


Table 2.15
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death^a Rates by State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	186.89	0.11		
High Five States				
Kentucky	219.31	1.02	(01)	17.35 ^c
Louisiana	217.01	1.01	(02)	16.12 ^c
West Virginia	211.40	1.39	(03)	13.12 ^c
Mississippi	211.34	1.22	(04)	13.09 ^c
Tennessee	208.83	0.83	(05)	11.74 ^c
Low Five States				
New Mexico	165.19	1.33	(47)	-11.61 ^c
Arizona	163.86	0.75	(48)	-12.32 ^c
Colorado	163.46	0.93	(49)	-12.53 ^c
Hawaii	150.28	1.47	(50)	-19.59 ^c
Utah	139.24	1.26	(51)	-25.49 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	203.72	0.92	(09)	9.01 ^c	Montana	182.67	1.88	(33)	-2.26
Alaska	182.04	3.24	(35)	-2.60	Nebraska	178.17	1.38	(39)	-4.67
Arizona	163.86	0.75	(48)	-12.32 ^c	Nevada	192.97	1.37	(21)	3.26
Arkansas	204.98	1.16	(06)	9.68 ^c	New Hampshire	190.62	1.70	(25)	2.00
California	169.70	0.33	(46)	-9.19 ^c	New Jersey	189.44	0.64	(27)	1.37
Colorado	163.46	0.93	(49)	-12.53 ^c	New Mexico	165.19	1.33	(47)	-11.61 ^c
Connecticut	182.60	0.98	(34)	-2.29	New York	176.13	0.42	(43)	-5.76 ^c
Delaware	198.65	2.13	(13)	6.29 ^c	North Carolina	194.29	0.68	(18)	3.96
D.C.	204.92	2.68	(07)	9.65 ^c	North Dakota	175.50	2.20	(44)	-6.10 ^c
Florida	177.25	0.40	(40)	-5.16 ^c	Ohio	202.29	0.57	(11)	8.24 ^c
Georgia	190.29	0.72	(26)	1.82	Oklahoma	198.10	1.03	(14)	6.00 ^c
Hawaii	150.28	1.47	(50)	-19.59 ^c	Oregon	189.25	1.00	(28)	1.26
Idaho	171.63	1.62	(45)	-8.17 ^c	Pennsylvania	196.43	0.51	(16)	5.11 ^c
Illinois	194.48	0.56	(17)	4.06	Rhode Island	191.35	1.79	(23)	2.39
Indiana	202.73	0.80	(10)	8.48 ^c	South Carolina	197.69	0.96	(15)	5.78 ^c
Iowa	184.30	1.04	(30)	-1.39	South Dakota	178.78	2.03	(38)	-4.34
Kansas	184.69	1.14	(29)	-1.18	Tennessee	208.83	0.83	(05)	11.74 ^c
Kentucky	219.31	1.02	(01)	17.35 ^c	Texas	181.67	0.44	(36)	-2.79
Louisiana	217.01	1.01	(02)	16.12 ^c	Utah	139.24	1.26	(51)	-25.49 ^c
Maine	204.15	1.63	(08)	9.23 ^c	Vermont	179.35	2.32	(37)	-4.03
Maryland	193.33	0.86	(20)	3.45	Virginia	191.95	0.74	(22)	2.71
Massachusetts	190.93	0.74	(24)	2.16	Washington	183.37	0.79	(32)	-1.88
Michigan	193.36	0.61	(19)	3.46	West Virginia	211.40	1.39	(03)	13.12 ^c
Minnesota	176.60	0.84	(42)	-5.50 ^c	Wisconsin	184.26	0.80	(31)	-1.41
Mississippi	211.34	1.22	(04)	13.09 ^c	Wyoming	176.74	2.65	(41)	-5.43 ^c
Missouri	200.08	0.81	(12)	7.06 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 2.16
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death^a Rates by State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	229.90	0.19		
High Five States				
Kentucky	280.75	1.82	(01)	22.12 ^c
Mississippi	280.13	2.22	(02)	21.85 ^c
Louisiana	278.60	1.80	(03)	21.18 ^c
D.C.	270.20	4.96	(04)	17.53 ^c
Tennessee	268.04	1.49	(05)	16.59 ^c
Low Five States				
New Mexico	199.19	2.22	(47)	-13.36 ^c
Arizona	196.85	1.23	(48)	-14.38 ^c
Colorado	195.98	1.59	(49)	-14.76 ^c
Hawaii	186.15	2.45	(50)	-19.03 ^c
Utah	167.02	2.10	(51)	-27.35 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	267.72	1.68	(06)	16.45 ^c	Montana	214.93	3.09	(41)	-6.51 ^c
Alaska	217.01	5.48	(37)	-5.61 ^c	Nebraska	220.47	2.37	(35)	-4.10
Arizona	196.85	1.23	(48)	-14.38 ^c	Nevada	223.33	2.21	(32)	-2.86
Arkansas	261.64	2.03	(08)	13.80 ^c	New Hampshire	233.22	2.95	(25)	1.44
California	202.23	0.55	(46)	-12.04 ^c	New Jersey	226.70	1.11	(29)	-1.39
Colorado	195.98	1.59	(49)	-14.76 ^c	New Mexico	199.19	2.22	(47)	-13.36 ^c
Connecticut	223.45	1.69	(31)	-2.81	New York	211.70	0.72	(43)	-7.92 ^c
Delaware	245.99	3.68	(16)	7.00 ^c	North Carolina	248.29	1.22	(14)	8.00 ^c
D.C.	270.20	4.96	(04)	17.53 ^c	North Dakota	214.10	3.66	(42)	-6.87 ^c
Florida	215.23	0.66	(40)	-6.38 ^c	Ohio	251.92	1.01	(11)	9.58 ^c
Georgia	245.93	1.33	(17)	6.97 ^c	Oklahoma	247.61	1.79	(15)	7.70 ^c
Hawaii	186.15	2.45	(50)	-19.03 ^c	Oregon	223.21	1.65	(33)	-2.91
Idaho	205.51	2.67	(45)	-10.61 ^c	Pennsylvania	243.19	0.89	(18)	5.78 ^c
Illinois	240.52	0.98	(20)	4.62	Rhode Island	240.44	3.14	(21)	4.58
Indiana	253.01	1.41	(10)	10.05 ^c	South Carolina	256.16	1.73	(09)	11.42 ^c
Iowa	229.10	1.78	(26)	-0.35	South Dakota	221.84	3.42	(34)	-3.51
Kansas	227.13	1.94	(28)	-1.21	Tennessee	268.04	1.49	(05)	16.59 ^c
Kentucky	280.75	1.82	(01)	22.12 ^c	Texas	227.34	0.77	(27)	-1.11
Louisiana	278.60	1.80	(03)	21.18 ^c	Utah	167.02	2.10	(51)	-27.35 ^c
Maine	251.33	2.81	(12)	9.32 ^c	Vermont	216.22	3.96	(38)	-5.95 ^c
Maryland	236.80	1.51	(22)	3.00	Virginia	241.35	1.32	(19)	4.98
Massachusetts	235.43	1.30	(24)	2.40	Washington	217.27	1.32	(36)	-5.49 ^c
Michigan	236.17	1.06	(23)	2.73	West Virginia	263.08	2.42	(07)	14.43 ^c
Minnesota	215.45	1.43	(39)	-6.29 ^c	Wisconsin	226.33	1.36	(30)	-1.55
Mississippi	280.13	2.22	(02)	21.85 ^c	Wyoming	206.71	4.36	(44)	-10.09 ^c
Missouri	249.40	1.40	(13)	8.48 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 2.17
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death^a Rates by State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	157.85	0.14		
High Five States				
Kentucky	178.71	1.22	(01)	13.22 ^c
West Virginia	175.95	1.70	(02)	11.47 ^c
Louisiana	175.78	1.20	(03)	11.36 ^c
Maine	172.37	2.01	(04)	9.20 ^c
Ohio	170.15	0.70	(05)	7.80 ^c
Low Five States				
Colorado	142.30	1.15	(47)	-9.85 ^c
New Mexico	140.20	1.65	(48)	-11.18 ^c
Arizona	138.42	0.93	(49)	-12.31 ^c
Hawaii	122.38	1.80	(50)	-22.47 ^c
Utah	118.70	1.56	(51)	-24.80 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	161.55	1.09	(23)	2.34	Montana	159.96	2.39	(26)	1.34
Alaska	155.23	4.00	(33)	-1.65	Nebraska	149.73	1.71	(41)	-5.14 ^c
Arizona	138.42	0.93	(49)	-12.31 ^c	Nevada	168.23	1.73	(08)	6.58 ^c
Arkansas	165.16	1.40	(17)	4.63	New Hampshire	163.06	2.10	(22)	3.31
California	147.64	0.41	(45)	-6.47 ^c	New Jersey	166.18	0.79	(11)	5.28 ^c
Colorado	142.30	1.15	(47)	-9.85 ^c	New Mexico	140.20	1.65	(48)	-11.18 ^c
Connecticut	156.84	1.20	(30)	-0.64	New York	153.87	0.51	(38)	-2.52
Delaware	165.83	2.59	(13)	5.06 ^c	North Carolina	158.93	0.81	(28)	0.68
D.C.	164.33	3.13	(19)	4.11	North Dakota	149.32	2.76	(42)	-5.40 ^c
Florida	147.93	0.50	(44)	-6.28 ^c	Ohio	170.15	0.70	(05)	7.80 ^c
Georgia	155.13	0.85	(35)	-1.72	Oklahoma	163.71	1.26	(20)	3.71
Hawaii	122.38	1.80	(50)	-22.47 ^c	Oregon	165.52	1.26	(14)	4.87
Idaho	146.21	2.02	(46)	-7.37 ^c	Pennsylvania	166.30	0.63	(10)	5.35 ^c
Illinois	165.32	0.68	(15)	4.74	Rhode Island	161.18	2.18	(24)	2.11
Indiana	170.08	0.98	(06)	7.75 ^c	South Carolina	158.29	1.14	(29)	0.28
Iowa	154.64	1.28	(36)	-2.03	South Dakota	148.11	2.51	(43)	-6.17 ^c
Kansas	156.53	1.41	(32)	-0.83	Tennessee	169.53	0.99	(07)	7.40 ^c
Kentucky	178.71	1.22	(01)	13.22 ^c	Texas	150.26	0.53	(40)	-4.80
Louisiana	175.78	1.20	(03)	11.36 ^c	Utah	118.70	1.56	(51)	-24.80 ^c
Maine	172.37	2.01	(04)	9.20 ^c	Vermont	155.14	2.88	(34)	-1.72
Maryland	165.26	1.04	(16)	4.70	Virginia	159.88	0.89	(27)	1.29
Massachusetts	163.48	0.91	(21)	3.57	Washington	160.16	0.99	(25)	1.47
Michigan	164.88	0.76	(18)	4.46	West Virginia	175.95	1.70	(02)	11.47 ^c
Minnesota	151.68	1.04	(39)	-3.91	Wisconsin	156.57	0.98	(31)	-0.81
Mississippi	165.98	1.43	(12)	5.15 ^c	Wyoming	154.37	3.36	(37)	-2.20
Missouri	167.07	0.98	(09)	5.84 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 3.15
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.33	0.02		
High Five States				
Tennessee	5.53	0.13	(01)	27.47 ^c
Washington	5.46	0.13	(02)	26.01 ^c
Idaho	5.46	0.29	(03)	25.93 ^c
Oregon	5.43	0.17	(04)	25.29 ^c
Kansas	5.36	0.20	(05)	23.65 ^c
Low Five States				
New Mexico	3.78	0.20	(47)	-12.89
New York	3.74	0.06	(48)	-13.75
New Jersey	3.67	0.09	(49)	-15.44 ^c
D.C.	3.15	0.34	(50)	-27.21 ^c
Hawaii	2.76	0.20	(51)	-36.34 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.29	0.13	(34)	-0.96	Montana	4.90	0.31	(13)	13.07
Alaska	4.24	0.45	(37)	-2.10	Nebraska	4.79	0.23	(16)	10.41
Arizona	4.10	0.12	(41)	-5.52	Nevada	4.26	0.20	(36)	-1.62
Arkansas	4.90	0.18	(14)	12.96	New Hampshire	4.99	0.28	(11)	15.02 ^c
California	4.31	0.05	(33)	-0.57	New Jersey	3.67	0.09	(49)	-15.44 ^c
Colorado	4.40	0.15	(30)	1.49	New Mexico	3.78	0.20	(47)	-12.89
Connecticut	4.12	0.15	(40)	-4.95	New York	3.74	0.06	(48)	-13.75
Delaware	4.41	0.32	(29)	1.63	North Carolina	4.27	0.10	(35)	-1.39
D.C.	3.15	0.34	(50)	-27.21 ^c	North Dakota	5.02	0.39	(09)	15.80 ^c
Florida	4.06	0.06	(42)	-6.31	Ohio	4.48	0.09	(27)	3.44
Georgia	3.82	0.10	(46)	-11.78	Oklahoma	4.85	0.16	(15)	11.81
Hawaii	2.76	0.20	(51)	-36.34 ^c	Oregon	5.43	0.17	(04)	25.29 ^c
Idaho	5.46	0.29	(03)	25.93 ^c	Pennsylvania	4.12	0.08	(39)	-4.89
Illinois	3.88	0.08	(44)	-10.41	Rhode Island	4.97	0.29	(12)	14.61
Indiana	4.64	0.12	(19)	7.06	South Carolina	4.53	0.15	(24)	4.54
Iowa	5.30	0.18	(06)	22.27 ^c	South Dakota	5.17	0.36	(07)	19.38 ^c
Kansas	5.36	0.20	(05)	23.65 ^c	Tennessee	5.53	0.13	(01)	27.47 ^c
Kentucky	4.34	0.14	(32)	0.20	Texas	4.42	0.07	(28)	1.87
Louisiana	4.63	0.15	(21)	6.72	Utah	4.67	0.22	(18)	7.77
Maine	5.08	0.26	(08)	17.29 ^c	Vermont	4.49	0.37	(26)	3.62
Maryland	3.93	0.12	(43)	-9.31	Virginia	3.83	0.10	(45)	-11.64
Massachusetts	4.17	0.11	(38)	-3.72	Washington	5.46	0.13	(02)	26.01 ^c
Michigan	4.72	0.10	(17)	8.92	West Virginia	4.60	0.21	(22)	6.17
Minnesota	4.37	0.13	(31)	0.87	Wisconsin	4.63	0.13	(20)	6.84
Mississippi	5.02	0.19	(10)	15.80 ^c	Wyoming	4.60	0.43	(23)	6.12
Missouri	4.51	0.12	(25)	3.98					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 3.16
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.30	0.03		
High Five States				
South Dakota	7.26	0.62	(01)	37.05 ^c
Washington	6.68	0.22	(02)	26.15 ^c
Kansas	6.64	0.33	(03)	25.45 ^c
Oregon	6.56	0.27	(04)	23.79 ^c
Maine	6.54	0.44	(05)	23.52 ^c
Low Five States				
New Mexico	4.63	0.33	(47)	-12.61
Georgia	4.63	0.17	(48)	-12.67
New Jersey	4.56	0.15	(49)	-13.98
D.C.	3.83	0.56	(50)	-27.61 ^c
Hawaii	3.35	0.33	(51)	-36.72 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.29	0.23	(31)	-0.06	Montana	5.69	0.48	(17)	7.42
Alaska	5.03	0.73	(37)	-5.09	Nebraska	5.64	0.37	(20)	6.42
Arizona	4.90	0.19	(42)	-7.52	Nevada	4.87	0.31	(43)	-8.04
Arkansas	5.76	0.30	(15)	8.75	New Hampshire	6.13	0.45	(10)	15.79 ^c
California	5.31	0.09	(29)	0.17	New Jersey	4.56	0.15	(49)	-13.98
Colorado	5.23	0.24	(34)	-1.16	New Mexico	4.63	0.33	(47)	-12.61
Connecticut	4.82	0.24	(44)	-9.03	New York	4.74	0.10	(45)	-10.58
Delaware	5.55	0.53	(23)	4.71	North Carolina	5.25	0.17	(32)	-0.86
D.C.	3.83	0.56	(50)	-27.61 ^c	North Dakota	5.64	0.61	(19)	6.47
Florida	4.91	0.10	(40)	-7.30	Ohio	5.47	0.14	(26)	3.24
Georgia	4.63	0.17	(48)	-12.67	Oklahoma	5.98	0.27	(11)	12.84
Hawaii	3.35	0.33	(51)	-36.72 ^c	Oregon	6.56	0.27	(04)	23.79 ^c
Idaho	6.24	0.44	(08)	17.91 ^c	Pennsylvania	4.97	0.13	(38)	-6.20
Illinois	4.67	0.13	(46)	-11.77	Rhode Island	6.20	0.49	(09)	16.99 ^c
Indiana	5.57	0.20	(21)	5.22	South Carolina	5.45	0.24	(27)	2.89
Iowa	6.32	0.29	(07)	19.27 ^c	South Dakota	7.26	0.62	(01)	37.05 ^c
Kansas	6.64	0.33	(03)	25.45 ^c	Tennessee	6.42	0.22	(06)	21.13 ^c
Kentucky	5.15	0.23	(35)	-2.72	Texas	5.41	0.11	(28)	2.18
Louisiana	5.93	0.25	(13)	11.88	Utah	5.64	0.36	(18)	6.48
Maine	6.54	0.44	(05)	23.52 ^c	Vermont	5.51	0.59	(24)	4.09
Maryland	4.90	0.20	(41)	-7.42	Virginia	4.95	0.18	(39)	-6.60
Massachusetts	5.25	0.19	(33)	-0.95	Washington	6.68	0.22	(02)	26.15 ^c
Michigan	5.84	0.16	(14)	10.30	West Virginia	5.73	0.35	(16)	8.14
Minnesota	5.31	0.21	(30)	0.17	Wisconsin	5.56	0.21	(22)	5.03
Mississippi	5.93	0.31	(12)	12.00	Wyoming	5.15	0.66	(36)	-2.81
Missouri	5.50	0.20	(25)	3.77					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 3.17
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.53	0.02		
High Five States				
Tennessee	4.76	0.17	(01)	34.78 ^c
Idaho	4.64	0.37	(02)	31.46 ^c
Washington	4.46	0.17	(03)	26.38 ^c
Iowa	4.46	0.23	(04)	26.27 ^c
Oregon	4.42	0.21	(05)	25.29 ^c
Low Five States				
Virginia	2.97	0.12	(47)	-15.91 ^c
New York	2.94	0.07	(48)	-16.77 ^c
New Jersey	2.93	0.11	(49)	-17.05 ^c
D.C.	2.57	0.41	(50)	-27.34 ^c
Hawaii	2.28	0.26	(51)	-35.42 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.45	0.16	(37)	-2.29	Montana	4.18	0.40	(09)	18.25 ^c
Alaska	3.51	0.59	(32)	-0.70	Nebraska	4.00	0.29	(13)	13.31
Arizona	3.40	0.15	(39)	-3.74	Nevada	3.68	0.25	(22)	4.15
Arkansas	4.17	0.23	(10)	18.07 ^c	New Hampshire	3.94	0.34	(14)	11.51
California	3.48	0.06	(36)	-1.46	New Jersey	2.93	0.11	(49)	-17.05 ^c
Colorado	3.64	0.18	(25)	3.06	New Mexico	3.03	0.25	(46)	-14.06
Connecticut	3.50	0.19	(34)	-0.95	New York	2.94	0.07	(48)	-16.77 ^c
Delaware	3.50	0.39	(33)	-0.90	North Carolina	3.49	0.12	(35)	-1.15
D.C.	2.57	0.41	(50)	-27.34 ^c	North Dakota	4.42	0.49	(06)	25.21 ^c
Florida	3.33	0.08	(41)	-5.80	Ohio	3.67	0.11	(24)	3.79
Georgia	3.16	0.12	(44)	-10.49	Oklahoma	3.83	0.20	(18)	8.58
Hawaii	2.28	0.26	(51)	-35.42 ^c	Oregon	4.42	0.21	(05)	25.29 ^c
Idaho	4.64	0.37	(02)	31.46 ^c	Pennsylvania	3.44	0.10	(38)	-2.66
Illinois	3.27	0.10	(43)	-7.54	Rhode Island	4.01	0.36	(12)	13.70
Indiana	3.87	0.15	(15)	9.71	South Carolina	3.84	0.18	(17)	8.79
Iowa	4.46	0.23	(04)	26.27 ^c	South Dakota	3.39	0.39	(40)	-3.99
Kansas	4.31	0.24	(07)	21.93 ^c	Tennessee	4.76	0.17	(01)	34.78 ^c
Kentucky	3.63	0.18	(26)	2.72	Texas	3.59	0.08	(28)	1.77
Louisiana	3.62	0.17	(27)	2.49	Utah	3.76	0.28	(21)	6.42
Maine	3.87	0.31	(16)	9.58	Vermont	3.55	0.45	(31)	0.39
Maryland	3.15	0.15	(45)	-10.75	Virginia	2.97	0.12	(47)	-15.91 ^c
Massachusetts	3.29	0.13	(42)	-6.94	Washington	4.46	0.17	(03)	26.38 ^c
Michigan	3.77	0.12	(20)	6.77	West Virginia	3.58	0.25	(29)	1.48
Minnesota	3.55	0.16	(30)	0.67	Wisconsin	3.83	0.16	(19)	8.47
Mississippi	4.22	0.23	(08)	19.38 ^c	Wyoming	4.09	0.56	(11)	15.78 ^c
Missouri	3.67	0.15	(23)	3.94					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 4.22
Cancer of the Female Breast (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	24.51	0.05		
High Five States				
Louisiana	28.93	0.49	(01)	18.05 ^c
D.C.	28.85	1.32	(02)	17.72 ^c
New Jersey	27.43	0.33	(03)	11.91 ^c
Ohio	27.13	0.28	(04)	10.71 ^c
Maryland	26.77	0.42	(05)	9.24
Low Five States				
Colorado	22.22	0.45	(47)	-9.33
Minnesota	22.17	0.40	(48)	-9.55
Arizona	21.75	0.37	(49)	-11.25 ^c
Alaska	21.65	1.41	(50)	-11.66 ^c
Hawaii	17.69	0.70	(51)	-27.83 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	25.11	0.43	(12)	2.44	Montana	23.06	0.91	(37)	-5.92
Alaska	21.65	1.41	(50)	-11.66 ^c	Nebraska	22.90	0.67	(41)	-6.58
Arizona	21.75	0.37	(49)	-11.25 ^c	Nevada	24.36	0.66	(22)	-0.60
Arkansas	24.27	0.54	(25)	-0.95	New Hampshire	23.27	0.79	(34)	-5.05
California	23.17	0.16	(35)	-5.44	New Jersey	27.43	0.33	(03)	11.91 ^c
Colorado	22.22	0.45	(47)	-9.33	New Mexico	22.48	0.66	(45)	-8.27
Connecticut	24.35	0.48	(24)	-0.63	New York	24.67	0.21	(19)	0.67
Delaware	24.01	1.00	(27)	-2.02	North Carolina	25.10	0.32	(13)	2.40
D.C.	28.85	1.32	(02)	17.72 ^c	North Dakota	23.01	1.11	(38)	-6.11
Florida	22.64	0.20	(44)	-7.63	Ohio	27.13	0.28	(04)	10.71 ^c
Georgia	24.48	0.33	(21)	-0.12	Oklahoma	25.06	0.50	(14)	2.26
Hawaii	17.69	0.70	(51)	-27.83 ^c	Oregon	23.90	0.48	(28)	-2.50
Idaho	22.29	0.79	(46)	-9.05	Pennsylvania	26.42	0.26	(06)	7.81
Illinois	25.72	0.27	(11)	4.92	Rhode Island	23.14	0.83	(36)	-5.60
Indiana	24.80	0.37	(18)	1.20	South Carolina	24.99	0.45	(16)	1.97
Iowa	22.86	0.50	(42)	-6.72	South Dakota	22.92	1.01	(40)	-6.46
Kansas	24.64	0.56	(20)	0.53	Tennessee	25.91	0.39	(10)	5.71
Kentucky	24.84	0.45	(17)	1.35	Texas	23.42	0.21	(30)	-4.46
Louisiana	28.93	0.49	(01)	18.05 ^c	Utah	23.75	0.70	(29)	-3.08
Maine	23.36	0.74	(32)	-4.69	Vermont	22.93	1.10	(39)	-6.45
Maryland	26.77	0.42	(05)	9.24	Virginia	26.00	0.36	(09)	6.10
Massachusetts	24.17	0.35	(26)	-1.37	Washington	23.34	0.38	(33)	-4.75
Michigan	25.06	0.30	(15)	2.25	West Virginia	24.35	0.64	(23)	-0.63
Minnesota	22.17	0.40	(48)	-9.55	Wisconsin	23.39	0.38	(31)	-4.56
Mississippi	26.39	0.57	(07)	7.66	Wyoming	22.65	1.29	(43)	-7.59
Missouri	26.25	0.39	(08)	7.12					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.



Table 5.13
Cancer of the Cervix Uteri (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.46	0.02		
High Five States				
Mississippi	3.98	0.23	(01)	62.05 ^c
Arkansas	3.46	0.21	(02)	40.60 ^c
D.C.	3.38	0.47	(03)	37.44 ^c
Texas	3.14	0.08	(04)	27.76 ^c
West Virginia	3.13	0.25	(05)	27.32 ^c
Low Five States				
South Dakota	1.64	0.29	(47)	-33.06 ^c
Wyoming	1.59	0.34	(48)	-35.13 ^c
Utah	1.56	0.18	(49)	-36.42 ^c
Massachusetts	1.49	0.09	(50)	-39.41 ^c
North Dakota	1.48	0.29	(51)	-39.96 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.03	0.15	(07)	23.11 ^c	Montana	1.98	0.28	(35)	-19.54 ^c
Alaska	1.94	0.40	(37)	-20.87 ^c	Nebraska	1.96	0.21	(36)	-20.42 ^c
Arizona	2.14	0.12	(30)	-13.09	Nevada	2.76	0.22	(11)	12.41
Arkansas	3.46	0.21	(02)	40.60 ^c	New Hampshire	2.25	0.25	(28)	-8.55
California	2.35	0.05	(24)	-4.22	New Jersey	2.65	0.10	(17)	7.65
Colorado	1.79	0.12	(43)	-27.16 ^c	New Mexico	2.54	0.23	(19)	3.51
Connecticut	1.75	0.13	(44)	-28.83 ^c	New York	2.57	0.07	(18)	4.53
Delaware	2.49	0.33	(21)	1.52	North Carolina	2.42	0.10	(23)	-1.54
D.C.	3.38	0.47	(03)	37.44 ^c	North Dakota	1.48	0.29	(51)	-39.96 ^c
Florida	2.75	0.07	(12)	11.71	Ohio	2.46	0.09	(22)	0.21
Georgia	2.71	0.11	(13)	10.44	Oklahoma	2.79	0.18	(09)	13.35
Hawaii	2.10	0.25	(31)	-14.58	Oregon	1.94	0.14	(38)	-21.21 ^c
Idaho	1.92	0.24	(39)	-21.93 ^c	Pennsylvania	2.34	0.08	(25)	-4.88
Illinois	2.77	0.09	(10)	12.91	Rhode Island	2.03	0.26	(33)	-17.25 ^c
Indiana	2.54	0.12	(20)	3.45	South Carolina	2.69	0.15	(15)	9.55
Iowa	2.29	0.17	(27)	-6.95	South Dakota	1.64	0.29	(47)	-33.06 ^c
Kansas	2.33	0.18	(26)	-5.00	Tennessee	2.89	0.13	(08)	17.62 ^c
Kentucky	2.71	0.16	(14)	10.29	Texas	3.14	0.08	(04)	27.76 ^c
Louisiana	3.11	0.16	(06)	26.62 ^c	Utah	1.56	0.18	(49)	-36.42 ^c
Maine	1.92	0.22	(40)	-21.96 ^c	Vermont	1.71	0.31	(45)	-30.59 ^c
Maryland	2.24	0.12	(29)	-8.96	Virginia	2.09	0.10	(32)	-14.90
Massachusetts	1.49	0.09	(50)	-39.41 ^c	Washington	1.81	0.11	(42)	-26.17 ^c
Michigan	2.02	0.09	(34)	-17.76 ^c	West Virginia	3.13	0.25	(05)	27.32 ^c
Minnesota	1.66	0.11	(46)	-32.58 ^c	Wisconsin	1.86	0.11	(41)	-24.42 ^c
Mississippi	3.98	0.23	(01)	62.05 ^c	Wyoming	1.59	0.34	(48)	-35.13 ^c
Missouri	2.66	0.13	(16)	8.40					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 6.22
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	18.19	0.03		
High Five States				
Louisiana	21.71	0.32	(01)	19.40 ^c
Kentucky	21.45	0.32	(02)	17.97 ^c
West Virginia	21.43	0.44	(03)	17.87 ^c
D.C.	20.90	0.85	(04)	14.95 ^c
Mississippi	20.68	0.38	(05)	13.73 ^c
Low Five States				
Washington	16.07	0.23	(47)	-11.62 ^c
Arizona	15.67	0.23	(48)	-13.82 ^c
Hawaii	15.53	0.47	(49)	-14.58 ^c
Idaho	15.35	0.48	(50)	-15.60 ^c
Utah	13.41	0.39	(51)	-26.28 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	18.87	0.28	(20)	3.74	Montana	16.83	0.57	(41)	-7.44
Alaska	16.73	1.01	(42)	-8.01	Nebraska	19.57	0.45	(12)	7.59
Arizona	15.67	0.23	(48)	-13.82 ^c	Nevada	19.60	0.44	(11)	7.78
Arkansas	19.53	0.36	(13)	7.37	New Hampshire	18.00	0.52	(32)	-1.04
California	16.25	0.10	(46)	-10.62 ^c	New Jersey	20.05	0.21	(08)	10.27 ^c
Colorado	16.86	0.30	(40)	-7.28	New Mexico	16.38	0.42	(45)	-9.90
Connecticut	17.08	0.30	(38)	-6.06	New York	18.37	0.13	(28)	1.03
Delaware	18.91	0.66	(18)	4.00	North Carolina	17.73	0.21	(35)	-2.51
D.C.	20.90	0.85	(04)	14.95 ^c	North Dakota	18.46	0.70	(25)	1.49
Florida	16.42	0.12	(44)	-9.70	Ohio	20.06	0.18	(07)	10.33 ^c
Georgia	17.94	0.22	(33)	-1.36	Oklahoma	18.89	0.32	(19)	3.88
Hawaii	15.53	0.47	(49)	-14.58 ^c	Oregon	16.90	0.30	(39)	-7.08
Idaho	15.35	0.48	(50)	-15.60 ^c	Pennsylvania	19.84	0.16	(10)	9.12
Illinois	20.13	0.18	(06)	10.72 ^c	Rhode Island	18.24	0.54	(30)	0.30
Indiana	19.92	0.25	(09)	9.54	South Carolina	18.66	0.30	(22)	2.64
Iowa	19.02	0.33	(17)	4.56	South Dakota	18.16	0.64	(31)	-0.13
Kansas	18.44	0.36	(26)	1.41	Tennessee	19.17	0.25	(16)	5.41
Kentucky	21.45	0.32	(02)	17.97 ^c	Texas	17.46	0.14	(36)	-4.01
Louisiana	21.71	0.32	(01)	19.40 ^c	Utah	13.41	0.39	(51)	-26.28 ^c
Maine	18.73	0.49	(21)	3.01	Vermont	18.57	0.74	(23)	2.11
Maryland	19.30	0.27	(14)	6.11	Virginia	18.29	0.23	(29)	0.55
Massachusetts	18.54	0.23	(24)	1.93	Washington	16.07	0.23	(47)	-11.62 ^c
Michigan	18.38	0.19	(27)	1.08	West Virginia	21.43	0.44	(03)	17.87 ^c
Minnesota	16.48	0.25	(43)	-9.40	Wisconsin	17.16	0.24	(37)	-5.65
Mississippi	20.68	0.38	(05)	13.73 ^c	Wyoming	17.91	0.85	(34)	-1.52
Missouri	19.22	0.25	(15)	5.69					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 6.23
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	21.94	0.06		
High Five States				
Louisiana	27.56	0.57	(01)	25.62 ^c
Kentucky	26.20	0.56	(02)	19.38 ^c
D.C.	26.16	1.55	(03)	19.21 ^c
West Virginia	25.98	0.76	(04)	18.39 ^c
Mississippi	25.01	0.66	(05)	13.96 ^c
Low Five States				
Arizona	19.03	0.38	(47)	-13.28 ^c
Washington	18.72	0.39	(48)	-14.68 ^c
Wyoming	18.66	1.32	(49)	-14.98 ^c
Idaho	17.64	0.79	(50)	-19.59 ^c
Utah	15.46	0.64	(51)	-29.54 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	24.19	0.51	(09)	10.25 ^c	Montana	19.77	0.94	(41)	-9.92
Alaska	19.70	1.67	(42)	-10.21 ^c	Nebraska	24.03	0.78	(12)	9.53
Arizona	19.03	0.38	(47)	-13.28 ^c	Nevada	23.11	0.72	(19)	5.31
Arkansas	23.96	0.62	(13)	9.21	New Hampshire	21.96	0.91	(28)	0.06
California	19.22	0.17	(46)	-12.39 ^c	New Jersey	24.12	0.36	(11)	9.91
Colorado	19.80	0.51	(40)	-9.75	New Mexico	19.88	0.70	(39)	-9.42
Connecticut	20.15	0.51	(37)	-8.18	New York	22.10	0.23	(26)	0.73
Delaware	23.22	1.13	(18)	5.84	North Carolina	21.61	0.36	(34)	-1.50
D.C.	26.16	1.55	(03)	19.21 ^c	North Dakota	21.68	1.16	(33)	-1.19
Florida	19.52	0.20	(44)	-11.05 ^c	Ohio	24.16	0.31	(10)	10.12 ^c
Georgia	22.20	0.40	(25)	1.17	Oklahoma	23.49	0.55	(15)	7.03
Hawaii	20.06	0.80	(38)	-8.57	Oregon	19.55	0.49	(43)	-10.92 ^c
Idaho	17.64	0.79	(50)	-19.59 ^c	Pennsylvania	24.56	0.28	(08)	11.92 ^c
Illinois	24.91	0.32	(06)	13.54 ^c	Rhode Island	22.32	0.96	(24)	1.71
Indiana	24.91	0.44	(07)	13.52 ^c	South Carolina	22.62	0.51	(21)	3.11
Iowa	23.00	0.57	(20)	4.84	South Dakota	21.74	1.07	(32)	-0.94
Kansas	21.95	0.60	(29)	0.05	Tennessee	23.66	0.45	(14)	7.82
Kentucky	26.20	0.56	(02)	19.38 ^c	Texas	21.52	0.24	(35)	-1.92
Louisiana	27.56	0.57	(01)	25.62 ^c	Utah	15.46	0.64	(51)	-29.54 ^c
Maine	21.88	0.84	(31)	-0.30	Vermont	22.08	1.27	(27)	0.64
Maryland	23.35	0.47	(16)	6.41	Virginia	22.59	0.40	(22)	2.94
Massachusetts	22.33	0.40	(23)	1.76	Washington	18.72	0.39	(48)	-14.68 ^c
Michigan	21.88	0.32	(30)	-0.28	West Virginia	25.98	0.76	(04)	18.39 ^c
Minnesota	19.27	0.43	(45)	-12.18 ^c	Wisconsin	20.75	0.42	(36)	-5.46
Mississippi	25.01	0.66	(05)	13.96 ^c	Wyoming	18.66	1.32	(49)	-14.98 ^c
Missouri	23.33	0.43	(17)	6.32					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 6.24
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	15.37	0.04		
High Five States				
Kentucky	18.21	0.38	(01)	18.50 ^c
West Virginia	18.07	0.53	(02)	17.61 ^c
Louisiana	17.52	0.38	(03)	14.03 ^c
Mississippi	17.49	0.46	(04)	13.83 ^c
D.C.	17.45	1.01	(05)	13.55 ^c
Low Five States				
New Mexico	13.57	0.51	(47)	-11.67 ^c
Idaho	13.44	0.61	(48)	-12.57 ^c
Arizona	12.93	0.28	(49)	-15.89 ^c
Utah	11.74	0.49	(50)	-23.62 ^c
Hawaii	11.72	0.55	(51)	-23.76 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	15.18	0.33	(30)	-1.24	Montana	14.48	0.71	(39)	-5.77
Alaska	14.13	1.25	(43)	-8.03	Nebraska	16.31	0.55	(13)	6.11
Arizona	12.93	0.28	(49)	-15.89 ^c	Nevada	16.52	0.55	(11)	7.51
Arkansas	16.10	0.43	(18)	4.79	New Hampshire	15.14	0.63	(32)	-1.48
California	13.93	0.12	(44)	-9.36	New Jersey	17.19	0.25	(06)	11.85 ^c
Colorado	14.54	0.36	(38)	-5.40	New Mexico	13.57	0.51	(47)	-11.67 ^c
Connecticut	14.92	0.36	(36)	-2.90	New York	15.77	0.16	(22)	2.63
Delaware	15.51	0.79	(28)	0.90	North Carolina	14.92	0.25	(35)	-2.89
D.C.	17.45	1.01	(05)	13.55 ^c	North Dakota	16.14	0.87	(17)	5.01
Florida	13.85	0.15	(46)	-9.85	Ohio	17.13	0.22	(07)	11.47 ^c
Georgia	15.13	0.27	(33)	-1.54	Oklahoma	15.44	0.38	(29)	0.48
Hawaii	11.72	0.55	(51)	-23.76 ^c	Oregon	14.81	0.37	(37)	-3.61
Idaho	13.44	0.61	(48)	-12.57 ^c	Pennsylvania	16.54	0.19	(10)	7.65
Illinois	16.83	0.21	(09)	9.52	Rhode Island	15.53	0.66	(27)	1.04
Indiana	16.31	0.30	(12)	6.14	South Carolina	15.65	0.36	(26)	1.83
Iowa	16.06	0.40	(19)	4.52	South Dakota	15.13	0.77	(34)	-1.54
Kansas	15.72	0.43	(24)	2.28	Tennessee	15.91	0.30	(20)	3.55
Kentucky	18.21	0.38	(01)	18.50 ^c	Texas	14.32	0.16	(42)	-6.82
Louisiana	17.52	0.38	(03)	14.03 ^c	Utah	11.74	0.49	(50)	-23.62 ^c
Maine	16.28	0.60	(15)	5.97	Vermont	15.72	0.90	(25)	2.27
Maryland	16.31	0.32	(14)	6.11	Virginia	15.17	0.27	(31)	-1.31
Massachusetts	15.86	0.27	(21)	3.21	Washington	13.88	0.29	(45)	-9.68
Michigan	15.73	0.23	(23)	2.36	West Virginia	18.07	0.53	(02)	17.61 ^c
Minnesota	14.43	0.31	(41)	-6.14	Wisconsin	14.48	0.29	(40)	-5.81
Mississippi	17.49	0.46	(04)	13.83 ^c	Wyoming	17.08	1.11	(08)	11.15 ^c
Missouri	16.17	0.30	(16)	5.21					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 7.13
Cancer of the Corpus and Uterus, NOS (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.14	0.02		
High Five States				
D.C.	6.32	0.62	(01)	52.51 ^c
Vermont	5.21	0.53	(02)	25.85 ^c
North Dakota	5.17	0.51	(03)	24.77 ^c
New Jersey	5.17	0.14	(04)	24.67 ^c
New York	5.01	0.09	(05)	20.91 ^c
Low Five States				
Arkansas	3.28	0.20	(47)	-20.90 ^c
Wyoming	3.18	0.49	(48)	-23.37 ^c
Nevada	2.99	0.23	(49)	-27.72 ^c
Arizona	2.89	0.13	(50)	-30.19 ^c
Alaska	2.60	0.53	(51)	-37.20 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.46	0.16	(43)	-16.56 ^c	Montana	4.93	0.42	(06)	19.04 ^c
Alaska	2.60	0.53	(51)	-37.20 ^c	Nebraska	4.93	0.31	(07)	19.02 ^c
Arizona	2.89	0.13	(50)	-30.19 ^c	Nevada	2.99	0.23	(49)	-27.72 ^c
Arkansas	3.28	0.20	(47)	-20.90 ^c	New Hampshire	3.62	0.31	(37)	-12.54
California	3.93	0.07	(29)	-5.14	New Jersey	5.17	0.14	(04)	24.67 ^c
Colorado	3.31	0.17	(46)	-20.09 ^c	New Mexico	3.82	0.27	(32)	-7.77
Connecticut	4.36	0.20	(20)	5.16	New York	5.01	0.09	(05)	20.91 ^c
Delaware	4.84	0.44	(09)	16.88 ^c	North Carolina	4.00	0.13	(27)	-3.54
D.C.	6.32	0.62	(01)	52.51 ^c	North Dakota	5.17	0.51	(03)	24.77 ^c
Florida	3.59	0.08	(39)	-13.30	Ohio	4.45	0.11	(16)	7.47
Georgia	3.58	0.13	(40)	-13.66	Oklahoma	4.08	0.20	(23)	-1.43
Hawaii	3.57	0.31	(41)	-13.79	Oregon	3.83	0.19	(31)	-7.51
Idaho	3.35	0.30	(45)	-19.14 ^c	Pennsylvania	4.91	0.11	(08)	18.50 ^c
Illinois	4.49	0.11	(13)	8.29	Rhode Island	3.67	0.33	(34)	-11.47
Indiana	4.71	0.16	(10)	13.63	South Carolina	4.03	0.18	(26)	-2.64
Iowa	4.66	0.22	(12)	12.57	South Dakota	3.83	0.40	(30)	-7.47
Kansas	3.62	0.21	(36)	-12.54	Tennessee	3.74	0.15	(33)	-9.79
Kentucky	3.50	0.17	(42)	-15.42 ^c	Texas	3.62	0.08	(38)	-12.56
Louisiana	3.65	0.17	(35)	-11.99	Utah	3.42	0.27	(44)	-17.54 ^c
Maine	4.40	0.32	(18)	6.15	Vermont	5.21	0.53	(02)	25.85 ^c
Maryland	4.42	0.17	(17)	6.62	Virginia	4.04	0.14	(25)	-2.40
Massachusetts	4.37	0.15	(19)	5.42	Washington	4.06	0.16	(24)	-2.07
Michigan	4.70	0.13	(11)	13.50	West Virginia	4.22	0.26	(22)	1.83
Minnesota	4.45	0.18	(15)	7.48	Wisconsin	4.45	0.17	(14)	7.51
Mississippi	3.97	0.22	(28)	-4.26	Wyoming	3.18	0.49	(48)	-23.37 ^c
Missouri	4.30	0.16	(21)	3.67					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 8.15<h2>
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.40	0.02		
High Five States				
Maine	6.25	0.28	(01)	42.10 ^c
New Hampshire	6.05	0.30	(02)	37.67 ^c
D.C.	6.00	0.46	(03)	36.42 ^c
Wyoming	5.85	0.48	(04)	33.08 ^c
Massachusetts	5.48	0.13	(05)	24.60 ^c
Low Five States				
Texas	3.74	0.06	(47)	-14.95
Arkansas	3.73	0.16	(48)	-15.29 ^c
California	3.69	0.05	(49)	-16.05 ^c
Utah	3.16	0.19	(50)	-28.23 ^c
Hawaii	2.75	0.20	(51)	-37.55 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.96	0.13	(44)	-9.87	Montana	4.76	0.30	(16)	8.29
Alaska	5.06	0.52	(11)	15.15 ^c	Nebraska	4.46	0.22	(31)	1.44
Arizona	4.11	0.12	(39)	-6.62	Nevada	4.46	0.20	(30)	1.50
Arkansas	3.73	0.16	(48)	-15.29 ^c	New Hampshire	6.05	0.30	(02)	37.67 ^c
California	3.69	0.05	(49)	-16.05 ^c	New Jersey	4.30	0.10	(35)	-2.13
Colorado	4.04	0.14	(42)	-8.04	New Mexico	4.04	0.21	(43)	-8.16
Connecticut	4.48	0.15	(29)	1.96	New York	4.37	0.07	(34)	-0.67
Delaware	5.03	0.34	(12)	14.37	North Carolina	4.24	0.10	(37)	-3.60
D.C.	6.00	0.46	(03)	36.42 ^c	North Dakota	3.95	0.33	(45)	-10.10
Florida	4.08	0.06	(40)	-7.26	Ohio	5.18	0.09	(08)	17.69 ^c
Georgia	4.26	0.11	(36)	-3.20	Oklahoma	4.08	0.15	(41)	-7.31
Hawaii	2.75	0.20	(51)	-37.55 ^c	Oregon	5.06	0.16	(10)	15.16 ^c
Idaho	4.61	0.26	(20)	4.76	Pennsylvania	5.12	0.08	(09)	16.45 ^c
Illinois	4.58	0.09	(22)	4.04	Rhode Island	5.46	0.30	(06)	24.09 ^c
Indiana	4.63	0.12	(19)	5.35	South Carolina	4.78	0.15	(15)	8.74
Iowa	4.84	0.17	(14)	10.00	South Dakota	4.54	0.33	(24)	3.22
Kansas	3.77	0.16	(46)	-14.22	Tennessee	4.11	0.12	(38)	-6.58
Kentucky	4.50	0.14	(27)	2.28	Texas	3.74	0.06	(47)	-14.95
Louisiana	4.42	0.14	(33)	0.56	Utah	3.16	0.19	(50)	-28.23 ^c
Maine	6.25	0.28	(01)	42.10 ^c	Vermont	4.76	0.38	(17)	8.21
Maryland	4.54	0.13	(25)	3.17	Virginia	4.43	0.11	(32)	0.75
Massachusetts	5.48	0.13	(05)	24.60 ^c	Washington	4.65	0.13	(18)	5.64
Michigan	5.02	0.10	(13)	14.12	West Virginia	4.58	0.21	(21)	4.07
Minnesota	4.49	0.13	(28)	2.13	Wisconsin	5.23	0.13	(07)	19.04 ^c
Mississippi	4.52	0.18	(26)	2.72	Wyoming	5.85	0.48	(04)	33.08 ^c
Missouri	4.56	0.12	(23)	3.72					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 8.16
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.82	0.03		
High Five States				
Maine	11.59	0.59	(01)	48.23 ^c
D.C.	10.99	0.97	(02)	40.51 ^c
New Hampshire	10.73	0.62	(03)	37.30 ^c
Massachusetts	10.14	0.26	(04)	29.69 ^c
Wyoming	10.02	0.94	(05)	28.11 ^c
Low Five States				
Texas	6.70	0.13	(47)	-14.29
Arkansas	6.51	0.31	(48)	-16.74 ^c
California	6.38	0.10	(49)	-18.40 ^c
Utah	5.86	0.38	(50)	-24.99 ^c
Hawaii	4.87	0.39	(51)	-37.74 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.32	0.27	(40)	-6.38	Montana	7.61	0.57	(32)	-2.69
Alaska	8.18	1.02	(21)	4.68	Nebraska	8.13	0.45	(26)	4.03
Arizona	7.29	0.23	(41)	-6.74	Nevada	7.57	0.39	(34)	-3.24
Arkansas	6.51	0.31	(48)	-16.74 ^c	New Hampshire	10.73	0.62	(03)	37.30 ^c
California	6.38	0.10	(49)	-18.40 ^c	New Jersey	7.52	0.20	(36)	-3.86
Colorado	7.26	0.29	(42)	-7.15	New Mexico	7.13	0.41	(44)	-8.82
Connecticut	8.19	0.32	(19)	4.71	New York	7.57	0.13	(33)	-3.13
Delaware	9.07	0.69	(10)	15.98 ^c	North Carolina	7.47	0.20	(38)	-4.46
D.C.	10.99	0.97	(02)	40.51 ^c	North Dakota	7.16	0.67	(43)	-8.41
Florida	7.08	0.12	(45)	-9.46	Ohio	9.49	0.19	(07)	21.32 ^c
Georgia	7.47	0.22	(37)	-4.41	Oklahoma	7.56	0.30	(35)	-3.31
Hawaii	4.87	0.39	(51)	-37.74 ^c	Oregon	8.82	0.32	(13)	12.81
Idaho	8.11	0.52	(27)	3.76	Pennsylvania	9.31	0.17	(08)	19.07 ^c
Illinois	8.19	0.18	(20)	4.69	Rhode Island	9.60	0.62	(06)	22.83 ^c
Indiana	8.36	0.25	(17)	6.88	South Carolina	8.59	0.30	(14)	9.82
Iowa	8.84	0.35	(12)	13.07	South Dakota	8.26	0.66	(18)	5.69
Kansas	7.02	0.34	(46)	-10.23	Tennessee	7.44	0.24	(39)	-4.89
Kentucky	8.17	0.30	(22)	4.56	Texas	6.70	0.13	(47)	-14.29
Louisiana	8.06	0.30	(29)	3.12	Utah	5.86	0.38	(50)	-24.99 ^c
Maine	11.59	0.59	(01)	48.23 ^c	Vermont	8.15	0.76	(24)	4.18
Maryland	8.01	0.27	(30)	2.51	Virginia	7.75	0.23	(31)	-0.92
Massachusetts	10.14	0.26	(04)	29.69 ^c	Washington	8.16	0.25	(23)	4.43
Michigan	8.87	0.20	(11)	13.49	West Virginia	8.13	0.42	(25)	4.04
Minnesota	8.07	0.27	(28)	3.20	Wisconsin	9.30	0.27	(09)	18.89 ^c
Mississippi	8.43	0.37	(15)	7.77	Wyoming	10.02	0.94	(05)	28.11 ^c
Missouri	8.37	0.25	(16)	7.00					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 8.17
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.71	0.01		
High Five States				
D.C.	2.34	0.37	(01)	37.07 ^c
Rhode Island	2.34	0.26	(02)	36.64 ^c
New Hampshire	2.33	0.25	(03)	36.12 ^c
Montana	2.32	0.29	(04)	35.95 ^c
Alaska	2.26	0.49	(05)	32.35 ^c
Low Five States				
Alabama	1.35	0.10	(47)	-21.24 ^c
Oklahoma	1.27	0.11	(48)	-25.68 ^c
Kansas	1.18	0.12	(49)	-31.14 ^c
Hawaii	0.95	0.16	(50)	-44.44 ^c
Utah	0.83	0.13	(51)	-51.71 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.35	0.10	(47)	-21.24 ^c	Montana	2.32	0.29	(04)	35.95 ^c
Alaska	2.26	0.49	(05)	32.35 ^c	Nebraska	1.55	0.17	(34)	-9.03
Arizona	1.40	0.09	(46)	-17.96 ^c	Nevada	1.63	0.17	(29)	-4.43
Arkansas	1.46	0.13	(41)	-14.69	New Hampshire	2.33	0.25	(03)	36.12 ^c
California	1.58	0.04	(32)	-7.42	New Jersey	1.84	0.08	(21)	7.41
Colorado	1.45	0.12	(42)	-14.99	New Mexico	1.48	0.17	(38)	-13.68
Connecticut	1.66	0.12	(28)	-2.98	New York	1.96	0.06	(12)	14.67
Delaware	1.88	0.28	(17)	10.16	North Carolina	1.73	0.08	(26)	1.44
D.C.	2.34	0.37	(01)	37.07 ^c	North Dakota	1.45	0.27	(43)	-15.42 ^c
Florida	1.58	0.05	(33)	-7.55	Ohio	1.93	0.07	(13)	12.98
Georgia	1.84	0.09	(20)	7.45	Oklahoma	1.27	0.11	(48)	-25.68 ^c
Hawaii	0.95	0.16	(50)	-44.44 ^c	Oregon	2.01	0.14	(08)	17.53 ^c
Idaho	1.47	0.20	(39)	-13.88	Pennsylvania	1.92	0.07	(14)	12.14
Illinois	1.84	0.07	(19)	7.90	Rhode Island	2.34	0.26	(02)	36.64 ^c
Indiana	1.73	0.10	(25)	1.49	South Carolina	1.82	0.12	(23)	6.47
Iowa	1.70	0.13	(27)	-0.56	South Dakota	1.44	0.25	(44)	-16.01 ^c
Kansas	1.18	0.12	(49)	-31.14 ^c	Tennessee	1.50	0.09	(37)	-12.08
Kentucky	1.53	0.11	(36)	-10.37	Texas	1.41	0.05	(45)	-17.58 ^c
Louisiana	1.55	0.11	(35)	-9.23	Utah	0.83	0.13	(51)	-51.71 ^c
Maine	1.90	0.21	(15)	11.28	Vermont	2.12	0.34	(07)	23.80 ^c
Maryland	1.90	0.11	(16)	11.06	Virginia	1.83	0.10	(22)	7.06
Massachusetts	1.99	0.10	(10)	16.61 ^c	Washington	1.84	0.11	(18)	7.92
Michigan	2.00	0.08	(09)	16.95 ^c	West Virginia	1.76	0.17	(24)	2.70
Minnesota	1.61	0.11	(31)	-5.53	Wisconsin	1.97	0.11	(11)	15.46 ^c
Mississippi	1.47	0.14	(40)	-14.21	Wyoming	2.24	0.41	(06)	31.30 ^c
Missouri	1.62	0.10	(30)	-5.38					



Age-Adjusted Death Rates per 100,000 Quantile Interval

- 2.01 - 2.34
- 1.89 - 2.00
- 1.74 - 1.88
- 1.56 - 1.73
- 1.45 - 1.55
- 0.83 - 1.44

U.S. Rate: 1.71

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.



Table 9.15
Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.44	0.01		
High Five States				
Mississippi	0.60	0.07	(01)	36.00 ^c
Arkansas	0.57	0.06	(02)	29.27 ^c
Oklahoma	0.56	0.06	(03)	27.12 ^c
Louisiana	0.56	0.05	(04)	26.35 ^c
Pennsylvania	0.54	0.03	(05)	21.09 ^c
Low Five States				
Idaho	0.36	0.07	(47)	-19.38 ^c
Utah	0.35	0.06	(48)	-20.02 ^c
Arizona	0.35	0.03	(49)	-21.55 ^c
Hawaii	0.34	0.07	(50)	-23.20 ^c
New Hampshire	0.32	0.07	(51)	-27.58 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.45	0.04	(20)	2.34	Montana	0.43	0.09	(30)	-3.81
Alaska	0.49	0.16	(10)	10.97	Nebraska	0.47	0.07	(12)	6.83
Arizona	0.35	0.03	(49)	-21.55 ^c	Nevada	0.39	0.06	(38)	-11.59
Arkansas	0.57	0.06	(02)	29.27 ^c	New Hampshire	0.32	0.07	(51)	-27.58 ^c
California	0.44	0.02	(26)	-0.95	New Jersey	0.51	0.03	(08)	15.18 ^c
Colorado	0.37	0.04	(46)	-16.06 ^c	New Mexico	0.39	0.06	(40)	-12.18
Connecticut	0.39	0.05	(41)	-12.50	New York	0.44	0.02	(23)	0.35
Delaware	0.43	0.10	(31)	-4.00	North Carolina	0.44	0.03	(25)	-0.82
D.C.	0.44	0.12	(29)	-1.38	North Dakota	0.42	0.12	(32)	-6.15
Florida	0.39	0.02	(39)	-11.78	Ohio	0.49	0.03	(11)	10.57
Georgia	0.38	0.03	(43)	-13.24	Oklahoma	0.56	0.06	(03)	27.12 ^c
Hawaii	0.34	0.07	(50)	-23.20 ^c	Oregon	0.46	0.05	(19)	2.77
Idaho	0.36	0.07	(47)	-19.38 ^c	Pennsylvania	0.54	0.03	(05)	21.09 ^c
Illinois	0.45	0.03	(22)	0.77	Rhode Island	0.47	0.09	(13)	6.01
Indiana	0.39	0.04	(42)	-12.62	South Carolina	0.40	0.04	(37)	-10.48
Iowa	0.44	0.05	(27)	-1.03	South Dakota	0.51	0.12	(07)	16.15 ^c
Kansas	0.41	0.05	(33)	-6.75	Tennessee	0.44	0.04	(28)	-1.09
Kentucky	0.52	0.05	(06)	18.23 ^c	Texas	0.47	0.02	(14)	5.67
Louisiana	0.56	0.05	(04)	26.35 ^c	Utah	0.35	0.06	(48)	-20.02 ^c
Maine	0.37	0.07	(45)	-15.98 ^c	Vermont	0.38	0.11	(44)	-14.04
Maryland	0.45	0.04	(21)	1.11	Virginia	0.40	0.03	(35)	-9.17
Massachusetts	0.41	0.03	(34)	-8.20	Washington	0.40	0.04	(36)	-9.56
Michigan	0.46	0.03	(17)	4.00	West Virginia	0.50	0.07	(09)	12.58
Minnesota	0.46	0.04	(18)	3.25	Wisconsin	0.47	0.04	(15)	5.34
Mississippi	0.60	0.07	(01)	36.00 ^c	Wyoming	0.46	0.13	(16)	4.77
Missouri	0.44	0.04	(24)	-0.41					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 9.16
Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.55	0.01		
High Five States				
Mississippi	0.75	0.11	(01)	36.87 ^c
Arkansas	0.72	0.10	(02)	32.25 ^c
Kentucky	0.72	0.09	(03)	31.98 ^c
Pennsylvania	0.67	0.05	(04)	23.37 ^c
Louisiana	0.67	0.09	(05)	23.00 ^c
Low Five States				
New Mexico	0.42	0.10	(47)	-23.88 ^c
Nevada	0.41	0.09	(48)	-24.17 ^c
Hawaii	0.36	0.11	(49)	-34.31 ^c
Alaska	0.30	0.12	(50)	-44.34 ^c
Vermont	0.28	0.14	(51)	-48.73 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.58	0.07	(17)	5.32	Montana	0.57	0.16	(18)	4.50
Alaska	0.30	0.12	(50)	-44.34 ^c	Nebraska	0.63	0.12	(06)	14.33
Arizona	0.47	0.06	(39)	-13.28	Nevada	0.41	0.09	(48)	-24.17 ^c
Arkansas	0.72	0.10	(02)	32.25 ^c	New Hampshire	0.42	0.13	(45)	-22.67 ^c
California	0.55	0.03	(19)	-0.16	New Jersey	0.59	0.05	(13)	7.34
Colorado	0.51	0.08	(28)	-7.41	New Mexico	0.42	0.10	(47)	-23.88 ^c
Connecticut	0.59	0.09	(12)	8.26	New York	0.54	0.04	(20)	-0.85
Delaware	0.51	0.17	(27)	-5.87	North Carolina	0.61	0.06	(09)	10.99
D.C.	0.50	0.19	(32)	-8.57	North Dakota	0.52	0.18	(26)	-5.67
Florida	0.51	0.03	(29)	-7.44	Ohio	0.61	0.05	(08)	12.40
Georgia	0.43	0.05	(44)	-20.77 ^c	Oklahoma	0.62	0.09	(07)	13.02
Hawaii	0.36	0.11	(49)	-34.31 ^c	Oregon	0.52	0.08	(24)	-4.64
Idaho	0.42	0.12	(46)	-23.31 ^c	Pennsylvania	0.67	0.05	(04)	23.37 ^c
Illinois	0.58	0.05	(14)	6.37	Rhode Island	0.60	0.15	(11)	8.98
Indiana	0.54	0.06	(22)	-2.13	South Carolina	0.52	0.08	(25)	-4.73
Iowa	0.44	0.08	(43)	-18.99 ^c	South Dakota	0.54	0.17	(21)	-2.03
Kansas	0.50	0.09	(34)	-9.00	Tennessee	0.46	0.06	(41)	-16.64 ^c
Kentucky	0.72	0.09	(03)	31.98 ^c	Texas	0.58	0.04	(16)	5.63
Louisiana	0.67	0.09	(05)	23.00 ^c	Utah	0.50	0.11	(31)	-8.50
Maine	0.47	0.12	(40)	-13.71	Vermont	0.28	0.14	(51)	-48.73 ^c
Maryland	0.49	0.06	(37)	-9.58	Virginia	0.48	0.05	(38)	-13.10
Massachusetts	0.53	0.06	(23)	-2.91	Washington	0.49	0.06	(36)	-9.54
Michigan	0.58	0.05	(15)	6.05	West Virginia	0.50	0.11	(30)	-8.17
Minnesota	0.50	0.07	(33)	-8.77	Wisconsin	0.60	0.07	(10)	9.20
Mississippi	0.75	0.11	(01)	36.87 ^c	Wyoming	0.44	0.21	(42)	-18.65 ^c
Missouri	0.50	0.06	(35)	-9.20					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.



Table 9.17
Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.36	0.01		
High Five States				
Alaska	0.62	0.26	(01)	73.31 ^c
Oklahoma	0.53	0.07	(02)	46.82 ^c
Vermont	0.51	0.18	(03)	41.23 ^c
Mississippi	0.51	0.08	(04)	41.00 ^c
South Dakota	0.50	0.16	(05)	40.34 ^c
Low Five States				
Indiana	0.29	0.04	(47)	-20.33 ^c
Colorado	0.25	0.05	(48)	-30.13 ^c
Utah	0.24	0.07	(49)	-32.66 ^c
Arizona	0.23	0.04	(50)	-35.64 ^c
Connecticut	0.22	0.05	(51)	-39.43 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.35	0.05	(28)	-1.54	Montana	0.29	0.10	(45)	-19.24 ^c
Alaska	0.62	0.26	(01)	73.31 ^c	Nebraska	0.34	0.08	(32)	-4.35
Arizona	0.23	0.04	(50)	-35.64 ^c	Nevada	0.35	0.08	(30)	-1.63
Arkansas	0.43	0.07	(12)	19.24 ^c	New Hampshire	0.29	0.09	(44)	-18.70 ^c
California	0.35	0.02	(29)	-1.63	New Jersey	0.45	0.04	(09)	25.85 ^c
Colorado	0.25	0.05	(48)	-30.13 ^c	New Mexico	0.36	0.08	(25)	0.84
Connecticut	0.22	0.05	(51)	-39.43 ^c	New York	0.37	0.03	(22)	2.99
Delaware	0.37	0.13	(21)	3.13	North Carolina	0.30	0.04	(42)	-16.85 ^c
D.C.	0.34	0.14	(33)	-4.51	North Dakota	0.33	0.15	(36)	-7.42
Florida	0.29	0.02	(46)	-19.41 ^c	Ohio	0.39	0.03	(17)	9.82
Georgia	0.34	0.04	(34)	-5.50	Oklahoma	0.53	0.07	(02)	46.82 ^c
Hawaii	0.32	0.09	(39)	-10.97	Oregon	0.40	0.06	(15)	11.51
Idaho	0.32	0.10	(38)	-10.89	Pennsylvania	0.43	0.03	(13)	18.89 ^c
Illinois	0.35	0.03	(31)	-3.11	Rhode Island	0.38	0.11	(19)	6.23
Indiana	0.29	0.04	(47)	-20.33 ^c	South Carolina	0.30	0.05	(43)	-17.80 ^c
Iowa	0.43	0.07	(11)	20.50 ^c	South Dakota	0.50	0.16	(05)	40.34 ^c
Kansas	0.36	0.07	(26)	-0.34	Tennessee	0.42	0.05	(14)	17.81 ^c
Kentucky	0.37	0.06	(23)	2.51	Texas	0.37	0.03	(20)	3.81
Louisiana	0.49	0.06	(06)	36.41 ^c	Utah	0.24	0.07	(49)	-32.66 ^c
Maine	0.31	0.09	(41)	-14.98	Vermont	0.51	0.18	(03)	41.23 ^c
Maryland	0.40	0.05	(16)	11.49	Virginia	0.34	0.04	(35)	-6.30
Massachusetts	0.31	0.04	(40)	-14.75	Washington	0.32	0.04	(37)	-10.01
Michigan	0.37	0.04	(24)	2.17	West Virginia	0.48	0.09	(08)	32.52 ^c
Minnesota	0.44	0.06	(10)	22.71 ^c	Wisconsin	0.36	0.05	(27)	-0.66
Mississippi	0.51	0.08	(04)	41.00 ^c	Wyoming	0.49	0.18	(07)	35.15 ^c
Missouri	0.38	0.05	(18)	6.57					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 11.15
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006


Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.12	0.02		
High Five States				
Louisiana	5.26	0.16	(01)	27.71 ^c
Oklahoma	5.25	0.17	(02)	27.32 ^c
North Dakota	5.13	0.37	(03)	24.43 ^c
Indiana	4.92	0.12	(04)	19.35 ^c
Nebraska	4.84	0.23	(05)	17.42 ^c
Low Five States				
Florida	3.51	0.06	(47)	-14.94
New York	3.47	0.06	(48)	-15.83 ^c
Utah	3.44	0.20	(49)	-16.53 ^c
D.C.	3.05	0.33	(50)	-26.02 ^c
Hawaii	2.45	0.19	(51)	-40.63 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.01	0.13	(32)	-2.59	Montana	4.35	0.29	(22)	5.49
Alaska	3.91	0.47	(36)	-5.14	Nebraska	4.84	0.23	(05)	17.42 ^c
Arizona	3.96	0.12	(35)	-3.79	Nevada	3.90	0.19	(37)	-5.36
Arkansas	4.44	0.17	(18)	7.71	New Hampshire	4.11	0.25	(30)	-0.23
California	3.65	0.05	(45)	-11.51	New Jersey	3.85	0.09	(39)	-6.56
Colorado	3.83	0.14	(40)	-7.13	New Mexico	4.45	0.22	(17)	8.06
Connecticut	3.51	0.14	(46)	-14.87	New York	3.47	0.06	(48)	-15.83 ^c
Delaware	3.71	0.29	(43)	-9.90	North Carolina	4.22	0.10	(26)	2.48
D.C.	3.05	0.33	(50)	-26.02 ^c	North Dakota	5.13	0.37	(03)	24.43 ^c
Florida	3.51	0.06	(47)	-14.94	Ohio	4.52	0.09	(15)	9.58
Georgia	4.12	0.11	(29)	0.05	Oklahoma	5.25	0.17	(02)	27.32 ^c
Hawaii	2.45	0.19	(51)	-40.63 ^c	Oregon	3.98	0.14	(34)	-3.50
Idaho	4.21	0.25	(27)	2.23	Pennsylvania	4.26	0.08	(24)	3.29
Illinois	4.34	0.08	(23)	5.25	Rhode Island	4.68	0.28	(10)	13.50
Indiana	4.92	0.12	(04)	19.35 ^c	South Carolina	4.17	0.14	(28)	1.11
Iowa	4.78	0.17	(07)	15.99 ^c	South Dakota	4.79	0.33	(06)	16.14 ^c
Kansas	4.58	0.18	(13)	11.23	Tennessee	4.41	0.12	(19)	6.90
Kentucky	4.59	0.15	(11)	11.46	Texas	4.76	0.07	(09)	15.45 ^c
Louisiana	5.26	0.16	(01)	27.71 ^c	Utah	3.44	0.20	(49)	-16.53 ^c
Maine	4.37	0.24	(21)	6.04	Vermont	3.90	0.34	(38)	-5.38
Maryland	3.75	0.12	(42)	-8.98	Virginia	3.99	0.11	(33)	-3.30
Massachusetts	3.80	0.10	(41)	-7.90	Washington	4.11	0.12	(31)	-0.28
Michigan	4.25	0.09	(25)	3.16	West Virginia	4.50	0.20	(16)	9.18
Minnesota	4.40	0.13	(20)	6.66	Wisconsin	4.55	0.13	(14)	10.51
Mississippi	4.58	0.18	(12)	11.24	Wyoming	3.71	0.38	(44)	-10.01
Missouri	4.76	0.12	(08)	15.52 ^c					



Age-Adjusted Death Rates per 100,000 Quantile Interval



U.S. Rate: 4.12

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 11.16
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.96	0.03		
High Five States				
Louisiana	7.53	0.29	(01)	26.50 ^c
Oklahoma	7.40	0.30	(02)	24.26 ^c
North Dakota	7.20	0.67	(03)	20.95 ^c
South Dakota	6.96	0.60	(04)	16.81 ^c
Indiana	6.96	0.23	(05)	16.80 ^c
Low Five States				
Connecticut	4.96	0.25	(47)	-16.70 ^c
Vermont	4.95	0.58	(48)	-16.82 ^c
Utah	4.75	0.35	(49)	-20.23 ^c
D.C.	4.46	0.62	(50)	-25.09 ^c
Hawaii	3.44	0.33	(51)	-42.26 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.76	0.24	(34)	-3.25	Montana	6.36	0.53	(21)	6.87
Alaska	5.89	0.86	(31)	-1.06	Nebraska	6.57	0.41	(12)	10.24
Arizona	5.60	0.21	(39)	-5.96	Nevada	5.55	0.34	(41)	-6.84
Arkansas	6.39	0.31	(20)	7.38	New Hampshire	5.49	0.45	(42)	-7.78
California	5.28	0.09	(44)	-11.37	New Jersey	5.67	0.17	(38)	-4.81
Colorado	5.72	0.27	(37)	-4.03	New Mexico	6.22	0.39	(22)	4.44
Connecticut	4.96	0.25	(47)	-16.70 ^c	New York	5.07	0.11	(46)	-14.85
Delaware	5.56	0.55	(40)	-6.64	North Carolina	6.48	0.20	(15)	8.80
D.C.	4.46	0.62	(50)	-25.09 ^c	North Dakota	7.20	0.67	(03)	20.95 ^c
Florida	5.16	0.10	(45)	-13.27	Ohio	6.48	0.16	(14)	8.82
Georgia	6.04	0.21	(26)	1.47	Oklahoma	7.40	0.30	(02)	24.26 ^c
Hawaii	3.44	0.33	(51)	-42.26 ^c	Oregon	5.74	0.26	(36)	-3.54
Idaho	6.00	0.45	(29)	0.83	Pennsylvania	6.00	0.14	(30)	0.82
Illinois	6.45	0.16	(17)	8.35	Rhode Island	6.71	0.52	(10)	12.74
Indiana	6.96	0.23	(05)	16.80 ^c	South Carolina	6.08	0.26	(25)	2.12
Iowa	6.90	0.31	(06)	15.92 ^c	South Dakota	6.96	0.60	(04)	16.81 ^c
Kansas	6.61	0.33	(11)	10.96	Tennessee	6.42	0.23	(18)	7.76
Kentucky	6.90	0.28	(07)	15.91 ^c	Texas	6.77	0.13	(09)	13.65
Louisiana	7.53	0.29	(01)	26.50 ^c	Utah	4.75	0.35	(49)	-20.23 ^c
Maine	6.18	0.43	(23)	3.76	Vermont	4.95	0.58	(48)	-16.82 ^c
Maryland	5.46	0.22	(43)	-8.25	Virginia	6.01	0.20	(28)	0.94
Massachusetts	5.81	0.20	(33)	-2.51	Washington	5.75	0.21	(35)	-3.50
Michigan	6.01	0.17	(27)	0.95	West Virginia	6.15	0.36	(24)	3.34
Minnesota	6.42	0.24	(19)	7.73	Wisconsin	6.48	0.23	(16)	8.75
Mississippi	6.55	0.33	(13)	9.97	Wyoming	5.82	0.71	(32)	-2.29
Missouri	6.88	0.23	(08)	15.51 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 11.17
Cancer of the Kidney And Renal Pelvis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.71	0.02		
High Five States				
North Dakota	3.58	0.43	(01)	31.90 ^c
Louisiana	3.52	0.17	(02)	29.56 ^c
Oklahoma	3.48	0.18	(03)	28.28 ^c
Nebraska	3.47	0.26	(04)	28.05 ^c
Indiana	3.37	0.14	(05)	24.37 ^c
Low Five States				
Florida	2.15	0.06	(47)	-20.66 ^c
Alaska	2.02	0.47	(48)	-25.49 ^c
D.C.	1.90	0.34	(49)	-30.12 ^c
Wyoming	1.77	0.35	(50)	-34.91 ^c
Hawaii	1.62	0.21	(51)	-40.26 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.68	0.14	(32)	-1.36	Montana	2.76	0.31	(28)	1.76
Alaska	2.02	0.47	(48)	-25.49 ^c	Nebraska	3.47	0.26	(04)	28.05 ^c
Arizona	2.59	0.13	(33)	-4.53	Nevada	2.46	0.21	(39)	-9.45
Arkansas	2.87	0.18	(23)	5.94	New Hampshire	3.13	0.29	(10)	15.31 ^c
California	2.38	0.05	(43)	-12.37	New Jersey	2.54	0.10	(36)	-6.56
Colorado	2.37	0.15	(44)	-12.53	New Mexico	3.00	0.24	(19)	10.45
Connecticut	2.47	0.15	(38)	-9.11	New York	2.31	0.06	(46)	-15.03 ^c
Delaware	2.42	0.31	(41)	-10.76	North Carolina	2.57	0.10	(34)	-5.29
D.C.	1.90	0.34	(49)	-30.12 ^c	North Dakota	3.58	0.43	(01)	31.90 ^c
Florida	2.15	0.06	(47)	-20.66 ^c	Ohio	3.09	0.09	(12)	13.81
Georgia	2.80	0.11	(25)	3.03	Oklahoma	3.48	0.18	(03)	28.28 ^c
Hawaii	1.62	0.21	(51)	-40.26 ^c	Oregon	2.55	0.15	(35)	-5.92
Idaho	2.69	0.27	(31)	-0.93	Pennsylvania	2.93	0.08	(20)	7.92
Illinois	2.75	0.09	(29)	1.47	Rhode Island	3.26	0.31	(06)	20.21 ^c
Indiana	3.37	0.14	(05)	24.37 ^c	South Carolina	2.74	0.15	(30)	1.06
Iowa	3.11	0.18	(11)	14.64	South Dakota	3.08	0.36	(13)	13.33
Kansas	3.03	0.20	(17)	11.60	Tennessee	2.89	0.13	(22)	6.38
Kentucky	2.86	0.15	(24)	5.48	Texas	3.19	0.08	(07)	17.70 ^c
Louisiana	3.52	0.17	(02)	29.56 ^c	Utah	2.31	0.22	(45)	-14.72
Maine	3.01	0.27	(18)	11.07	Vermont	3.05	0.41	(15)	12.23
Maryland	2.50	0.13	(37)	-7.71	Virginia	2.45	0.11	(40)	-9.88
Massachusetts	2.39	0.11	(42)	-11.78	Washington	2.77	0.13	(26)	2.21
Michigan	2.91	0.10	(21)	7.38	West Virginia	3.15	0.22	(09)	16.05 ^c
Minnesota	2.76	0.14	(27)	1.82	Wisconsin	3.04	0.14	(16)	11.88
Mississippi	3.07	0.19	(14)	13.07	Wyoming	1.77	0.35	(50)	-34.91 ^c
Missouri	3.19	0.14	(08)	17.64 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.



Table 12.15
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.25	0.01		
High Five States				
D.C.	2.48	0.30	(01)	97.67 ^c
West Virginia	1.88	0.13	(02)	49.85 ^c
Louisiana	1.77	0.09	(03)	40.75 ^c
Delaware	1.65	0.19	(04)	31.31 ^c
Kentucky	1.63	0.09	(05)	29.88 ^c
Low Five States				
Montana	0.81	0.13	(47)	-35.76 ^c
Colorado	0.79	0.06	(48)	-36.96 ^c
Hawaii	0.71	0.10	(49)	-43.71 ^c
Alaska	0.53	0.17	(50)	-58.00 ^c
Utah	0.49	0.07	(51)	-61.22 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.45	0.08	(09)	15.84 ^c	Montana	0.81	0.13	(47)	-35.76 ^c
Alaska	0.53	0.17	(50)	-58.00 ^c	Nebraska	0.91	0.10	(42)	-27.76 ^c
Arizona	0.95	0.06	(40)	-24.50 ^c	Nevada	1.08	0.10	(31)	-14.15
Arkansas	1.28	0.09	(22)	1.94	New Hampshire	1.02	0.13	(36)	-18.80 ^c
California	0.98	0.02	(39)	-21.86 ^c	New Jersey	1.32	0.05	(18)	5.13
Colorado	0.79	0.06	(48)	-36.96 ^c	New Mexico	1.05	0.11	(33)	-16.11 ^c
Connecticut	1.15	0.08	(28)	-8.22	New York	1.34	0.04	(16)	7.01
Delaware	1.65	0.19	(04)	31.31 ^c	North Carolina	1.42	0.06	(12)	13.47
D.C.	2.48	0.30	(01)	97.67 ^c	North Dakota	1.00	0.17	(37)	-19.94 ^c
Florida	1.47	0.04	(08)	17.43 ^c	Ohio	1.43	0.05	(11)	13.78
Georgia	1.39	0.06	(14)	10.88	Oklahoma	1.20	0.08	(27)	-3.98
Hawaii	0.71	0.10	(49)	-43.71 ^c	Oregon	0.92	0.07	(41)	-27.03 ^c
Idaho	0.89	0.12	(43)	-29.20 ^c	Pennsylvania	1.35	0.04	(15)	7.60
Illinois	1.30	0.05	(19)	3.85	Rhode Island	1.26	0.15	(24)	0.10
Indiana	1.21	0.06	(26)	-3.70	South Carolina	1.48	0.08	(07)	17.83 ^c
Iowa	1.07	0.08	(32)	-14.60	South Dakota	0.81	0.14	(46)	-35.30 ^c
Kansas	1.04	0.09	(35)	-16.97 ^c	Tennessee	1.55	0.07	(06)	23.32 ^c
Kentucky	1.63	0.09	(05)	29.88 ^c	Texas	1.30	0.04	(20)	3.40
Louisiana	1.77	0.09	(03)	40.75 ^c	Utah	0.49	0.07	(51)	-61.22 ^c
Maine	1.12	0.12	(29)	-10.36	Vermont	1.11	0.18	(30)	-11.16
Maryland	1.45	0.07	(10)	15.40 ^c	Virginia	1.23	0.06	(25)	-1.70
Massachusetts	1.29	0.06	(21)	2.90	Washington	0.99	0.06	(38)	-21.17 ^c
Michigan	1.40	0.05	(13)	11.47	West Virginia	1.88	0.13	(02)	49.85 ^c
Minnesota	0.85	0.06	(45)	-32.26 ^c	Wisconsin	1.05	0.06	(34)	-16.16 ^c
Mississippi	1.33	0.10	(17)	6.12	Wyoming	0.85	0.18	(44)	-32.15 ^c
Missouri	1.28	0.06	(23)	1.91					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.



Table 12.16
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	2.27	0.02		
High Five States				
D.C.	5.14	0.68	(01)	126.35 ^c
Louisiana	3.25	0.19	(02)	43.19 ^c
West Virginia	3.18	0.26	(03)	39.96 ^c
Kentucky	2.99	0.18	(04)	31.56 ^c
Delaware	2.90	0.38	(05)	27.84 ^c
Low Five States				
Colorado	1.37	0.13	(47)	-39.52 ^c
North Dakota	1.37	0.29	(48)	-39.64 ^c
Montana	1.08	0.21	(49)	-52.42 ^c
Alaska	0.95	0.33	(50)	-58.30 ^c
Utah	0.84	0.15	(51)	-63.11 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.68	0.16	(11)	17.79 ^c	Montana	1.08	0.21	(49)	-52.42 ^c
Alaska	0.95	0.33	(50)	-58.30 ^c	Nebraska	1.69	0.21	(37)	-25.40 ^c
Arizona	1.66	0.11	(40)	-26.85 ^c	Nevada	1.74	0.19	(36)	-23.30 ^c
Arkansas	2.40	0.19	(20)	5.79	New Hampshire	1.68	0.25	(38)	-26.20 ^c
California	1.81	0.05	(35)	-20.18 ^c	New Jersey	2.46	0.11	(17)	8.28
Colorado	1.37	0.13	(47)	-39.52 ^c	New Mexico	1.86	0.21	(31)	-18.09 ^c
Connecticut	2.15	0.16	(26)	-5.30	New York	2.48	0.08	(16)	9.13
Delaware	2.90	0.38	(05)	27.84 ^c	North Carolina	2.75	0.13	(08)	21.21 ^c
D.C.	5.14	0.68	(01)	126.35 ^c	North Dakota	1.37	0.29	(48)	-39.64 ^c
Florida	2.59	0.07	(12)	14.19	Ohio	2.58	0.10	(14)	13.69
Georgia	2.59	0.13	(13)	14.07	Oklahoma	2.04	0.16	(28)	-10.00
Hawaii	1.38	0.21	(46)	-39.29 ^c	Oregon	1.48	0.13	(43)	-34.79 ^c
Idaho	1.38	0.22	(45)	-39.20 ^c	Pennsylvania	2.43	0.09	(19)	7.05
Illinois	2.34	0.09	(22)	2.90	Rhode Island	2.25	0.30	(25)	-0.80
Indiana	2.15	0.13	(27)	-5.31	South Carolina	2.87	0.18	(06)	26.47 ^c
Iowa	1.97	0.16	(30)	-13.24	South Dakota	1.54	0.28	(41)	-32.06 ^c
Kansas	1.82	0.17	(34)	-19.83 ^c	Tennessee	2.85	0.15	(07)	25.38 ^c
Kentucky	2.99	0.18	(04)	31.56 ^c	Texas	2.44	0.08	(18)	7.36
Louisiana	3.25	0.19	(02)	43.19 ^c	Utah	0.84	0.15	(51)	-63.11 ^c
Maine	2.00	0.24	(29)	-12.06	Vermont	1.86	0.34	(32)	-18.13 ^c
Maryland	2.68	0.16	(10)	17.92 ^c	Virginia	2.32	0.13	(23)	2.15
Massachusetts	2.28	0.13	(24)	0.15	Washington	1.67	0.11	(39)	-26.37 ^c
Michigan	2.48	0.11	(15)	9.36	West Virginia	3.18	0.26	(03)	39.96 ^c
Minnesota	1.52	0.12	(42)	-33.19 ^c	Wisconsin	1.84	0.12	(33)	-19.17 ^c
Mississippi	2.71	0.21	(09)	19.36 ^c	Wyoming	1.44	0.35	(44)	-36.60 ^c
Missouri	2.34	0.13	(21)	3.03					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 12.17
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.47	0.01		
High Five States				
West Virginia	0.84	0.12	(01)	77.60 ^c
D.C.	0.67	0.20	(02)	42.23 ^c
North Dakota	0.67	0.19	(03)	42.23 ^c
Louisiana	0.64	0.07	(04)	35.10 ^c
Delaware	0.62	0.16	(05)	31.43 ^c
Low Five States				
Nebraska	0.31	0.08	(47)	-35.47 ^c
Utah	0.22	0.07	(48)	-53.28 ^c
South Dakota	0.16	0.08	(49)	-65.49 ^c
Hawaii	0.16	0.07	(50)	-66.35 ^c
Alaska	0.13	0.13	(51)	-73.45 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.53	0.06	(19)	11.86	Montana	0.61	0.15	(06)	28.00 ^c
Alaska	0.13	0.13	(51)	-73.45 ^c	Nebraska	0.31	0.08	(47)	-35.47 ^c
Arizona	0.34	0.05	(43)	-28.10 ^c	Nevada	0.51	0.10	(23)	7.76
Arkansas	0.40	0.07	(38)	-14.90	New Hampshire	0.51	0.12	(21)	8.52
California	0.35	0.02	(42)	-26.79 ^c	New Jersey	0.49	0.04	(25)	3.78
Colorado	0.34	0.06	(44)	-29.06 ^c	New Mexico	0.37	0.09	(40)	-21.12 ^c
Connecticut	0.41	0.06	(36)	-12.71	New York	0.51	0.03	(24)	7.59
Delaware	0.62	0.16	(05)	31.43 ^c	North Carolina	0.48	0.04	(26)	1.96
D.C.	0.67	0.20	(02)	42.23 ^c	North Dakota	0.67	0.19	(03)	42.23 ^c
Florida	0.54	0.03	(16)	14.78	Ohio	0.55	0.04	(12)	16.52 ^c
Georgia	0.52	0.05	(20)	8.89	Oklahoma	0.55	0.07	(13)	16.03 ^c
Hawaii	0.16	0.07	(50)	-66.35 ^c	Oregon	0.45	0.07	(28)	-4.39
Idaho	0.55	0.12	(11)	17.00 ^c	Pennsylvania	0.55	0.04	(14)	15.92 ^c
Illinois	0.51	0.04	(22)	8.21	Rhode Island	0.57	0.13	(09)	20.26 ^c
Indiana	0.54	0.06	(18)	13.47	South Carolina	0.41	0.06	(37)	-13.64
Iowa	0.39	0.07	(39)	-16.84 ^c	South Dakota	0.16	0.08	(49)	-65.49 ^c
Kansas	0.43	0.07	(32)	-9.34	Tennessee	0.54	0.06	(17)	14.14
Kentucky	0.60	0.07	(07)	26.89 ^c	Texas	0.43	0.03	(33)	-9.36
Louisiana	0.64	0.07	(04)	35.10 ^c	Utah	0.22	0.07	(48)	-53.28 ^c
Maine	0.42	0.10	(35)	-11.04	Vermont	0.44	0.15	(30)	-7.24
Maryland	0.56	0.06	(10)	17.40 ^c	Virginia	0.42	0.05	(34)	-10.72
Massachusetts	0.57	0.05	(08)	20.53 ^c	Washington	0.44	0.05	(29)	-6.54
Michigan	0.55	0.04	(15)	15.13 ^c	West Virginia	0.84	0.12	(01)	77.60 ^c
Minnesota	0.33	0.05	(45)	-31.29 ^c	Wisconsin	0.43	0.05	(31)	-9.06
Mississippi	0.31	0.06	(46)	-35.40 ^c	Wyoming	0.35	0.16	(41)	-26.53 ^c
Missouri	0.47	0.05	(27)	0.05					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 13.23
Leukemia

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	7.30	0.02		
High Five States				
Idaho	8.40	0.36	(01)	15.12 ^c
Wisconsin	8.26	0.17	(02)	13.22
Iowa	8.22	0.22	(03)	12.60
Missouri	8.03	0.16	(04)	10.09
Arkansas	8.03	0.23	(05)	10.09
Low Five States				
Utah	6.81	0.28	(47)	-6.60
California	6.80	0.07	(48)	-6.79
Arizona	6.43	0.15	(49)	-11.93
New Mexico	6.35	0.26	(50)	-12.93
Hawaii	5.32	0.28	(51)	-27.10 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.42	0.18	(24)	1.74	Montana	7.54	0.38	(22)	3.30
Alaska	7.06	0.65	(38)	-3.20	Nebraska	7.43	0.28	(23)	1.87
Arizona	6.43	0.15	(49)	-11.93	Nevada	6.86	0.26	(43)	-5.91
Arkansas	8.03	0.23	(05)	10.09	New Hampshire	7.22	0.33	(28)	-1.03
California	6.80	0.07	(48)	-6.79	New Jersey	7.26	0.13	(27)	-0.51
Colorado	6.86	0.19	(44)	-5.99	New Mexico	6.35	0.26	(50)	-12.93
Connecticut	7.08	0.19	(36)	-2.91	New York	6.85	0.08	(45)	-6.08
Delaware	7.61	0.42	(17)	4.35	North Carolina	7.10	0.13	(33)	-2.61
D.C.	6.83	0.49	(46)	-6.45	North Dakota	7.10	0.44	(34)	-2.62
Florida	7.06	0.08	(39)	-3.24	Ohio	7.73	0.11	(15)	5.98
Georgia	7.07	0.14	(37)	-3.10	Oklahoma	7.78	0.21	(12)	6.70
Hawaii	5.32	0.28	(51)	-27.10 ^c	Oregon	7.42	0.20	(25)	1.71
Idaho	8.40	0.36	(01)	15.12 ^c	Pennsylvania	7.55	0.10	(21)	3.54
Illinois	7.78	0.11	(13)	6.59	Rhode Island	7.05	0.34	(40)	-3.30
Indiana	7.83	0.16	(10)	7.28	South Carolina	7.20	0.19	(29)	-1.29
Iowa	8.22	0.22	(03)	12.60	South Dakota	7.73	0.42	(14)	5.99
Kansas	7.64	0.23	(16)	4.72	Tennessee	7.56	0.16	(20)	3.56
Kentucky	7.61	0.19	(18)	4.34	Texas	7.19	0.09	(30)	-1.47
Louisiana	7.11	0.18	(31)	-2.55	Utah	6.81	0.28	(47)	-6.60
Maine	7.81	0.32	(11)	7.03	Vermont	7.90	0.49	(07)	8.29
Maryland	7.10	0.17	(35)	-2.67	Virginia	7.11	0.14	(32)	-2.55
Massachusetts	6.95	0.14	(41)	-4.77	Washington	7.56	0.16	(19)	3.56
Michigan	7.89	0.12	(08)	8.12	West Virginia	7.42	0.26	(26)	1.66
Minnesota	7.85	0.18	(09)	7.54	Wisconsin	8.26	0.17	(02)	13.22
Mississippi	7.97	0.24	(06)	9.28	Wyoming	6.92	0.53	(42)	-5.11
Missouri	8.03	0.16	(04)	10.09					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 13.24
Leukemia

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	9.81	0.04		
High Five States				
Maine	11.29	0.60	(01)	15.12 ^c
Iowa	11.20	0.40	(02)	14.13
Missouri	11.05	0.30	(03)	12.60
Wisconsin	11.02	0.30	(04)	12.30
Oklahoma	10.87	0.38	(05)	10.82
Low Five States				
Nevada	8.86	0.44	(47)	-9.72
New Mexico	8.35	0.46	(48)	-14.90
D.C.	8.34	0.87	(49)	-14.99
Arizona	8.34	0.25	(50)	-15.01 ^c
Hawaii	7.25	0.48	(51)	-26.13 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.92	0.33	(27)	1.17	Montana	10.08	0.68	(23)	2.75
Alaska	9.60	1.19	(34)	-2.09	Nebraska	10.19	0.51	(21)	3.90
Arizona	8.34	0.25	(50)	-15.01 ^c	Nevada	8.86	0.44	(47)	-9.72
Arkansas	10.81	0.42	(06)	10.21	New Hampshire	9.97	0.62	(25)	1.64
California	9.00	0.12	(46)	-8.28	New Jersey	9.76	0.23	(30)	-0.48
Colorado	9.08	0.34	(45)	-7.42	New Mexico	8.35	0.46	(48)	-14.90
Connecticut	9.54	0.35	(38)	-2.77	New York	9.24	0.15	(42)	-5.81
Delaware	9.87	0.75	(28)	0.62	North Carolina	9.72	0.24	(32)	-0.87
D.C.	8.34	0.87	(49)	-14.99	North Dakota	10.73	0.82	(09)	9.34
Florida	9.38	0.14	(40)	-4.38	Ohio	10.36	0.21	(16)	5.58
Georgia	9.44	0.26	(39)	-3.80	Oklahoma	10.87	0.38	(05)	10.82
Hawaii	7.25	0.48	(51)	-26.13 ^c	Oregon	10.06	0.35	(24)	2.56
Idaho	10.37	0.60	(15)	5.66	Pennsylvania	10.13	0.18	(22)	3.30
Illinois	10.73	0.21	(08)	9.42	Rhode Island	9.17	0.61	(44)	-6.57
Indiana	10.71	0.29	(10)	9.17	South Carolina	9.94	0.35	(26)	1.28
Iowa	11.20	0.40	(02)	14.13	South Dakota	10.74	0.75	(07)	9.47
Kansas	10.24	0.41	(19)	4.43	Tennessee	10.69	0.30	(11)	8.93
Kentucky	10.24	0.35	(20)	4.34	Texas	9.80	0.16	(29)	-0.07
Louisiana	9.58	0.34	(35)	-2.33	Utah	9.36	0.50	(41)	-4.62
Maine	11.29	0.60	(01)	15.12 ^c	Vermont	10.63	0.89	(13)	8.38
Maryland	9.69	0.31	(33)	-1.24	Virginia	9.55	0.26	(37)	-2.65
Massachusetts	9.18	0.26	(43)	-6.43	Washington	10.30	0.29	(18)	5.05
Michigan	10.53	0.23	(14)	7.35	West Virginia	9.73	0.47	(31)	-0.82
Minnesota	10.66	0.32	(12)	8.64	Wisconsin	11.02	0.30	(04)	12.30
Mississippi	10.33	0.43	(17)	5.33	Wyoming	9.56	0.95	(36)	-2.56
Missouri	11.05	0.30	(03)	12.60					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 13.25
Leukemia

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.51	0.03		
High Five States				
Idaho	6.78	0.43	(01)	23.06 ^c
Mississippi	6.42	0.28	(02)	16.60 ^c
Wisconsin	6.26	0.20	(03)	13.63
Iowa	6.14	0.25	(04)	11.45
Delaware	6.13	0.50	(05)	11.31
Low Five States				
Arizona	4.84	0.17	(47)	-12.24
Utah	4.83	0.31	(48)	-12.29
Wyoming	4.78	0.58	(49)	-13.29
North Dakota	4.53	0.46	(50)	-17.77 ^c
Hawaii	3.72	0.32	(51)	-32.57 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.73	0.20	(18)	3.92	Montana	5.64	0.45	(21)	2.36
Alaska	5.06	0.73	(45)	-8.21	Nebraska	5.45	0.32	(30)	-1.08
Arizona	4.84	0.17	(47)	-12.24	Nevada	5.21	0.31	(42)	-5.47
Arkansas	6.10	0.27	(06)	10.65	New Hampshire	5.37	0.38	(35)	-2.52
California	5.17	0.08	(44)	-6.15	New Jersey	5.51	0.14	(24)	0.01
Colorado	5.33	0.22	(39)	-3.35	New Mexico	4.88	0.31	(46)	-11.52
Connecticut	5.48	0.22	(27)	-0.59	New York	5.22	0.09	(41)	-5.18
Delaware	6.13	0.50	(05)	11.31	North Carolina	5.33	0.15	(38)	-3.33
D.C.	5.69	0.58	(19)	3.26	North Dakota	4.53	0.46	(50)	-17.77 ^c
Florida	5.21	0.09	(43)	-5.52	Ohio	5.95	0.13	(08)	8.06
Georgia	5.49	0.16	(26)	-0.38	Oklahoma	5.61	0.23	(22)	1.73
Hawaii	3.72	0.32	(51)	-32.57 ^c	Oregon	5.45	0.23	(31)	-1.17
Idaho	6.78	0.43	(01)	23.06 ^c	Pennsylvania	5.78	0.12	(16)	4.97
Illinois	5.81	0.13	(14)	5.40	Rhode Island	5.67	0.41	(20)	2.89
Indiana	5.94	0.18	(09)	7.80	South Carolina	5.36	0.21	(36)	-2.71
Iowa	6.14	0.25	(04)	11.45	South Dakota	5.38	0.47	(34)	-2.41
Kansas	5.82	0.27	(13)	5.58	Tennessee	5.52	0.18	(23)	0.13
Kentucky	5.81	0.22	(15)	5.36	Texas	5.33	0.10	(37)	-3.19
Louisiana	5.38	0.21	(33)	-2.32	Utah	4.83	0.31	(48)	-12.29
Maine	5.40	0.36	(32)	-2.05	Vermont	5.90	0.56	(12)	7.09
Maryland	5.31	0.19	(40)	-3.59	Virginia	5.46	0.16	(29)	-0.99
Massachusetts	5.48	0.17	(28)	-0.60	Washington	5.51	0.18	(25)	-0.09
Michigan	5.97	0.14	(07)	8.36	West Virginia	5.76	0.31	(17)	4.51
Minnesota	5.92	0.20	(11)	7.50	Wisconsin	6.26	0.20	(03)	13.63
Mississippi	6.42	0.28	(02)	16.60 ^c	Wyoming	4.78	0.58	(49)	-13.29
Missouri	5.94	0.18	(10)	7.73					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.



Table 14.15
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.11	0.02		
High Five States				
Hawaii	7.27	0.32	(01)	42.21 ^c
Texas	6.81	0.08	(02)	33.17 ^c
Louisiana	6.72	0.18	(03)	31.47 ^c
California	6.33	0.06	(04)	23.69 ^c
New Mexico	6.20	0.26	(05)	21.15 ^c
Low Five States				
Nebraska	3.76	0.20	(47)	-26.43 ^c
North Dakota	3.75	0.32	(48)	-26.70 ^c
Iowa	3.66	0.15	(49)	-28.41 ^c
Vermont	3.52	0.32	(50)	-31.13 ^c
Utah	3.19	0.19	(51)	-37.55 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.21	0.15	(13)	1.80	Montana	3.81	0.27	(46)	-25.59 ^c
Alaska	6.14	0.58	(06)	19.98 ^c	Nebraska	3.76	0.20	(47)	-26.43 ^c
Arizona	4.87	0.13	(22)	-4.84	Nevada	5.65	0.23	(09)	10.40
Arkansas	5.50	0.19	(10)	7.51	New Hampshire	4.47	0.26	(32)	-12.66
California	6.33	0.06	(04)	23.69 ^c	New Jersey	4.97	0.10	(17)	-2.79
Colorado	4.31	0.15	(38)	-15.65 ^c	New Mexico	6.20	0.26	(05)	21.15 ^c
Connecticut	4.71	0.16	(27)	-7.92	New York	5.20	0.07	(14)	1.70
Delaware	4.64	0.33	(29)	-9.20	North Carolina	4.47	0.10	(33)	-12.67
D.C.	5.97	0.46	(08)	16.74 ^c	North Dakota	3.75	0.32	(48)	-26.70 ^c
Florida	4.93	0.07	(19)	-3.66	Ohio	4.39	0.08	(37)	-14.08
Georgia	4.42	0.11	(34)	-13.52	Oklahoma	4.47	0.16	(31)	-12.50
Hawaii	7.27	0.32	(01)	42.21 ^c	Oregon	4.41	0.15	(36)	-13.84
Idaho	3.99	0.25	(44)	-22.03 ^c	Pennsylvania	4.70	0.08	(28)	-8.13
Illinois	5.06	0.09	(15)	-1.11	Rhode Island	5.24	0.30	(12)	2.46
Indiana	4.61	0.12	(30)	-9.87	South Carolina	4.84	0.15	(23)	-5.40
Iowa	3.66	0.15	(49)	-28.41 ^c	South Dakota	3.87	0.30	(45)	-24.42 ^c
Kansas	4.07	0.17	(42)	-20.43 ^c	Tennessee	4.84	0.13	(24)	-5.44
Kentucky	4.93	0.15	(20)	-3.66	Texas	6.81	0.08	(02)	33.17 ^c
Louisiana	6.72	0.18	(03)	31.47 ^c	Utah	3.19	0.19	(51)	-37.55 ^c
Maine	4.16	0.23	(41)	-18.61 ^c	Vermont	3.52	0.32	(50)	-31.13 ^c
Maryland	4.83	0.13	(25)	-5.51	Virginia	4.41	0.11	(35)	-13.70
Massachusetts	5.03	0.12	(16)	-1.55	Washington	5.41	0.13	(11)	5.86
Michigan	4.93	0.10	(18)	-3.62	West Virginia	4.29	0.20	(40)	-16.18 ^c
Minnesota	4.31	0.13	(39)	-15.75 ^c	Wisconsin	4.78	0.13	(26)	-6.61
Mississippi	6.11	0.21	(07)	19.54 ^c	Wyoming	4.05	0.40	(43)	-20.73 ^c
Missouri	4.87	0.13	(21)	-4.73					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 14.16
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.50	0.03		
High Five States				
Hawaii	10.48	0.58	(01)	39.70 ^c
Texas	10.05	0.16	(02)	33.95 ^c
Louisiana	9.99	0.33	(03)	33.17 ^c
D.C.	9.37	0.89	(04)	24.83 ^c
Mississippi	9.16	0.39	(05)	22.12 ^c
Low Five States				
Wyoming	5.09	0.66	(47)	-32.17 ^c
Montana	5.07	0.46	(48)	-32.47 ^c
Vermont	4.69	0.56	(49)	-37.55 ^c
North Dakota	4.64	0.54	(50)	-38.17 ^c
Utah	4.21	0.32	(51)	-43.90 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.47	0.27	(17)	-0.47	Montana	5.07	0.46	(48)	-32.47 ^c
Alaska	8.89	1.07	(07)	18.56 ^c	Nebraska	5.59	0.37	(44)	-25.51 ^c
Arizona	7.02	0.23	(27)	-6.44	Nevada	7.86	0.39	(11)	4.80
Arkansas	7.54	0.34	(13)	0.52	New Hampshire	6.73	0.49	(30)	-10.32
California	8.98	0.11	(06)	19.71 ^c	New Jersey	7.47	0.20	(16)	-0.38
Colorado	6.04	0.27	(40)	-19.47 ^c	New Mexico	8.75	0.45	(08)	16.57 ^c
Connecticut	7.38	0.30	(20)	-1.66	New York	7.83	0.13	(12)	4.33
Delaware	7.49	0.62	(15)	-0.12	North Carolina	6.66	0.19	(32)	-11.25
D.C.	9.37	0.89	(04)	24.83 ^c	North Dakota	4.64	0.54	(50)	-38.17 ^c
Florida	7.45	0.12	(18)	-0.72	Ohio	6.50	0.16	(36)	-13.40
Georgia	6.60	0.21	(34)	-11.99	Oklahoma	6.52	0.28	(35)	-13.06
Hawaii	10.48	0.58	(01)	39.70 ^c	Oregon	6.18	0.26	(38)	-17.60 ^c
Idaho	5.75	0.44	(43)	-23.40 ^c	Pennsylvania	7.07	0.15	(25)	-5.73
Illinois	7.39	0.17	(19)	-1.55	Rhode Island	8.23	0.57	(09)	9.75
Indiana	6.60	0.22	(33)	-11.99	South Carolina	7.31	0.28	(22)	-2.57
Iowa	5.25	0.27	(46)	-30.00 ^c	South Dakota	5.41	0.53	(45)	-27.91 ^c
Kansas	5.97	0.31	(41)	-20.43 ^c	Tennessee	7.17	0.24	(24)	-4.48
Kentucky	6.96	0.28	(28)	-7.20	Texas	10.05	0.16	(02)	33.95 ^c
Louisiana	9.99	0.33	(03)	33.17 ^c	Utah	4.21	0.32	(51)	-43.90 ^c
Maine	6.14	0.43	(39)	-18.17 ^c	Vermont	4.69	0.56	(49)	-37.55 ^c
Maryland	7.32	0.25	(21)	-2.47	Virginia	6.68	0.21	(31)	-10.90
Massachusetts	8.07	0.23	(10)	7.50	Washington	7.51	0.23	(14)	0.07
Michigan	7.02	0.18	(26)	-6.42	West Virginia	5.91	0.36	(42)	-21.19 ^c
Minnesota	6.31	0.24	(37)	-15.96 ^c	Wisconsin	6.93	0.23	(29)	-7.66
Mississippi	9.16	0.39	(05)	22.12 ^c	Wyoming	5.09	0.66	(47)	-32.17 ^c
Missouri	7.28	0.23	(23)	-3.01					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 14.17
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.16	0.02		
High Five States				
Hawaii	4.43	0.34	(01)	40.23 ^c
Texas	4.17	0.09	(02)	31.79 ^c
Louisiana	4.11	0.18	(03)	29.95 ^c
California	4.07	0.07	(04)	28.83 ^c
New Mexico	3.98	0.28	(05)	25.94 ^c
Low Five States				
Idaho	2.44	0.26	(47)	-22.92 ^c
Iowa	2.41	0.16	(48)	-23.85 ^c
Utah	2.29	0.22	(49)	-27.60 ^c
Nebraska	2.26	0.21	(50)	-28.56 ^c
Delaware	2.24	0.30	(51)	-29.12 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.44	0.16	(11)	8.66	Montana	2.64	0.31	(40)	-16.40 ^c
Alaska	3.76	0.62	(08)	18.99 ^c	Nebraska	2.26	0.21	(50)	-28.56 ^c
Arizona	3.01	0.14	(19)	-4.85	Nevada	3.54	0.25	(10)	11.91
Arkansas	3.81	0.21	(07)	20.35 ^c	New Hampshire	2.59	0.26	(43)	-18.06 ^c
California	4.07	0.07	(04)	28.83 ^c	New Jersey	3.02	0.11	(18)	-4.54
Colorado	2.89	0.16	(27)	-8.59	New Mexico	3.98	0.28	(05)	25.94 ^c
Connecticut	2.65	0.16	(39)	-16.16 ^c	New York	3.13	0.07	(16)	-0.95
Delaware	2.24	0.30	(51)	-29.12 ^c	North Carolina	2.74	0.11	(35)	-13.36
D.C.	3.34	0.44	(12)	5.52	North Dakota	2.99	0.39	(20)	-5.43
Florida	2.76	0.07	(32)	-12.61	Ohio	2.75	0.09	(34)	-12.97
Georgia	2.74	0.11	(36)	-13.43	Oklahoma	2.76	0.16	(33)	-12.71
Hawaii	4.43	0.34	(01)	40.23 ^c	Oregon	2.87	0.17	(29)	-9.35
Idaho	2.44	0.26	(47)	-22.92 ^c	Pennsylvania	2.83	0.08	(31)	-10.39
Illinois	3.22	0.09	(15)	1.97	Rhode Island	2.95	0.29	(24)	-6.78
Indiana	2.99	0.13	(21)	-5.56	South Carolina	2.84	0.15	(30)	-10.31
Iowa	2.41	0.16	(48)	-23.85 ^c	South Dakota	2.56	0.33	(44)	-19.18 ^c
Kansas	2.50	0.18	(45)	-20.99 ^c	Tennessee	2.95	0.13	(25)	-6.80
Kentucky	3.33	0.17	(13)	5.31	Texas	4.17	0.09	(02)	31.79 ^c
Louisiana	4.11	0.18	(03)	29.95 ^c	Utah	2.29	0.22	(49)	-27.60 ^c
Maine	2.60	0.25	(42)	-17.86 ^c	Vermont	2.44	0.36	(46)	-22.84 ^c
Maryland	2.88	0.14	(28)	-8.88	Virginia	2.64	0.11	(41)	-16.52 ^c
Massachusetts	2.67	0.11	(38)	-15.68 ^c	Washington	3.59	0.15	(09)	13.54
Michigan	3.23	0.11	(14)	2.29	West Virginia	2.98	0.22	(22)	-5.84
Minnesota	2.67	0.14	(37)	-15.65 ^c	Wisconsin	2.95	0.13	(23)	-6.65
Mississippi	3.83	0.22	(06)	21.29 ^c	Wyoming	3.03	0.47	(17)	-4.05
Missouri	2.94	0.13	(26)	-7.02					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 15.21
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	53.38	0.06		
High Five States				
Kentucky	77.68	0.60	(01)	45.54 ^c
Arkansas	68.68	0.67	(02)	28.67 ^c
Tennessee	68.38	0.47	(03)	28.10 ^c
West Virginia	68.27	0.79	(04)	27.90 ^c
Mississippi	67.26	0.69	(05)	26.01 ^c
Low Five States				
California	42.84	0.17	(47)	-19.75 ^c
Colorado	39.99	0.46	(48)	-25.09 ^c
New Mexico	37.37	0.63	(49)	-29.98 ^c
Hawaii	37.22	0.73	(50)	-30.27 ^c
Utah	24.42	0.53	(51)	-54.24 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	63.19	0.51	(09)	18.37 ^c	Montana	51.10	1.00	(33)	-4.27
Alaska	52.68	1.73	(29)	-1.31	Nebraska	48.33	0.73	(40)	-9.45 ^c
Arizona	45.33	0.39	(45)	-15.08 ^c	Nevada	58.11	0.74	(16)	8.87 ^c
Arkansas	68.68	0.67	(02)	28.67 ^c	New Hampshire	53.95	0.91	(24)	1.08
California	42.84	0.17	(47)	-19.75 ^c	New Jersey	49.23	0.33	(35)	-7.78 ^c
Colorado	39.99	0.46	(48)	-25.09 ^c	New Mexico	37.37	0.63	(49)	-29.98 ^c
Connecticut	48.64	0.51	(39)	-8.87 ^c	New York	46.28	0.21	(42)	-13.30 ^c
Delaware	61.37	1.18	(12)	14.97 ^c	North Carolina	59.17	0.37	(15)	10.86 ^c
D.C.	50.80	1.34	(34)	-4.83	North Dakota	45.41	1.13	(44)	-14.92 ^c
Florida	53.30	0.22	(28)	-0.15	Ohio	60.43	0.31	(13)	13.21 ^c
Georgia	57.82	0.40	(17)	8.32 ^c	Oklahoma	63.52	0.58	(07)	19.00 ^c
Hawaii	37.22	0.73	(50)	-30.27 ^c	Oregon	54.85	0.54	(21)	2.76
Idaho	43.64	0.82	(46)	-18.25 ^c	Pennsylvania	53.79	0.27	(25)	0.78
Illinois	54.56	0.30	(23)	2.21	Rhode Island	53.79	0.96	(26)	0.76
Indiana	63.34	0.45	(08)	18.66 ^c	South Carolina	59.73	0.53	(14)	11.91 ^c
Iowa	52.32	0.56	(31)	-1.98	South Dakota	49.07	1.07	(36)	-8.07 ^c
Kansas	54.74	0.62	(22)	2.55	Tennessee	68.38	0.47	(03)	28.10 ^c
Kentucky	77.68	0.60	(01)	45.54 ^c	Texas	52.11	0.24	(32)	-2.38
Louisiana	65.66	0.55	(06)	23.01 ^c	Utah	24.42	0.53	(51)	-54.24 ^c
Maine	61.88	0.90	(11)	15.93 ^c	Vermont	49.02	1.21	(37)	-8.16 ^c
Maryland	55.30	0.46	(20)	3.61	Virginia	56.50	0.40	(18)	5.85 ^c
Massachusetts	53.34	0.39	(27)	-0.06	Washington	52.51	0.42	(30)	-1.62
Michigan	56.29	0.33	(19)	5.45 ^c	West Virginia	68.27	0.79	(04)	27.90 ^c
Minnesota	46.17	0.43	(43)	-13.50 ^c	Wisconsin	48.74	0.41	(38)	-8.69 ^c
Mississippi	67.26	0.69	(05)	26.01 ^c	Wyoming	46.52	1.36	(41)	-12.85 ^c
Missouri	63.10	0.45	(10)	18.22 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 15.22
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	70.49	0.11		
High Five States				
Kentucky	107.59	1.11	(01)	52.63 ^c
Mississippi	100.18	1.31	(02)	42.12 ^c
Tennessee	97.83	0.89	(03)	38.79 ^c
Arkansas	96.90	1.22	(04)	37.46 ^c
Alabama	93.42	0.97	(05)	32.53 ^c
Low Five States				
California	53.04	0.28	(47)	-24.76 ^c
Hawaii	50.48	1.28	(48)	-28.39 ^c
Colorado	49.66	0.80	(49)	-29.55 ^c
New Mexico	47.26	1.08	(50)	-32.95 ^c
Utah	32.76	0.92	(51)	-53.53 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	93.42	0.97	(05)	32.53 ^c	Montana	61.70	1.64	(38)	-12.47 ^c
Alaska	63.63	2.87	(34)	-9.74 ^c	Nebraska	65.84	1.29	(31)	-6.60 ^c
Arizona	56.44	0.65	(45)	-19.94 ^c	Nevada	66.10	1.17	(30)	-6.22 ^c
Arkansas	96.90	1.22	(04)	37.46 ^c	New Hampshire	66.91	1.56	(29)	-5.08 ^c
California	53.04	0.28	(47)	-24.76 ^c	New Jersey	62.70	0.58	(37)	-11.05 ^c
Colorado	49.66	0.80	(49)	-29.55 ^c	New Mexico	47.26	1.08	(50)	-32.95 ^c
Connecticut	61.19	0.88	(39)	-13.20 ^c	New York	59.50	0.38	(42)	-15.59 ^c
Delaware	78.65	2.05	(16)	11.57 ^c	North Carolina	83.71	0.70	(13)	18.75 ^c
D.C.	74.34	2.57	(18)	5.47 ^c	North Dakota	60.48	1.95	(41)	-14.20 ^c
Florida	68.27	0.37	(27)	-3.15	Ohio	81.55	0.57	(14)	15.69 ^c
Georgia	84.07	0.77	(12)	19.27 ^c	Oklahoma	85.94	1.04	(10)	21.92 ^c
Hawaii	50.48	1.28	(48)	-28.39 ^c	Oregon	65.68	0.89	(32)	-6.83 ^c
Idaho	54.59	1.37	(46)	-22.56 ^c	Pennsylvania	72.45	0.48	(22)	2.78
Illinois	72.59	0.53	(21)	2.98	Rhode Island	70.51	1.70	(26)	0.02
Indiana	85.27	0.81	(11)	20.97 ^c	South Carolina	86.20	0.99	(08)	22.28 ^c
Iowa	72.08	1.00	(23)	2.26	South Dakota	65.43	1.85	(33)	-7.18 ^c
Kansas	72.65	1.10	(20)	3.07	Tennessee	97.83	0.89	(03)	38.79 ^c
Kentucky	107.59	1.11	(01)	52.63 ^c	Texas	70.84	0.43	(25)	0.50
Louisiana	92.92	1.03	(06)	31.81 ^c	Utah	32.76	0.92	(51)	-53.53 ^c
Maine	79.16	1.56	(15)	12.30 ^c	Vermont	60.67	2.07	(40)	-13.94 ^c
Maryland	71.54	0.82	(24)	1.49	Virginia	76.44	0.73	(17)	8.43 ^c
Massachusetts	66.98	0.69	(28)	-4.98	Washington	63.56	0.71	(35)	-9.83 ^c
Michigan	73.48	0.59	(19)	4.25	West Virginia	92.76	1.41	(07)	31.59 ^c
Minnesota	58.66	0.74	(43)	-16.78 ^c	Wisconsin	62.93	0.71	(36)	-10.72 ^c
Mississippi	100.18	1.31	(02)	42.12 ^c	Wyoming	57.28	2.28	(44)	-18.74 ^c
Missouri	85.95	0.81	(09)	21.93 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 15.23
Cancer of the Lung And Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	40.88	0.07		
High Five States				
Kentucky	56.40	0.69	(01)	37.96 ^c
Nevada	51.18	0.95	(02)	25.19 ^c
West Virginia	50.41	0.91	(03)	23.32 ^c
Maine	49.44	1.08	(04)	20.94 ^c
Delaware	48.52	1.40	(05)	18.69 ^c
Low Five States				
North Dakota	34.19	1.35	(47)	-16.37 ^c
Colorado	33.17	0.56	(48)	-18.85 ^c
New Mexico	29.69	0.76	(49)	-27.36 ^c
Hawaii	26.44	0.84	(50)	-35.33 ^c
Utah	17.62	0.61	(51)	-56.90 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	41.75	0.55	(27)	2.13	Montana	42.97	1.25	(21)	5.13 ^c
Alaska	43.41	2.13	(20)	6.18 ^c	Nebraska	35.50	0.84	(43)	-13.17 ^c
Arizona	36.30	0.48	(42)	-11.19 ^c	Nevada	51.18	0.95	(02)	25.19 ^c
Arkansas	47.62	0.75	(07)	16.49 ^c	New Hampshire	44.72	1.11	(14)	9.40 ^c
California	35.33	0.20	(44)	-13.58 ^c	New Jersey	40.07	0.39	(33)	-1.98
Colorado	33.17	0.56	(48)	-18.85 ^c	New Mexico	29.69	0.76	(49)	-27.36 ^c
Connecticut	40.10	0.61	(32)	-1.91	New York	37.19	0.25	(40)	-9.03 ^c
Delaware	48.52	1.40	(05)	18.69 ^c	North Carolina	41.76	0.41	(26)	2.16
D.C.	35.04	1.45	(46)	-14.29 ^c	North Dakota	34.19	1.35	(47)	-16.37 ^c
Florida	41.20	0.26	(28)	0.79	Ohio	45.54	0.36	(13)	11.41 ^c
Georgia	39.50	0.43	(34)	-3.37	Oklahoma	46.92	0.67	(09)	14.78 ^c
Hawaii	26.44	0.84	(50)	-35.33 ^c	Oregon	46.77	0.67	(10)	14.43 ^c
Idaho	35.07	1.00	(45)	-14.21 ^c	Pennsylvania	40.73	0.31	(29)	-0.36
Illinois	41.99	0.35	(24)	2.71	Rhode Island	42.60	1.13	(22)	4.20
Indiana	47.96	0.52	(06)	17.33 ^c	South Carolina	40.49	0.57	(31)	-0.94
Iowa	38.07	0.65	(38)	-6.86 ^c	South Dakota	36.89	1.27	(41)	-9.75 ^c
Kansas	41.82	0.74	(25)	2.29	Tennessee	47.39	0.52	(08)	15.93 ^c
Kentucky	56.40	0.69	(01)	37.96 ^c	Texas	38.30	0.27	(36)	-6.30 ^c
Louisiana	46.05	0.62	(12)	12.66 ^c	Utah	17.62	0.61	(51)	-56.90 ^c
Maine	49.44	1.08	(04)	20.94 ^c	Vermont	40.72	1.49	(30)	-0.38
Maryland	43.80	0.54	(18)	7.15 ^c	Virginia	42.24	0.46	(23)	3.34
Massachusetts	44.21	0.48	(16)	8.15 ^c	Washington	44.39	0.52	(15)	8.58 ^c
Michigan	43.94	0.39	(17)	7.49 ^c	West Virginia	50.41	0.91	(03)	23.32 ^c
Minnesota	37.35	0.52	(39)	-8.64 ^c	Wisconsin	38.74	0.50	(35)	-5.22 ^c
Mississippi	43.73	0.73	(19)	6.97 ^c	Wyoming	38.12	1.68	(37)	-6.75 ^c
Missouri	46.61	0.52	(11)	14.03 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 16.15
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.68	0.01		
High Five States				
Utah	3.46	0.20	(01)	29.10 ^c
New Hampshire	3.41	0.23	(02)	27.16 ^c
Oklahoma	3.37	0.14	(03)	25.68 ^c
Wyoming	3.32	0.36	(04)	24.05 ^c
Idaho	3.31	0.23	(05)	23.71 ^c
Low Five States				
New York	2.19	0.05	(47)	-18.32 ^c
Alaska	2.08	0.31	(48)	-22.50 ^c
Hawaii	1.76	0.16	(49)	-34.31 ^c
North Dakota	1.70	0.22	(50)	-36.73 ^c
D.C.	1.43	0.22	(51)	-46.57 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.80	0.11	(29)	4.61	Montana	2.86	0.24	(24)	6.77
Alaska	2.08	0.31	(48)	-22.50 ^c	Nebraska	2.91	0.18	(17)	8.43
Arizona	2.83	0.10	(25)	5.76	Nevada	2.86	0.16	(23)	6.86
Arkansas	2.49	0.13	(39)	-7.17	New Hampshire	3.41	0.23	(02)	27.16 ^c
California	2.57	0.04	(34)	-4.11	New Jersey	2.66	0.08	(31)	-0.64
Colorado	2.83	0.12	(26)	5.71	New Mexico	2.65	0.17	(32)	-1.20
Connecticut	2.83	0.12	(27)	5.62	New York	2.19	0.05	(47)	-18.32 ^c
Delaware	2.94	0.26	(14)	9.75	North Carolina	2.91	0.08	(16)	8.51
D.C.	1.43	0.22	(51)	-46.57 ^c	North Dakota	1.70	0.22	(50)	-36.73 ^c
Florida	2.89	0.05	(20)	7.91	Ohio	2.63	0.07	(33)	-1.92
Georgia	2.48	0.08	(40)	-7.32	Oklahoma	3.37	0.14	(03)	25.68 ^c
Hawaii	1.76	0.16	(49)	-34.31 ^c	Oregon	3.13	0.13	(08)	16.99 ^c
Idaho	3.31	0.23	(05)	23.71 ^c	Pennsylvania	2.90	0.06	(18)	8.06
Illinois	2.53	0.06	(37)	-5.53	Rhode Island	3.09	0.23	(10)	15.32 ^c
Indiana	2.89	0.10	(21)	7.88	South Carolina	2.54	0.11	(35)	-5.06
Iowa	2.46	0.12	(41)	-8.03	South Dakota	2.43	0.24	(42)	-9.32
Kansas	2.80	0.14	(28)	4.68	Tennessee	2.91	0.10	(15)	8.60
Kentucky	3.15	0.12	(06)	17.44 ^c	Texas	2.52	0.05	(38)	-6.02
Louisiana	2.22	0.10	(46)	-17.02 ^c	Utah	3.46	0.20	(01)	29.10 ^c
Maine	2.97	0.20	(13)	10.68	Vermont	3.00	0.30	(11)	11.86
Maryland	2.78	0.10	(30)	3.82	Virginia	2.89	0.09	(22)	7.85
Massachusetts	2.97	0.09	(12)	10.73	Washington	2.89	0.10	(19)	8.03
Michigan	2.40	0.07	(43)	-10.62	West Virginia	3.14	0.17	(07)	17.34 ^c
Minnesota	2.26	0.09	(44)	-15.51 ^c	Wisconsin	2.54	0.09	(36)	-5.31
Mississippi	2.23	0.13	(45)	-16.69 ^c	Wyoming	3.32	0.36	(04)	24.05 ^c
Missouri	3.11	0.10	(09)	16.04 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 16.16
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	3.94	0.02		
High Five States				
Utah	5.09	0.36	(01)	29.30 ^c
New Hampshire	5.09	0.42	(02)	29.19 ^c
Oklahoma	4.98	0.25	(03)	26.52 ^c
Idaho	4.91	0.40	(04)	24.62 ^c
Delaware	4.86	0.51	(05)	23.31 ^c
Low Five States				
New York	3.03	0.09	(47)	-23.05 ^c
Alaska	2.71	0.49	(48)	-31.18 ^c
Hawaii	2.52	0.28	(49)	-36.13 ^c
D.C.	2.38	0.47	(50)	-39.60 ^c
North Dakota	2.33	0.38	(51)	-40.81 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.08	0.20	(26)	3.62	Montana	3.58	0.40	(42)	-9.19
Alaska	2.71	0.49	(48)	-31.18 ^c	Nebraska	4.28	0.33	(21)	8.71
Arizona	4.36	0.18	(16)	10.66	Nevada	4.30	0.29	(19)	9.11
Arkansas	3.62	0.24	(39)	-8.02	New Hampshire	5.09	0.42	(02)	29.19 ^c
California	3.90	0.08	(31)	-0.94	New Jersey	3.95	0.14	(30)	0.21
Colorado	4.10	0.22	(25)	4.11	New Mexico	3.69	0.30	(35)	-6.41
Connecticut	4.40	0.24	(12)	11.82	New York	3.03	0.09	(47)	-23.05 ^c
Delaware	4.86	0.51	(05)	23.31 ^c	North Carolina	4.19	0.15	(23)	6.33
D.C.	2.38	0.47	(50)	-39.60 ^c	North Dakota	2.33	0.38	(51)	-40.81 ^c
Florida	4.32	0.09	(18)	9.60	Ohio	3.84	0.12	(33)	-2.50
Georgia	3.62	0.16	(40)	-8.05	Oklahoma	4.98	0.25	(03)	26.52 ^c
Hawaii	2.52	0.28	(49)	-36.13 ^c	Oregon	4.56	0.23	(08)	15.78 ^c
Idaho	4.91	0.40	(04)	24.62 ^c	Pennsylvania	4.20	0.12	(22)	6.58
Illinois	3.61	0.12	(41)	-8.30	Rhode Island	4.78	0.44	(06)	21.27 ^c
Indiana	4.14	0.18	(24)	5.15	South Carolina	3.86	0.21	(32)	-1.90
Iowa	3.39	0.21	(44)	-13.99	South Dakota	3.17	0.41	(46)	-19.56 ^c
Kansas	4.06	0.26	(27)	3.19	Tennessee	4.39	0.19	(14)	11.35
Kentucky	4.71	0.23	(07)	19.71 ^c	Texas	3.74	0.10	(34)	-4.93
Louisiana	3.44	0.20	(43)	-12.78	Utah	5.09	0.36	(01)	29.30 ^c
Maine	4.37	0.37	(15)	11.02	Vermont	4.47	0.56	(10)	13.56
Maryland	4.30	0.20	(20)	9.09	Virginia	4.06	0.17	(28)	3.14
Massachusetts	4.32	0.17	(17)	9.70	Washington	4.05	0.18	(29)	2.79
Michigan	3.65	0.13	(36)	-7.33	West Virginia	4.52	0.31	(09)	14.77
Minnesota	3.17	0.17	(45)	-19.54 ^c	Wisconsin	3.63	0.17	(38)	-7.79
Mississippi	3.65	0.25	(37)	-7.39	Wyoming	4.40	0.63	(13)	11.64
Missouri	4.44	0.18	(11)	12.82					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 16.17
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.74	0.01		
High Five States				
Wyoming	2.42	0.43	(01)	39.40 ^c
Montana	2.25	0.29	(02)	29.69 ^c
Oklahoma	2.10	0.15	(03)	20.69 ^c
Missouri	2.08	0.11	(04)	19.90 ^c
New Hampshire	2.08	0.24	(05)	19.76 ^c
Low Five States				
Louisiana	1.31	0.10	(47)	-24.87 ^c
Mississippi	1.23	0.12	(48)	-29.06 ^c
North Dakota	1.17	0.25	(49)	-32.79 ^c
Hawaii	1.09	0.17	(50)	-37.08 ^c
D.C.	0.89	0.23	(51)	-48.96 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.89	0.12	(20)	8.91	Montana	2.25	0.29	(02)	29.69 ^c
Alaska	1.32	0.34	(46)	-23.90 ^c	Nebraska	1.90	0.20	(19)	9.14
Arizona	1.55	0.10	(43)	-10.65	Nevada	1.60	0.17	(39)	-7.95
Arkansas	1.59	0.14	(40)	-8.41	New Hampshire	2.08	0.24	(05)	19.76 ^c
California	1.57	0.04	(42)	-9.86	New Jersey	1.74	0.08	(29)	0.08
Colorado	1.85	0.13	(24)	6.54	New Mexico	1.83	0.19	(26)	5.58
Connecticut	1.72	0.13	(31)	-1.27	New York	1.62	0.05	(37)	-6.96
Delaware	1.53	0.25	(44)	-11.79	North Carolina	1.98	0.09	(12)	13.84
D.C.	0.89	0.23	(51)	-48.96 ^c	North Dakota	1.17	0.25	(49)	-32.79 ^c
Florida	1.74	0.06	(27)	0.30	Ohio	1.74	0.07	(28)	0.29
Georgia	1.67	0.09	(34)	-3.94	Oklahoma	2.10	0.15	(03)	20.69 ^c
Hawaii	1.09	0.17	(50)	-37.08 ^c	Oregon	2.01	0.14	(10)	15.59 ^c
Idaho	1.88	0.23	(21)	8.30	Pennsylvania	1.95	0.07	(15)	12.09
Illinois	1.72	0.07	(30)	-0.91	Rhode Island	1.91	0.24	(16)	10.20
Indiana	1.97	0.11	(13)	13.25	South Carolina	1.58	0.11	(41)	-9.16
Iowa	1.70	0.14	(32)	-2.04	South Dakota	1.85	0.29	(25)	6.22
Kansas	1.87	0.16	(22)	7.44	Tennessee	1.86	0.11	(23)	6.98
Kentucky	1.96	0.13	(14)	12.80	Texas	1.61	0.05	(38)	-7.46
Louisiana	1.31	0.10	(47)	-24.87 ^c	Utah	2.08	0.21	(06)	19.74 ^c
Maine	1.91	0.22	(18)	9.79	Vermont	1.91	0.32	(17)	9.88
Maryland	1.70	0.10	(33)	-2.19	Virginia	2.07	0.10	(07)	18.94 ^c
Massachusetts	2.05	0.10	(08)	17.86 ^c	Washington	1.99	0.11	(11)	14.67
Michigan	1.48	0.07	(45)	-15.00 ^c	West Virginia	2.03	0.19	(09)	16.79 ^c
Minnesota	1.62	0.11	(36)	-6.94	Wisconsin	1.66	0.10	(35)	-4.68
Mississippi	1.23	0.12	(48)	-29.06 ^c	Wyoming	2.42	0.43	(01)	39.40 ^c
Missouri	2.08	0.11	(04)	19.90 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 18.15
Myeloma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.62	0.02		
High Five States				
D.C.	5.30	0.43	(01)	46.52 ^c
Alabama	4.40	0.14	(02)	21.46 ^c
South Carolina	4.26	0.14	(03)	17.75 ^c
South Dakota	4.22	0.31	(04)	16.54 ^c
Louisiana	4.21	0.14	(05)	16.30 ^c
Low Five States				
Vermont	3.20	0.31	(47)	-11.69
Arizona	3.17	0.10	(48)	-12.31
Nevada	3.10	0.17	(49)	-14.46
Florida	3.09	0.05	(50)	-14.65
Hawaii	2.25	0.18	(51)	-37.71 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.40	0.14	(02)	21.46 ^c	Montana	3.77	0.27	(21)	4.21
Alaska	3.30	0.43	(43)	-8.78	Nebraska	3.34	0.19	(40)	-7.75
Arizona	3.17	0.10	(48)	-12.31	Nevada	3.10	0.17	(49)	-14.46
Arkansas	3.64	0.16	(26)	0.49	New Hampshire	3.83	0.24	(19)	5.70
California	3.33	0.05	(41)	-8.01	New Jersey	3.58	0.09	(30)	-0.96
Colorado	3.51	0.14	(35)	-3.10	New Mexico	3.35	0.19	(38)	-7.39
Connecticut	3.59	0.14	(29)	-0.67	New York	3.20	0.06	(46)	-11.53
Delaware	3.29	0.27	(44)	-8.95	North Carolina	4.16	0.10	(06)	15.01 ^c
D.C.	5.30	0.43	(01)	46.52 ^c	North Dakota	3.89	0.32	(15)	7.49
Florida	3.09	0.05	(50)	-14.65	Ohio	3.92	0.08	(14)	8.44
Georgia	3.87	0.10	(16)	6.98	Oklahoma	3.26	0.13	(45)	-9.80
Hawaii	2.25	0.18	(51)	-37.71 ^c	Oregon	3.94	0.14	(13)	8.85
Idaho	3.55	0.23	(33)	-1.91	Pennsylvania	3.57	0.07	(32)	-1.24
Illinois	3.75	0.08	(22)	3.59	Rhode Island	3.35	0.24	(39)	-7.44
Indiana	3.74	0.11	(23)	3.21	South Carolina	4.26	0.14	(03)	17.75 ^c
Iowa	3.54	0.14	(34)	-2.17	South Dakota	4.22	0.31	(04)	16.54 ^c
Kansas	3.98	0.17	(12)	10.01	Tennessee	4.15	0.12	(07)	14.61
Kentucky	3.84	0.14	(18)	6.24	Texas	3.62	0.06	(28)	-0.08
Louisiana	4.21	0.14	(05)	16.30 ^c	Utah	3.31	0.20	(42)	-8.55
Maine	3.63	0.22	(27)	0.18	Vermont	3.20	0.31	(47)	-11.69
Maryland	4.05	0.12	(08)	11.98	Virginia	4.00	0.11	(10)	10.53
Massachusetts	3.49	0.10	(36)	-3.64	Washington	3.73	0.11	(24)	2.93
Michigan	4.00	0.09	(09)	10.66	West Virginia	3.72	0.18	(25)	2.85
Minnesota	3.85	0.12	(17)	6.28	Wisconsin	3.58	0.11	(31)	-1.01
Mississippi	3.99	0.17	(11)	10.27	Wyoming	3.47	0.37	(37)	-4.01
Missouri	3.78	0.11	(20)	4.41					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 18.16
Myeloma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	4.51	0.03		
High Five States				
D.C.	6.52	0.78	(01)	44.39 ^c
Louisiana	5.49	0.25	(02)	21.64 ^c
South Dakota	5.34	0.53	(03)	18.36 ^c
Alabama	5.29	0.24	(04)	17.19 ^c
Michigan	5.22	0.16	(05)	15.73 ^c
Low Five States				
Arizona	4.06	0.18	(47)	-9.98
Florida	3.96	0.09	(48)	-12.23
Nevada	3.74	0.28	(49)	-17.19 ^c
Alaska	2.99	0.59	(50)	-33.75 ^c
Hawaii	2.85	0.30	(51)	-36.84 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.29	0.24	(04)	17.19 ^c	Montana	5.07	0.47	(09)	12.29
Alaska	2.99	0.59	(50)	-33.75 ^c	Nebraska	4.11	0.32	(44)	-8.90
Arizona	4.06	0.18	(47)	-9.98	Nevada	3.74	0.28	(49)	-17.19 ^c
Arkansas	4.57	0.27	(28)	1.36	New Hampshire	4.44	0.40	(33)	-1.55
California	4.13	0.08	(42)	-8.37	New Jersey	4.39	0.15	(36)	-2.74
Colorado	4.51	0.24	(31)	-0.08	New Mexico	4.39	0.33	(35)	-2.74
Connecticut	4.21	0.23	(40)	-6.73	New York	4.07	0.10	(46)	-9.84
Delaware	4.11	0.48	(43)	-8.89	North Carolina	5.02	0.17	(11)	11.25
D.C.	6.52	0.78	(01)	44.39 ^c	North Dakota	5.09	0.56	(08)	12.76
Florida	3.96	0.09	(48)	-12.23	Ohio	4.76	0.14	(22)	5.59
Georgia	4.69	0.18	(23)	3.86	Oklahoma	4.28	0.24	(39)	-5.25
Hawaii	2.85	0.30	(51)	-36.84 ^c	Oregon	4.85	0.24	(19)	7.37
Idaho	4.52	0.40	(30)	0.10	Pennsylvania	4.32	0.12	(38)	-4.19
Illinois	4.53	0.13	(29)	0.31	Rhode Island	4.09	0.41	(45)	-9.37
Indiana	4.95	0.20	(14)	9.77	South Carolina	5.17	0.25	(07)	14.65
Iowa	4.48	0.25	(32)	-0.77	South Dakota	5.34	0.53	(03)	18.36 ^c
Kansas	4.93	0.29	(15)	9.14	Tennessee	5.20	0.21	(06)	15.28 ^c
Kentucky	4.98	0.24	(12)	10.33	Texas	4.43	0.11	(34)	-1.90
Louisiana	5.49	0.25	(02)	21.64 ^c	Utah	4.14	0.33	(41)	-8.26
Maine	4.88	0.40	(17)	8.19	Vermont	4.58	0.59	(27)	1.43
Maryland	4.91	0.22	(16)	8.90	Virginia	5.04	0.19	(10)	11.59
Massachusetts	4.36	0.18	(37)	-3.29	Washington	4.87	0.20	(18)	7.82
Michigan	5.22	0.16	(05)	15.73 ^c	West Virginia	4.78	0.33	(20)	5.97
Minnesota	4.98	0.22	(13)	10.31	Wisconsin	4.58	0.19	(26)	1.51
Mississippi	4.61	0.28	(25)	2.14	Wyoming	4.65	0.65	(24)	3.12
Missouri	4.77	0.19	(21)	5.78					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 18.17
Myeloma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.98	0.02		
High Five States				
D.C.	4.62	0.52	(01)	54.73 ^c
Alabama	3.77	0.16	(02)	26.52 ^c
South Carolina	3.66	0.17	(03)	22.84 ^c
North Carolina	3.56	0.12	(04)	19.31 ^c
Mississippi	3.50	0.20	(05)	17.47 ^c
Low Five States				
Wyoming	2.48	0.43	(47)	-16.83 ^c
Arizona	2.47	0.12	(48)	-17.23 ^c
Florida	2.41	0.06	(49)	-19.24 ^c
Vermont	2.32	0.35	(50)	-22.39 ^c
Hawaii	1.78	0.22	(51)	-40.39 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.77	0.16	(02)	26.52 ^c	Montana	2.77	0.31	(38)	-7.14
Alaska	3.46	0.60	(07)	16.14 ^c	Nebraska	2.79	0.23	(37)	-6.40
Arizona	2.47	0.12	(48)	-17.23 ^c	Nevada	2.54	0.21	(46)	-14.86
Arkansas	2.94	0.18	(28)	-1.57	New Hampshire	3.27	0.30	(14)	9.61
California	2.74	0.06	(39)	-8.17	New Jersey	3.04	0.11	(24)	1.77
Colorado	2.80	0.16	(36)	-6.15	New Mexico	2.55	0.22	(44)	-14.51
Connecticut	3.17	0.17	(18)	6.22	New York	2.62	0.07	(43)	-12.12
Delaware	2.72	0.33	(40)	-8.94	North Carolina	3.56	0.12	(04)	19.31 ^c
D.C.	4.62	0.52	(01)	54.73 ^c	North Dakota	2.94	0.38	(27)	-1.48
Florida	2.41	0.06	(49)	-19.24 ^c	Ohio	3.37	0.10	(09)	12.88
Georgia	3.32	0.13	(12)	11.19	Oklahoma	2.55	0.16	(45)	-14.58
Hawaii	1.78	0.22	(51)	-40.39 ^c	Oregon	3.25	0.18	(15)	9.07
Idaho	2.88	0.29	(34)	-3.60	Pennsylvania	3.06	0.08	(21)	2.60
Illinois	3.21	0.09	(17)	7.74	Rhode Island	2.90	0.29	(32)	-2.94
Indiana	2.92	0.13	(30)	-2.19	South Carolina	3.66	0.17	(03)	22.84 ^c
Iowa	2.92	0.17	(29)	-2.09	South Dakota	3.23	0.36	(16)	8.18
Kansas	3.32	0.20	(11)	11.22	Tennessee	3.42	0.14	(08)	14.52
Kentucky	3.06	0.16	(22)	2.47	Texas	3.02	0.08	(25)	1.14
Louisiana	3.33	0.16	(10)	11.67	Utah	2.69	0.24	(42)	-9.68
Maine	2.72	0.25	(41)	-8.96	Vermont	2.32	0.35	(50)	-22.39 ^c
Maryland	3.47	0.15	(06)	16.44 ^c	Virginia	3.29	0.13	(13)	10.36
Massachusetts	2.87	0.12	(35)	-3.83	Washington	2.91	0.13	(31)	-2.53
Michigan	3.13	0.10	(19)	4.94	West Virginia	2.96	0.22	(26)	-0.81
Minnesota	3.04	0.15	(23)	1.85	Wisconsin	2.89	0.13	(33)	-3.19
Mississippi	3.50	0.20	(05)	17.47 ^c	Wyoming	2.48	0.43	(47)	-16.83 ^c
Missouri	3.09	0.13	(20)	3.65					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 19.19
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	7.11	0.02		
High Five States				
Michigan	8.01	0.13	(01)	12.79
West Virginia	7.99	0.27	(02)	12.43
Oregon	7.93	0.20	(03)	11.61
Indiana	7.85	0.16	(04)	10.44
Iowa	7.82	0.21	(05)	10.13
Low Five States				
Georgia	6.44	0.13	(47)	-9.40
D.C.	6.29	0.47	(48)	-11.41
Nevada	6.25	0.25	(49)	-12.02
New Mexico	6.08	0.26	(50)	-14.43
Hawaii	5.68	0.29	(51)	-19.99 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.11	0.17	(25)	0.05	Montana	7.08	0.37	(27)	-0.31
Alaska	6.46	0.61	(46)	-9.09	Nebraska	7.43	0.28	(16)	4.55
Arizona	6.64	0.15	(40)	-6.59	Nevada	6.25	0.25	(49)	-12.02
Arkansas	6.98	0.21	(31)	-1.82	New Hampshire	7.38	0.34	(19)	3.93
California	6.70	0.07	(37)	-5.71	New Jersey	7.37	0.13	(21)	3.69
Colorado	6.64	0.19	(41)	-6.60	New Mexico	6.08	0.26	(50)	-14.43
Connecticut	6.99	0.19	(30)	-1.59	New York	6.60	0.08	(45)	-7.16
Delaware	6.87	0.40	(33)	-3.34	North Carolina	6.93	0.13	(32)	-2.40
D.C.	6.29	0.47	(48)	-11.41	North Dakota	7.00	0.44	(29)	-1.45
Florida	6.79	0.08	(35)	-4.38	Ohio	7.52	0.11	(13)	5.88
Georgia	6.44	0.13	(47)	-9.40	Oklahoma	7.35	0.20	(22)	3.49
Hawaii	5.68	0.29	(51)	-19.99 ^c	Oregon	7.93	0.20	(03)	11.61
Idaho	7.09	0.33	(26)	-0.16	Pennsylvania	7.76	0.10	(06)	9.27
Illinois	7.39	0.11	(18)	3.96	Rhode Island	7.13	0.34	(24)	0.38
Indiana	7.85	0.16	(04)	10.44	South Carolina	6.64	0.18	(39)	-6.53
Iowa	7.82	0.21	(05)	10.13	South Dakota	6.77	0.39	(36)	-4.73
Kansas	7.51	0.23	(14)	5.67	Tennessee	7.66	0.16	(08)	7.78
Kentucky	7.60	0.19	(09)	7.01	Texas	6.83	0.09	(34)	-3.82
Louisiana	7.76	0.19	(07)	9.25	Utah	6.63	0.28	(43)	-6.74
Maine	7.41	0.31	(17)	4.25	Vermont	7.00	0.46	(28)	-1.45
Maryland	6.62	0.16	(44)	-6.86	Virginia	6.69	0.14	(38)	-5.80
Massachusetts	7.37	0.14	(20)	3.76	Washington	7.57	0.16	(11)	6.60
Michigan	8.01	0.13	(01)	12.79	West Virginia	7.99	0.27	(02)	12.43
Minnesota	7.57	0.17	(12)	6.50	Wisconsin	7.50	0.16	(15)	5.59
Mississippi	6.63	0.22	(42)	-6.70	Wyoming	7.58	0.55	(10)	6.65
Missouri	7.28	0.15	(23)	2.42					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 19.20
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	8.98	0.04		
High Five States				
West Virginia	0.48	(01)	12.96	
Indiana ^b	0.28	(02)	12.68	
Michigan ^b	0.22	(03)	11.69	
Iowa ^b	0.37	(04)	10.92	
Minnesota	0.31	(05)	10.40	
Low Five States				
Alaska ^b	1.06	(47)	-11.11	
Wyoming ^b	0.87	(48)	-11.48	
New Mexico	0.43	(49)	-16.37 ^c	
Hawaii ^b	0.48	(50)	-20.50 ^c	
Nevada ^b	0.39	(51)	-20.73 ^c	

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	8.83	0.30	(29)	-1.67	Montana	8.80	0.63	(30)	-2.03
Alaska	7.98	1.06	(47)	-11.11	Nebraska	9.11	0.48	(27)	1.43
Arizona	8.15	0.25	(46)	-9.27	Nevada	7.12	0.39	(51)	-20.73 ^c
Arkansas	9.27	0.38	(22)	3.24	New Hampshire	9.15	0.59	(25)	1.88
California	8.50	0.11	(38)	-5.37	New Jersey	9.49	0.23	(16)	5.64
Colorado	8.47	0.33	(39)	-5.74	New Mexico	7.51	0.43	(49)	-16.37 ^c
Connecticut	9.10	0.34	(28)	1.34	New York	8.28	0.14	(43)	-7.84
Delaware	9.11	0.72	(26)	1.46	North Carolina	8.63	0.23	(33)	-3.93
D.C.	9.57	0.92	(15)	6.58	North Dakota	8.53	0.73	(37)	-5.05
Florida	8.56	0.13	(36)	-4.73	Ohio	9.67	0.20	(11)	7.67
Georgia	8.27	0.25	(44)	-7.98	Oklahoma	9.62	0.35	(14)	7.06
Hawaii	7.14	0.48	(50)	-20.50 ^c	Oregon	9.74	0.35	(08)	8.39
Idaho	8.57	0.54	(35)	-4.63	Pennsylvania	9.91	0.18	(06)	10.31
Illinois	9.39	0.19	(17)	4.57	Rhode Island	9.28	0.62	(21)	3.26
Indiana	10.12	0.28	(02)	12.68	South Carolina	8.26	0.31	(45)	-8.09
Iowa	9.96	0.37	(04)	10.92	South Dakota	8.57	0.67	(34)	-4.61
Kansas	9.66	0.40	(13)	7.52	Tennessee	9.73	0.29	(09)	8.35
Kentucky	9.84	0.35	(07)	9.60	Texas	8.45	0.15	(40)	-5.92
Louisiana	9.73	0.34	(10)	8.33	Utah	8.41	0.47	(41)	-6.34
Maine	9.32	0.55	(20)	3.75	Vermont	9.23	0.83	(24)	2.73
Maryland	8.65	0.29	(32)	-3.66	Virginia	8.30	0.24	(42)	-7.57
Massachusetts	9.34	0.26	(19)	4.00	Washington	9.67	0.28	(12)	7.61
Michigan	10.03	0.22	(03)	11.69	West Virginia	10.15	0.48	(01)	12.96
Minnesota	9.92	0.31	(05)	10.40	Wisconsin	9.38	0.28	(18)	4.44
Mississippi	8.67	0.39	(31)	-3.46	Wyoming	7.95	0.87	(48)	-11.48
Missouri	9.24	0.27	(23)	2.86					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 19.21
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.71	0.03		
High Five States				
Wyoming	7.30	0.73	(01)	27.75 ^c
Michigan	6.57	0.15	(02)	14.92
Oregon	6.53	0.25	(03)	14.25
Louisiana	6.38	0.23	(04)	11.70
West Virginia	6.34	0.32	(05)	10.92
Low Five States				
Mississippi	5.14	0.25	(47)	-9.99
Utah	5.11	0.32	(48)	-10.54
New Mexico	4.96	0.31	(49)	-13.12
Hawaii	4.46	0.34	(50)	-21.90 ^c
D.C.	3.99	0.48	(51)	-30.22 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.82	0.20	(23)	1.82	Montana	5.76	0.45	(25)	0.74
Alaska	5.29	0.76	(42)	-7.48	Nebraska	6.16	0.34	(11)	7.76
Arizona	5.39	0.18	(36)	-5.67	Nevada	5.40	0.31	(35)	-5.56
Arkansas	5.27	0.25	(43)	-7.73	New Hampshire	6.16	0.41	(10)	7.81
California	5.32	0.08	(40)	-6.90	New Jersey	5.84	0.15	(21)	2.16
Colorado	5.24	0.22	(44)	-8.28	New Mexico	4.96	0.31	(49)	-13.12
Connecticut	5.54	0.22	(31)	-3.06	New York	5.37	0.10	(37)	-6.03
Delaware	5.29	0.46	(41)	-7.39	North Carolina	5.69	0.15	(27)	-0.50
D.C.	3.99	0.48	(51)	-30.22 ^c	North Dakota	5.70	0.52	(26)	-0.28
Florida	5.35	0.09	(38)	-6.30	Ohio	5.96	0.13	(19)	4.33
Georgia	5.18	0.16	(45)	-9.38	Oklahoma	5.61	0.23	(29)	-1.81
Hawaii	4.46	0.34	(50)	-21.90 ^c	Oregon	6.53	0.25	(03)	14.25
Idaho	5.77	0.40	(24)	0.99	Pennsylvania	6.25	0.12	(07)	9.34
Illinois	5.99	0.13	(17)	4.84	Rhode Island	5.59	0.39	(30)	-2.27
Indiana	6.24	0.18	(08)	9.23	South Carolina	5.47	0.21	(34)	-4.34
Iowa	6.31	0.25	(06)	10.47	South Dakota	5.34	0.46	(39)	-6.52
Kansas	6.06	0.27	(15)	6.07	Tennessee	6.18	0.19	(09)	8.19
Kentucky	6.03	0.22	(16)	5.46	Texas	5.62	0.10	(28)	-1.73
Louisiana	6.38	0.23	(04)	11.70	Utah	5.11	0.32	(48)	-10.54
Maine	6.06	0.37	(14)	6.12	Vermont	5.49	0.54	(32)	-3.86
Maryland	5.15	0.18	(46)	-9.84	Virginia	5.48	0.16	(33)	-4.13
Massachusetts	6.08	0.17	(13)	6.45	Washington	5.99	0.19	(18)	4.84
Michigan	6.57	0.15	(02)	14.92	West Virginia	6.34	0.32	(05)	10.92
Minnesota	5.82	0.20	(22)	1.89	Wisconsin	6.10	0.19	(12)	6.83
Mississippi	5.14	0.25	(47)	-9.99	Wyoming	7.30	0.73	(01)	27.75 ^c
Missouri	5.91	0.18	(20)	3.42					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 20.15
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.58	0.01		
High Five States				
D.C.	3.93	0.37	(01)	52.52 ^c
Mississippi	3.48	0.15	(02)	35.04 ^c
Louisiana	3.37	0.12	(03)	30.58 ^c
South Carolina	3.10	0.12	(04)	20.36 ^c
Tennessee	3.08	0.10	(05)	19.24 ^c
Low Five States				
Montana	2.11	0.20	(47)	-18.26 ^c
Nebraska	2.03	0.15	(48)	-21.18 ^c
Connecticut	2.01	0.10	(49)	-22.03 ^c
New Mexico	1.96	0.14	(50)	-23.94 ^c
Utah	1.69	0.14	(51)	-34.46 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.00	0.11	(08)	16.47 ^c	Montana	2.11	0.20	(47)	-18.26 ^c
Alaska	2.85	0.39	(12)	10.36	Nebraska	2.03	0.15	(48)	-21.18 ^c
Arizona	2.14	0.09	(46)	-17.11 ^c	Nevada	2.82	0.16	(14)	9.17
Arkansas	3.07	0.14	(06)	18.87 ^c	New Hampshire	2.96	0.21	(10)	14.70
California	2.66	0.04	(21)	3.07	New Jersey	2.38	0.07	(35)	-7.91
Colorado	2.17	0.11	(43)	-15.87 ^c	New Mexico	1.96	0.14	(50)	-23.94 ^c
Connecticut	2.01	0.10	(49)	-22.03 ^c	New York	2.29	0.05	(38)	-11.26
Delaware	2.72	0.25	(17)	5.37	North Carolina	2.78	0.08	(15)	7.67
D.C.	3.93	0.37	(01)	52.52 ^c	North Dakota	2.47	0.26	(32)	-4.05
Florida	2.77	0.05	(16)	7.54	Ohio	2.64	0.07	(24)	2.26
Georgia	2.82	0.09	(13)	9.37	Oklahoma	2.60	0.12	(25)	0.86
Hawaii	3.05	0.21	(07)	18.20 ^c	Oregon	2.57	0.12	(26)	-0.53
Idaho	2.64	0.20	(23)	2.42	Pennsylvania	2.34	0.06	(37)	-9.24
Illinois	2.51	0.06	(30)	-2.81	Rhode Island	2.14	0.19	(45)	-16.93 ^c
Indiana	2.27	0.08	(40)	-12.05	South Carolina	3.10	0.12	(04)	20.36 ^c
Iowa	2.52	0.12	(29)	-2.44	South Dakota	2.18	0.22	(42)	-15.63 ^c
Kansas	2.21	0.12	(41)	-14.13	Tennessee	3.08	0.10	(05)	19.24 ^c
Kentucky	2.86	0.12	(11)	11.03	Texas	2.65	0.05	(22)	2.88
Louisiana	3.37	0.12	(03)	30.58 ^c	Utah	1.69	0.14	(51)	-34.46 ^c
Maine	2.71	0.19	(18)	4.96	Vermont	2.40	0.27	(34)	-7.07
Maryland	2.67	0.10	(20)	3.69	Virginia	2.34	0.08	(36)	-9.12
Massachusetts	2.55	0.09	(27)	-1.27	Washington	3.00	0.10	(09)	16.22 ^c
Michigan	2.54	0.07	(28)	-1.53	West Virginia	2.69	0.16	(19)	4.48
Minnesota	2.27	0.09	(39)	-11.99	Wisconsin	2.50	0.09	(31)	-3.00
Mississippi	3.48	0.15	(02)	35.04 ^c	Wyoming	2.16	0.29	(44)	-16.07 ^c
Missouri	2.42	0.09	(33)	-6.33					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 20.16
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	3.94	0.02		
High Five States				
D.C.	6.92	0.77	(01)	75.41 ^c
Mississippi	5.85	0.30	(02)	48.33 ^c
Louisiana	5.56	0.24	(03)	40.88 ^c
Arkansas	5.08	0.28	(04)	28.81 ^c
South Carolina	5.00	0.23	(05)	26.89 ^c
Low Five States				
Colorado	2.96	0.19	(47)	-24.90 ^c
New Mexico	2.95	0.26	(48)	-25.27 ^c
Alaska	2.58	0.49	(49)	-34.46 ^c
Wyoming	2.57	0.46	(50)	-34.93 ^c
Utah	2.37	0.24	(51)	-39.81 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.75	0.22	(07)	20.37 ^c	Montana	3.11	0.36	(43)	-21.05 ^c
Alaska	2.58	0.49	(49)	-34.46 ^c	Nebraska	3.00	0.27	(46)	-23.99 ^c
Arizona	3.30	0.16	(40)	-16.29 ^c	Nevada	4.05	0.29	(21)	2.61
Arkansas	5.08	0.28	(04)	28.81 ^c	New Hampshire	4.18	0.38	(17)	5.91
California	3.96	0.08	(22)	0.49	New Jersey	3.59	0.14	(35)	-8.96
Colorado	2.96	0.19	(47)	-24.90 ^c	New Mexico	2.95	0.26	(48)	-25.27 ^c
Connecticut	3.26	0.20	(41)	-17.34 ^c	New York	3.52	0.09	(36)	-10.72
Delaware	4.42	0.48	(10)	12.07	North Carolina	4.12	0.15	(18)	4.58
D.C.	6.92	0.77	(01)	75.41 ^c	North Dakota	3.03	0.43	(44)	-23.28 ^c
Florida	4.24	0.09	(14)	7.41	Ohio	4.08	0.12	(20)	3.54
Georgia	4.33	0.17	(12)	9.70	Oklahoma	3.92	0.22	(24)	-0.63
Hawaii	4.48	0.37	(09)	13.73	Oregon	3.62	0.21	(34)	-8.08
Idaho	3.83	0.36	(27)	-2.99	Pennsylvania	3.68	0.11	(30)	-6.59
Illinois	3.80	0.12	(28)	-3.52	Rhode Island	3.02	0.35	(45)	-23.43 ^c
Indiana	3.52	0.16	(37)	-10.72	South Carolina	5.00	0.23	(05)	26.89 ^c
Iowa	3.67	0.22	(31)	-6.86	South Dakota	3.17	0.40	(42)	-19.49 ^c
Kansas	3.34	0.23	(39)	-15.37 ^c	Tennessee	4.86	0.19	(06)	23.27 ^c
Kentucky	4.53	0.22	(08)	15.01 ^c	Texas	4.23	0.10	(15)	7.18
Louisiana	5.56	0.24	(03)	40.88 ^c	Utah	2.37	0.24	(51)	-39.81 ^c
Maine	4.09	0.35	(19)	3.71	Vermont	3.86	0.51	(26)	-2.05
Maryland	4.24	0.19	(13)	7.53	Virginia	3.70	0.16	(29)	-6.19
Massachusetts	3.94	0.16	(23)	-0.01	Washington	4.35	0.18	(11)	10.26
Michigan	3.86	0.13	(25)	-1.99	West Virginia	4.20	0.30	(16)	6.60
Minnesota	3.35	0.17	(38)	-15.06 ^c	Wisconsin	3.65	0.17	(32)	-7.36
Mississippi	5.85	0.30	(02)	48.33 ^c	Wyoming	2.57	0.46	(50)	-34.93 ^c
Missouri	3.64	0.16	(33)	-7.78					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.



Table 20.17
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.46	0.01		
High Five States				
Alaska	2.96	0.56	(01)	102.89 ^c
North Dakota	2.05	0.33	(02)	40.76 ^c
New Hampshire	1.91	0.23	(03)	31.14 ^c
Washington	1.86	0.11	(04)	27.92 ^c
D.C.	1.76	0.33	(05)	20.80 ^c
Low Five States				
Arizona	1.13	0.08	(47)	-22.45 ^c
Vermont	1.11	0.24	(48)	-23.56 ^c
New Mexico	1.08	0.14	(49)	-25.64 ^c
Utah	1.08	0.15	(50)	-25.65 ^c
Connecticut	1.05	0.10	(51)	-27.87 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.64	0.11	(11)	12.67	Montana	1.23	0.21	(45)	-15.35 ^c
Alaska	2.96	0.56	(01)	102.89 ^c	Nebraska	1.23	0.16	(46)	-15.54 ^c
Arizona	1.13	0.08	(47)	-22.45 ^c	Nevada	1.71	0.17	(07)	17.04 ^c
Arkansas	1.41	0.13	(32)	-3.32	New Hampshire	1.91	0.23	(03)	31.14 ^c
California	1.60	0.04	(15)	9.63	New Jersey	1.41	0.07	(31)	-2.95
Colorado	1.55	0.12	(17)	6.59	New Mexico	1.08	0.14	(49)	-25.64 ^c
Connecticut	1.05	0.10	(51)	-27.87 ^c	New York	1.31	0.05	(39)	-9.99
Delaware	1.33	0.23	(38)	-8.83	North Carolina	1.64	0.08	(10)	12.75
D.C.	1.76	0.33	(05)	20.80 ^c	North Dakota	2.05	0.33	(02)	40.76 ^c
Florida	1.51	0.05	(23)	3.87	Ohio	1.45	0.06	(28)	-0.30
Georgia	1.64	0.09	(12)	12.53	Oklahoma	1.53	0.12	(22)	4.85
Hawaii	1.75	0.21	(06)	20.15 ^c	Oregon	1.65	0.12	(09)	12.88
Idaho	1.59	0.21	(16)	9.19	Pennsylvania	1.26	0.06	(42)	-13.31
Illinois	1.44	0.06	(30)	-1.21	Rhode Island	1.44	0.21	(29)	-0.92
Indiana	1.25	0.08	(44)	-14.26	South Carolina	1.53	0.11	(21)	4.93
Iowa	1.55	0.13	(20)	6.18	South Dakota	1.26	0.22	(41)	-13.26
Kansas	1.31	0.13	(40)	-10.14	Tennessee	1.62	0.10	(14)	11.11
Kentucky	1.50	0.11	(25)	3.08	Texas	1.37	0.05	(36)	-5.99
Louisiana	1.62	0.12	(13)	11.36	Utah	1.08	0.15	(50)	-25.65 ^c
Maine	1.55	0.19	(18)	6.57	Vermont	1.11	0.24	(48)	-23.56 ^c
Maryland	1.41	0.10	(33)	-3.62	Virginia	1.26	0.08	(43)	-13.42
Massachusetts	1.48	0.09	(26)	1.37	Washington	1.86	0.11	(04)	27.92 ^c
Michigan	1.46	0.07	(27)	0.19	West Virginia	1.37	0.15	(37)	-6.11
Minnesota	1.39	0.10	(35)	-4.52	Wisconsin	1.55	0.10	(19)	6.39
Mississippi	1.51	0.13	(24)	3.85	Wyoming	1.70	0.35	(08)	16.64 ^c
Missouri	1.40	0.09	(34)	-4.27					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 21.13
Cancer of the Ovary (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	8.76	0.03		
High Five States				
Washington	10.15	0.25	(01)	15.95 ^c
Oregon	9.96	0.31	(02)	13.76
Wyoming	9.85	0.85	(03)	12.55
Montana	9.61	0.59	(04)	9.80
Alabama	9.45	0.26	(05)	7.93
Low Five States				
Texas	8.02	0.12	(47)	-8.46
D.C.	7.95	0.69	(48)	-9.21
Mississippi	7.91	0.31	(49)	-9.67
Hawaii	6.78	0.43	(50)	-22.51 ^c
Alaska	6.05	0.76	(51)	-30.93 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.45	0.26	(05)	7.93	Montana	9.61	0.59	(04)	9.80
Alaska	6.05	0.76	(51)	-30.93 ^c	Nebraska	8.75	0.42	(26)	-0.08
Arizona	8.50	0.23	(35)	-2.89	Nevada	8.68	0.39	(30)	-0.81
Arkansas	8.25	0.31	(41)	-5.82	New Hampshire	8.82	0.49	(21)	0.75
California	8.80	0.10	(23)	0.48	New Jersey	9.24	0.19	(13)	5.51
Colorado	9.21	0.29	(14)	5.13	New Mexico	8.16	0.40	(42)	-6.86
Connecticut	8.46	0.28	(37)	-3.35	New York	8.82	0.12	(22)	0.70
Delaware	8.44	0.59	(38)	-3.63	North Carolina	8.62	0.19	(32)	-1.59
D.C.	7.95	0.69	(48)	-9.21	North Dakota	8.13	0.65	(43)	-7.20
Florida	8.04	0.12	(46)	-8.23	Ohio	8.79	0.16	(24)	0.42
Georgia	8.71	0.20	(29)	-0.54	Oklahoma	8.57	0.29	(34)	-2.15
Hawaii	6.78	0.43	(50)	-22.51 ^c	Oregon	9.96	0.31	(02)	13.76
Idaho	8.47	0.49	(36)	-3.24	Pennsylvania	9.28	0.15	(10)	5.95
Illinois	8.83	0.16	(20)	0.81	Rhode Island	9.31	0.53	(09)	6.38
Indiana	9.33	0.23	(08)	6.51	South Carolina	8.11	0.26	(44)	-7.39
Iowa	9.14	0.31	(15)	4.42	South Dakota	9.24	0.63	(12)	5.55
Kansas	9.25	0.34	(11)	5.69	Tennessee	9.14	0.23	(16)	4.34
Kentucky	8.68	0.27	(31)	-0.88	Texas	8.02	0.12	(47)	-8.46
Louisiana	8.36	0.26	(40)	-4.53	Utah	8.72	0.42	(28)	-0.43
Maine	8.72	0.45	(27)	-0.37	Vermont	8.06	0.66	(45)	-7.90
Maryland	8.57	0.24	(33)	-2.13	Virginia	8.92	0.21	(19)	1.91
Massachusetts	9.13	0.22	(17)	4.29	Washington	10.15	0.25	(01)	15.95 ^c
Michigan	9.09	0.18	(18)	3.83	West Virginia	9.35	0.40	(06)	6.80
Minnesota	8.76	0.25	(25)	0.04	Wisconsin	9.33	0.24	(07)	6.55
Mississippi	7.91	0.31	(49)	-9.67	Wyoming	9.85	0.85	(03)	12.55
Missouri	8.42	0.22	(39)	-3.79					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 22.15
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	10.65	0.03		
High Five States				
Louisiana	12.19	0.24	(01)	14.44
D.C.	12.18	0.65	(02)	14.32
Connecticut	11.73	0.25	(03)	10.10
Mississippi	11.51	0.28	(04)	8.00
Massachusetts	11.46	0.18	(05)	7.60
Low Five States				
Nebraska	9.71	0.32	(47)	-8.87
Vermont	9.37	0.53	(48)	-12.04
West Virginia	9.29	0.29	(49)	-12.77
Utah	9.26	0.33	(50)	-13.14
Arizona	9.11	0.18	(51)	-14.51

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	10.54	0.21	(31)	-1.11	Montana	10.26	0.44	(40)	-3.75
Alaska	10.46	0.78	(32)	-1.87	Nebraska	9.71	0.32	(47)	-8.87
Arizona	9.11	0.18	(51)	-14.51	Nevada	10.77	0.32	(24)	1.09
Arkansas	10.69	0.27	(28)	0.37	New Hampshire	11.12	0.41	(11)	4.40
California	10.28	0.08	(39)	-3.55	New Jersey	11.15	0.16	(10)	4.64
Colorado	9.86	0.23	(46)	-7.48	New Mexico	10.19	0.33	(41)	-4.39
Connecticut	11.73	0.25	(03)	10.10	New York	10.93	0.10	(18)	2.54
Delaware	10.11	0.48	(42)	-5.08	North Carolina	10.94	0.16	(17)	2.68
D.C.	12.18	0.65	(02)	14.32	North Dakota	10.41	0.54	(34)	-2.31
Florida	9.91	0.09	(45)	-7.03	Ohio	10.79	0.13	(23)	1.30
Georgia	10.44	0.17	(33)	-2.06	Oklahoma	9.95	0.23	(44)	-6.57
Hawaii	10.62	0.39	(30)	-0.35	Oregon	11.00	0.24	(15)	3.27
Idaho	10.82	0.40	(21)	1.59	Pennsylvania	11.23	0.12	(09)	5.37
Illinois	11.32	0.13	(08)	6.29	Rhode Island	10.29	0.41	(38)	-3.45
Indiana	11.10	0.19	(13)	4.22	South Carolina	10.71	0.22	(26)	0.54
Iowa	10.38	0.25	(36)	-2.63	South Dakota	10.35	0.49	(37)	-2.87
Kansas	10.69	0.27	(29)	0.33	Tennessee	10.88	0.19	(20)	2.08
Kentucky	10.70	0.22	(27)	0.39	Texas	10.04	0.10	(43)	-5.80
Louisiana	12.19	0.24	(01)	14.44	Utah	9.26	0.33	(50)	-13.14
Maine	11.41	0.38	(07)	7.08	Vermont	9.37	0.53	(48)	-12.04
Maryland	11.45	0.21	(06)	7.46	Virginia	10.97	0.18	(16)	2.98
Massachusetts	11.46	0.18	(05)	7.60	Washington	10.74	0.19	(25)	0.83
Michigan	11.06	0.15	(14)	3.78	West Virginia	9.29	0.29	(49)	-12.77
Minnesota	10.39	0.20	(35)	-2.52	Wisconsin	11.11	0.20	(12)	4.28
Mississippi	11.51	0.28	(04)	8.00	Wyoming	10.89	0.66	(19)	2.23
Missouri	10.82	0.19	(22)	1.51					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 22.16
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	12.28	0.04		
High Five States				
D.C.	14.95	1.16	(01)	21.75 ^c
Connecticut	13.87	0.42	(02)	13.02 ^c
Louisiana	13.80	0.40	(03)	12.44 ^c
Massachusetts	13.50	0.31	(04)	10.01 ^c
Mississippi	13.24	0.48	(05)	7.87
Low Five States				
Colorado	11.06	0.37	(47)	-9.89
West Virginia	10.88	0.49	(48)	-11.37 ^c
Utah	10.84	0.53	(49)	-11.66 ^c
Vermont	10.69	0.86	(50)	-12.92 ^c
Arizona	10.59	0.28	(51)	-13.75 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	12.66	0.36	(17)	3.11	Montana	11.87	0.72	(32)	-3.33
Alaska	12.21	1.30	(28)	-0.55	Nebraska	11.92	0.55	(31)	-2.91
Arizona	10.59	0.28	(51)	-13.75 ^c	Nevada	12.28	0.51	(26)	0.04
Arkansas	12.64	0.44	(19)	2.99	New Hampshire	11.58	0.65	(41)	-5.69
California	11.54	0.13	(42)	-5.97	New Jersey	12.66	0.26	(16)	3.12
Colorado	11.06	0.37	(47)	-9.89	New Mexico	11.32	0.52	(44)	-7.80
Connecticut	13.87	0.42	(02)	13.02 ^c	New York	12.45	0.17	(22)	1.40
Delaware	11.52	0.78	(43)	-6.17	North Carolina	12.97	0.27	(09)	5.62
D.C.	14.95	1.16	(01)	21.75 ^c	North Dakota	11.68	0.85	(39)	-4.87
Florida	11.60	0.15	(40)	-5.53	Ohio	12.50	0.22	(20)	1.82
Georgia	12.28	0.30	(27)	0.02	Oklahoma	11.80	0.39	(33)	-3.91
Hawaii	11.77	0.61	(34)	-4.13	Oregon	12.36	0.39	(24)	0.67
Idaho	11.29	0.62	(45)	-8.03	Pennsylvania	13.17	0.21	(07)	7.24
Illinois	13.19	0.23	(06)	7.48	Rhode Island	11.77	0.69	(35)	-4.15
Indiana	13.06	0.32	(08)	6.36	South Carolina	12.33	0.38	(25)	0.48
Iowa	11.70	0.40	(38)	-4.73	South Dakota	11.27	0.77	(46)	-8.21
Kansas	12.49	0.45	(21)	1.70	Tennessee	12.65	0.32	(18)	3.08
Kentucky	12.41	0.38	(23)	1.12	Texas	11.70	0.17	(37)	-4.67
Louisiana	13.80	0.40	(03)	12.44 ^c	Utah	10.84	0.53	(49)	-11.66 ^c
Maine	12.94	0.63	(10)	5.37	Vermont	10.69	0.86	(50)	-12.92 ^c
Maryland	12.90	0.35	(11)	5.07	Virginia	12.71	0.30	(15)	3.51
Massachusetts	13.50	0.31	(04)	10.01 ^c	Washington	12.05	0.31	(30)	-1.86
Michigan	12.88	0.24	(12)	4.88	West Virginia	10.88	0.49	(48)	-11.37 ^c
Minnesota	11.72	0.33	(36)	-4.53	Wisconsin	12.73	0.32	(14)	3.66
Mississippi	13.24	0.48	(05)	7.87	Wyoming	12.11	1.05	(29)	-1.39
Missouri	12.73	0.31	(13)	3.66					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 22.17
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	9.33	0.03		
High Five States				
Louisiana	10.84	0.30	(01)	16.24 ^c
New Hampshire	10.79	0.54	(02)	15.65 ^c
Maryland	10.31	0.26	(03)	10.60
Idaho	10.26	0.53	(04)	10.02
Mississippi	10.18	0.35	(05)	9.21
Low Five States				
Vermont	8.23	0.66	(47)	-11.74
Nebraska	8.14	0.40	(48)	-12.73
West Virginia	8.01	0.36	(49)	-14.07
Utah	7.90	0.41	(50)	-15.31 ^c
Arizona	7.80	0.22	(51)	-16.35 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	8.93	0.25	(40)	-4.20	Montana	8.74	0.55	(43)	-6.25
Alaska	9.09	0.99	(37)	-2.54	Nebraska	8.14	0.40	(48)	-12.73
Arizona	7.80	0.22	(51)	-16.35 ^c	Nevada	9.38	0.41	(25)	0.63
Arkansas	9.13	0.33	(36)	-2.08	New Hampshire	10.79	0.54	(02)	15.65 ^c
California	9.22	0.10	(32)	-1.15	New Jersey	9.97	0.19	(09)	6.89
Colorado	8.81	0.29	(41)	-5.58	New Mexico	9.20	0.42	(34)	-1.31
Connecticut	10.09	0.30	(08)	8.16	New York	9.72	0.13	(14)	4.28
Delaware	8.80	0.59	(42)	-5.67	North Carolina	9.30	0.19	(28)	-0.26
D.C.	10.16	0.77	(06)	8.96	North Dakota	9.08	0.67	(38)	-2.67
Florida	8.46	0.12	(46)	-9.33	Ohio	9.40	0.16	(24)	0.84
Georgia	9.06	0.21	(39)	-2.85	Oklahoma	8.48	0.28	(45)	-9.09
Hawaii	9.63	0.50	(17)	3.26	Oregon	9.84	0.31	(13)	5.49
Idaho	10.26	0.53	(04)	10.02	Pennsylvania	9.69	0.15	(16)	3.88
Illinois	9.86	0.17	(12)	5.75	Rhode Island	9.33	0.52	(27)	0.04
Indiana	9.55	0.23	(21)	2.44	South Carolina	9.46	0.28	(23)	1.44
Iowa	9.28	0.31	(30)	-0.51	South Dakota	9.58	0.63	(19)	2.75
Kansas	9.17	0.33	(35)	-1.63	Tennessee	9.35	0.23	(26)	0.31
Kentucky	9.29	0.28	(29)	-0.37	Texas	8.67	0.13	(44)	-7.01
Louisiana	10.84	0.30	(01)	16.24 ^c	Utah	7.90	0.41	(50)	-15.31 ^c
Maine	10.10	0.48	(07)	8.26	Vermont	8.23	0.66	(47)	-11.74
Maryland	10.31	0.26	(03)	10.60	Virginia	9.58	0.22	(20)	2.72
Massachusetts	9.95	0.22	(10)	6.68	Washington	9.62	0.24	(18)	3.15
Michigan	9.53	0.18	(22)	2.22	West Virginia	8.01	0.36	(49)	-14.07
Minnesota	9.23	0.25	(31)	-1.00	Wisconsin	9.71	0.24	(15)	4.13
Mississippi	10.18	0.35	(05)	9.21	Wyoming	9.91	0.85	(11)	6.27
Missouri	9.22	0.23	(33)	-1.16					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 23.13
Cancer of the Prostate (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	25.64	0.07		
High Five States				
D.C.	43.31	2.08	(01)	68.91 ^c
Mississippi	34.20	0.83	(02)	33.35 ^c
Alabama	31.24	0.61	(03)	21.81 ^c
South Carolina	30.62	0.65	(04)	19.43 ^c
Louisiana	30.42	0.64	(05)	18.64 ^c
Low Five States				
West Virginia	23.88	0.77	(47)	-6.86
Kansas	23.52	0.64	(48)	-8.27
Arizona	22.10	0.42	(49)	-13.83 ^c
Florida	21.33	0.21	(50)	-16.84 ^c
Hawaii	17.42	0.76	(51)	-32.07 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	31.24	0.61	(03)	21.81 ^c	Montana	28.37	1.16	(10)	10.65 ^c
Alaska	24.23	2.09	(43)	-5.52	Nebraska	24.51	0.81	(39)	-4.41
Arizona	22.10	0.42	(49)	-13.83 ^c	Nevada	25.42	0.81	(36)	-0.87
Arkansas	27.49	0.69	(15)	7.22	New Hampshire	27.23	1.07	(17)	6.18
California	24.24	0.20	(42)	-5.47	New Jersey	24.47	0.38	(40)	-4.59
Colorado	25.54	0.61	(33)	-0.39	New Mexico	26.31	0.85	(26)	2.61
Connecticut	26.58	0.60	(25)	3.64	New York	24.60	0.26	(38)	-4.08
Delaware	26.80	1.28	(22)	4.50	North Carolina	28.91	0.45	(08)	12.72 ^c
D.C.	43.31	2.08	(01)	68.91 ^c	North Dakota	27.86	1.32	(13)	8.65
Florida	21.33	0.21	(50)	-16.84 ^c	Ohio	27.00	0.35	(18)	5.28
Georgia	29.75	0.51	(06)	16.00 ^c	Oklahoma	24.06	0.59	(45)	-6.18
Hawaii	17.42	0.76	(51)	-32.07 ^c	Oregon	26.91	0.59	(20)	4.95
Idaho	28.19	1.03	(12)	9.92	Pennsylvania	26.00	0.30	(31)	1.39
Illinois	26.95	0.34	(19)	5.10	Rhode Island	25.46	1.04	(35)	-0.71
Indiana	26.23	0.48	(27)	2.27	South Carolina	30.62	0.65	(04)	19.43 ^c
Iowa	26.67	0.62	(23)	3.99	South Dakota	27.42	1.21	(16)	6.91
Kansas	23.52	0.64	(48)	-8.27	Tennessee	28.35	0.52	(11)	10.54 ^c
Kentucky	26.64	0.60	(24)	3.88	Texas	24.27	0.27	(41)	-5.37
Louisiana	30.42	0.64	(05)	18.64 ^c	Utah	25.97	0.87	(32)	1.29
Maine	26.21	0.95	(28)	2.22	Vermont	26.20	1.45	(29)	2.19
Maryland	28.41	0.55	(09)	10.80 ^c	Virginia	28.94	0.49	(07)	12.84 ^c
Massachusetts	25.49	0.44	(34)	-0.61	Washington	26.13	0.48	(30)	1.88
Michigan	24.80	0.36	(37)	-3.29	West Virginia	23.88	0.77	(47)	-6.86
Minnesota	26.83	0.52	(21)	4.62	Wisconsin	27.76	0.49	(14)	8.24
Mississippi	34.20	0.83	(02)	33.35 ^c	Wyoming	24.09	1.58	(44)	-6.05
Missouri	24.03	0.46	(46)	-6.29					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 24.15
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006


Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.95	0.02		
High Five States				
Hawaii	7.17	0.32	(01)	81.42 ^c
D.C.	5.65	0.44	(02)	42.95 ^c
Louisiana	4.92	0.15	(03)	24.40 ^c
Alaska	4.89	0.52	(04)	23.70 ^c
Rhode Island	4.86	0.28	(05)	22.97 ^c
Low Five States				
Iowa	2.81	0.13	(47)	-29.02 ^c
Idaho	2.71	0.20	(48)	-31.48 ^c
Utah	2.69	0.17	(49)	-32.00 ^c
Nebraska	2.54	0.16	(50)	-35.88 ^c
Vermont	2.44	0.27	(51)	-38.37 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.89	0.13	(18)	-1.63	Montana	3.14	0.25	(41)	-20.65 ^c
Alaska	4.89	0.52	(04)	23.70 ^c	Nebraska	2.54	0.16	(50)	-35.88 ^c
Arizona	3.29	0.11	(36)	-16.84 ^c	Nevada	3.82	0.19	(21)	-3.29
Arkansas	3.70	0.16	(25)	-6.29	New Hampshire	2.99	0.21	(44)	-24.32 ^c
California	4.78	0.05	(09)	20.94 ^c	New Jersey	4.80	0.10	(07)	21.46 ^c
Colorado	3.26	0.13	(37)	-17.45 ^c	New Mexico	4.80	0.23	(08)	21.41 ^c
Connecticut	4.52	0.15	(10)	14.36	New York	4.85	0.07	(06)	22.68 ^c
Delaware	3.84	0.30	(19)	-2.97	North Carolina	3.64	0.09	(26)	-7.94
D.C.	5.65	0.44	(02)	42.95 ^c	North Dakota	4.01	0.34	(17)	1.54
Florida	3.56	0.06	(27)	-9.98	Ohio	3.50	0.08	(29)	-11.57
Georgia	3.83	0.10	(20)	-3.03	Oklahoma	3.23	0.13	(38)	-18.42 ^c
Hawaii	7.17	0.32	(01)	81.42 ^c	Oregon	3.14	0.13	(40)	-20.62 ^c
Idaho	2.71	0.20	(48)	-31.48 ^c	Pennsylvania	3.80	0.07	(22)	-3.95
Illinois	4.34	0.08	(13)	9.69	Rhode Island	4.86	0.28	(05)	22.97 ^c
Indiana	3.07	0.10	(43)	-22.41 ^c	South Carolina	4.39	0.14	(11)	10.94
Iowa	2.81	0.13	(47)	-29.02 ^c	South Dakota	3.32	0.28	(34)	-16.13 ^c
Kansas	2.90	0.14	(46)	-26.56 ^c	Tennessee	3.47	0.11	(31)	-12.17
Kentucky	3.43	0.13	(32)	-13.18	Texas	4.29	0.07	(14)	8.59
Louisiana	4.92	0.15	(03)	24.40 ^c	Utah	2.69	0.17	(49)	-32.00 ^c
Maine	3.33	0.21	(33)	-15.67 ^c	Vermont	2.44	0.27	(51)	-38.37 ^c
Maryland	4.02	0.12	(16)	1.65	Virginia	3.76	0.10	(23)	-4.83
Massachusetts	4.11	0.11	(15)	3.99	Washington	3.48	0.11	(30)	-12.01
Michigan	3.74	0.09	(24)	-5.51	West Virginia	3.51	0.18	(28)	-11.19
Minnesota	3.10	0.11	(42)	-21.55 ^c	Wisconsin	3.19	0.10	(39)	-19.30 ^c
Mississippi	4.38	0.18	(12)	10.89	Wyoming	2.91	0.34	(45)	-26.28 ^c
Missouri	3.30	0.10	(35)	-16.49 ^c					



Age-Adjusted Death Rates per 100,000 Quantile Interval



U.S. Rate: 3.95

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.



Table 24.16
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.48	0.03		
High Five States				
Hawaii	9.71	0.56	(01)	77.08 ^c
D.C.	8.08	0.85	(02)	47.32 ^c
Rhode Island	7.67	0.56	(03)	39.93 ^c
Louisiana	7.09	0.29	(04)	29.19 ^c
New Mexico	6.94	0.41	(05)	26.54 ^c
Low Five States				
Wyoming	3.95	0.60	(47)	-28.06 ^c
Idaho	3.85	0.36	(48)	-29.88 ^c
Nebraska	3.48	0.30	(49)	-36.47 ^c
Vermont	3.40	0.50	(50)	-37.94 ^c
Utah	3.28	0.29	(51)	-40.17 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.41	0.24	(17)	-1.27	Montana	4.37	0.44	(42)	-20.33 ^c
Alaska	6.37	0.89	(09)	16.08 ^c	Nebraska	3.48	0.30	(49)	-36.47 ^c
Arizona	4.41	0.18	(40)	-19.64 ^c	Nevada	4.96	0.33	(27)	-9.60
Arkansas	5.01	0.28	(25)	-8.68	New Hampshire	4.63	0.41	(35)	-15.63 ^c
California	6.43	0.10	(08)	17.32 ^c	New Jersey	6.75	0.19	(06)	23.00 ^c
Colorado	4.47	0.24	(38)	-18.53 ^c	New Mexico	6.94	0.41	(05)	26.54 ^c
Connecticut	6.33	0.28	(10)	15.49 ^c	New York	6.69	0.13	(07)	22.04 ^c
Delaware	5.31	0.54	(21)	-3.16	North Carolina	5.16	0.18	(24)	-6.00
D.C.	8.08	0.85	(02)	47.32 ^c	North Dakota	5.22	0.57	(23)	-4.85
Florida	4.88	0.10	(30)	-11.08	Ohio	4.94	0.14	(29)	-9.90
Georgia	5.31	0.19	(20)	-3.11	Oklahoma	4.45	0.24	(39)	-18.88 ^c
Hawaii	9.71	0.56	(01)	77.08 ^c	Oregon	4.04	0.22	(45)	-26.30 ^c
Idaho	3.85	0.36	(48)	-29.88 ^c	Pennsylvania	5.41	0.13	(18)	-1.40
Illinois	6.11	0.16	(13)	11.46	Rhode Island	7.67	0.56	(03)	39.93 ^c
Indiana	4.32	0.18	(44)	-21.17 ^c	South Carolina	6.23	0.27	(12)	13.65
Iowa	3.95	0.23	(46)	-27.99 ^c	South Dakota	4.63	0.50	(34)	-15.57 ^c
Kansas	4.35	0.27	(43)	-20.65 ^c	Tennessee	4.72	0.20	(31)	-13.94
Kentucky	4.96	0.24	(28)	-9.63	Texas	5.92	0.12	(15)	7.92
Louisiana	7.09	0.29	(04)	29.19 ^c	Utah	3.28	0.29	(51)	-40.17 ^c
Maine	4.61	0.38	(36)	-16.02 ^c	Vermont	3.40	0.50	(50)	-37.94 ^c
Maryland	5.33	0.23	(19)	-2.78	Virginia	5.52	0.20	(16)	0.69
Massachusetts	6.00	0.21	(14)	9.48	Washington	4.65	0.19	(33)	-15.17 ^c
Michigan	5.26	0.16	(22)	-4.04	West Virginia	5.00	0.34	(26)	-8.89
Minnesota	4.37	0.20	(41)	-20.29 ^c	Wisconsin	4.57	0.19	(37)	-16.59 ^c
Mississippi	6.30	0.33	(11)	14.86	Wyoming	3.95	0.60	(47)	-28.06 ^c
Missouri	4.66	0.19	(32)	-15.11 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.



Table 24.17
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.83	0.02		
High Five States				
Hawaii	5.19	0.37	(01)	83.53 ^c
D.C.	3.87	0.47	(02)	36.76 ^c
New York	3.59	0.08	(03)	26.84 ^c
California	3.56	0.06	(04)	26.00 ^c
Alaska	3.55	0.60	(05)	25.47 ^c
Low Five States				
Nebraska	1.85	0.19	(47)	-34.56 ^c
Kansas	1.77	0.15	(48)	-37.34 ^c
Vermont	1.74	0.31	(49)	-38.40 ^c
Idaho	1.73	0.22	(50)	-38.82 ^c
New Hampshire	1.69	0.21	(51)	-40.31 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.77	0.14	(20)	-1.99	Montana	2.18	0.28	(42)	-22.89 ^c
Alaska	3.55	0.60	(05)	25.47 ^c	Nebraska	1.85	0.19	(47)	-34.56 ^c
Arizona	2.41	0.12	(33)	-14.93	Nevada	2.83	0.23	(17)	0.03
Arkansas	2.73	0.18	(21)	-3.35	New Hampshire	1.69	0.21	(51)	-40.31 ^c
California	3.56	0.06	(04)	26.00 ^c	New Jersey	3.44	0.11	(06)	21.73 ^c
Colorado	2.38	0.15	(34)	-15.78 ^c	New Mexico	3.15	0.25	(09)	11.38
Connecticut	3.16	0.17	(08)	11.88	New York	3.59	0.08	(03)	26.84 ^c
Delaware	2.72	0.33	(22)	-3.69	North Carolina	2.54	0.10	(28)	-10.15
D.C.	3.87	0.47	(02)	36.76 ^c	North Dakota	3.00	0.39	(14)	6.24
Florida	2.51	0.06	(29)	-11.37	Ohio	2.48	0.08	(30)	-12.26
Georgia	2.81	0.11	(18)	-0.80	Oklahoma	2.26	0.15	(39)	-20.21 ^c
Hawaii	5.19	0.37	(01)	83.53 ^c	Oregon	2.44	0.15	(32)	-13.61
Idaho	1.73	0.22	(50)	-38.82 ^c	Pennsylvania	2.65	0.08	(23)	-6.46
Illinois	3.08	0.09	(11)	9.00	Rhode Island	3.02	0.29	(13)	6.96
Indiana	2.17	0.11	(43)	-23.23 ^c	South Carolina	2.96	0.15	(15)	4.64
Iowa	1.96	0.14	(46)	-30.85 ^c	South Dakota	2.30	0.32	(37)	-18.54 ^c
Kansas	1.77	0.15	(48)	-37.34 ^c	Tennessee	2.58	0.12	(25)	-8.63
Kentucky	2.32	0.14	(36)	-17.94 ^c	Texas	3.07	0.08	(12)	8.40
Louisiana	3.32	0.16	(07)	17.54 ^c	Utah	2.19	0.21	(40)	-22.40 ^c
Maine	2.38	0.23	(35)	-15.95 ^c	Vermont	1.74	0.31	(49)	-38.40 ^c
Maryland	3.09	0.14	(10)	9.19	Virginia	2.57	0.11	(27)	-9.02
Massachusetts	2.79	0.12	(19)	-1.19	Washington	2.59	0.12	(24)	-8.42
Michigan	2.58	0.09	(26)	-8.76	West Virginia	2.48	0.20	(31)	-12.34
Minnesota	2.19	0.12	(41)	-22.40 ^c	Wisconsin	2.17	0.11	(44)	-23.24 ^c
Mississippi	2.94	0.19	(16)	4.10	Wyoming	2.08	0.39	(45)	-26.40 ^c
Missouri	2.29	0.11	(38)	-18.93 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 25.13
Cancer of the Testis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.25	0.01		
High Five States				
North Dakota	0.42	0.16	(01)	65.48 ^c
Idaho	0.42	0.11	(02)	64.87 ^c
Oregon	0.36	0.06	(03)	40.97 ^c
Rhode Island	0.35	0.12	(04)	37.96 ^c
D.C.	0.35	0.16	(05)	37.49 ^c
Low Five States				
Wisconsin	0.16	0.03	(47)	-38.44 ^c
Delaware	0.16	0.09	(48)	-38.84 ^c
Montana	0.14	0.08	(49)	-46.22 ^c
South Dakota	0.10	0.07	(50)	-59.06 ^c
Alaska	0.00	.	(51)	-100.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.24	0.05	(27)	-4.08	Montana	0.14	0.08	(49)	-46.22 ^c
Alaska	0.00	.	(51)	-100.0 ^c	Nebraska	0.33	0.09	(07)	29.50 ^c
Arizona	0.28	0.04	(15)	11.16	Nevada	0.34	0.08	(06)	35.15 ^c
Arkansas	0.20	0.06	(41)	-20.70 ^c	New Hampshire	0.33	0.10	(08)	29.46 ^c
California	0.32	0.02	(10)	26.73 ^c	New Jersey	0.19	0.03	(42)	-24.03 ^c
Colorado	0.24	0.05	(30)	-5.32	New Mexico	0.23	0.07	(34)	-9.13
Connecticut	0.27	0.06	(19)	6.12	New York	0.25	0.02	(25)	-3.50
Delaware	0.16	0.09	(48)	-38.84 ^c	North Carolina	0.27	0.04	(20)	6.11
D.C.	0.35	0.16	(05)	37.49 ^c	North Dakota	0.42	0.16	(01)	65.48 ^c
Florida	0.24	0.02	(26)	-3.98	Ohio	0.27	0.03	(18)	6.31
Georgia	0.16	0.03	(46)	-35.48 ^c	Oklahoma	0.23	0.05	(33)	-8.74
Hawaii	0.25	0.09	(24)	-2.28	Oregon	0.36	0.06	(03)	40.97 ^c
Idaho	0.42	0.11	(02)	64.87 ^c	Pennsylvania	0.25	0.03	(23)	0.36
Illinois	0.23	0.03	(36)	-11.21	Rhode Island	0.35	0.12	(04)	37.96 ^c
Indiana	0.23	0.04	(35)	-9.42	South Carolina	0.17	0.04	(44)	-31.96 ^c
Iowa	0.29	0.06	(13)	12.69	South Dakota	0.10	0.07	(50)	-59.06 ^c
Kansas	0.27	0.06	(17)	7.51	Tennessee	0.26	0.04	(22)	1.86
Kentucky	0.16	0.04	(45)	-35.08 ^c	Texas	0.28	0.02	(16)	8.53
Louisiana	0.28	0.05	(14)	11.87	Utah	0.22	0.07	(37)	-12.16
Maine	0.24	0.09	(29)	-4.90	Vermont	0.24	0.12	(31)	-6.12
Maryland	0.29	0.05	(11)	14.45	Virginia	0.18	0.03	(43)	-30.52 ^c
Massachusetts	0.22	0.04	(38)	-12.79	Washington	0.22	0.04	(39)	-14.14
Michigan	0.24	0.03	(32)	-6.60	West Virginia	0.26	0.08	(21)	2.17
Minnesota	0.22	0.04	(40)	-14.71	Wisconsin	0.16	0.03	(47)	-38.44 ^c
Mississippi	0.33	0.07	(09)	28.25 ^c	Wyoming	0.24	0.14	(28)	-4.66
Missouri	0.29	0.05	(12)	12.92					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 26.15
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.47	0.01		
High Five States				
New Mexico	0.65	0.08	(01)	37.04 ^c
Idaho	0.58	0.09	(02)	21.73 ^c
Kansas	0.56	0.06	(03)	18.16 ^c
Kentucky	0.55	0.05	(04)	17.30 ^c
Arizona	0.55	0.04	(05)	17.18 ^c
Low Five States				
Alabama	0.37	0.04	(47)	-21.01 ^c
Delaware	0.36	0.09	(48)	-23.19 ^c
D.C.	0.32	0.11	(49)	-32.03 ^c
Alaska	0.25	0.13	(50)	-47.32 ^c
Vermont	0.22	0.08	(51)	-53.56 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.37	0.04	(47)	-21.01 ^c	Montana	0.44	0.09	(34)	-6.38
Alaska	0.25	0.13	(50)	-47.32 ^c	Nebraska	0.38	0.06	(45)	-19.45 ^c
Arizona	0.55	0.04	(05)	17.18 ^c	Nevada	0.39	0.06	(42)	-17.74 ^c
Arkansas	0.43	0.05	(36)	-9.42	New Hampshire	0.45	0.08	(30)	-4.98
California	0.52	0.02	(11)	9.47	New Jersey	0.52	0.03	(10)	10.11
Colorado	0.49	0.05	(18)	4.23	New Mexico	0.65	0.08	(01)	37.04 ^c
Connecticut	0.53	0.05	(07)	12.13	New York	0.49	0.02	(20)	3.45
Delaware	0.36	0.09	(48)	-23.19 ^c	North Carolina	0.40	0.03	(41)	-15.31 ^c
D.C.	0.32	0.11	(49)	-32.03 ^c	North Dakota	0.51	0.12	(14)	8.36
Florida	0.40	0.02	(38)	-14.87	Ohio	0.49	0.03	(19)	3.49
Georgia	0.38	0.03	(46)	-19.83 ^c	Oklahoma	0.44	0.05	(32)	-6.00
Hawaii	0.53	0.09	(08)	11.30	Oregon	0.41	0.05	(37)	-14.38
Idaho	0.58	0.09	(02)	21.73 ^c	Pennsylvania	0.46	0.02	(26)	-1.91
Illinois	0.51	0.03	(12)	8.65	Rhode Island	0.45	0.08	(29)	-4.82
Indiana	0.46	0.04	(27)	-3.10	South Carolina	0.39	0.04	(44)	-18.06 ^c
Iowa	0.51	0.05	(17)	7.53	South Dakota	0.54	0.11	(06)	13.66
Kansas	0.56	0.06	(03)	18.16 ^c	Tennessee	0.48	0.04	(22)	0.73
Kentucky	0.55	0.05	(04)	17.30 ^c	Texas	0.51	0.02	(15)	8.16
Louisiana	0.51	0.05	(13)	8.60	Utah	0.39	0.07	(43)	-17.97 ^c
Maine	0.44	0.07	(35)	-7.22	Vermont	0.22	0.08	(51)	-53.56 ^c
Maryland	0.53	0.04	(09)	11.00	Virginia	0.40	0.03	(40)	-15.17 ^c
Massachusetts	0.44	0.04	(33)	-6.15	Washington	0.45	0.04	(31)	-5.70
Michigan	0.51	0.03	(16)	8.07	West Virginia	0.40	0.06	(39)	-15.06 ^c
Minnesota	0.46	0.04	(28)	-3.83	Wisconsin	0.47	0.04	(23)	0.17
Mississippi	0.47	0.06	(25)	-1.68	Wyoming	0.49	0.14	(21)	2.93
Missouri	0.47	0.04	(24)	-0.86					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 26.16
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.46	0.01		
High Five States				
New Mexico	0.66	0.13	(01)	43.69 ^c
South Dakota	0.62	0.18	(02)	36.82 ^c
Kansas	0.58	0.10	(03)	27.30 ^c
Iowa	0.55	0.09	(04)	21.16 ^c
Maine	0.55	0.13	(05)	21.09 ^c
Low Five States				
Montana	0.32	0.12	(47)	-29.64 ^c
Nebraska	0.32	0.09	(48)	-29.80 ^c
Alaska	0.31	0.26	(49)	-31.92 ^c
Minnesota	0.30	0.05	(50)	-34.08 ^c
Vermont	0.29	0.14	(51)	-36.81 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.40	0.07	(35)	-11.34	Montana	0.32	0.12	(47)	-29.64 ^c
Alaska	0.31	0.26	(49)	-31.92 ^c	Nebraska	0.32	0.09	(48)	-29.80 ^c
Arizona	0.53	0.06	(09)	16.47 ^c	Nevada	0.38	0.09	(40)	-16.26 ^c
Arkansas	0.44	0.08	(27)	-4.43	New Hampshire	0.41	0.12	(34)	-10.16
California	0.48	0.03	(16)	6.32	New Jersey	0.47	0.05	(23)	2.43
Colorado	0.51	0.08	(14)	12.27	New Mexico	0.66	0.13	(01)	43.69 ^c
Connecticut	0.48	0.08	(17)	6.27	New York	0.47	0.03	(22)	2.87
Delaware	0.36	0.14	(44)	-21.48 ^c	North Carolina	0.47	0.05	(21)	3.24
D.C.	0.39	0.20	(39)	-14.60	North Dakota	0.42	0.16	(29)	-7.78
Florida	0.40	0.03	(38)	-13.03	Ohio	0.54	0.05	(07)	18.66 ^c
Georgia	0.37	0.05	(43)	-19.57 ^c	Oklahoma	0.47	0.08	(18)	3.69
Hawaii	0.49	0.13	(15)	7.04	Oregon	0.42	0.07	(33)	-8.44
Idaho	0.54	0.13	(08)	17.48 ^c	Pennsylvania	0.46	0.04	(24)	-0.09
Illinois	0.43	0.04	(28)	-5.59	Rhode Island	0.44	0.13	(26)	-2.98
Indiana	0.52	0.06	(10)	13.73	South Carolina	0.42	0.07	(31)	-8.26
Iowa	0.55	0.09	(04)	21.16 ^c	South Dakota	0.62	0.18	(02)	36.82 ^c
Kansas	0.58	0.10	(03)	27.30 ^c	Tennessee	0.42	0.06	(30)	-8.02
Kentucky	0.55	0.08	(06)	20.22 ^c	Texas	0.52	0.04	(11)	13.49
Louisiana	0.38	0.06	(42)	-17.06 ^c	Utah	0.47	0.11	(19)	3.67
Maine	0.55	0.13	(05)	21.09 ^c	Vermont	0.29	0.14	(51)	-36.81 ^c
Maryland	0.51	0.07	(12)	12.60	Virginia	0.40	0.05	(36)	-11.88
Massachusetts	0.40	0.05	(37)	-12.54	Washington	0.33	0.05	(46)	-28.10 ^c
Michigan	0.51	0.05	(13)	12.28	West Virginia	0.34	0.09	(45)	-24.90 ^c
Minnesota	0.30	0.05	(50)	-34.08 ^c	Wisconsin	0.38	0.05	(41)	-16.70 ^c
Mississippi	0.44	0.09	(25)	-2.94	Wyoming	0.42	0.19	(32)	-8.36
Missouri	0.47	0.06	(20)	3.35					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 26.17
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.48	0.01		
High Five States				
New Mexico	0.63	0.11	(01)	30.97 ^c
Louisiana	0.60	0.07	(02)	24.74 ^c
Idaho	0.59	0.13	(03)	23.96 ^c
Illinois	0.58	0.04	(04)	20.19 ^c
Hawaii	0.57	0.12	(05)	19.42 ^c
Low Five States				
Alabama	0.34	0.05	(47)	-29.77 ^c
Utah	0.30	0.08	(48)	-36.66 ^c
D.C.	0.27	0.12	(49)	-44.16 ^c
Alaska	0.24	0.17	(50)	-50.67 ^c
Vermont	0.17	0.10	(51)	-64.52 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.34	0.05	(47)	-29.77 ^c	Montana	0.52	0.13	(18)	7.92
Alaska	0.24	0.17	(50)	-50.67 ^c	Nebraska	0.40	0.09	(39)	-16.77 ^c
Arizona	0.57	0.06	(06)	19.23 ^c	Nevada	0.39	0.08	(41)	-18.06 ^c
Arkansas	0.41	0.07	(35)	-14.94	New Hampshire	0.48	0.11	(24)	-0.74
California	0.54	0.02	(13)	13.13	New Jersey	0.55	0.05	(08)	15.12 ^c
Colorado	0.47	0.07	(25)	-1.97	New Mexico	0.63	0.11	(01)	30.97 ^c
Connecticut	0.55	0.07	(10)	14.65	New York	0.50	0.03	(21)	3.97
Delaware	0.36	0.12	(44)	-25.09 ^c	North Carolina	0.34	0.04	(46)	-29.31 ^c
D.C.	0.27	0.12	(49)	-44.16 ^c	North Dakota	0.53	0.16	(16)	10.06
Florida	0.40	0.03	(38)	-15.93 ^c	Ohio	0.45	0.04	(31)	-6.94
Georgia	0.39	0.04	(42)	-19.30 ^c	Oklahoma	0.41	0.06	(34)	-13.60
Hawaii	0.57	0.12	(05)	19.42 ^c	Oregon	0.40	0.06	(37)	-15.92 ^c
Idaho	0.59	0.13	(03)	23.96 ^c	Pennsylvania	0.46	0.03	(28)	-3.89
Illinois	0.58	0.04	(04)	20.19 ^c	Rhode Island	0.42	0.10	(33)	-13.18
Indiana	0.41	0.05	(36)	-15.13 ^c	South Carolina	0.37	0.05	(43)	-23.69 ^c
Iowa	0.46	0.07	(29)	-3.94	South Dakota	0.46	0.14	(30)	-4.04
Kansas	0.55	0.08	(12)	14.02	Tennessee	0.50	0.05	(20)	5.01
Kentucky	0.55	0.07	(11)	14.21	Texas	0.51	0.03	(19)	5.46
Louisiana	0.60	0.07	(02)	24.74 ^c	Utah	0.30	0.08	(48)	-36.66 ^c
Maine	0.34	0.09	(45)	-28.83 ^c	Vermont	0.17	0.10	(51)	-64.52 ^c
Maryland	0.52	0.06	(17)	9.27	Virginia	0.39	0.04	(40)	-17.86 ^c
Massachusetts	0.47	0.05	(26)	-2.10	Washington	0.54	0.06	(14)	12.44
Michigan	0.50	0.04	(22)	3.20	West Virginia	0.44	0.08	(32)	-8.70
Minnesota	0.55	0.06	(09)	14.68	Wisconsin	0.53	0.06	(15)	11.15
Mississippi	0.48	0.08	(23)	0.22	Wyoming	0.56	0.20	(07)	16.33 ^c
Missouri	0.46	0.05	(27)	-3.83					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.



Table 27.15
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.34	0.02		
High Five States				
Delaware	5.91	0.37	(01)	36.10 ^c
Maine	5.64	0.27	(02)	29.78 ^c
Vermont	5.63	0.41	(03)	29.58 ^c
Nevada	5.38	0.23	(04)	23.84 ^c
Rhode Island	5.29	0.29	(05)	21.73 ^c
Low Five States				
Alabama	3.75	0.13	(47)	-13.58
Mississippi	3.67	0.16	(48)	-15.38 ^c
New Mexico	3.62	0.20	(49)	-16.65 ^c
Utah	3.43	0.20	(50)	-21.07 ^c
Hawaii	2.56	0.19	(51)	-40.98 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.75	0.13	(47)	-13.58	Montana	4.32	0.29	(26)	-0.61
Alaska	4.59	0.55	(17)	5.77	Nebraska	3.80	0.20	(45)	-12.47
Arizona	3.99	0.12	(40)	-8.22	Nevada	5.38	0.23	(04)	23.84 ^c
Arkansas	3.81	0.16	(44)	-12.29	New Hampshire	4.86	0.27	(12)	11.87
California	4.15	0.05	(31)	-4.32	New Jersey	4.87	0.10	(11)	12.18
Colorado	4.04	0.15	(38)	-6.88	New Mexico	3.62	0.20	(49)	-16.65 ^c
Connecticut	4.89	0.16	(10)	12.71	New York	4.33	0.07	(25)	-0.24
Delaware	5.91	0.37	(01)	36.10 ^c	North Carolina	4.09	0.10	(35)	-5.83
D.C.	5.10	0.42	(06)	17.43 ^c	North Dakota	4.05	0.32	(37)	-6.67
Florida	4.36	0.06	(24)	0.45	Ohio	5.02	0.09	(07)	15.64 ^c
Georgia	3.78	0.10	(46)	-12.96	Oklahoma	4.10	0.15	(34)	-5.70
Hawaii	2.56	0.19	(51)	-40.98 ^c	Oregon	4.97	0.16	(09)	14.46
Idaho	4.20	0.25	(28)	-3.22	Pennsylvania	4.71	0.08	(14)	8.41
Illinois	4.41	0.08	(22)	1.62	Rhode Island	5.29	0.29	(05)	21.73 ^c
Indiana	4.44	0.12	(21)	2.14	South Carolina	3.94	0.14	(42)	-9.34
Iowa	4.12	0.15	(33)	-5.09	South Dakota	4.37	0.31	(23)	0.53
Kansas	3.97	0.16	(41)	-8.60	Tennessee	4.14	0.12	(32)	-4.67
Kentucky	4.58	0.15	(18)	5.45	Texas	3.82	0.07	(43)	-12.03
Louisiana	4.08	0.14	(36)	-6.11	Utah	3.43	0.20	(50)	-21.07 ^c
Maine	5.64	0.27	(02)	29.78 ^c	Vermont	5.63	0.41	(03)	29.58 ^c
Maryland	4.58	0.13	(19)	5.37	Virginia	4.23	0.11	(27)	-2.54
Massachusetts	5.01	0.12	(08)	15.33 ^c	Washington	4.53	0.12	(20)	4.35
Michigan	4.61	0.10	(16)	6.23	West Virginia	4.66	0.21	(15)	7.21
Minnesota	4.19	0.13	(29)	-3.46	Wisconsin	4.82	0.13	(13)	11.08
Mississippi	3.67	0.16	(48)	-15.38 ^c	Wyoming	4.03	0.40	(39)	-7.20
Missouri	4.16	0.12	(30)	-4.16					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 27.16
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.53	0.04		
High Five States				
Vermont	10.31	0.89	(01)	36.93 ^c
Delaware	10.10	0.76	(02)	34.09 ^c
Rhode Island	9.78	0.64	(03)	29.84 ^c
Maine	9.46	0.56	(04)	25.68 ^c
Ohio	9.08	0.20	(05)	20.55 ^c
Low Five States				
Texas	6.58	0.14	(47)	-12.60
Utah	6.42	0.42	(48)	-14.67
Arkansas	6.37	0.32	(49)	-15.40 ^c
New Mexico	5.65	0.38	(50)	-24.98 ^c
Hawaii	4.10	0.37	(51)	-45.57 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.87	0.28	(41)	-8.78	Montana	7.44	0.58	(28)	-1.22
Alaska	8.60	1.17	(10)	14.27	Nebraska	6.69	0.42	(44)	-11.09
Arizona	6.67	0.23	(45)	-11.43	Nevada	8.98	0.46	(06)	19.29 ^c
Arkansas	6.37	0.32	(49)	-15.40 ^c	New Hampshire	8.86	0.59	(08)	17.67 ^c
California	7.14	0.11	(35)	-5.22	New Jersey	8.36	0.22	(12)	11.06
Colorado	7.13	0.32	(36)	-5.34	New Mexico	5.65	0.38	(50)	-24.98 ^c
Connecticut	8.76	0.34	(09)	16.37 ^c	New York	7.52	0.14	(24)	-0.15
Delaware	10.10	0.76	(02)	34.09 ^c	North Carolina	7.02	0.21	(37)	-6.77
D.C.	7.74	0.86	(19)	2.79	North Dakota	7.49	0.68	(26)	-0.58
Florida	7.28	0.12	(31)	-3.29	Ohio	9.08	0.20	(05)	20.55 ^c
Georgia	6.61	0.23	(46)	-12.15	Oklahoma	6.92	0.31	(40)	-8.15
Hawaii	4.10	0.37	(51)	-45.57 ^c	Oregon	8.40	0.33	(11)	11.60
Idaho	7.38	0.52	(29)	-1.92	Pennsylvania	8.30	0.17	(14)	10.29
Illinois	7.66	0.18	(21)	1.72	Rhode Island	9.78	0.64	(03)	29.84 ^c
Indiana	7.55	0.25	(23)	0.33	South Carolina	6.92	0.30	(38)	-8.03
Iowa	7.28	0.32	(32)	-3.35	South Dakota	8.29	0.66	(15)	10.07
Kansas	7.22	0.35	(33)	-4.06	Tennessee	7.22	0.26	(34)	-4.16
Kentucky	7.97	0.32	(17)	5.84	Texas	6.58	0.14	(47)	-12.60
Louisiana	6.92	0.30	(39)	-8.07	Utah	6.42	0.42	(48)	-14.67
Maine	9.46	0.56	(04)	25.68 ^c	Vermont	10.31	0.89	(01)	36.93 ^c
Maryland	7.81	0.28	(18)	3.69	Virginia	7.48	0.24	(27)	-0.61
Massachusetts	8.86	0.26	(07)	17.69 ^c	Washington	7.73	0.25	(20)	2.60
Michigan	8.22	0.20	(16)	9.16	West Virginia	7.58	0.42	(22)	0.61
Minnesota	7.33	0.27	(30)	-2.68	Wisconsin	8.34	0.27	(13)	10.76
Mississippi	6.81	0.36	(42)	-9.51	Wyoming	6.79	0.82	(43)	-9.87
Missouri	7.51	0.25	(25)	-0.23					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table 27.17
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2002-2006

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.24	0.02		
High Five States				
D.C.	3.39	0.43	(01)	51.09 ^c
Maine	2.98	0.26	(02)	32.95 ^c
Delaware	2.87	0.33	(03)	27.82 ^c
West Virginia	2.69	0.20	(04)	19.98 ^c
New Jersey	2.67	0.10	(05)	19.02 ^c
Low Five States				
South Dakota	1.71	0.26	(47)	-23.85 ^c
North Dakota	1.62	0.27	(48)	-27.70 ^c
Alaska	1.44	0.40	(49)	-35.73 ^c
Hawaii	1.41	0.19	(50)	-37.33 ^c
Utah	1.21	0.16	(51)	-46.08 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.85	0.11	(44)	-17.44 ^c	Montana	2.08	0.26	(32)	-7.36
Alaska	1.44	0.40	(49)	-35.73 ^c	Nebraska	1.94	0.18	(42)	-13.74
Arizona	1.96	0.11	(41)	-12.64	Nevada	2.53	0.21	(11)	12.62
Arkansas	2.07	0.15	(35)	-7.70	New Hampshire	2.20	0.24	(24)	-1.83
California	2.11	0.05	(31)	-6.16	New Jersey	2.67	0.10	(05)	19.02 ^c
Colorado	2.00	0.14	(39)	-10.67	New Mexico	2.07	0.20	(34)	-7.54
Connecticut	2.48	0.15	(13)	10.37	New York	2.35	0.06	(18)	4.91
Delaware	2.87	0.33	(03)	27.82 ^c	North Carolina	2.27	0.09	(22)	1.13
D.C.	3.39	0.43	(01)	51.09 ^c	North Dakota	1.62	0.27	(48)	-27.70 ^c
Florida	2.17	0.06	(29)	-3.47	Ohio	2.49	0.08	(12)	10.88
Georgia	2.06	0.10	(36)	-8.26	Oklahoma	2.18	0.14	(27)	-2.94
Hawaii	1.41	0.19	(50)	-37.33 ^c	Oregon	2.55	0.15	(09)	13.50
Idaho	1.79	0.22	(45)	-20.20 ^c	Pennsylvania	2.44	0.07	(14)	8.98
Illinois	2.40	0.08	(17)	6.96	Rhode Island	2.59	0.26	(07)	15.52 ^c
Indiana	2.44	0.11	(15)	8.72	South Carolina	2.01	0.13	(37)	-10.33
Iowa	2.18	0.14	(26)	-2.85	South Dakota	1.71	0.26	(47)	-23.85 ^c
Kansas	1.89	0.15	(43)	-15.65 ^c	Tennessee	2.17	0.11	(28)	-3.14
Kentucky	2.42	0.14	(16)	7.74	Texas	2.01	0.06	(38)	-10.61
Louisiana	2.28	0.13	(21)	1.79	Utah	1.21	0.16	(51)	-46.08 ^c
Maine	2.98	0.26	(02)	32.95 ^c	Vermont	2.32	0.34	(19)	3.42
Maryland	2.55	0.13	(08)	13.61	Virginia	2.19	0.10	(25)	-2.22
Massachusetts	2.64	0.11	(06)	17.47 ^c	Washington	2.30	0.12	(20)	2.31
Michigan	2.25	0.09	(23)	0.45	West Virginia	2.69	0.20	(04)	19.98 ^c
Minnesota	2.16	0.12	(30)	-3.78	Wisconsin	2.53	0.12	(10)	12.79
Mississippi	1.78	0.14	(46)	-20.46 ^c	Wyoming	2.08	0.39	(33)	-7.45
Missouri	1.99	0.10	(40)	-11.12					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.