


Table 2.18
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death^a Rates by State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	168.51	0.10		
High Five States				
Kentucky	202.63	0.92	01(01,02)	20.25 ^c
Mississippi	198.61	1.13	02(01,03)	17.86 ^c
West Virginia	196.13	1.30	03(02,05)	16.39 ^c
Louisiana	193.45	0.91	04(03,06)	14.80 ^c
Arkansas	192.16	1.07	05(03,07)	14.04 ^c
Low Five States				
Arizona	149.36	0.65	47(46,48)	-11.37 ^c
New Mexico	147.97	1.16	48(47,49)	-12.19 ^c
Colorado	145.38	0.79	49(48,49)	-13.73 ^c
Hawaii	137.76	1.32	50(50,50)	-18.25 ^c
Utah	128.08	1.10	51(51,51)	-23.99 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	188.11	0.84	08(06,09)	11.63 ^c	Montana	159.04	1.64	41(35,45)	-5.62 ^c
Alaska	176.26	2.81	18(10,29)	4.60	Nebraska	164.13	1.27	34(30,39)	-2.60
Arizona	149.36	0.65	47(46,48)	-11.37 ^c	Nevada	169.37	1.14	27(21,33)	0.51
Arkansas	192.16	1.07	05(03,07)	14.04 ^c	New Hampshire	168.88	1.51	29(20,34)	0.22
California	152.56	0.29	46(44,46)	-9.47 ^c	New Jersey	163.75	0.58	35(33,37)	-2.82
Colorado	145.38	0.79	49(48,49)	-13.73 ^c	New Mexico	147.97	1.16	48(47,49)	-12.19 ^c
Connecticut	156.53	0.87	44(41,45)	-7.11 ^c	New York	160.06	0.38	40(37,42)	-5.02 ^c
Delaware	176.85	1.84	16(12,22)	4.95	North Carolina	173.44	0.58	20(18,22)	2.92
D.C.	182.30	2.52	11(07,19)	8.19 ^c	North Dakota	154.67	1.98	45(39,47)	-8.21 ^c
Florida	160.32	0.36	39(37,42)	-4.86	Ohio	184.06	0.53	10(09,12)	9.23 ^c
Georgia	171.62	0.63	21(20,26)	1.84	Oklahoma	189.09	0.96	07(05,09)	12.21 ^c
Hawaii	137.76	1.32	50(50,50)	-18.25 ^c	Oregon	169.69	0.88	25(21,32)	0.70
Idaho	156.71	1.40	43(38,45)	-7.00 ^c	Pennsylvania	176.84	0.47	17(14,19)	4.94
Illinois	176.19	0.51	19(15,20)	4.56	Rhode Island	171.30	1.66	22(18,32)	1.66
Indiana	185.24	0.73	09(08,11)	9.93 ^c	South Carolina	180.04	0.84	14(11,16)	6.84 ^c
Iowa	169.70	0.97	24(21,32)	0.71	South Dakota	162.53	1.85	36(31,43)	-3.55
Kansas	168.43	1.04	30(22,34)	-0.05	Tennessee	190.80	0.74	06(04,08)	13.23 ^c
Kentucky	202.63	0.92	01(01,02)	20.25 ^c	Texas	161.79	0.38	37(35,39)	-3.99
Louisiana	193.45	0.91	04(03,06)	14.80 ^c	Utah	128.08	1.10	51(51,51)	-23.99 ^c
Maine	181.37	1.46	13(09,16)	7.63 ^c	Vermont	171.19	2.15	23(16,34)	1.59
Maryland	168.23	0.75	31(23,33)	-0.16	Virginia	169.09	0.64	28(22,32)	0.34
Massachusetts	166.84	0.67	32(28,34)	-0.99	Washington	165.67	0.69	33(30,35)	-1.69
Michigan	177.09	0.56	15(14,19)	5.09 ^c	West Virginia	196.13	1.30	03(02,05)	16.39 ^c
Minnesota	161.00	0.75	38(35,42)	-4.46	Wisconsin	169.65	0.73	26(21,31)	0.68
Mississippi	198.61	1.13	02(01,03)	17.86 ^c	Wyoming	158.09	2.33	42(34,46)	-6.18 ^c
Missouri	182.09	0.73	12(10,14)	8.06 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 2.19
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death^a Rates by State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	203.98	0.17		
High Five States				
Mississippi	256.97	2.01	01(01,02)	25.98 ^c
Kentucky	250.53	1.59	02(01,02)	22.82 ^c
Arkansas	242.54	1.84	03(03,07)	18.91 ^c
Alabama	242.28	1.49	04(03,07)	18.78 ^c
Louisiana	241.91	1.58	05(03,07)	18.59 ^c
Low Five States				
New Mexico	178.19	1.92	47(44,48)	-12.64 ^c
Arizona	177.02	1.05	48(46,48)	-13.21 ^c
Colorado	169.98	1.31	49(49,50)	-16.67 ^c
Hawaii	168.80	2.18	50(49,50)	-17.25 ^c
Utah	152.11	1.80	51(51,51)	-25.43 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	242.28	1.49	04(03,07)	18.78 ^c	Montana	184.67	2.62	44(39,48)	-9.47 ^c
Alaska	206.54	4.65	23(13,38)	1.26	Nebraska	198.52	2.12	32(25,39)	-2.68
Arizona	177.02	1.05	48(46,48)	-13.21 ^c	Nevada	197.13	1.85	34(27,40)	-3.36
Arkansas	242.54	1.84	03(03,07)	18.91 ^c	New Hampshire	203.41	2.52	26(21,35)	-0.28
California	180.35	0.48	46(44,47)	-11.58 ^c	New Jersey	194.74	0.98	36(33,40)	-4.53
Colorado	169.98	1.31	49(49,50)	-16.67 ^c	New Mexico	178.19	1.92	47(44,48)	-12.64 ^c
Connecticut	186.64	1.46	43(40,45)	-8.50 ^c	New York	190.68	0.65	41(38,43)	-6.52 ^c
Delaware	212.41	3.09	19(12,27)	4.14	North Carolina	218.21	1.02	14(12,17)	6.98 ^c
D.C.	216.33	4.34	15(09,27)	6.06 ^c	North Dakota	191.77	3.31	40(29,45)	-5.99 ^c
Florida	193.13	0.58	39(36,41)	-5.32 ^c	Ohio	223.47	0.89	11(10,13)	9.55 ^c
Georgia	215.66	1.12	16(13,20)	5.73 ^c	Oklahoma	230.50	1.62	08(08,10)	13.00 ^c
Hawaii	168.80	2.18	50(49,50)	-17.25 ^c	Oregon	200.33	1.44	31(25,35)	-1.79
Idaho	186.82	2.29	42(38,46)	-8.41 ^c	Pennsylvania	214.05	0.80	17(15,21)	4.94
Illinois	211.89	0.87	21(16,22)	3.88	Rhode Island	212.75	2.86	18(12,26)	4.30
Indiana	228.04	1.25	09(08,11)	11.80 ^c	South Carolina	227.95	1.47	10(08,11)	11.75 ^c
Iowa	206.62	1.61	22(20,29)	1.30	South Dakota	193.49	3.03	37(27,44)	-5.14 ^c
Kansas	202.12	1.73	29(22,34)	-0.91	Tennessee	241.18	1.30	06(03,07)	18.24 ^c
Kentucky	250.53	1.59	02(01,02)	22.82 ^c	Texas	198.42	0.65	33(29,35)	-2.73
Louisiana	241.91	1.58	05(03,07)	18.59 ^c	Utah	152.11	1.80	51(51,51)	-25.43 ^c
Maine	220.57	2.44	12(10,18)	8.13 ^c	Vermont	202.89	3.56	27(19,39)	-0.53
Maryland	202.41	1.28	28(23,32)	-0.77	Virginia	206.21	1.10	24(21,28)	1.09
Massachusetts	200.35	1.13	30(26,34)	-1.78	Washington	196.28	1.14	35(31,39)	-3.77
Michigan	212.11	0.94	20(16,22)	3.99	West Virginia	239.84	2.18	07(03,07)	17.58 ^c
Minnesota	193.35	1.25	38(34,41)	-5.21 ^c	Wisconsin	205.22	1.21	25(21,29)	0.61
Mississippi	256.97	2.01	01(01,02)	25.98 ^c	Wyoming	184.22	3.77	45(36,48)	-9.69 ^c
Missouri	219.59	1.23	13(11,16)	7.65 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 2.20
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death^a Rates by State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	143.35	0.12		
High Five States				
Kentucky	168.96	1.13	01(01,02)	17.86 ^c
West Virginia	164.35	1.61	02(01,04)	14.65 ^c
D.C.	161.10	3.12	03(01,14)	12.38 ^c
Oklahoma	158.75	1.19	04(03,11)	10.74 ^c
Louisiana	158.74	1.10	05(02,10)	10.73 ^c
Low Five States				
North Dakota	127.79	2.46	47(42,49)	-10.86 ^c
Arizona	127.68	0.82	48(45,49)	-10.93 ^c
New Mexico	125.04	1.44	49(46,49)	-12.77 ^c
Hawaii	114.21	1.64	50(50,51)	-20.33 ^c
Utah	109.81	1.38	51(50,51)	-23.40 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	150.20	1.00	18(12,21)	4.78	Montana	139.08	2.10	38(24,44)	-2.98
Alaska	152.04	3.53	15(03,35)	6.06 ^c	Nebraska	139.55	1.59	37(26,43)	-2.66
Arizona	127.68	0.82	48(45,49)	-10.93 ^c	Nevada	146.97	1.45	21(16,33)	2.52
Arkansas	155.20	1.31	11(04,17)	8.26 ^c	New Hampshire	144.21	1.88	27(18,39)	0.60
California	132.86	0.36	44(43,46)	-7.32 ^c	New Jersey	143.73	0.72	30(22,35)	0.26
Colorado	127.94	0.99	46(45,49)	-10.75 ^c	New Mexico	125.04	1.44	49(46,49)	-12.77 ^c
Connecticut	136.40	1.08	41(36,44)	-4.85	New York	139.87	0.48	35(34,40)	-2.43
Delaware	151.53	2.29	16(05,25)	5.70 ^c	North Carolina	142.90	0.70	33(24,36)	-0.32
D.C.	161.10	3.12	03(01,14)	12.38 ^c	North Dakota	127.79	2.46	47(42,49)	-10.86 ^c
Florida	134.86	0.45	43(40,44)	-5.93 ^c	Ohio	156.76	0.65	07(04,12)	9.35 ^c
Georgia	142.02	0.75	34(26,38)	-0.93	Oklahoma	158.75	1.19	04(03,11)	10.74 ^c
Hawaii	114.21	1.64	50(50,51)	-20.33 ^c	Oregon	147.37	1.12	20(17,28)	2.80
Idaho	132.77	1.76	45(39,47)	-7.38 ^c	Pennsylvania	151.50	0.59	17(12,19)	5.68 ^c
Illinois	152.28	0.64	14(11,18)	6.23 ^c	Rhode Island	144.22	2.04	26(17,39)	0.60
Indiana	155.32	0.90	10(05,14)	8.35 ^c	South Carolina	146.47	1.01	22(18,30)	2.17
Iowa	143.89	1.21	29(20,36)	0.38	South Dakota	139.87	2.34	36(21,44)	-2.43
Kansas	143.97	1.30	28(19,37)	0.43	Tennessee	155.35	0.89	09(05,14)	8.37 ^c
Kentucky	168.96	1.13	01(01,02)	17.86 ^c	Texas	135.13	0.47	42(39,44)	-5.74 ^c
Louisiana	158.74	1.10	05(02,10)	10.73 ^c	Utah	109.81	1.38	51(50,51)	-23.40 ^c
Maine	153.02	1.81	12(05,20)	6.74 ^c	Vermont	147.83	2.69	19(09,37)	3.12
Maryland	145.44	0.93	23(18,33)	1.46	Virginia	143.40	0.79	32(23,36)	0.03
Massachusetts	144.62	0.83	25(20,34)	0.89	Washington	143.44	0.87	31(22,36)	0.06
Michigan	152.57	0.70	13(10,18)	6.43 ^c	West Virginia	164.35	1.61	02(01,04)	14.65 ^c
Minnesota	138.29	0.94	40(35,41)	-3.53	Wisconsin	144.70	0.91	24(19,34)	0.94
Mississippi	158.16	1.34	06(03,12)	10.33 ^c	Wyoming	138.57	2.99	39(20,46)	-3.34
Missouri	155.41	0.91	08(05,14)	8.41 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 3.18
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.32	0.02		
High Five States				
South Dakota	5.59	0.35	01(01,21)	29.49 ^c
Vermont	5.38	0.39	02(01,34)	24.59 ^c
Kansas	5.13	0.18	03(01,20)	18.94 ^c
Iowa	5.13	0.17	04(01,20)	18.83 ^c
Oregon	5.11	0.16	05(01,19)	18.51 ^c
Low Five States				
New York	3.79	0.06	47(43,50)	-12.21
New Jersey	3.75	0.09	48(42,50)	-13.22
New Mexico	3.55	0.18	49(40,50)	-17.85 ^c
D.C.	3.28	0.34	50(26,51)	-24.05 ^c
Hawaii	2.59	0.18	51(50,51)	-40.09 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	5.01	0.14	08(01,20)	16.00 ^c	Montana	4.53	0.28	21(02,48)	4.92
Alaska	4.79	0.43	14(01,49)	10.97	Nebraska	5.08	0.23	06(01,27)	17.79 ^c
Arizona	4.22	0.11	36(19,46)	-2.28	Nevada	4.03	0.17	44(20,50)	-6.57
Arkansas	4.87	0.17	10(02,29)	12.86	New Hampshire	4.70	0.25	18(01,45)	8.93
California	4.29	0.05	30(23,40)	-0.62	New Jersey	3.75	0.09	48(42,50)	-13.22
Colorado	4.50	0.14	23(09,41)	4.37	New Mexico	3.55	0.18	49(40,50)	-17.85 ^c
Connecticut	4.20	0.15	37(18,47)	-2.71	New York	3.79	0.06	47(43,50)	-12.21
Delaware	4.18	0.29	38(06,50)	-3.11	North Carolina	4.39	0.09	27(16,40)	1.66
D.C.	3.28	0.34	50(26,51)	-24.05 ^c	North Dakota	4.28	0.34	31(02,50)	-0.77
Florida	4.11	0.06	40(31,46)	-4.82	Ohio	4.46	0.08	25(15,35)	3.26
Georgia	4.07	0.09	42(27,47)	-5.71	Oklahoma	4.53	0.15	20(08,40)	4.99
Hawaii	2.59	0.18	51(50,51)	-40.09 ^c	Oregon	5.11	0.16	05(01,19)	18.51 ^c
Idaho	4.76	0.24	16(01,43)	10.37	Pennsylvania	4.26	0.08	34(22,44)	-1.36
Illinois	4.13	0.08	39(27,46)	-4.25	Rhode Island	4.02	0.26	45(13,50)	-6.76
Indiana	4.51	0.11	22(11,37)	4.52	South Carolina	4.32	0.13	29(15,45)	0.05
Iowa	5.13	0.17	04(01,20)	18.83 ^c	South Dakota	5.59	0.35	01(01,21)	29.49 ^c
Kansas	5.13	0.18	03(01,20)	18.94 ^c	Tennessee	4.86	0.12	11(03,23)	12.53
Kentucky	4.71	0.14	17(04,32)	9.06	Texas	4.07	0.06	41(32,46)	-5.59
Louisiana	4.28	0.14	33(16,46)	-0.84	Utah	4.34	0.19	28(09,47)	0.54
Maine	4.87	0.25	09(01,38)	12.95	Vermont	5.38	0.39	02(01,34)	24.59 ^c
Maryland	4.06	0.12	43(25,48)	-5.91	Virginia	3.97	0.10	46(32,49)	-8.02
Massachusetts	4.28	0.11	32(18,45)	-0.78	Washington	5.05	0.12	07(01,18)	17.07 ^c
Michigan	4.69	0.09	19(07,27)	8.74	West Virginia	4.22	0.19	35(12,49)	-2.19
Minnesota	4.49	0.13	24(10,40)	4.06	Wisconsin	4.85	0.12	12(03,24)	12.46
Mississippi	4.78	0.18	15(02,34)	10.78	Wyoming	4.80	0.41	13(01,48)	11.14
Missouri	4.45	0.12	26(13,40)	3.01					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.



- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 3.19
Cancer of the Brain and Other Nervous System (Invasive)
Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	5.27	0.03		
High Five States				
South Dakota	6.72	0.55	01(01,36)	27.52 ^c
Idaho	6.70	0.42	02(01,23)	27.23 ^c
Vermont	6.66	0.63	03(01,43)	26.50 ^c
Maine	6.46	0.42	04(01,32)	22.58 ^c
Alaska	6.44	0.76	05(01,49)	22.29 ^c
Low Five States				
New Jersey	4.71	0.15	47(32,50)	-10.61
New York	4.58	0.10	48(41,50)	-13.10
D.C.	4.21	0.60	49(09,51)	-20.00 ^c
New Mexico	4.08	0.28	50(40,51)	-22.48 ^c
Hawaii	3.01	0.28	51(50,51)	-42.89 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	6.15	0.23	10(01,23)	16.68 ^c	Montana	5.05	0.43	37(05,50)	-4.06
Alaska	6.44	0.76	05(01,49)	22.29 ^c	Nebraska	6.21	0.38	09(01,36)	17.88 ^c
Arizona	4.95	0.17	43(22,49)	-6.10	Nevada	4.99	0.28	42(14,50)	-5.31
Arkansas	6.06	0.29	13(01,32)	15.07 ^c	New Hampshire	6.09	0.43	12(01,42)	15.64 ^c
California	5.23	0.08	30(22,41)	-0.77	New Jersey	4.71	0.15	47(32,50)	-10.61
Colorado	5.47	0.22	21(09,44)	3.92	New Mexico	4.08	0.28	50(40,51)	-22.48 ^c
Connecticut	5.04	0.24	40(15,49)	-4.24	New York	4.58	0.10	48(41,50)	-13.10
Delaware	4.84	0.47	45(06,50)	-8.10	North Carolina	5.46	0.16	22(12,40)	3.72
D.C.	4.21	0.60	49(09,51)	-20.00 ^c	North Dakota	5.19	0.55	32(01,50)	-1.41
Florida	5.05	0.10	38(25,45)	-4.10	Ohio	5.44	0.14	25(13,39)	3.19
Georgia	5.12	0.16	34(19,47)	-2.88	Oklahoma	5.46	0.24	23(08,45)	3.70
Hawaii	3.01	0.28	51(50,51)	-42.89 ^c	Oregon	6.29	0.25	08(01,22)	19.37 ^c
Idaho	6.70	0.42	02(01,23)	27.23 ^c	Pennsylvania	5.13	0.12	33(21,45)	-2.66
Illinois	5.09	0.13	36(21,46)	-3.40	Rhode Island	5.05	0.44	39(04,50)	-4.18
Indiana	5.56	0.19	20(09,39)	5.60	South Carolina	5.57	0.22	19(07,41)	5.81
Iowa	6.35	0.28	07(01,22)	20.48 ^c	South Dakota	6.72	0.55	01(01,36)	27.52 ^c
Kansas	6.39	0.30	06(01,24)	21.29 ^c	Tennessee	5.72	0.19	15(06,34)	8.54
Kentucky	5.45	0.23	24(09,45)	3.49	Texas	4.93	0.10	44(30,47)	-6.41
Louisiana	5.26	0.23	28(12,47)	-0.05	Utah	5.27	0.31	27(07,49)	-0.03
Maine	6.46	0.42	04(01,32)	22.58 ^c	Vermont	6.66	0.63	03(01,43)	26.50 ^c
Maryland	5.10	0.20	35(16,48)	-3.25	Virginia	4.71	0.16	46(31,50)	-10.50
Massachusetts	5.23	0.18	29(14,46)	-0.74	Washington	6.14	0.19	11(02,22)	16.61 ^c
Michigan	5.69	0.15	16(08,32)	7.97	West Virginia	5.00	0.31	41(12,50)	-5.12
Minnesota	5.64	0.21	17(07,38)	7.13	Wisconsin	5.82	0.20	14(04,31)	10.53
Mississippi	5.64	0.29	18(04,45)	7.07	Wyoming	5.23	0.61	31(01,50)	-0.80
Missouri	5.32	0.19	26(12,45)	1.01					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 3.20
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.51	0.02		
High Five States				
South Dakota	4.57	0.45	01(01,37)	30.13 ^c
Wyoming	4.39	0.54	02(01,48)	25.20 ^c
Vermont	4.22	0.47	03(01,48)	20.41 ^c
Mississippi	4.15	0.22	04(01,28)	18.24 ^c
Tennessee	4.11	0.15	05(01,22)	17.22 ^c
Low Five States				
New Mexico	3.06	0.23	47(17,51)	-12.75
Idaho	2.97	0.26	48(18,51)	-15.31 ^c
New Jersey	2.96	0.11	49(38,50)	-15.59 ^c
D.C.	2.63	0.40	50(12,51)	-25.14 ^c
Hawaii	2.21	0.24	51(47,51)	-37.11 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.05	0.17	12(01,26)	15.37 ^c	Montana	4.02	0.37	14(01,47)	14.42
Alaska	3.36	0.49	36(01,51)	-4.19	Nebraska	4.07	0.28	07(01,40)	15.96 ^c
Arizona	3.57	0.14	23(10,44)	1.66	Nevada	3.16	0.21	44(16,50)	-9.83
Arkansas	3.93	0.21	15(01,38)	12.02	New Hampshire	3.51	0.30	28(02,50)	-0.07
California	3.49	0.06	32(19,40)	-0.61	New Jersey	2.96	0.11	49(38,50)	-15.59 ^c
Colorado	3.68	0.17	19(05,42)	4.87	New Mexico	3.06	0.23	47(17,51)	-12.75
Connecticut	3.47	0.18	33(11,48)	-1.13	New York	3.16	0.07	45(33,49)	-9.95
Delaware	3.72	0.37	17(01,50)	6.03	North Carolina	3.52	0.11	26(14,43)	0.43
D.C.	2.63	0.40	50(12,51)	-25.14 ^c	North Dakota	3.49	0.43	31(01,51)	-0.47
Florida	3.28	0.07	40(27,47)	-6.52	Ohio	3.60	0.10	22(12,39)	2.72
Georgia	3.23	0.11	43(24,49)	-7.91	Oklahoma	3.67	0.18	20(04,44)	4.64
Hawaii	2.21	0.24	51(47,51)	-37.11 ^c	Oregon	4.06	0.19	08(01,30)	15.67 ^c
Idaho	2.97	0.26	48(18,51)	-15.31 ^c	Pennsylvania	3.53	0.09	25(15,42)	0.56
Illinois	3.34	0.10	39(21,47)	-4.70	Rhode Island	3.14	0.31	46(07,51)	-10.40
Indiana	3.62	0.14	21(09,43)	3.28	South Carolina	3.25	0.15	41(17,50)	-7.39
Iowa	4.04	0.21	13(01,33)	15.23 ^c	South Dakota	4.57	0.45	01(01,37)	30.13 ^c
Kansas	4.05	0.22	11(01,36)	15.49 ^c	Tennessee	4.11	0.15	05(01,22)	17.22 ^c
Kentucky	4.06	0.18	09(01,28)	15.62 ^c	Texas	3.35	0.07	37(24,45)	-4.42
Louisiana	3.46	0.16	34(13,48)	-1.26	Utah	3.51	0.24	27(04,49)	-0.05
Maine	3.50	0.29	29(02,50)	-0.24	Vermont	4.22	0.47	03(01,48)	20.41 ^c
Maryland	3.24	0.14	42(20,49)	-7.74	Virginia	3.35	0.12	38(19,48)	-4.49
Massachusetts	3.50	0.13	30(13,46)	-0.28	Washington	4.07	0.15	06(01,23)	16.12 ^c
Michigan	3.85	0.12	16(04,29)	9.58	West Virginia	3.54	0.25	24(03,49)	0.82
Minnesota	3.46	0.15	35(12,47)	-1.38	Wisconsin	4.05	0.16	10(01,25)	15.54 ^c
Mississippi	4.15	0.22	04(01,28)	18.24 ^c	Wyoming	4.39	0.54	02(01,48)	25.20 ^c
Missouri	3.69	0.14	18(06,40)	5.14					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 4.23
Cancer of the Female Breast (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	21.52	0.05		
High Five States				
D.C.	29.33	1.34	01(01,01)	36.25 ^c
Louisiana	24.53	0.44	02(02,06)	13.99 ^c
Mississippi	24.09	0.53	03(02,17)	11.93 ^c
New Jersey	23.38	0.29	04(02,15)	8.64
Ohio	23.32	0.25	05(03,15)	8.37
Low Five States				
Rhode Island	19.40	0.76	47(22,50)	-9.86
Vermont	19.08	0.98	48(19,50)	-11.37 ^c
North Dakota	19.01	0.96	49(21,50)	-11.65 ^c
Maine	18.84	0.65	50(30,50)	-12.48 ^c
Hawaii	14.86	0.61	51(51,51)	-30.94 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	22.20	0.39	20(05,28)	3.13	Montana	20.87	0.83	28(05,50)	-3.01
Alaska	21.09	1.28	25(02,50)	-2.01	Nebraska	19.78	0.61	40(23,50)	-8.10
Arizona	19.65	0.32	43(30,50)	-8.71	Nevada	23.08	0.57	07(02,25)	7.24
Arkansas	22.27	0.50	18(04,30)	3.47	New Hampshire	19.62	0.69	45(21,50)	-8.82
California	20.89	0.14	27(23,35)	-2.95	New Jersey	23.38	0.29	04(02,15)	8.64
Colorado	19.47	0.38	46(31,50)	-9.55	New Mexico	19.91	0.58	38(23,50)	-7.49
Connecticut	19.83	0.42	39(26,50)	-7.86	New York	21.22	0.19	24(21,32)	-1.40
Delaware	21.76	0.88	22(02,47)	1.09	North Carolina	21.70	0.28	23(13,30)	0.81
D.C.	29.33	1.34	01(01,01)	36.25 ^c	North Dakota	19.01	0.96	49(21,50)	-11.65 ^c
Florida	20.57	0.18	33(25,40)	-4.45	Ohio	23.32	0.25	05(03,15)	8.37
Georgia	22.93	0.30	09(03,21)	6.54	Oklahoma	23.22	0.46	06(02,22)	7.89
Hawaii	14.86	0.61	51(51,51)	-30.94 ^c	Oregon	20.47	0.42	34(22,47)	-4.91
Idaho	20.77	0.70	29(10,49)	-3.49	Pennsylvania	22.78	0.23	11(04,21)	5.85
Illinois	22.69	0.25	12(04,22)	5.44	Rhode Island	19.40	0.76	47(22,50)	-9.86
Indiana	22.30	0.34	17(05,26)	3.61	South Carolina	22.49	0.40	13(04,26)	4.52
Iowa	19.71	0.46	41(26,50)	-8.41	South Dakota	20.06	0.90	37(11,50)	-6.78
Kansas	20.57	0.50	32(19,48)	-4.43	Tennessee	22.49	0.34	14(04,25)	4.51
Kentucky	22.39	0.41	16(04,27)	4.03	Texas	20.66	0.18	31(24,38)	-3.99
Louisiana	24.53	0.44	02(02,06)	13.99 ^c	Utah	20.73	0.60	30(15,48)	-3.66
Maine	18.84	0.65	50(30,50)	-12.48 ^c	Vermont	19.08	0.98	48(19,50)	-11.37 ^c
Maryland	22.98	0.37	08(03,22)	6.75	Virginia	22.21	0.31	19(07,26)	3.20
Massachusetts	19.67	0.31	42(31,50)	-8.60	Washington	20.16	0.33	36(26,47)	-6.31
Michigan	22.47	0.27	15(05,24)	4.40	West Virginia	22.16	0.60	21(03,34)	2.98
Minnesota	19.64	0.35	44(30,50)	-8.73	Wisconsin	20.92	0.35	26(20,40)	-2.82
Mississippi	24.09	0.53	03(02,17)	11.93 ^c	Wyoming	20.23	1.13	35(04,50)	-6.01
Missouri	22.79	0.35	10(03,23)	5.90					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 5.14
Cancer of the Cervix Uteri (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.30	0.02		
High Five States				
West Virginia	3.45	0.26	01(01,12)	50.09 ^c
Mississippi	3.44	0.21	02(01,09)	49.63 ^c
Arkansas	3.43	0.21	03(01,10)	49.12 ^c
Alabama	3.12	0.16	04(01,13)	35.99 ^c
Kentucky	2.99	0.16	05(01,16)	30.01 ^c
Low Five States				
Utah	1.43	0.16	47(33,51)	-37.89 ^c
Massachusetts	1.33	0.08	48(40,51)	-41.94 ^c
Rhode Island	1.27	0.21	49(32,51)	-44.77 ^c
North Dakota	1.24	0.27	50(26,51)	-46.11 ^c
Vermont	1.18	0.25	51(30,51)	-48.52 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.12	0.16	04(01,13)	35.99 ^c	Montana	1.82	0.26	35(13,51)	-20.64 ^c
Alaska	1.92	0.35	34(04,51)	-16.39 ^c	Nebraska	2.04	0.21	31(10,45)	-11.14
Arizona	2.09	0.11	26(17,38)	-8.91	Nevada	2.13	0.17	23(12,41)	-7.20
Arkansas	3.43	0.21	03(01,10)	49.12 ^c	New Hampshire	1.59	0.20	42(22,51)	-30.83 ^c
California	2.26	0.05	22(17,29)	-1.77	New Jersey	2.34	0.10	21(13,30)	1.83
Colorado	1.54	0.11	43(33,50)	-33.03 ^c	New Mexico	2.06	0.20	29(12,44)	-10.19
Connecticut	1.60	0.12	41(30,50)	-30.56 ^c	New York	2.34	0.06	20(14,27)	1.98
Delaware	2.61	0.32	15(01,40)	13.55	North Carolina	2.08	0.09	27(20,37)	-9.61
D.C.	2.92	0.43	07(01,38)	27.08 ^c	North Dakota	1.24	0.27	50(26,51)	-46.11 ^c
Florida	2.74	0.07	09(05,17)	19.39 ^c	Ohio	2.52	0.09	17(09,24)	9.65
Georgia	2.61	0.10	13(07,21)	13.70	Oklahoma	2.91	0.18	08(02,20)	26.47 ^c
Hawaii	2.06	0.24	30(08,47)	-10.52	Oregon	1.78	0.13	38(24,47)	-22.58 ^c
Idaho	1.67	0.21	40(21,51)	-27.24 ^c	Pennsylvania	2.12	0.08	24(19,35)	-7.82
Illinois	2.42	0.08	19(11,26)	5.41	Rhode Island	1.27	0.21	49(32,51)	-44.77 ^c
Indiana	2.46	0.12	18(08,28)	6.90	South Carolina	2.67	0.15	12(04,23)	16.26 ^c
Iowa	2.01	0.16	32(16,43)	-12.65	South Dakota	1.69	0.28	39(14,51)	-26.37 ^c
Kansas	1.94	0.16	33(17,45)	-15.37 ^c	Tennessee	2.68	0.12	11(05,21)	16.59 ^c
Kentucky	2.99	0.16	05(01,16)	30.01 ^c	Texas	2.73	0.07	10(06,17)	18.72 ^c
Louisiana	2.96	0.16	06(02,16)	28.88 ^c	Utah	1.43	0.16	47(33,51)	-37.89 ^c
Maine	1.44	0.19	46(29,51)	-37.27 ^c	Vermont	1.18	0.25	51(30,51)	-48.52 ^c
Maryland	2.07	0.11	28(18,39)	-9.79	Virginia	1.80	0.09	36(27,44)	-21.56 ^c
Massachusetts	1.33	0.08	48(40,51)	-41.94 ^c	Washington	1.79	0.10	37(26,44)	-21.91 ^c
Michigan	2.12	0.09	25(18,36)	-7.91	West Virginia	3.45	0.26	01(01,12)	50.09 ^c
Minnesota	1.45	0.10	44(36,51)	-36.78 ^c	Wisconsin	1.45	0.10	45(37,51)	-37.08 ^c
Mississippi	3.44	0.21	02(01,09)	49.63 ^c	Wyoming	2.60	0.42	16(01,46)	13.32
Missouri	2.61	0.13	14(06,23)	13.60					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 6.25
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	15.10	0.03		
High Five States				
Mississippi	19.46	0.35	01(01,03)	28.92 ^c
West Virginia	18.78	0.40	02(01,06)	24.39 ^c
Arkansas	18.16	0.33	03(01,08)	20.27 ^c
Louisiana	18.08	0.28	04(02,07)	19.78 ^c
Kentucky	17.61	0.27	05(02,10)	16.62 ^c
Low Five States				
Minnesota	13.49	0.22	47(36,49)	-10.60 ^c
Arizona	13.24	0.19	48(40,49)	-12.31 ^c
Colorado	12.95	0.23	49(43,50)	-14.20 ^c
Connecticut	12.20	0.24	50(48,51)	-19.16 ^c
Utah	11.50	0.33	51(50,51)	-23.82 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	16.89	0.25	09(05,17)	11.90 ^c	Montana	13.95	0.48	40(22,49)	-7.62
Alaska	15.64	0.83	21(03,47)	3.61	Nebraska	16.27	0.40	15(05,26)	7.78
Arizona	13.24	0.19	48(40,49)	-12.31 ^c	Nevada	16.97	0.36	08(03,19)	12.43 ^c
Arkansas	18.16	0.33	03(01,08)	20.27 ^c	New Hampshire	13.70	0.43	43(26,49)	-9.27
California	13.86	0.09	41(35,46)	-8.19	New Jersey	15.65	0.18	20(15,25)	3.64
Colorado	12.95	0.23	49(43,50)	-14.20 ^c	New Mexico	14.31	0.36	34(22,47)	-5.19
Connecticut	12.20	0.24	50(48,51)	-19.16 ^c	New York	14.63	0.12	29(25,36)	-3.07
Delaware	14.09	0.52	38(20,49)	-6.68	North Carolina	14.27	0.17	36(28,43)	-5.48
D.C.	16.69	0.76	10(02,32)	10.59 ^c	North Dakota	15.56	0.62	22(06,42)	3.08
Florida	13.96	0.10	39(33,45)	-7.50	Ohio	16.59	0.16	11(07,18)	9.91
Georgia	15.48	0.19	23(16,28)	2.54	Oklahoma	17.27	0.29	06(03,14)	14.42 ^c
Hawaii	13.68	0.41	44(27,49)	-9.36	Oregon	14.56	0.26	30(23,42)	-3.53
Idaho	13.52	0.41	46(29,50)	-10.42 ^c	Pennsylvania	16.33	0.14	14(09,20)	8.16
Illinois	16.20	0.15	17(10,21)	7.31	Rhode Island	14.72	0.48	28(15,47)	-2.52
Indiana	16.24	0.22	16(08,21)	7.60	South Carolina	15.79	0.25	18(11,27)	4.57
Iowa	16.43	0.30	12(06,22)	8.84	South Dakota	15.71	0.57	19(06,38)	4.07
Kansas	15.30	0.31	24(14,34)	1.36	Tennessee	17.11	0.22	07(04,13)	13.34 ^c
Kentucky	17.61	0.27	05(02,10)	16.62 ^c	Texas	15.07	0.12	26(21,31)	-0.16
Louisiana	18.08	0.28	04(02,07)	19.78 ^c	Utah	11.50	0.33	51(50,51)	-23.82 ^c
Maine	14.44	0.41	32(20,48)	-4.34	Vermont	14.35	0.62	33(16,49)	-4.96
Maryland	14.86	0.22	27(21,37)	-1.55	Virginia	14.52	0.19	31(25,40)	-3.81
Massachusetts	13.77	0.19	42(33,48)	-8.82	Washington	13.55	0.20	45(36,49)	-10.25 ^c
Michigan	15.22	0.16	25(19,30)	0.82	West Virginia	18.78	0.40	02(01,06)	24.39 ^c
Minnesota	13.49	0.22	47(36,49)	-10.60 ^c	Wisconsin	14.17	0.21	37(28,45)	-6.16
Mississippi	19.46	0.35	01(01,03)	28.92 ^c	Wyoming	14.27	0.70	35(14,50)	-5.47
Missouri	16.35	0.22	13(07,21)	8.34					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 6.26
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	18.13	0.05		
High Five States				
Mississippi	23.82	0.61	01(01,04)	31.38 ^c
West Virginia	23.12	0.68	02(01,06)	27.55 ^c
Louisiana	22.10	0.48	03(01,09)	21.93 ^c
Arkansas	22.01	0.56	04(01,10)	21.38 ^c
Kentucky	21.15	0.46	05(02,14)	16.64 ^c
Low Five States				
Arizona	15.52	0.31	47(39,50)	-14.37 ^c
New Hampshire	15.28	0.70	48(29,51)	-15.72 ^c
Colorado	14.83	0.38	49(43,51)	-18.22 ^c
Connecticut	14.32	0.40	50(46,51)	-21.01 ^c
Utah	13.05	0.52	51(49,51)	-28.03 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	21.14	0.44	06(02,13)	16.58 ^c	Montana	16.46	0.77	41(21,50)	-9.20
Alaska	18.20	1.35	24(03,49)	0.41	Nebraska	18.75	0.65	19(08,37)	3.41
Arizona	15.52	0.31	47(39,50)	-14.37 ^c	Nevada	20.44	0.59	09(03,22)	12.74 ^c
Arkansas	22.01	0.56	04(01,10)	21.38 ^c	New Hampshire	15.28	0.70	48(29,51)	-15.72 ^c
California	16.37	0.14	42(35,45)	-9.73	New Jersey	18.88	0.30	18(12,28)	4.12
Colorado	14.83	0.38	49(43,51)	-18.22 ^c	New Mexico	17.58	0.60	30(15,44)	-3.02
Connecticut	14.32	0.40	50(46,51)	-21.01 ^c	New York	17.40	0.20	31(25,38)	-4.01
Delaware	17.92	0.90	28(08,47)	-1.14	North Carolina	17.61	0.29	29(21,39)	-2.85
D.C.	18.03	1.24	27(04,49)	-0.54	North Dakota	18.30	1.02	22(05,47)	0.93
Florida	16.77	0.17	38(30,43)	-7.52	Ohio	20.42	0.27	10(05,15)	12.65 ^c
Georgia	19.44	0.33	15(09,24)	7.22	Oklahoma	20.99	0.49	07(02,16)	15.75 ^c
Hawaii	17.30	0.69	33(16,47)	-4.59	Oregon	16.92	0.42	36(24,46)	-6.67
Idaho	16.31	0.67	43(22,50)	-10.06 ^c	Pennsylvania	19.58	0.24	14(09,21)	8.01
Illinois	19.63	0.26	13(08,22)	8.25	Rhode Island	16.65	0.80	39(18,50)	-8.14
Indiana	19.79	0.37	12(07,22)	9.17	South Carolina	19.31	0.43	17(08,27)	6.49
Iowa	19.44	0.50	16(07,28)	7.21	South Dakota	18.72	0.94	20(05,44)	3.28
Kansas	18.11	0.52	26(13,39)	-0.08	Tennessee	20.95	0.38	08(03,13)	15.54 ^c
Kentucky	21.15	0.46	05(02,14)	16.64 ^c	Texas	18.65	0.20	21(16,28)	2.89
Louisiana	22.10	0.48	03(01,09)	21.93 ^c	Utah	13.05	0.52	51(49,51)	-28.03 ^c
Maine	17.00	0.69	35(18,48)	-6.23	Vermont	16.24	1.02	44(15,51)	-10.41 ^c
Maryland	18.12	0.38	25(16,36)	-0.04	Virginia	17.30	0.32	32(23,42)	-4.58
Massachusetts	16.53	0.33	40(29,47)	-8.83	Washington	15.63	0.32	46(38,49)	-13.78 ^c
Michigan	18.29	0.28	23(17,32)	0.90	West Virginia	23.12	0.68	02(01,06)	27.55 ^c
Minnesota	15.68	0.35	45(37,49)	-13.53 ^c	Wisconsin	16.81	0.35	37(26,46)	-7.26
Mississippi	23.82	0.61	01(01,04)	31.38 ^c	Wyoming	17.25	1.13	34(07,50)	-4.84
Missouri	19.85	0.37	11(06,22)	9.50					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 6.27
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	12.69	0.04		
High Five States				
Mississippi	16.14	0.43	01(01,05)	27.17 ^c
D.C.	15.76	0.98	02(01,25)	24.13 ^c
West Virginia	15.38	0.49	03(01,12)	21.13 ^c
Arkansas	15.08	0.41	04(01,13)	18.78 ^c
Louisiana	14.97	0.34	05(01,12)	17.96 ^c
Low Five States				
Idaho	11.14	0.51	47(25,51)	-12.23 ^c
Delaware	10.98	0.61	48(24,51)	-13.53 ^c
Hawaii	10.75	0.50	49(31,51)	-15.35 ^c
Connecticut	10.56	0.30	50(43,51)	-16.84 ^c
Utah	10.06	0.42	51(45,51)	-20.72 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	13.68	0.30	14(06,26)	7.80	Montana	11.71	0.60	40(15,51)	-7.74
Alaska	13.28	1.04	20(01,50)	4.60	Nebraska	14.26	0.50	08(02,26)	12.35 ^c
Arizona	11.28	0.24	46(35,50)	-11.14 ^c	Nevada	13.87	0.45	11(03,30)	9.23
Arkansas	15.08	0.41	04(01,13)	18.78 ^c	New Hampshire	12.53	0.55	30(09,48)	-1.28
California	11.85	0.11	36(32,45)	-6.63	New Jersey	13.26	0.21	21(11,28)	4.47
Colorado	11.46	0.29	45(30,50)	-9.71	New Mexico	11.70	0.44	41(22,50)	-7.82
Connecticut	10.56	0.30	50(43,51)	-16.84 ^c	New York	12.54	0.14	29(21,36)	-1.19
Delaware	10.98	0.61	48(24,51)	-13.53 ^c	North Carolina	11.74	0.20	39(31,47)	-7.54
D.C.	15.76	0.98	02(01,25)	24.13 ^c	North Dakota	13.30	0.77	19(02,47)	4.74
Florida	11.61	0.13	43(34,48)	-8.58	Ohio	13.69	0.19	13(08,23)	7.80
Georgia	12.59	0.22	28(19,38)	-0.80	Oklahoma	14.31	0.36	07(03,21)	12.69 ^c
Hawaii	10.75	0.50	49(31,51)	-15.35 ^c	Oregon	12.63	0.32	27(15,41)	-0.52
Idaho	11.14	0.51	47(25,51)	-12.23 ^c	Pennsylvania	13.80	0.17	12(07,21)	8.73
Illinois	13.63	0.19	15(09,23)	7.34	Rhode Island	13.33	0.60	18(04,43)	4.99
Indiana	13.41	0.26	17(08,28)	5.66	South Carolina	12.93	0.30	23(12,36)	1.83
Iowa	14.10	0.37	10(03,23)	11.05 ^c	South Dakota	13.04	0.69	22(04,47)	2.69
Kansas	12.87	0.38	24(10,40)	1.42	Tennessee	14.23	0.27	09(04,19)	12.10 ^c
Kentucky	14.77	0.33	06(02,14)	16.36 ^c	Texas	12.20	0.14	34(26,40)	-3.88
Louisiana	14.97	0.34	05(01,12)	17.96 ^c	Utah	10.06	0.42	51(45,51)	-20.72 ^c
Maine	12.45	0.51	32(10,47)	-1.92	Vermont	12.69	0.78	26(04,50)	-0.01
Maryland	12.46	0.27	31(19,42)	-1.87	Virginia	12.34	0.23	33(21,42)	-2.83
Massachusetts	11.76	0.23	38(28,48)	-7.36	Washington	11.77	0.25	37(28,48)	-7.32
Michigan	12.84	0.20	25(17,34)	1.13	West Virginia	15.38	0.49	03(01,12)	21.13 ^c
Minnesota	11.69	0.27	42(28,49)	-7.94	Wisconsin	12.09	0.26	35(23,46)	-4.78
Mississippi	16.14	0.43	01(01,05)	27.17 ^c	Wyoming	11.59	0.87	44(08,51)	-8.72
Missouri	13.61	0.27	16(07,25)	7.19					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 7.14
Cancer of the Corpus and Uterus, NOS (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.46	0.02		
High Five States				
D.C.	7.78	0.68	01(01,02)	74.57 ^c
New Jersey	5.46	0.14	02(02,10)	22.45 ^c
New York	5.45	0.09	03(02,08)	22.21 ^c
Vermont	5.42	0.51	04(01,38)	21.66 ^c
Maryland	5.35	0.18	05(02,12)	19.95 ^c
Low Five States				
Nevada	3.43	0.22	47(32,51)	-22.97 ^c
Alabama	3.42	0.15	48(37,51)	-23.33 ^c
Arkansas	3.24	0.19	49(38,51)	-27.41 ^c
North Dakota	3.17	0.39	50(22,51)	-28.97 ^c
Wyoming	2.97	0.44	51(22,51)	-33.43 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.42	0.15	48(37,51)	-23.33 ^c	Montana	4.11	0.36	34(06,50)	-7.67
Alaska	3.66	0.55	44(04,51)	-17.79 ^c	Nebraska	4.46	0.29	20(04,45)	0.10
Arizona	3.49	0.14	46(37,51)	-21.60 ^c	Nevada	3.43	0.22	47(32,51)	-22.97 ^c
Arkansas	3.24	0.19	49(38,51)	-27.41 ^c	New Hampshire	4.60	0.33	14(02,44)	3.26
California	4.35	0.07	24(16,34)	-2.48	New Jersey	5.46	0.14	02(02,10)	22.45 ^c
Colorado	3.83	0.17	39(24,49)	-14.16	New Mexico	3.52	0.24	45(27,51)	-21.02 ^c
Connecticut	4.46	0.20	22(08,39)	0.02	New York	5.45	0.09	03(02,08)	22.21 ^c
Delaware	4.99	0.42	10(02,40)	11.91	North Carolina	4.33	0.12	25(13,38)	-2.89
D.C.	7.78	0.68	01(01,02)	74.57 ^c	North Dakota	3.17	0.39	50(22,51)	-28.97 ^c
Florida	4.08	0.08	36(25,40)	-8.44	Ohio	4.80	0.11	11(06,22)	7.80
Georgia	4.24	0.13	31(15,40)	-4.75	Oklahoma	3.84	0.18	38(22,49)	-13.92
Hawaii	4.46	0.33	21(03,45)	0.06	Oregon	4.32	0.19	27(10,41)	-3.02
Idaho	4.18	0.31	33(07,49)	-6.24	Pennsylvania	5.20	0.11	06(02,12)	16.66 ^c
Illinois	5.13	0.12	07(03,13)	15.21 ^c	Rhode Island	4.53	0.36	19(02,46)	1.65
Indiana	4.63	0.15	12(08,32)	3.89	South Carolina	4.63	0.18	13(07,34)	3.87
Iowa	5.11	0.23	08(02,22)	14.63	South Dakota	4.54	0.43	18(02,48)	1.77
Kansas	4.33	0.23	26(09,43)	-2.91	Tennessee	3.81	0.14	40(28,48)	-14.52
Kentucky	3.72	0.17	43(29,50)	-16.47 ^c	Texas	3.76	0.08	42(34,47)	-15.62 ^c
Louisiana	3.77	0.17	41(27,49)	-15.38 ^c	Utah	4.09	0.27	35(11,49)	-8.33
Maine	4.31	0.30	29(06,47)	-3.26	Vermont	5.42	0.51	04(01,38)	21.66 ^c
Maryland	5.35	0.18	05(02,12)	19.95 ^c	Virginia	4.58	0.14	15(09,32)	2.83
Massachusetts	4.55	0.15	17(09,34)	2.08	Washington	4.28	0.15	30(13,40)	-3.88
Michigan	5.09	0.13	09(03,15)	14.22	West Virginia	4.32	0.26	28(08,45)	-3.17
Minnesota	4.23	0.16	32(13,42)	-5.17	Wisconsin	4.56	0.16	16(09,34)	2.29
Mississippi	4.00	0.21	37(15,48)	-10.32	Wyoming	2.97	0.44	51(22,51)	-33.43 ^c
Missouri	4.37	0.15	23(11,39)	-1.97					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 8.18
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.14	0.02		
High Five States				
Maine	5.69	0.26	01(01,07)	37.49 ^c
New Hampshire	5.53	0.27	02(01,12)	33.61 ^c
Wyoming	5.17	0.42	03(01,41)	24.85 ^c
West Virginia	5.01	0.21	04(01,23)	21.02 ^c
Ohio	5.00	0.09	05(02,12)	20.85 ^c
Low Five States				
New Mexico	3.46	0.18	47(37,51)	-16.51 ^c
California	3.41	0.04	48(46,50)	-17.57 ^c
Texas	3.35	0.05	49(46,51)	-18.98 ^c
Utah	3.10	0.17	50(46,51)	-25.12 ^c
Hawaii	3.01	0.19	51(46,51)	-27.20 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.01	0.12	40(22,46)	-3.20	Montana	4.20	0.26	30(05,47)	1.58
Alaska	4.42	0.43	20(01,50)	6.79	Nebraska	4.63	0.21	15(03,37)	11.95
Arizona	4.09	0.11	35(20,45)	-1.18	Nevada	4.67	0.19	14(03,33)	12.71
Arkansas	4.11	0.16	34(15,46)	-0.82	New Hampshire	5.53	0.27	02(01,12)	33.61 ^c
California	3.41	0.04	48(46,50)	-17.57 ^c	New Jersey	3.94	0.09	42(28,46)	-4.81
Colorado	3.88	0.13	45(27,47)	-6.20	New Mexico	3.46	0.18	47(37,51)	-16.51 ^c
Connecticut	4.03	0.14	38(19,47)	-2.66	New York	3.94	0.06	43(31,46)	-4.90
Delaware	4.32	0.28	27(03,47)	4.38	North Carolina	3.89	0.09	44(30,47)	-6.00
D.C.	4.12	0.38	32(03,51)	-0.52	North Dakota	4.25	0.34	28(03,49)	2.76
Florida	3.97	0.06	41(30,46)	-4.15	Ohio	5.00	0.09	05(02,12)	20.85 ^c
Georgia	3.78	0.09	46(34,47)	-8.71	Oklahoma	4.34	0.14	25(10,41)	4.90
Hawaii	3.01	0.19	51(46,51)	-27.20 ^c	Oregon	4.41	0.14	22(09,39)	6.48
Idaho	4.05	0.22	37(11,47)	-2.08	Pennsylvania	4.81	0.08	08(03,18)	16.19 ^c
Illinois	4.42	0.08	19(12,32)	6.80	Rhode Island	4.67	0.28	13(02,43)	12.74
Indiana	4.91	0.12	06(02,18)	18.70 ^c	South Carolina	4.11	0.12	33(18,46)	-0.61
Iowa	4.61	0.16	16(04,32)	11.49	South Dakota	4.83	0.32	07(01,42)	16.68 ^c
Kansas	4.40	0.17	23(07,42)	6.20	Tennessee	4.14	0.11	31(19,45)	-0.01
Kentucky	4.35	0.13	24(10,39)	5.12	Texas	3.35	0.05	49(46,51)	-18.98 ^c
Louisiana	4.42	0.14	21(08,38)	6.79	Utah	3.10	0.17	50(46,51)	-25.12 ^c
Maine	5.69	0.26	01(01,07)	37.49 ^c	Vermont	4.77	0.36	10(01,45)	15.18 ^c
Maryland	4.03	0.12	39(22,46)	-2.73	Virginia	4.08	0.10	36(22,46)	-1.46
Massachusetts	4.74	0.11	12(03,23)	14.43	Washington	4.48	0.11	17(09,32)	8.23
Michigan	4.80	0.09	09(03,19)	15.92 ^c	West Virginia	5.01	0.21	04(01,23)	21.02 ^c
Minnesota	4.45	0.12	18(08,34)	7.61	Wisconsin	4.75	0.12	11(03,23)	14.73
Mississippi	4.22	0.16	29(11,46)	2.00	Wyoming	5.17	0.42	03(01,41)	24.85 ^c
Missouri	4.33	0.11	26(12,39)	4.56					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 8.19
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	7.37	0.03		
High Five States				
Maine	10.27	0.51	01(01,09)	39.24 ^c
New Hampshire	9.80	0.54	02(01,15)	32.92 ^c
West Virginia	9.09	0.41	03(01,20)	23.22 ^c
Ohio	9.07	0.18	04(02,12)	22.96 ^c
Wyoming	8.91	0.81	05(01,45)	20.87 ^c
Low Five States				
Texas	6.01	0.11	47(45,51)	-18.46 ^c
New Mexico	5.95	0.34	48(37,51)	-19.31 ^c
California	5.95	0.09	49(45,51)	-19.35 ^c
Hawaii	5.54	0.39	50(43,51)	-24.83 ^c
Utah	5.45	0.33	51(45,51)	-26.06 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	7.20	0.25	32(18,46)	-2.38	Montana	7.07	0.50	39(10,50)	-4.16
Alaska	7.07	0.81	40(02,51)	-4.17	Nebraska	8.36	0.43	14(02,36)	13.40
Arizona	7.09	0.21	37(21,46)	-3.79	Nevada	7.90	0.35	18(05,42)	7.13
Arkansas	7.27	0.31	30(14,46)	-1.36	New Hampshire	9.80	0.54	02(01,15)	32.92 ^c
California	5.95	0.09	49(45,51)	-19.35 ^c	New Jersey	7.04	0.18	41(25,46)	-4.58
Colorado	6.92	0.25	44(23,47)	-6.10	New Mexico	5.95	0.34	48(37,51)	-19.31 ^c
Connecticut	7.14	0.28	35(18,46)	-3.20	New York	7.00	0.12	43(29,46)	-5.13
Delaware	7.18	0.54	33(07,50)	-2.57	North Carolina	7.09	0.18	38(24,46)	-3.85
D.C.	6.63	0.75	46(06,51)	-10.05	North Dakota	7.80	0.67	22(02,49)	5.74
Florida	7.01	0.11	42(29,46)	-4.87	Ohio	9.07	0.18	04(02,12)	22.96 ^c
Georgia	6.78	0.19	45(30,47)	-8.04	Oklahoma	7.86	0.29	20(08,40)	6.53
Hawaii	5.54	0.39	50(43,51)	-24.83 ^c	Oregon	7.76	0.28	23(09,41)	5.24
Idaho	7.13	0.44	36(11,49)	-3.28	Pennsylvania	8.74	0.16	08(03,15)	18.56 ^c
Illinois	7.90	0.16	17(12,30)	7.16	Rhode Island	8.60	0.57	10(01,40)	16.66 ^c
Indiana	8.89	0.24	07(02,16)	20.51 ^c	South Carolina	7.22	0.25	31(17,46)	-2.05
Iowa	8.55	0.33	12(02,26)	15.95 ^c	South Dakota	8.90	0.64	06(01,38)	20.74 ^c
Kansas	8.08	0.34	15(04,36)	9.58	Tennessee	7.55	0.22	27(14,42)	2.44
Kentucky	7.88	0.27	19(09,38)	6.83	Texas	6.01	0.11	47(45,51)	-18.46 ^c
Louisiana	7.83	0.27	21(08,40)	6.25	Utah	5.45	0.33	51(45,51)	-26.06 ^c
Maine	10.27	0.51	01(01,09)	39.24 ^c	Vermont	8.05	0.69	16(01,49)	9.13
Maryland	7.15	0.23	34(19,46)	-2.98	Virginia	7.36	0.20	29(17,44)	-0.23
Massachusetts	8.72	0.23	09(02,18)	18.32 ^c	Washington	7.75	0.22	24(12,38)	5.14
Michigan	8.56	0.19	11(04,19)	16.03 ^c	West Virginia	9.09	0.41	03(01,20)	23.22 ^c
Minnesota	7.52	0.24	28(13,44)	1.95	Wisconsin	8.39	0.24	13(04,24)	13.75
Mississippi	7.60	0.33	25(10,45)	3.11	Wyoming	8.91	0.81	05(01,45)	20.87 ^c
Missouri	7.56	0.22	26(14,42)	2.56					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 8.20
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	1.52	0.01		
High Five States				
D.C.	2.33	0.37	01(01,45)	52.63 ^c
Vermont	2.04	0.32	02(01,49)	33.75 ^c
Minnesota	1.95	0.11	03(01,22)	27.72 ^c
New Hampshire	1.93	0.21	04(01,43)	26.90 ^c
Alaska	1.93	0.39	05(01,51)	26.60 ^c
Low Five States				
Kansas	1.31	0.12	47(14,51)	-14.07
North Dakota	1.20	0.24	48(04,51)	-21.42 ^c
Texas	1.18	0.04	49(43,51)	-22.92 ^c
Utah	1.05	0.13	50(33,51)	-31.00 ^c
Hawaii	0.87	0.14	51(42,51)	-42.90 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	1.47	0.10	35(10,48)	-3.88	Montana	1.65	0.23	21(01,50)	8.07
Alaska	1.93	0.39	05(01,51)	26.60 ^c	Nebraska	1.61	0.17	24(02,49)	5.56
Arizona	1.50	0.09	31(09,47)	-1.29	Nevada	1.72	0.15	14(01,46)	13.07
Arkansas	1.47	0.13	34(06,49)	-3.77	New Hampshire	1.93	0.21	04(01,43)	26.90 ^c
California	1.37	0.04	43(30,48)	-9.96	New Jersey	1.56	0.07	29(09,44)	2.14
Colorado	1.36	0.10	44(18,50)	-11.10	New Mexico	1.35	0.15	45(08,51)	-11.13
Connecticut	1.61	0.12	23(03,46)	5.57	New York	1.57	0.05	28(13,38)	3.05
Delaware	1.93	0.25	06(01,47)	26.31 ^c	North Carolina	1.37	0.07	41(23,49)	-9.90
D.C.	2.33	0.37	01(01,45)	52.63 ^c	North Dakota	1.20	0.24	48(04,51)	-21.42 ^c
Florida	1.43	0.04	36(24,46)	-6.37	Ohio	1.77	0.07	08(02,27)	16.16 ^c
Georgia	1.40	0.07	39(20,48)	-8.46	Oklahoma	1.41	0.11	38(10,50)	-7.63
Hawaii	0.87	0.14	51(42,51)	-42.90 ^c	Oregon	1.59	0.11	26(04,47)	4.23
Idaho	1.34	0.17	46(06,51)	-12.26	Pennsylvania	1.70	0.06	15(04,30)	11.69
Illinois	1.69	0.07	17(05,32)	10.69	Rhode Island	1.63	0.21	22(01,50)	7.20
Indiana	1.73	0.09	12(03,37)	13.26	South Carolina	1.59	0.10	27(05,46)	4.21
Iowa	1.38	0.12	40(11,50)	-9.27	South Dakota	1.53	0.24	30(01,51)	0.37
Kansas	1.31	0.12	47(14,51)	-14.07	Tennessee	1.37	0.08	42(21,49)	-9.96
Kentucky	1.42	0.10	37(11,49)	-6.84	Texas	1.18	0.04	49(43,51)	-22.92 ^c
Louisiana	1.65	0.11	20(03,44)	8.22	Utah	1.05	0.13	50(33,51)	-31.00 ^c
Maine	1.84	0.20	07(01,46)	20.92 ^c	Vermont	2.04	0.32	02(01,49)	33.75 ^c
Maryland	1.60	0.10	25(05,45)	5.07	Virginia	1.47	0.08	33(12,47)	-3.69
Massachusetts	1.66	0.09	18(04,39)	8.82	Washington	1.75	0.10	10(03,34)	15.00 ^c
Michigan	1.74	0.07	11(03,30)	14.39	West Virginia	1.66	0.16	19(02,48)	8.79
Minnesota	1.95	0.11	03(01,22)	27.72 ^c	Wisconsin	1.72	0.10	13(03,38)	13.11
Mississippi	1.50	0.13	32(06,49)	-1.69	Wyoming	1.77	0.33	09(01,51)	16.08 ^c
Missouri	1.69	0.09	16(03,38)	10.78					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 9.18
Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.36	0.00		
High Five States				
Vermont	0.64	0.14	01(01,43)	77.81 ^c
Louisiana	0.47	0.05	02(01,28)	32.63 ^c
West Virginia	0.44	0.07	03(01,43)	23.49 ^c
Mississippi	0.44	0.05	04(01,42)	23.42 ^c
New Hampshire	0.43	0.08	05(01,46)	19.17 ^c
Low Five States				
Arizona	0.29	0.03	42(15,46)	-18.22 ^c
Colorado	0.28	0.03	43(14,46)	-22.76 ^c
Kentucky	0.27	0.04	44(15,46)	-23.40 ^c
Utah	0.26	0.05	45(07,46)	-26.38 ^c
Hawaii	0.21	0.05	46(13,46)	-41.54 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.38	0.04	13(02,43)	6.99	Montana	0.34	0.08	35(01,46)	-5.86
Alaska	-	-	-(-,-)	-	Nebraska	0.36	0.06	26(01,46)	0.93
Arizona	0.29	0.03	42(15,46)	-18.22 ^c	Nevada	0.32	0.05	38(03,46)	-9.47
Arkansas	0.40	0.05	10(01,44)	12.08	New Hampshire	0.43	0.08	05(01,46)	19.17 ^c
California	0.38	0.01	15(06,32)	5.98	New Jersey	0.32	0.03	37(10,45)	-9.44
Colorado	0.28	0.03	43(14,46)	-22.76 ^c	New Mexico	0.37	0.06	18(01,46)	4.63
Connecticut	0.32	0.04	39(05,46)	-11.30	New York	0.36	0.02	27(08,39)	0.91
Delaware	0.37	0.09	17(01,46)	4.78	North Carolina	0.34	0.03	34(08,44)	-5.01
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.35	0.02	28(09,40)	-0.95	Ohio	0.36	0.02	25(05,41)	1.07
Georgia	0.36	0.03	24(04,42)	1.10	Oklahoma	0.40	0.05	09(01,43)	13.18
Hawaii	0.21	0.05	46(13,46)	-41.54 ^c	Oregon	0.36	0.04	23(02,45)	1.69
Idaho	0.41	0.07	08(01,46)	13.80	Pennsylvania	0.37	0.02	19(05,38)	3.50
Illinois	0.31	0.02	41(18,45)	-13.81	Rhode Island	0.41	0.08	07(01,46)	14.99
Indiana	0.35	0.03	31(04,44)	-1.50	South Carolina	0.33	0.04	36(05,46)	-6.73
Iowa	0.36	0.05	22(02,45)	1.92	South Dakota	-	-	-(-,-)	-
Kansas	0.37	0.05	20(02,45)	2.66	Tennessee	0.42	0.04	06(01,35)	18.15 ^c
Kentucky	0.27	0.04	44(15,46)	-23.40 ^c	Texas	0.39	0.02	11(05,34)	8.17
Louisiana	0.47	0.05	02(01,28)	32.63 ^c	Utah	0.26	0.05	45(07,46)	-26.38 ^c
Maine	0.38	0.07	16(01,46)	5.22	Vermont	0.64	0.14	01(01,43)	77.81 ^c
Maryland	0.35	0.03	32(04,44)	-1.63	Virginia	0.35	0.03	29(05,43)	-0.95
Massachusetts	0.34	0.03	33(05,45)	-3.94	Washington	0.31	0.03	40(10,46)	-12.56
Michigan	0.38	0.03	12(03,39)	7.58	West Virginia	0.44	0.07	03(01,43)	23.49 ^c
Minnesota	0.35	0.04	30(04,45)	-1.44	Wisconsin	0.38	0.04	14(02,42)	6.93
Mississippi	0.44	0.05	04(01,42)	23.42 ^c	Wyoming	-	-	-(-,-)	-
Missouri	0.37	0.03	21(03,43)	2.42					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 9.19
Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.45	0.01		
High Five States				
New Hampshire	0.62	0.15	01(01,42)	37.91 ^c
Rhode Island	0.59	0.15	02(01,42)	30.82 ^c
Tennessee	0.58	0.06	03(01,30)	29.31 ^c
Nebraska	0.56	0.11	04(01,41)	23.99 ^c
Mississippi	0.56	0.09	05(01,40)	23.29 ^c
Low Five States				
Connecticut	0.35	0.06	38(06,42)	-21.67 ^c
Arizona	0.34	0.05	39(15,42)	-23.56 ^c
Utah	0.33	0.08	40(04,42)	-26.57 ^c
Kentucky	0.32	0.06	41(14,42)	-29.85 ^c
Colorado	0.28	0.05	42(24,42)	-38.91 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.44	0.06	28(02,41)	-1.84	Montana	-	-	-(-,-)	-
Alaska	-	-	-(-,-)	-	Nebraska	0.56	0.11	04(01,41)	23.99 ^c
Arizona	0.34	0.05	39(15,42)	-23.56 ^c	Nevada	0.41	0.08	34(02,42)	-8.58
Arkansas	0.46	0.08	19(01,42)	1.88	New Hampshire	0.62	0.15	01(01,42)	37.91 ^c
California	0.50	0.02	11(04,28)	11.29	New Jersey	0.38	0.04	36(10,42)	-14.69
Colorado	0.28	0.05	42(24,42)	-38.91 ^c	New Mexico	0.54	0.11	07(01,42)	19.06 ^c
Connecticut	0.35	0.06	38(06,42)	-21.67 ^c	New York	0.45	0.03	26(07,38)	-1.28
Delaware	-	-	-(-,-)	-	North Carolina	0.45	0.05	23(04,40)	-0.70
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.44	0.03	30(09,38)	-2.62	Ohio	0.45	0.04	22(04,38)	0.16
Georgia	0.51	0.05	09(01,36)	13.32	Oklahoma	0.47	0.07	15(01,41)	5.24
Hawaii	-	-	-(-,-)	-	Oregon	0.50	0.07	13(01,40)	10.14
Idaho	0.45	0.11	24(01,42)	-1.03	Pennsylvania	0.42	0.04	32(08,40)	-6.79
Illinois	0.39	0.04	35(12,41)	-14.56	Rhode Island	0.59	0.15	02(01,42)	30.82 ^c
Indiana	0.46	0.06	21(02,40)	1.74	South Carolina	0.46	0.06	20(02,41)	1.83
Iowa	0.47	0.08	17(01,42)	4.93	South Dakota	-	-	-(-,-)	-
Kansas	0.44	0.08	31(01,42)	-3.36	Tennessee	0.58	0.06	03(01,30)	29.31 ^c
Kentucky	0.32	0.06	41(14,42)	-29.85 ^c	Texas	0.52	0.03	08(02,26)	16.36 ^c
Louisiana	0.55	0.07	06(01,38)	20.94 ^c	Utah	0.33	0.08	40(04,42)	-26.57 ^c
Maine	0.50	0.12	10(01,42)	11.45	Vermont	-	-	-(-,-)	-
Maryland	0.42	0.06	33(04,42)	-7.62	Virginia	0.50	0.05	12(01,38)	10.72
Massachusetts	0.45	0.05	25(03,40)	-1.05	Washington	0.36	0.05	37(11,42)	-20.49 ^c
Michigan	0.48	0.05	14(02,37)	7.30	West Virginia	0.47	0.10	16(01,42)	4.96
Minnesota	0.44	0.06	27(02,41)	-1.65	Wisconsin	0.47	0.06	18(02,40)	3.59
Mississippi	0.56	0.09	05(01,40)	23.29 ^c	Wyoming	-	-	-(-,-)	-
Missouri	0.44	0.06	29(03,41)	-2.10					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 9.20
Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.28	0.01		
High Five States				
Louisiana	0.42	0.06	01(01,24)	51.42 ^c
West Virginia	0.41	0.09	02(01,34)	47.88 ^c
Mississippi	0.36	0.07	03(01,34)	30.09 ^c
Oklahoma	0.36	0.06	04(01,34)	29.74 ^c
Arkansas	0.36	0.07	05(01,34)	29.00 ^c
Low Five States				
Arizona	0.25	0.04	31(05,35)	-12.02
Nevada	0.25	0.06	32(02,35)	-12.18
Illinois	0.25	0.03	33(08,35)	-12.44
Kentucky	0.23	0.04	34(04,35)	-18.98 ^c
South Carolina	0.22	0.04	35(06,35)	-23.08 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.33	0.05	07(01,34)	16.66 ^c	Montana	-	-	-(-,-)	-
Alaska	-	-	-(-,-)	-	Nebraska	-	-	-(-,-)	-
Arizona	0.25	0.04	31(05,35)	-12.02	Nevada	0.25	0.06	32(02,35)	-12.18
Arkansas	0.36	0.07	05(01,34)	29.00 ^c	New Hampshire	-	-	-(-,-)	-
California	0.28	0.02	16(07,31)	-0.24	New Jersey	0.27	0.03	24(04,35)	-2.83
Colorado	0.27	0.05	25(02,35)	-2.92	New Mexico	-	-	-(-,-)	-
Connecticut	0.28	0.05	19(02,35)	-0.61	New York	0.29	0.02	12(04,31)	4.94
Delaware	-	-	-(-,-)	-	North Carolina	0.25	0.03	29(06,35)	-10.93
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.28	0.02	17(05,33)	-0.26	Ohio	0.29	0.03	14(03,34)	1.88
Georgia	0.26	0.03	26(04,35)	-6.28	Oklahoma	0.36	0.06	04(01,34)	29.74 ^c
Hawaii	-	-	-(-,-)	-	Oregon	0.25	0.05	28(03,35)	-9.93
Idaho	-	-	-(-,-)	-	Pennsylvania	0.33	0.03	06(01,27)	17.62 ^c
Illinois	0.25	0.03	33(08,35)	-12.44	Rhode Island	-	-	-(-,-)	-
Indiana	0.28	0.04	20(03,35)	-1.28	South Carolina	0.22	0.04	35(06,35)	-23.08 ^c
Iowa	0.28	0.06	18(01,35)	-0.50	South Dakota	-	-	-(-,-)	-
Kansas	0.31	0.07	08(01,35)	11.97	Tennessee	0.28	0.04	15(02,35)	1.63
Kentucky	0.23	0.04	34(04,35)	-18.98 ^c	Texas	0.27	0.02	22(07,33)	-1.93
Louisiana	0.42	0.06	01(01,24)	51.42 ^c	Utah	-	-	-(-,-)	-
Maine	-	-	-(-,-)	-	Vermont	-	-	-(-,-)	-
Maryland	0.29	0.04	13(02,35)	2.72	Virginia	0.25	0.03	30(06,35)	-11.66
Massachusetts	0.26	0.04	27(04,35)	-7.98	Washington	0.28	0.04	21(03,35)	-1.36
Michigan	0.30	0.03	11(02,33)	5.80	West Virginia	0.41	0.09	02(01,34)	47.88 ^c
Minnesota	0.27	0.04	23(02,35)	-1.96	Wisconsin	0.31	0.04	09(01,34)	10.14
Mississippi	0.36	0.07	03(01,34)	30.09 ^c	Wyoming	-	-	-(-,-)	-
Missouri	0.31	0.04	10(01,35)	9.59					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 11.18
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.90	0.02		
High Five States				
Oklahoma	5.38	0.16	01(01,04)	38.22 ^c
Louisiana	5.00	0.15	02(01,09)	28.31 ^c
Alaska	4.78	0.46	03(01,42)	22.61 ^c
Arkansas	4.72	0.17	04(02,19)	21.13 ^c
Kentucky	4.62	0.14	05(02,20)	18.57 ^c
Low Five States				
New York	3.23	0.05	47(41,49)	-17.01 ^c
Wyoming	3.11	0.32	48(22,51)	-20.04 ^c
Connecticut	3.10	0.12	49(40,51)	-20.54 ^c
Hawaii	2.68	0.18	50(46,51)	-31.07 ^c
D.C.	2.68	0.30	51(36,51)	-31.15 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.04	0.12	24(13,35)	3.84	Montana	3.42	0.24	42(20,51)	-12.31
Alaska	4.78	0.46	03(01,42)	22.61 ^c	Nebraska	4.46	0.21	14(02,30)	14.60
Arizona	3.71	0.10	34(24,43)	-4.74	Nevada	3.64	0.17	37(21,48)	-6.49
Arkansas	4.72	0.17	04(02,19)	21.13 ^c	New Hampshire	3.79	0.22	31(11,48)	-2.80
California	3.50	0.04	41(34,45)	-10.07	New Jersey	3.33	0.08	44(37,49)	-14.61
Colorado	3.51	0.12	40(28,48)	-9.78	New Mexico	4.34	0.20	15(02,33)	11.54
Connecticut	3.10	0.12	49(40,51)	-20.54 ^c	New York	3.23	0.05	47(41,49)	-17.01 ^c
Delaware	3.77	0.27	33(07,49)	-3.20	North Carolina	3.90	0.09	29(20,37)	0.16
D.C.	2.68	0.30	51(36,51)	-31.15 ^c	North Dakota	4.08	0.33	23(02,48)	4.77
Florida	3.37	0.05	43(38,48)	-13.47	Ohio	4.20	0.08	19(12,28)	7.72
Georgia	3.54	0.09	39(30,46)	-9.19	Oklahoma	5.38	0.16	01(01,04)	38.22 ^c
Hawaii	2.68	0.18	50(46,51)	-31.07 ^c	Oregon	3.91	0.13	28(15,40)	0.46
Idaho	3.99	0.22	25(05,44)	2.51	Pennsylvania	3.95	0.07	27(19,34)	1.52
Illinois	4.12	0.08	22(14,30)	5.68	Rhode Island	3.78	0.25	32(10,48)	-2.91
Indiana	4.51	0.11	12(03,21)	15.81 ^c	South Carolina	3.98	0.12	26(14,38)	2.06
Iowa	4.60	0.16	07(02,22)	17.97 ^c	South Dakota	4.53	0.31	10(01,36)	16.41 ^c
Kansas	4.49	0.17	13(02,27)	15.23 ^c	Tennessee	4.31	0.11	18(06,27)	10.67
Kentucky	4.62	0.14	05(02,20)	18.57 ^c	Texas	4.52	0.06	11(04,17)	16.02 ^c
Louisiana	5.00	0.15	02(01,09)	28.31 ^c	Utah	3.58	0.18	38(20,49)	-8.20
Maine	4.14	0.22	21(04,40)	6.24	Vermont	3.28	0.30	46(19,51)	-15.72 ^c
Maryland	3.67	0.11	36(25,44)	-5.82	Virginia	3.70	0.09	35(25,42)	-4.96
Massachusetts	3.28	0.09	45(37,50)	-15.70 ^c	Washington	3.80	0.10	30(21,41)	-2.51
Michigan	4.15	0.09	20(12,30)	6.66	West Virginia	4.61	0.20	06(02,25)	18.43 ^c
Minnesota	4.33	0.12	16(04,27)	11.27	Wisconsin	4.32	0.12	17(05,27)	10.81
Mississippi	4.55	0.17	09(02,24)	16.77 ^c	Wyoming	3.11	0.32	48(22,51)	-20.04 ^c
Missouri	4.57	0.12	08(02,19)	17.42 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 11.19
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	5.68	0.03		
High Five States				
Oklahoma	7.95	0.29	01(01,04)	39.93 ^c
Louisiana	6.93	0.26	02(01,19)	22.04 ^c
Iowa	6.91	0.29	03(01,21)	21.76 ^c
Mississippi	6.91	0.33	04(01,24)	21.73 ^c
Arkansas	6.76	0.30	05(01,25)	19.07 ^c
Low Five States				
Connecticut	4.67	0.23	47(35,51)	-17.83 ^c
Massachusetts	4.65	0.17	48(37,51)	-18.18 ^c
Wyoming	4.47	0.56	49(15,51)	-21.36 ^c
D.C.	4.38	0.61	50(13,51)	-22.93 ^c
Hawaii	4.14	0.34	51(38,51)	-27.03 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	6.02	0.23	21(05,36)	5.95	Montana	5.32	0.44	37(07,51)	-6.36
Alaska	6.19	0.78	18(01,51)	8.98	Nebraska	6.60	0.38	07(01,33)	16.16 ^c
Arizona	5.39	0.18	35(21,45)	-4.99	Nevada	5.13	0.29	38(20,50)	-9.73
Arkansas	6.76	0.30	05(01,25)	19.07 ^c	New Hampshire	5.91	0.42	25(02,47)	4.15
California	5.07	0.08	39(34,46)	-10.70	New Jersey	5.02	0.16	42(32,49)	-11.61
Colorado	4.94	0.22	43(30,50)	-12.95	New Mexico	5.79	0.34	28(05,46)	1.88
Connecticut	4.67	0.23	47(35,51)	-17.83 ^c	New York	4.89	0.10	45(36,49)	-13.82
Delaware	5.04	0.48	40(09,51)	-11.26	North Carolina	5.99	0.17	23(09,34)	5.46
D.C.	4.38	0.61	50(13,51)	-22.93 ^c	North Dakota	6.04	0.59	20(01,50)	6.37
Florida	4.90	0.09	44(37,49)	-13.62	Ohio	5.88	0.14	26(15,35)	3.48
Georgia	5.32	0.17	36(24,46)	-6.27	Oklahoma	7.95	0.29	01(01,04)	39.93 ^c
Hawaii	4.14	0.34	51(38,51)	-27.03 ^c	Oregon	5.69	0.24	29(12,42)	0.15
Idaho	5.64	0.39	31(05,49)	-0.74	Pennsylvania	5.59	0.13	33(21,39)	-1.52
Illinois	6.01	0.15	22(11,33)	5.91	Rhode Island	6.11	0.48	19(02,47)	7.52
Indiana	6.48	0.21	11(02,26)	14.04	South Carolina	6.25	0.24	17(03,33)	10.14
Iowa	6.91	0.29	03(01,21)	21.76 ^c	South Dakota	5.95	0.53	24(02,49)	4.70
Kansas	6.49	0.31	10(02,32)	14.24	Tennessee	6.57	0.21	08(02,24)	15.64 ^c
Kentucky	6.66	0.26	06(02,25)	17.31 ^c	Texas	6.43	0.12	12(04,22)	13.27
Louisiana	6.93	0.26	02(01,19)	22.04 ^c	Utah	5.04	0.32	41(21,51)	-11.33
Maine	6.32	0.41	16(02,39)	11.22	Vermont	4.70	0.53	46(13,51)	-17.30 ^c
Maryland	5.67	0.21	30(14,42)	-0.08	Virginia	5.52	0.18	34(20,42)	-2.77
Massachusetts	4.65	0.17	48(37,51)	-18.18 ^c	Washington	5.60	0.19	32(17,41)	-1.36
Michigan	5.87	0.16	27(14,36)	3.45	West Virginia	6.38	0.36	13(02,37)	12.32
Minnesota	6.38	0.22	14(03,30)	12.31	Wisconsin	6.32	0.21	15(03,29)	11.38
Mississippi	6.91	0.33	04(01,24)	21.73 ^c	Wyoming	4.47	0.56	49(15,51)	-21.36 ^c
Missouri	6.54	0.21	09(02,25)	15.13 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 11.20
Cancer of the Kidney And Renal Pelvis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.48	0.02		
High Five States				
Alaska	3.48	0.55	01(01,45)	40.07 ^c
Louisiana	3.44	0.16	02(01,10)	38.57 ^c
South Dakota	3.33	0.36	03(01,37)	34.04 ^c
Oklahoma	3.29	0.17	04(01,16)	32.45 ^c
West Virginia	3.23	0.22	05(01,23)	29.91 ^c
Low Five States				
Wyoming	1.95	0.35	47(06,51)	-21.68 ^c
Connecticut	1.88	0.13	48(33,51)	-24.36 ^c
Montana	1.71	0.23	49(25,51)	-30.98 ^c
Hawaii	1.49	0.18	50(39,51)	-39.92 ^c
D.C.	1.48	0.30	51(28,51)	-40.47 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	2.54	0.13	25(10,41)	2.24	Montana	1.71	0.23	49(25,51)	-30.98 ^c
Alaska	3.48	0.55	01(01,45)	40.07 ^c	Nebraska	2.64	0.21	22(03,45)	6.41
Arizona	2.26	0.11	37(22,47)	-8.99	Nevada	2.34	0.18	31(11,48)	-5.91
Arkansas	3.10	0.18	06(01,25)	24.71 ^c	New Hampshire	1.95	0.21	46(21,51)	-21.50 ^c
California	2.24	0.05	38(28,44)	-9.90	New Jersey	2.03	0.08	44(33,49)	-18.14 ^c
Colorado	2.32	0.13	32(17,47)	-6.51	New Mexico	3.10	0.22	07(01,28)	24.67 ^c
Connecticut	1.88	0.13	48(33,51)	-24.36 ^c	New York	2.00	0.06	45(38,49)	-19.35 ^c
Delaware	2.82	0.31	14(01,47)	13.58	North Carolina	2.39	0.09	29(18,42)	-3.76
D.C.	1.48	0.30	51(28,51)	-40.47 ^c	North Dakota	2.55	0.35	24(01,50)	2.54
Florida	2.11	0.05	42(33,47)	-15.03 ^c	Ohio	2.87	0.09	13(05,23)	15.49 ^c
Georgia	2.21	0.09	39(25,47)	-10.98	Oklahoma	3.29	0.17	04(01,16)	32.45 ^c
Hawaii	1.49	0.18	50(39,51)	-39.92 ^c	Oregon	2.46	0.14	27(11,44)	-1.08
Idaho	2.51	0.24	26(04,48)	1.10	Pennsylvania	2.67	0.08	21(11,29)	7.58
Illinois	2.68	0.08	20(10,29)	7.83	Rhode Island	2.18	0.25	41(09,51)	-12.29
Indiana	2.92	0.12	11(03,25)	17.68 ^c	South Carolina	2.28	0.13	35(19,47)	-8.01
Iowa	2.76	0.17	16(03,36)	11.08	South Dakota	3.33	0.36	03(01,37)	34.04 ^c
Kansas	2.87	0.19	12(02,32)	15.75 ^c	Tennessee	2.58	0.12	23(09,37)	4.09
Kentucky	3.01	0.15	09(01,25)	21.17 ^c	Texas	2.96	0.07	10(04,18)	19.38 ^c
Louisiana	3.44	0.16	02(01,10)	38.57 ^c	Utah	2.35	0.20	30(09,48)	-5.44
Maine	2.43	0.22	28(06,49)	-2.17	Vermont	2.07	0.31	43(07,51)	-16.54 ^c
Maryland	2.19	0.11	40(24,48)	-11.84	Virginia	2.29	0.10	34(22,46)	-7.98
Massachusetts	2.27	0.10	36(22,47)	-8.59	Washington	2.30	0.11	33(20,47)	-7.33
Michigan	2.80	0.09	15(06,27)	12.66	West Virginia	3.23	0.22	05(01,23)	29.91 ^c
Minnesota	2.68	0.13	19(06,35)	7.94	Wisconsin	2.68	0.12	18(07,33)	8.04
Mississippi	2.69	0.17	17(04,40)	8.47	Wyoming	1.95	0.35	47(06,51)	-21.68 ^c
Missouri	3.05	0.13	08(01,20)	22.81 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 12.18
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	1.05	0.01		
High Five States				
D.C.	1.69	0.24	01(01,28)	60.32 ^c
West Virginia	1.60	0.12	02(01,10)	52.11 ^c
Kentucky	1.45	0.08	03(01,13)	37.88 ^c
Tennessee	1.36	0.06	04(01,15)	29.47 ^c
Louisiana	1.33	0.07	05(01,22)	26.16 ^c
Low Five States				
Idaho	0.69	0.09	47(31,50)	-34.75 ^c
Wyoming	0.68	0.15	48(14,51)	-35.40 ^c
Colorado	0.66	0.05	49(39,50)	-37.37 ^c
South Dakota	0.65	0.12	50(27,51)	-38.43 ^c
Utah	0.34	0.06	51(50,51)	-67.50 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	1.31	0.07	06(02,23)	24.75 ^c	Montana	1.13	0.14	21(02,45)	7.07
Alaska	0.91	0.20	34(02,50)	-13.35	Nebraska	0.84	0.09	39(18,50)	-19.99 ^c
Arizona	0.79	0.05	42(32,49)	-24.41 ^c	Nevada	0.91	0.08	35(15,47)	-13.51
Arkansas	0.94	0.07	32(15,46)	-10.52	New Hampshire	1.04	0.12	28(03,47)	-0.68
California	0.78	0.02	43(36,47)	-25.38 ^c	New Jersey	1.14	0.05	19(08,30)	8.00
Colorado	0.66	0.05	49(39,50)	-37.37 ^c	New Mexico	0.88	0.09	37(15,50)	-16.55 ^c
Connecticut	0.86	0.06	38(25,48)	-17.75 ^c	New York	1.14	0.03	18(10,29)	8.07
Delaware	1.07	0.14	26(02,48)	1.62	North Carolina	1.11	0.05	24(10,32)	5.81
D.C.	1.69	0.24	01(01,28)	60.32 ^c	North Dakota	1.16	0.17	16(01,47)	9.98
Florida	1.16	0.03	15(09,27)	10.02	Ohio	1.31	0.04	07(03,16)	24.40 ^c
Georgia	1.07	0.05	27(12,35)	1.55	Oklahoma	0.97	0.07	30(15,44)	-7.93
Hawaii	0.82	0.10	41(18,50)	-22.21 ^c	Oregon	0.74	0.06	44(33,50)	-29.16 ^c
Idaho	0.69	0.09	47(31,50)	-34.75 ^c	Pennsylvania	1.12	0.04	22(10,30)	6.88
Illinois	1.18	0.04	12(06,26)	12.27	Rhode Island	1.13	0.14	20(02,44)	7.89
Indiana	1.15	0.06	17(06,31)	9.06	South Carolina	1.28	0.07	09(02,26)	21.52 ^c
Iowa	0.93	0.07	33(17,46)	-11.22	South Dakota	0.65	0.12	50(27,51)	-38.43 ^c
Kansas	0.74	0.07	45(31,50)	-29.92 ^c	Tennessee	1.36	0.06	04(01,15)	29.47 ^c
Kentucky	1.45	0.08	03(01,13)	37.88 ^c	Texas	1.11	0.03	25(12,30)	5.77
Louisiana	1.33	0.07	05(01,22)	26.16 ^c	Utah	0.34	0.06	51(50,51)	-67.50 ^c
Maine	1.31	0.12	08(01,32)	24.16 ^c	Vermont	1.16	0.18	14(01,47)	10.65
Maryland	1.17	0.06	13(05,30)	11.52	Virginia	1.00	0.05	29(16,39)	-4.59
Massachusetts	0.96	0.05	31(21,41)	-8.88	Washington	0.89	0.05	36(26,45)	-15.20 ^c
Michigan	1.20	0.05	11(05,26)	13.89	West Virginia	1.60	0.12	02(01,10)	52.11 ^c
Minnesota	0.71	0.05	46(36,50)	-32.05 ^c	Wisconsin	0.83	0.05	40(29,48)	-21.27 ^c
Mississippi	1.26	0.09	10(02,30)	19.69 ^c	Wyoming	0.68	0.15	48(14,51)	-35.40 ^c
Missouri	1.12	0.06	23(07,33)	6.15					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.19
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	1.88	0.02		
High Five States				
D.C.	3.08	0.51	01(01,33)	63.79 ^c
West Virginia	2.56	0.22	02(01,25)	36.17 ^c
Kentucky	2.48	0.15	03(01,19)	31.64 ^c
South Carolina	2.46	0.15	04(01,19)	30.72 ^c
Tennessee	2.43	0.13	05(01,18)	28.82 ^c
Low Five States				
Wyoming	1.21	0.29	47(12,51)	-35.67 ^c
South Dakota	1.20	0.24	48(21,51)	-36.28 ^c
Idaho	1.15	0.17	49(31,51)	-39.17 ^c
Colorado	1.08	0.10	50(41,50)	-42.45 ^c
Utah	0.56	0.11	51(49,51)	-70.05 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	2.40	0.14	06(01,22)	27.31 ^c	Montana	1.94	0.27	25(01,48)	3.07
Alaska	1.55	0.38	38(01,51)	-17.60 ^c	Nebraska	1.53	0.19	40(13,50)	-18.54 ^c
Arizona	1.37	0.09	43(32,49)	-26.97 ^c	Nevada	1.58	0.16	36(15,49)	-16.26 ^c
Arkansas	1.69	0.15	31(12,46)	-10.15	New Hampshire	1.84	0.24	28(02,48)	-2.22
California	1.45	0.04	42(33,46)	-22.78 ^c	New Jersey	1.96	0.10	24(09,33)	4.08
Colorado	1.08	0.10	50(41,50)	-42.45 ^c	New Mexico	1.56	0.17	37(14,49)	-17.23 ^c
Connecticut	1.59	0.13	35(20,48)	-15.39 ^c	New York	2.06	0.07	15(08,28)	9.45
Delaware	1.72	0.27	30(03,50)	-8.65	North Carolina	2.04	0.09	18(07,31)	8.11
D.C.	3.08	0.51	01(01,33)	63.79 ^c	North Dakota	2.04	0.34	17(01,49)	8.26
Florida	2.07	0.06	14(09,27)	9.93	Ohio	2.29	0.09	10(02,20)	21.66 ^c
Georgia	1.96	0.10	23(08,34)	4.20	Oklahoma	1.76	0.14	29(11,44)	-6.47
Hawaii	1.63	0.21	34(08,49)	-13.53	Oregon	1.33	0.12	44(30,50)	-29.38 ^c
Idaho	1.15	0.17	49(31,51)	-39.17 ^c	Pennsylvania	1.99	0.08	21(10,31)	5.50
Illinois	2.19	0.09	11(04,25)	16.46 ^c	Rhode Island	1.86	0.27	27(02,49)	-1.46
Indiana	1.99	0.11	22(07,34)	5.50	South Carolina	2.46	0.15	04(01,19)	30.72 ^c
Iowa	1.65	0.14	33(15,47)	-12.50	South Dakota	1.20	0.24	48(21,51)	-36.28 ^c
Kansas	1.28	0.14	46(31,50)	-31.87 ^c	Tennessee	2.43	0.13	05(01,18)	28.82 ^c
Kentucky	2.48	0.15	03(01,19)	31.64 ^c	Texas	2.05	0.06	16(08,28)	9.11
Louisiana	2.33	0.15	08(01,26)	23.99 ^c	Utah	0.56	0.11	51(49,51)	-70.05 ^c
Maine	2.35	0.24	07(01,32)	24.94 ^c	Vermont	2.01	0.34	19(01,49)	6.51
Maryland	2.15	0.13	12(03,30)	14.06	Virginia	1.88	0.10	26(11,37)	-0.41
Massachusetts	1.67	0.10	32(20,44)	-11.17	Washington	1.54	0.10	39(27,47)	-18.08 ^c
Michigan	2.11	0.09	13(05,28)	12.33	West Virginia	2.56	0.22	02(01,25)	36.17 ^c
Minnesota	1.32	0.10	45(33,50)	-29.71 ^c	Wisconsin	1.48	0.10	41(28,48)	-21.36 ^c
Mississippi	2.30	0.18	09(01,29)	22.22 ^c	Wyoming	1.21	0.29	47(12,51)	-35.67 ^c
Missouri	1.99	0.12	20(07,34)	5.57					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 12.20
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.39	0.01		
High Five States				
West Virginia	0.77	0.11	01(01,10)	96.77 ^c
Kentucky	0.65	0.07	02(01,13)	66.17 ^c
Rhode Island	0.65	0.14	03(01,36)	64.68 ^c
Delaware	0.55	0.13	04(01,40)	39.56 ^c
Louisiana	0.55	0.06	05(01,25)	38.77 ^c
Low Five States				
New Mexico	0.30	0.07	36(07,40)	-24.75 ^c
Kansas	0.29	0.06	37(11,40)	-27.07 ^c
California	0.26	0.02	38(30,40)	-33.53 ^c
Oregon	0.26	0.05	39(19,40)	-33.86 ^c
Minnesota	0.22	0.04	40(27,40)	-42.86 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.46	0.05	12(03,32)	16.27 ^c	Montana	-	-	-(-,-)	-
Alaska	-	-	-(-,-)	-	Nebraska	-	-	-(-,-)	-
Arizona	0.30	0.04	35(17,40)	-23.51 ^c	Nevada	0.33	0.07	30(06,40)	-16.92 ^c
Arkansas	0.31	0.06	34(09,40)	-21.77 ^c	New Hampshire	0.39	0.10	21(02,40)	0.11
California	0.26	0.02	38(30,40)	-33.53 ^c	New Jersey	0.51	0.04	08(02,21)	30.52 ^c
Colorado	0.32	0.05	31(10,40)	-18.86 ^c	New Mexico	0.30	0.07	36(07,40)	-24.75 ^c
Connecticut	0.31	0.05	33(11,40)	-21.59 ^c	New York	0.45	0.03	13(05,25)	15.48 ^c
Delaware	0.55	0.13	04(01,40)	39.56 ^c	North Carolina	0.40	0.04	20(07,34)	1.85
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.39	0.02	22(11,32)	-1.14	Ohio	0.53	0.04	06(02,18)	35.72 ^c
Georgia	0.38	0.04	23(08,36)	-4.52	Oklahoma	0.34	0.05	29(08,40)	-14.36
Hawaii	-	-	-(-,-)	-	Oregon	0.26	0.05	39(19,40)	-33.86 ^c
Idaho	-	-	-(-,-)	-	Pennsylvania	0.46	0.03	11(04,25)	17.32 ^c
Illinois	0.42	0.03	18(07,31)	5.74	Rhode Island	0.65	0.14	03(01,36)	64.68 ^c
Indiana	0.48	0.05	09(03,31)	21.06 ^c	South Carolina	0.35	0.05	28(08,40)	-11.22
Iowa	0.37	0.06	24(05,40)	-7.06	South Dakota	-	-	-(-,-)	-
Kansas	0.29	0.06	37(11,40)	-27.07 ^c	Tennessee	0.53	0.05	07(02,23)	34.78 ^c
Kentucky	0.65	0.07	02(01,13)	66.17 ^c	Texas	0.35	0.02	26(15,35)	-9.72
Louisiana	0.55	0.06	05(01,25)	38.77 ^c	Utah	-	-	-(-,-)	-
Maine	0.43	0.10	15(01,40)	10.71	Vermont	-	-	-(-,-)	-
Maryland	0.43	0.05	16(04,35)	8.33	Virginia	0.35	0.04	27(11,38)	-10.73
Massachusetts	0.41	0.04	19(05,34)	5.57	Washington	0.36	0.04	25(09,38)	-7.74
Michigan	0.47	0.04	10(03,26)	20.59 ^c	West Virginia	0.77	0.11	01(01,10)	96.77 ^c
Minnesota	0.22	0.04	40(27,40)	-42.86 ^c	Wisconsin	0.32	0.04	32(13,40)	-19.01 ^c
Mississippi	0.42	0.07	17(03,39)	8.06	Wyoming	-	-	-(-,-)	-
Missouri	0.45	0.05	14(03,32)	14.35					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 13.26
Leukemia

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	6.89	0.02		
High Five States				
Wyoming	8.42	0.54	01(01,28)	22.10 ^c
Wisconsin	7.96	0.16	02(01,11)	15.52 ^c
South Dakota	7.88	0.42	03(01,33)	14.37
Oklahoma	7.78	0.20	04(01,19)	12.91
West Virginia	7.68	0.26	05(01,27)	11.35
Low Five States				
Colorado	6.41	0.17	47(25,49)	-7.01
New Mexico	5.95	0.23	48(33,51)	-13.65
Utah	5.90	0.24	49(34,51)	-14.44
D.C.	5.61	0.45	50(23,51)	-18.57 ^c
Hawaii	5.34	0.26	51(47,51)	-22.51 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	7.49	0.17	11(02,25)	8.65	Montana	6.59	0.34	39(08,50)	-4.41
Alaska	6.72	0.57	32(01,51)	-2.58	Nebraska	6.79	0.26	27(08,48)	-1.47
Arizona	6.59	0.14	38(22,48)	-4.36	Nevada	6.74	0.23	29(11,48)	-2.23
Arkansas	7.57	0.22	09(01,26)	9.83	New Hampshire	6.73	0.30	30(06,49)	-2.36
California	6.52	0.06	41(30,47)	-5.43	New Jersey	6.46	0.12	46(28,48)	-6.35
Colorado	6.41	0.17	47(25,49)	-7.01	New Mexico	5.95	0.23	48(33,51)	-13.65
Connecticut	6.80	0.18	26(15,47)	-1.30	New York	6.52	0.08	43(29,48)	-5.44
Delaware	6.88	0.37	24(02,49)	-0.22	North Carolina	6.73	0.12	31(21,45)	-2.42
D.C.	5.61	0.45	50(23,51)	-18.57 ^c	North Dakota	6.67	0.41	34(03,50)	-3.22
Florida	6.51	0.07	44(29,47)	-5.60	Ohio	7.24	0.11	16(08,27)	5.06
Georgia	6.46	0.12	45(27,48)	-6.32	Oklahoma	7.78	0.20	04(01,19)	12.91
Hawaii	5.34	0.26	51(47,51)	-22.51 ^c	Oregon	6.97	0.18	22(09,43)	1.14
Idaho	6.67	0.29	35(09,50)	-3.31	Pennsylvania	7.25	0.10	15(08,26)	5.21
Illinois	7.44	0.11	13(04,22)	7.97	Rhode Island	6.66	0.33	36(07,50)	-3.40
Indiana	7.46	0.15	12(03,24)	8.15	South Carolina	6.71	0.17	33(18,47)	-2.72
Iowa	7.16	0.20	20(05,38)	3.85	South Dakota	7.88	0.42	03(01,33)	14.37
Kansas	7.67	0.22	06(01,24)	11.19	Tennessee	7.37	0.15	14(04,26)	6.87
Kentucky	7.63	0.18	08(01,22)	10.75	Texas	6.80	0.08	25(21,39)	-1.29
Louisiana	7.16	0.18	21(05,36)	3.84	Utah	5.90	0.24	49(34,51)	-14.44
Maine	7.19	0.30	19(02,46)	4.23	Vermont	6.65	0.43	37(02,51)	-3.49
Maryland	6.56	0.15	40(22,48)	-4.83	Virginia	6.52	0.13	42(25,48)	-5.44
Massachusetts	6.78	0.14	28(18,45)	-1.68	Washington	6.89	0.14	23(14,41)	0.00
Michigan	7.19	0.11	18(08,29)	4.30	West Virginia	7.68	0.26	05(01,27)	11.35
Minnesota	7.64	0.16	07(01,21)	10.77	Wisconsin	7.96	0.16	02(01,11)	15.52 ^c
Mississippi	7.50	0.22	10(01,28)	8.72	Wyoming	8.42	0.54	01(01,28)	22.10 ^c
Missouri	7.20	0.15	17(06,33)	4.45					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 13.27
Leukemia

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	9.27	0.04		
High Five States				
Wyoming	10.88	0.93	01(01,47)	17.44 ^c
Wisconsin	10.88	0.28	02(01,16)	17.37 ^c
Arkansas	10.87	0.40	03(01,20)	17.30 ^c
Kentucky	10.62	0.34	04(01,23)	14.63
South Dakota	10.61	0.73	05(01,44)	14.48
Low Five States				
Colorado	8.40	0.29	47(26,50)	-9.32
New Mexico	7.82	0.40	48(32,51)	-15.62 ^c
D.C.	7.70	0.84	49(08,51)	-16.89 ^c
Utah	7.64	0.41	50(36,51)	-17.59 ^c
Hawaii	6.97	0.45	51(45,51)	-24.78 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	10.41	0.32	07(01,26)	12.29	Montana	8.42	0.57	46(11,51)	-9.09
Alaska	9.27	1.02	27(01,51)	-0.01	Nebraska	8.96	0.46	35(07,50)	-3.32
Arizona	8.63	0.23	44(26,49)	-6.88	Nevada	9.13	0.40	32(07,48)	-1.47
Arkansas	10.87	0.40	03(01,20)	17.30 ^c	New Hampshire	8.95	0.54	36(05,50)	-3.36
California	8.49	0.10	45(36,48)	-8.34	New Jersey	8.80	0.21	40(23,48)	-5.03
Colorado	8.40	0.29	47(26,50)	-9.32	New Mexico	7.82	0.40	48(32,51)	-15.62 ^c
Connecticut	9.42	0.33	24(07,45)	1.69	New York	8.95	0.14	37(24,45)	-3.38
Delaware	9.82	0.68	16(01,49)	5.98	North Carolina	9.17	0.22	29(16,44)	-1.04
D.C.	7.70	0.84	49(08,51)	-16.89 ^c	North Dakota	9.22	0.73	28(01,51)	-0.53
Florida	8.71	0.12	43(30,47)	-6.00	Ohio	9.88	0.19	14(06,28)	6.58
Georgia	8.73	0.23	42(23,48)	-5.74	Oklahoma	10.48	0.35	06(01,25)	13.12
Hawaii	6.97	0.45	51(45,51)	-24.78 ^c	Oregon	9.44	0.32	23(07,44)	1.89
Idaho	8.95	0.51	38(06,50)	-3.44	Pennsylvania	9.65	0.17	20(10,32)	4.14
Illinois	10.05	0.19	13(04,26)	8.51	Rhode Island	9.16	0.60	30(02,50)	-1.10
Indiana	9.83	0.26	15(04,32)	6.06	South Carolina	9.28	0.31	26(10,46)	0.18
Iowa	9.66	0.35	19(03,43)	4.26	South Dakota	10.61	0.73	05(01,44)	14.48
Kansas	10.34	0.40	09(01,31)	11.61	Tennessee	10.17	0.27	12(02,27)	9.76
Kentucky	10.62	0.34	04(01,23)	14.63	Texas	9.08	0.14	34(22,43)	-2.05
Louisiana	9.68	0.33	18(04,41)	4.45	Utah	7.64	0.41	50(36,51)	-17.59 ^c
Maine	9.49	0.52	22(02,48)	2.40	Vermont	9.39	0.79	25(01,51)	1.39
Maryland	8.77	0.27	41(20,49)	-5.40	Virginia	8.87	0.23	39(21,48)	-4.24
Massachusetts	9.09	0.24	33(16,46)	-1.95	Washington	9.13	0.25	31(14,46)	-1.41
Michigan	9.65	0.20	21(08,34)	4.14	West Virginia	10.20	0.46	11(01,37)	10.10
Minnesota	10.39	0.29	08(01,24)	12.08	Wisconsin	10.88	0.28	02(01,16)	17.37 ^c
Mississippi	10.26	0.41	10(01,33)	10.68	Wyoming	10.88	0.93	01(01,47)	17.44 ^c
Missouri	9.79	0.26	17(05,34)	5.69					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 13.28
Leukemia

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	5.16	0.02		
High Five States				
Wyoming	6.55	0.65	01(01,43)	26.94 ^c
West Virginia	5.87	0.31	02(01,33)	13.84
Indiana	5.80	0.17	03(01,21)	12.50
Oklahoma	5.78	0.23	04(01,27)	12.01
Wisconsin	5.76	0.18	05(01,24)	11.75
Low Five States				
Alaska	4.52	0.63	47(01,51)	-12.36
Utah	4.51	0.28	48(17,51)	-12.56
New Mexico	4.42	0.27	49(21,51)	-14.24
D.C.	4.30	0.51	50(03,51)	-16.62 ^c
Hawaii	4.08	0.32	51(30,51)	-20.79 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	5.48	0.19	13(02,36)	6.23	Montana	5.11	0.41	27(01,51)	-0.89
Alaska	4.52	0.63	47(01,51)	-12.36	Nebraska	5.20	0.30	23(02,48)	0.80
Arizona	4.98	0.16	32(12,47)	-3.50	Nevada	4.71	0.26	44(11,51)	-8.64
Arkansas	5.16	0.24	25(03,48)	0.10	New Hampshire	5.18	0.36	24(01,50)	0.41
California	5.03	0.07	30(19,41)	-2.40	New Jersey	4.82	0.13	38(21,49)	-6.53
Colorado	4.87	0.19	36(13,49)	-5.52	New Mexico	4.42	0.27	49(21,51)	-14.24
Connecticut	4.90	0.20	34(13,50)	-4.99	New York	4.80	0.09	39(27,48)	-6.85
Delaware	4.67	0.40	45(03,51)	-9.34	North Carolina	5.09	0.13	28(12,44)	-1.34
D.C.	4.30	0.51	50(03,51)	-16.62 ^c	North Dakota	4.71	0.47	43(02,51)	-8.57
Florida	4.77	0.08	42(29,48)	-7.43	Ohio	5.38	0.12	14(05,31)	4.38
Georgia	4.87	0.14	37(19,48)	-5.60	Oklahoma	5.78	0.23	04(01,27)	12.01
Hawaii	4.08	0.32	51(30,51)	-20.79 ^c	Oregon	5.07	0.21	29(06,48)	-1.66
Idaho	4.78	0.33	41(04,51)	-7.22	Pennsylvania	5.55	0.11	11(03,24)	7.60
Illinois	5.64	0.12	08(02,22)	9.44	Rhode Island	4.87	0.37	35(02,51)	-5.51
Indiana	5.80	0.17	03(01,21)	12.50	South Carolina	4.92	0.19	33(12,49)	-4.59
Iowa	5.34	0.23	18(02,45)	3.47	South Dakota	5.71	0.47	06(01,48)	10.65
Kansas	5.70	0.26	07(01,33)	10.62	Tennessee	5.36	0.17	17(03,37)	3.98
Kentucky	5.56	0.21	10(01,35)	7.89	Texas	5.13	0.09	26(14,39)	-0.48
Louisiana	5.37	0.20	16(02,41)	4.10	Utah	4.51	0.28	48(17,51)	-12.56
Maine	5.33	0.34	19(01,49)	3.37	Vermont	4.52	0.47	46(02,51)	-12.34
Maryland	5.01	0.17	31(10,47)	-2.85	Virginia	4.80	0.15	40(21,49)	-6.87
Massachusetts	5.22	0.16	21(07,42)	1.26	Washington	5.20	0.17	22(06,43)	0.90
Michigan	5.37	0.13	15(05,34)	4.12	West Virginia	5.87	0.31	02(01,33)	13.84
Minnesota	5.59	0.19	09(01,30)	8.31	Wisconsin	5.76	0.18	05(01,24)	11.75
Mississippi	5.53	0.25	12(01,40)	7.23	Wyoming	6.55	0.65	01(01,43)	26.94 ^c
Missouri	5.28	0.17	20(04,41)	2.42					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.18
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	6.14	0.02		
High Five States				
D.C.	9.59	0.57	01(01,02)	56.31 ^c
Louisiana	7.86	0.18	02(02,06)	28.04 ^c
Texas	7.81	0.08	03(02,04)	27.25 ^c
Hawaii	7.70	0.31	04(01,08)	25.45 ^c
California	7.43	0.06	05(03,07)	21.11 ^c
Low Five States				
Montana	4.69	0.28	47(32,51)	-23.49 ^c
Idaho	4.60	0.23	48(36,51)	-25.07 ^c
South Dakota	4.58	0.31	49(32,51)	-25.34 ^c
Utah	4.55	0.20	50(38,51)	-25.82 ^c
North Dakota	3.76	0.31	51(46,51)	-38.80 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	6.34	0.15	14(07,22)	3.36	Montana	4.69	0.28	47(32,51)	-23.49 ^c
Alaska	5.92	0.47	21(05,48)	-3.50	Nebraska	4.81	0.22	45(33,50)	-21.58 ^c
Arizona	6.00	0.13	20(13,30)	-2.23	Nevada	5.66	0.20	29(14,41)	-7.82
Arkansas	6.56	0.20	11(06,20)	6.99	New Hampshire	5.00	0.25	42(25,50)	-18.46 ^c
California	7.43	0.06	05(03,07)	21.11 ^c	New Jersey	5.63	0.11	30(20,38)	-8.20
Colorado	5.27	0.15	37(26,46)	-14.07	New Mexico	6.86	0.25	07(04,17)	11.77
Connecticut	5.43	0.16	34(22,43)	-11.45	New York	6.05	0.07	18(14,25)	-1.41
Delaware	6.61	0.35	09(04,30)	7.67	North Carolina	5.71	0.10	27(19,36)	-7.01
D.C.	9.59	0.57	01(01,02)	56.31 ^c	North Dakota	3.76	0.31	51(46,51)	-38.80 ^c
Florida	5.84	0.07	24(18,31)	-4.82	Ohio	5.47	0.09	33(25,40)	-10.88
Georgia	5.75	0.11	26(17,35)	-6.25	Oklahoma	6.01	0.17	19(10,33)	-2.04
Hawaii	7.70	0.31	04(01,08)	25.45 ^c	Oregon	6.34	0.17	15(07,24)	3.29
Idaho	4.60	0.23	48(36,51)	-25.07 ^c	Pennsylvania	5.68	0.08	28(20,36)	-7.38
Illinois	5.86	0.09	22(16,31)	-4.41	Rhode Island	6.67	0.33	08(04,28)	8.67
Indiana	5.38	0.12	35(26,43)	-12.26	South Carolina	5.86	0.15	23(14,35)	-4.52
Iowa	4.75	0.16	46(37,50)	-22.55 ^c	South Dakota	4.58	0.31	49(32,51)	-25.34 ^c
Kansas	5.10	0.18	41(29,48)	-16.83 ^c	Tennessee	6.42	0.13	13(07,20)	4.59
Kentucky	5.82	0.15	25(14,36)	-5.09	Texas	7.81	0.08	03(02,04)	27.25 ^c
Louisiana	7.86	0.18	02(02,06)	28.04 ^c	Utah	4.55	0.20	50(38,51)	-25.82 ^c
Maine	5.12	0.24	39(22,50)	-16.54 ^c	Vermont	5.16	0.37	38(14,50)	-15.88 ^c
Maryland	6.25	0.14	16(08,24)	1.85	Virginia	5.60	0.11	31(20,38)	-8.67
Massachusetts	6.46	0.13	12(07,18)	5.32	Washington	6.57	0.14	10(06,17)	7.12
Michigan	5.55	0.10	32(23,38)	-9.51	West Virginia	4.92	0.20	44(32,50)	-19.85 ^c
Minnesota	5.00	0.13	43(35,48)	-18.47 ^c	Wisconsin	5.29	0.13	36(27,44)	-13.71
Mississippi	7.12	0.21	06(03,11)	15.98 ^c	Wyoming	5.12	0.41	40(12,51)	-16.60 ^c
Missouri	6.07	0.13	17(11,28)	-1.02					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 14.19
 Cancer of the Liver and Intrahepatic Bile Duct (Invasive)
 Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	9.07	0.03		
High Five States				
D.C.	14.97	1.08	01(01,02)	65.04 ^c
Louisiana	11.81	0.33	02(01,06)	30.21 ^c
Texas	11.54	0.15	03(02,05)	27.24 ^c
Hawaii	11.31	0.55	04(02,12)	24.73 ^c
California	10.87	0.11	05(04,08)	19.83 ^c
Low Five States				
Idaho	6.46	0.41	47(36,51)	-28.73 ^c
Montana	6.38	0.47	48(35,51)	-29.64 ^c
Utah	5.97	0.34	49(44,51)	-34.18 ^c
South Dakota	5.96	0.52	50(38,51)	-34.29 ^c
North Dakota	5.40	0.55	51(42,51)	-40.42 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	9.39	0.28	14(07,26)	3.54	Montana	6.38	0.47	48(35,51)	-29.64 ^c
Alaska	8.51	0.82	25(04,49)	-6.14	Nebraska	7.08	0.39	44(29,50)	-21.92 ^c
Arizona	8.41	0.22	30(17,37)	-7.31	Nevada	7.94	0.35	34(18,46)	-12.42
Arkansas	9.79	0.36	11(05,23)	7.93	New Hampshire	7.74	0.47	37(17,48)	-14.66
California	10.87	0.11	05(04,08)	19.83 ^c	New Jersey	8.49	0.20	28(17,36)	-6.44
Colorado	7.78	0.26	36(25,44)	-14.23	New Mexico	9.94	0.43	10(04,23)	9.57
Connecticut	8.50	0.30	26(14,39)	-6.34	New York	9.09	0.14	18(12,24)	0.18
Delaware	10.00	0.64	09(02,32)	10.20	North Carolina	8.75	0.19	21(14,33)	-3.57
D.C.	14.97	1.08	01(01,02)	65.04 ^c	North Dakota	5.40	0.55	51(42,51)	-40.42 ^c
Florida	8.73	0.12	22(16,30)	-3.73	Ohio	8.20	0.17	32(23,38)	-9.65
Georgia	8.63	0.21	23(15,35)	-4.90	Oklahoma	8.87	0.30	20(10,35)	-2.26
Hawaii	11.31	0.55	04(02,12)	24.73 ^c	Oregon	9.29	0.30	16(08,29)	2.45
Idaho	6.46	0.41	47(36,51)	-28.73 ^c	Pennsylvania	8.49	0.15	27(19,35)	-6.36
Illinois	8.52	0.17	24(18,35)	-6.03	Rhode Island	10.57	0.62	06(02,21)	16.54 ^c
Indiana	7.69	0.22	38(28,45)	-15.22 ^c	South Carolina	8.98	0.28	19(10,33)	-0.97
Iowa	6.88	0.29	46(36,50)	-24.13 ^c	South Dakota	5.96	0.52	50(38,51)	-34.29 ^c
Kansas	7.18	0.31	43(32,48)	-20.80 ^c	Tennessee	9.71	0.25	12(06,20)	7.02
Kentucky	8.41	0.28	31(15,39)	-7.33	Texas	11.54	0.15	03(02,05)	27.24 ^c
Louisiana	11.81	0.33	02(01,06)	30.21 ^c	Utah	5.97	0.34	49(44,51)	-34.18 ^c
Maine	7.49	0.44	39(21,48)	-17.37 ^c	Vermont	7.81	0.68	35(10,50)	-13.89
Maryland	9.40	0.26	13(07,25)	3.59	Virginia	8.43	0.21	29(17,37)	-7.03
Massachusetts	10.07	0.25	08(05,16)	11.05	Washington	9.36	0.24	15(08,24)	3.23
Michigan	8.07	0.18	33(24,40)	-11.06	West Virginia	7.29	0.37	41(27,49)	-19.63 ^c
Minnesota	7.24	0.23	42(34,48)	-20.16 ^c	Wisconsin	7.48	0.22	40(32,47)	-17.54 ^c
Mississippi	10.43	0.39	07(03,16)	15.01 ^c	Wyoming	6.96	0.70	45(16,51)	-23.24 ^c
Missouri	9.09	0.24	17(10,29)	0.19					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.20
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.65	0.02		
High Five States				
D.C.	5.35	0.56	01(01,16)	46.70 ^c
Texas	4.58	0.09	02(01,07)	25.42 ^c
Louisiana	4.57	0.19	03(01,09)	25.10 ^c
California	4.49	0.07	04(01,07)	22.96 ^c
Hawaii	4.39	0.31	05(01,23)	20.29 ^c
Low Five States				
Nebraska	2.91	0.23	47(16,51)	-20.19 ^c
Idaho	2.91	0.25	48(16,51)	-20.37 ^c
Vermont	2.88	0.38	49(07,51)	-21.08 ^c
New Hampshire	2.58	0.25	50(29,51)	-29.41 ^c
North Dakota	2.23	0.31	51(35,51)	-38.83 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.85	0.16	10(05,30)	5.43	Montana	3.08	0.31	41(08,51)	-15.60 ^c
Alaska	3.33	0.51	32(01,51)	-8.86	Nebraska	2.91	0.23	47(16,51)	-20.19 ^c
Arizona	3.85	0.14	09(06,26)	5.60	Nevada	3.54	0.22	21(06,47)	-3.06
Arkansas	3.82	0.20	11(04,36)	4.53	New Hampshire	2.58	0.25	50(29,51)	-29.41 ^c
California	4.49	0.07	04(01,07)	22.96 ^c	New Jersey	3.38	0.11	29(14,43)	-7.47
Colorado	3.06	0.15	42(22,50)	-16.18 ^c	New Mexico	4.06	0.26	08(01,33)	11.37
Connecticut	2.96	0.16	44(23,51)	-18.96 ^c	New York	3.59	0.08	18(10,31)	-1.60
Delaware	3.77	0.36	12(01,49)	3.38	North Carolina	3.21	0.11	39(21,47)	-12.04
D.C.	5.35	0.56	01(01,16)	46.70 ^c	North Dakota	2.23	0.31	51(35,51)	-38.83 ^c
Florida	3.34	0.07	31(19,42)	-8.42	Ohio	3.24	0.09	37(21,45)	-11.24
Georgia	3.41	0.12	27(12,42)	-6.67	Oklahoma	3.55	0.18	20(08,43)	-2.64
Hawaii	4.39	0.31	05(01,23)	20.29 ^c	Oregon	3.68	0.17	16(06,39)	0.84
Idaho	2.91	0.25	48(16,51)	-20.37 ^c	Pennsylvania	3.35	0.09	30(17,42)	-8.31
Illinois	3.69	0.10	14(08,28)	1.23	Rhode Island	3.62	0.33	17(02,49)	-0.69
Indiana	3.46	0.13	25(10,43)	-5.28	South Carolina	3.28	0.15	34(14,48)	-10.17
Iowa	2.93	0.17	46(25,51)	-19.85 ^c	South Dakota	3.32	0.36	33(04,51)	-9.08
Kansas	3.27	0.20	35(11,49)	-10.36	Tennessee	3.69	0.14	15(07,35)	1.05
Kentucky	3.59	0.16	19(07,41)	-1.63	Texas	4.58	0.09	02(01,07)	25.42 ^c
Louisiana	4.57	0.19	03(01,09)	25.10 ^c	Utah	3.22	0.24	38(08,50)	-11.67
Maine	3.19	0.26	40(08,50)	-12.63	Vermont	2.88	0.38	49(07,51)	-21.08 ^c
Maryland	3.71	0.15	13(07,35)	1.54	Virginia	3.24	0.12	36(18,47)	-11.13
Massachusetts	3.53	0.13	22(09,41)	-3.16	Washington	4.07	0.15	07(03,19)	11.42
Michigan	3.42	0.10	26(13,42)	-6.30	West Virginia	2.95	0.22	45(17,51)	-19.27 ^c
Minnesota	3.05	0.14	43(22,50)	-16.31 ^c	Wisconsin	3.40	0.14	28(12,45)	-6.95
Mississippi	4.37	0.22	06(01,16)	19.72 ^c	Wyoming	3.48	0.48	24(01,51)	-4.61
Missouri	3.50	0.14	23(10,42)	-4.10					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 15.24
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	45.97	0.05		
High Five States				
Kentucky	69.96	0.54	01(01,01)	52.19 ^c
Arkansas	61.86	0.60	02(02,04)	34.57 ^c
West Virginia	61.61	0.72	03(02,05)	34.03 ^c
Mississippi	60.00	0.62	04(02,06)	30.51 ^c
Tennessee	59.95	0.41	05(03,05)	30.41 ^c
Low Five States				
California	34.81	0.14	47(47,47)	-24.27 ^c
Colorado	33.11	0.38	48(48,50)	-27.98 ^c
Hawaii	32.60	0.64	49(48,50)	-29.08 ^c
New Mexico	32.12	0.54	50(48,50)	-30.13 ^c
Utah	20.35	0.44	51(51,51)	-55.74 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	57.29	0.46	07(06,10)	24.63 ^c	Montana	40.83	0.83	39(35,44)	-11.18 ^c
Alaska	50.78	1.52	17(11,27)	10.46 ^c	Nebraska	43.70	0.66	34(29,36)	-4.94
Arizona	37.46	0.32	46(44,46)	-18.50 ^c	Nevada	47.77	0.60	21(18,27)	3.92
Arkansas	61.86	0.60	02(02,04)	34.57 ^c	New Hampshire	47.54	0.80	23(17,29)	3.42
California	34.81	0.14	47(47,47)	-24.27 ^c	New Jersey	40.86	0.29	38(37,43)	-11.10 ^c
Colorado	33.11	0.38	48(48,50)	-27.98 ^c	New Mexico	32.12	0.54	50(48,50)	-30.13 ^c
Connecticut	40.63	0.44	41(36,44)	-11.62 ^c	New York	41.20	0.20	37(36,42)	-10.37 ^c
Delaware	52.19	0.99	13(11,18)	13.53 ^c	North Carolina	51.79	0.32	15(12,17)	12.66 ^c
D.C.	40.64	1.20	40(34,45)	-11.58 ^c	North Dakota	40.25	1.01	43(35,45)	-12.44 ^c
Florida	45.25	0.19	30(27,32)	-1.56	Ohio	54.26	0.29	11(11,13)	18.04 ^c
Georgia	48.96	0.33	19(17,22)	6.51 ^c	Oklahoma	57.87	0.53	06(05,08)	25.89 ^c
Hawaii	32.60	0.64	49(48,50)	-29.08 ^c	Oregon	45.72	0.46	28(25,33)	-0.53
Idaho	38.00	0.69	45(42,46)	-17.34 ^c	Pennsylvania	47.49	0.25	24(20,27)	3.31
Illinois	48.47	0.27	20(17,23)	5.43 ^c	Rhode Island	49.00	0.90	18(15,27)	6.60 ^c
Indiana	56.28	0.40	09(07,10)	22.42 ^c	South Carolina	52.01	0.45	14(12,17)	13.15 ^c
Iowa	47.15	0.51	26(20,28)	2.57	South Dakota	43.58	0.96	35(28,39)	-5.19 ^c
Kansas	47.41	0.55	25(18,28)	3.13	Tennessee	59.95	0.41	05(03,05)	30.41 ^c
Kentucky	69.96	0.54	01(01,01)	52.19 ^c	Texas	41.97	0.20	36(35,38)	-8.69 ^c
Louisiana	56.37	0.49	08(06,10)	22.62 ^c	Utah	20.35	0.44	51(51,51)	-55.74 ^c
Maine	53.71	0.79	12(10,15)	16.85 ^c	Vermont	47.55	1.13	22(17,32)	3.45
Maryland	44.55	0.39	32(28,35)	-3.09	Virginia	46.97	0.34	27(21,28)	2.18
Massachusetts	45.31	0.35	29(27,33)	-1.44	Washington	44.01	0.36	33(30,35)	-4.26
Michigan	50.93	0.30	16(14,18)	10.80 ^c	West Virginia	61.61	0.72	03(02,05)	34.03 ^c
Minnesota	40.40	0.38	42(37,44)	-12.11 ^c	Wisconsin	44.92	0.37	31(28,34)	-2.29
Mississippi	60.00	0.62	04(02,06)	30.51 ^c	Wyoming	38.53	1.15	44(37,46)	-16.19 ^c
Missouri	55.99	0.40	10(07,11)	21.80 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 15.25
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	57.80	0.09		
High Five States				
Kentucky	90.49	0.94	01(01,01)	56.55 ^c
Mississippi	85.45	1.14	02(02,03)	47.84 ^c
Arkansas	84.60	1.07	03(02,04)	46.36 ^c
Tennessee	80.67	0.74	04(04,06)	39.57 ^c
Alabama	80.30	0.84	05(04,06)	38.93 ^c
Low Five States				
Hawaii	41.88	1.09	47(44,50)	-27.54 ^c
California	41.88	0.23	48(46,48)	-27.55 ^c
New Mexico	39.04	0.90	49(48,50)	-32.46 ^c
Colorado	38.73	0.62	50(48,50)	-32.99 ^c
Utah	25.82	0.74	51(51,51)	-55.32 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	80.30	0.84	05(04,06)	38.93 ^c	Montana	46.14	1.30	43(40,47)	-20.18 ^c
Alaska	59.21	2.49	23(14,36)	2.43	Nebraska	55.11	1.12	29(23,38)	-4.67
Arizona	44.89	0.53	45(43,47)	-22.34 ^c	Nevada	54.23	0.96	33(25,38)	-6.17 ^c
Arkansas	84.60	1.07	03(02,04)	46.36 ^c	New Hampshire	55.76	1.31	26(22,37)	-3.53
California	41.88	0.23	48(46,48)	-27.55 ^c	New Jersey	50.24	0.50	40(37,41)	-13.08 ^c
Colorado	38.73	0.62	50(48,50)	-32.99 ^c	New Mexico	39.04	0.90	49(48,50)	-7.66 ^c
Connecticut	48.08	0.74	42(40,44)	-16.83 ^c	New York	50.58	0.33	39(37,41)	-12.50 ^c
Delaware	64.74	1.68	16(12,22)	12.01 ^c	North Carolina	70.04	0.57	11(10,13)	21.18 ^c
D.C.	51.39	2.11	38(24,44)	-11.10 ^c	North Dakota	53.37	1.74	35(23,41)	-7.66 ^c
Florida	56.09	0.31	25(24,30)	-2.97	Ohio	68.75	0.49	13(11,14)	18.94 ^c
Georgia	66.18	0.61	14(14,16)	14.49 ^c	Oklahoma	73.94	0.91	08(07,10)	27.93 ^c
Hawaii	41.88	1.09	47(44,50)	-27.54 ^c	Oregon	52.83	0.74	36(28,39)	-8.60 ^c
Idaho	44.89	1.11	46(42,48)	-22.34 ^c	Pennsylvania	60.71	0.42	18(17,23)	5.03 ^c
Illinois	60.22	0.46	21(18,24)	4.18	Rhode Island	60.51	1.53	19(15,26)	4.68
Indiana	73.23	0.70	09(07,10)	26.69 ^c	South Carolina	69.76	0.80	12(10,14)	20.69 ^c
Iowa	60.39	0.87	20(17,25)	4.47	South Dakota	55.17	1.61	28(21,39)	-4.55
Kansas	58.76	0.93	24(18,27)	1.66	Tennessee	80.67	0.74	04(04,06)	39.57 ^c
Kentucky	90.49	0.94	01(01,01)	56.55 ^c	Texas	54.26	0.34	31(27,36)	-6.13 ^c
Louisiana	75.07	0.87	07(06,09)	29.88 ^c	Utah	25.82	0.74	51(51,51)	-55.32 ^c
Maine	65.75	1.32	15(12,18)	13.76 ^c	Vermont	55.59	1.84	27(18,39)	-3.84
Maryland	54.24	0.66	32(26,37)	-6.17 ^c	Virginia	59.59	0.59	22(18,25)	3.09
Massachusetts	53.99	0.59	34(27,37)	-6.60 ^c	Washington	51.89	0.59	37(33,40)	-10.23 ^c
Michigan	62.79	0.51	17(15,19)	8.63 ^c	West Virginia	78.92	1.23	06(04,07)	36.54 ^c
Minnesota	48.47	0.62	41(39,44)	-16.15 ^c	Wisconsin	54.91	0.62	30(24,36)	-5.00 ^c
Mississippi	85.45	1.14	02(02,03)	47.84 ^c	Wyoming	46.11	1.88	44(38,48)	-20.22 ^c
Missouri	70.89	0.69	10(09,13)	22.65 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 15.26
Cancer of the Lung And Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	37.05	0.06		
High Five States				
Kentucky	54.84	0.64	01(01,01)	48.01 ^c
West Virginia	48.41	0.87	02(02,03)	30.67 ^c
Oklahoma	45.60	0.63	03(02,09)	23.08 ^c
Missouri	44.87	0.49	04(03,10)	21.11 ^c
Maine	44.52	0.97	05(02,15)	20.16 ^c
Low Five States				
California	29.49	0.17	47(45,48)	-20.41 ^c
Colorado	28.81	0.47	48(46,49)	-22.25 ^c
New Mexico	26.70	0.66	49(48,50)	-27.92 ^c
Hawaii	25.27	0.77	50(49,50)	-31.79 ^c
Utah	15.83	0.53	51(51,51)	-57.27 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	40.26	0.51	20(14,24)	8.67	Montana	36.56	1.07	32(21,41)	-1.33
Alaska	44.20	1.93	08(02,24)	19.31 ^c	Nebraska	35.00	0.80	36(30,43)	-5.54
Arizona	31.40	0.40	45(42,46)	-15.26 ^c	Nevada	42.45	0.78	12(05,20)	14.57 ^c
Arkansas	44.29	0.69	07(03,14)	19.56 ^c	New Hampshire	41.75	1.02	15(05,24)	12.68 ^c
California	29.49	0.17	47(45,48)	-20.41 ^c	New Jersey	34.32	0.35	40(35,42)	-7.37
Colorado	28.81	0.47	48(46,49)	-22.25 ^c	New Mexico	26.70	0.66	49(48,50)	-27.92 ^c
Connecticut	35.32	0.55	35(31,41)	-4.67	New York	34.60	0.24	38(35,42)	-6.61
Delaware	42.86	1.21	11(03,22)	15.69 ^c	North Carolina	38.38	0.36	25(21,31)	3.59
D.C.	33.17	1.42	41(29,48)	-10.46 ^c	North Dakota	30.51	1.21	46(40,49)	-17.65 ^c
Florida	36.49	0.23	34(29,36)	-1.51	Ohio	43.54	0.34	10(05,14)	17.52 ^c
Georgia	36.52	0.38	33(28,36)	-1.43	Oklahoma	45.60	0.63	03(02,09)	23.08 ^c
Hawaii	25.27	0.77	50(49,50)	-31.79 ^c	Oregon	40.33	0.59	19(13,24)	8.84
Idaho	32.25	0.87	44(37,46)	-12.94 ^c	Pennsylvania	37.93	0.29	27(23,32)	2.37
Illinois	40.02	0.33	21(16,23)	8.02	Rhode Island	41.34	1.10	17(06,25)	11.57 ^c
Indiana	43.55	0.48	09(04,14)	17.54 ^c	South Carolina	38.72	0.52	24(19,31)	4.52
Iowa	37.13	0.62	31(23,36)	0.22	South Dakota	34.83	1.17	37(25,44)	-6.00
Kansas	38.77	0.68	23(17,32)	4.65	Tennessee	44.49	0.47	06(03,11)	20.08 ^c
Kentucky	54.84	0.64	01(01,01)	48.01 ^c	Texas	32.62	0.23	42(40,45)	-11.94 ^c
Louisiana	42.20	0.57	13(08,19)	13.91 ^c	Utah	15.83	0.53	51(51,51)	-57.27 ^c
Maine	44.52	0.97	05(02,15)	20.16 ^c	Vermont	41.50	1.43	16(03,30)	12.02 ^c
Maryland	37.47	0.47	30(23,34)	1.14	Virginia	37.48	0.40	29(23,34)	1.17
Massachusetts	39.31	0.43	22(17,27)	6.12	Washington	37.99	0.45	26(22,33)	2.53
Michigan	42.08	0.37	14(10,18)	13.59 ^c	West Virginia	48.41	0.87	02(02,03)	30.67 ^c
Minnesota	34.53	0.47	39(34,42)	-6.80	Wisconsin	37.52	0.46	28(23,34)	1.28
Mississippi	41.13	0.68	18(10,22)	11.01 ^c	Wyoming	32.32	1.45	43(32,48)	-12.76 ^c
Missouri	44.87	0.49	04(03,10)	21.11 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 16.18
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.71	0.01		
High Five States				
Utah	3.73	0.18	01(01,07)	37.55 ^c
Idaho	3.57	0.21	02(01,21)	31.53 ^c
West Virginia	3.38	0.18	03(01,23)	24.66 ^c
Oregon	3.33	0.12	04(01,19)	22.72 ^c
Kentucky	3.22	0.12	05(01,24)	18.64 ^c
Low Five States				
New York	2.24	0.05	47(41,49)	-17.43 ^c
South Dakota	2.22	0.22	48(19,50)	-18.13 ^c
Alaska	2.14	0.30	49(10,51)	-21.04 ^c
Hawaii	1.58	0.14	50(48,51)	-41.55 ^c
D.C.	1.13	0.19	51(49,51)	-58.26 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	2.86	0.10	25(07,36)	5.47	Montana	3.04	0.23	14(01,42)	11.98
Alaska	2.14	0.30	49(10,51)	-21.04 ^c	Nebraska	3.06	0.18	12(02,36)	12.68
Arizona	3.02	0.09	15(04,30)	11.51	Nevada	2.90	0.15	22(04,40)	6.86
Arkansas	2.84	0.13	28(05,40)	4.67	New Hampshire	2.75	0.19	29(04,48)	1.46
California	2.50	0.04	38(31,43)	-7.66	New Jersey	2.40	0.07	42(32,48)	-11.62
Colorado	3.21	0.12	06(02,25)	18.21 ^c	New Mexico	2.68	0.16	31(09,46)	-1.00
Connecticut	2.44	0.11	41(27,49)	-10.05	New York	2.24	0.05	47(41,49)	-17.43 ^c
Delaware	2.88	0.24	24(02,47)	6.18	North Carolina	2.84	0.07	27(12,34)	4.88
D.C.	1.13	0.19	51(49,51)	-58.26 ^c	North Dakota	2.31	0.24	44(10,50)	-14.93
Florida	2.96	0.05	18(08,28)	9.20	Ohio	2.97	0.07	17(06,29)	9.66
Georgia	2.48	0.08	39(29,47)	-8.46	Oklahoma	3.08	0.12	10(02,31)	13.55
Hawaii	1.58	0.14	50(48,51)	-41.55 ^c	Oregon	3.33	0.12	04(01,19)	22.72 ^c
Idaho	3.57	0.21	02(01,21)	31.53 ^c	Pennsylvania	2.90	0.06	21(10,32)	6.88
Illinois	2.53	0.06	36(29,44)	-6.69	Rhode Island	2.53	0.20	37(07,49)	-6.77
Indiana	3.14	0.10	07(02,25)	15.62 ^c	South Carolina	2.73	0.10	30(14,42)	0.52
Iowa	2.92	0.13	20(04,38)	7.83	South Dakota	2.22	0.22	48(19,50)	-18.13 ^c
Kansas	3.06	0.14	11(02,33)	12.69	Tennessee	3.13	0.10	08(02,26)	15.39 ^c
Kentucky	3.22	0.12	05(01,24)	18.64 ^c	Texas	2.37	0.05	43(36,48)	-12.74
Louisiana	2.27	0.10	45(34,49)	-16.16 ^c	Utah	3.73	0.18	01(01,07)	37.55 ^c
Maine	3.04	0.19	13(02,40)	12.07	Vermont	2.64	0.27	34(02,49)	-2.66
Maryland	2.64	0.09	33(19,44)	-2.64	Virginia	2.85	0.08	26(10,35)	5.02
Massachusetts	2.93	0.09	19(07,33)	8.12	Washington	2.89	0.09	23(08,34)	6.46
Michigan	2.45	0.07	40(31,47)	-9.53	West Virginia	3.38	0.18	03(01,23)	24.66 ^c
Minnesota	2.57	0.10	35(23,45)	-5.08	Wisconsin	2.67	0.09	32(18,42)	-1.56
Mississippi	2.25	0.12	46(33,49)	-17.09 ^c	Wyoming	3.00	0.32	16(01,49)	10.58
Missouri	3.10	0.10	09(03,27)	14.19					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 16.19
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.06	0.02		
High Five States				
Utah	5.90	0.35	01(01,05)	45.59 ^c
West Virginia	5.15	0.32	02(01,25)	26.95 ^c
Tennessee	5.04	0.19	03(01,14)	24.26 ^c
Idaho	4.94	0.37	04(01,33)	21.92 ^c
Missouri	4.77	0.18	05(02,24)	17.57 ^c
Low Five States				
New York	3.25	0.08	47(41,49)	-19.84 ^c
South Dakota	2.78	0.36	48(31,50)	-31.50 ^c
Alaska	2.67	0.52	49(14,51)	-34.13 ^c
Hawaii	2.37	0.26	50(47,51)	-41.49 ^c
D.C.	1.18	0.30	51(50,51)	-70.93 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.42	0.20	14(03,35)	9.02	Montana	3.94	0.38	34(03,48)	-2.73
Alaska	2.67	0.52	49(14,51)	-34.13 ^c	Nebraska	4.11	0.31	28(03,46)	1.37
Arizona	4.42	0.17	15(04,33)	9.01	Nevada	4.36	0.27	19(02,42)	7.58
Arkansas	4.59	0.25	10(02,35)	13.16	New Hampshire	4.07	0.36	29(02,48)	0.43
California	3.78	0.07	36(29,43)	-6.74	New Jersey	3.56	0.13	43(30,48)	-12.11
Colorado	4.52	0.21	12(02,33)	11.52	New Mexico	3.95	0.29	32(05,48)	-2.52
Connecticut	3.35	0.19	46(31,49)	-17.43 ^c	New York	3.25	0.08	47(41,49)	-19.84 ^c
Delaware	3.95	0.42	33(02,49)	-2.60	North Carolina	4.40	0.15	17(05,32)	8.55
D.C.	1.18	0.30	51(50,51)	-70.93 ^c	North Dakota	3.51	0.45	45(05,50)	-13.51
Florida	4.41	0.09	16(07,28)	8.81	Ohio	4.40	0.13	18(06,31)	8.43
Georgia	3.96	0.15	31(16,43)	-2.29	Oklahoma	4.57	0.23	11(02,33)	12.78
Hawaii	2.37	0.26	50(47,51)	-41.49 ^c	Oregon	4.63	0.22	07(02,32)	14.09
Idaho	4.94	0.37	04(01,33)	21.92 ^c	Pennsylvania	4.27	0.11	23(09,33)	5.17
Illinois	3.74	0.11	38(28,45)	-7.90	Rhode Island	3.74	0.37	37(05,49)	-7.67
Indiana	4.51	0.17	13(03,32)	11.26	South Carolina	4.18	0.20	25(06,41)	3.02
Iowa	4.28	0.23	21(03,40)	5.54	South Dakota	2.78	0.36	48(31,50)	-31.50 ^c
Kansas	4.67	0.26	06(02,34)	15.24 ^c	Tennessee	5.04	0.19	03(01,14)	24.26 ^c
Kentucky	4.61	0.22	08(02,32)	13.62	Texas	3.62	0.09	40(32,46)	-10.71
Louisiana	3.52	0.19	44(27,48)	-13.24	Utah	5.90	0.35	01(01,05)	45.59 ^c
Maine	4.60	0.36	09(01,41)	13.39	Vermont	3.71	0.49	39(02,50)	-8.60
Maryland	4.17	0.19	26(06,40)	2.86	Virginia	4.31	0.16	20(05,35)	6.36
Massachusetts	4.28	0.16	22(06,37)	5.44	Washington	4.19	0.16	24(07,37)	3.35
Michigan	3.60	0.12	41(30,47)	-11.13	West Virginia	5.15	0.32	02(01,25)	26.95 ^c
Minnesota	3.84	0.17	35(17,46)	-5.44	Wisconsin	4.12	0.17	27(07,40)	1.60
Mississippi	3.56	0.24	42(22,49)	-12.10	Wyoming	3.97	0.55	30(01,49)	-2.00
Missouri	4.77	0.18	05(02,24)	17.57 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 16.20
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	1.69	0.01		
High Five States				
Idaho	2.35	0.24	01(01,33)	39.00 ^c
Oregon	2.29	0.14	02(01,20)	35.35 ^c
Nebraska	2.24	0.21	03(01,33)	32.39 ^c
Colorado	2.20	0.13	04(01,22)	29.85 ^c
Montana	2.19	0.28	05(01,46)	29.67 ^c
Low Five States				
Louisiana	1.38	0.10	47(28,51)	-18.23 ^c
Mississippi	1.35	0.13	48(26,51)	-20.07 ^c
North Dakota	1.32	0.25	49(06,51)	-21.93 ^c
D.C.	1.12	0.26	50(12,51)	-34.08 ^c
Hawaii	0.95	0.15	51(44,51)	-44.06 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	1.72	0.11	31(08,46)	1.68	Montana	2.19	0.28	05(01,46)	29.67 ^c
Alaska	1.76	0.38	27(01,51)	3.85	Nebraska	2.24	0.21	03(01,33)	32.39 ^c
Arizona	1.87	0.10	17(05,38)	10.50	Nevada	1.62	0.15	36(07,50)	-4.16
Arkansas	1.49	0.13	44(15,50)	-11.79	New Hampshire	1.78	0.21	25(01,50)	5.36
California	1.51	0.04	42(31,48)	-10.79	New Jersey	1.54	0.07	40(23,49)	-8.75
Colorado	2.20	0.13	04(01,22)	29.85 ^c	New Mexico	1.71	0.17	32(04,50)	0.81
Connecticut	1.80	0.12	24(05,46)	6.30	New York	1.51	0.05	43(30,48)	-10.85
Delaware	2.11	0.28	08(01,48)	24.49 ^c	North Carolina	1.75	0.08	28(11,41)	3.24
D.C.	1.12	0.26	50(12,51)	-34.08 ^c	North Dakota	1.32	0.25	49(06,51)	-21.93 ^c
Florida	1.80	0.05	22(11,34)	6.65	Ohio	1.91	0.07	13(06,31)	12.99
Georgia	1.42	0.08	45(31,50)	-16.06 ^c	Oklahoma	1.89	0.13	15(03,42)	11.79
Hawaii	0.95	0.15	51(44,51)	-44.06 ^c	Oregon	2.29	0.14	02(01,20)	35.35 ^c
Idaho	2.35	0.24	01(01,33)	39.00 ^c	Pennsylvania	1.88	0.07	16(07,32)	10.91
Illinois	1.65	0.07	35(17,45)	-2.44	Rhode Island	1.70	0.22	33(01,50)	0.62
Indiana	2.10	0.11	09(01,24)	24.38 ^c	South Carolina	1.67	0.11	34(10,48)	-1.33
Iowa	1.87	0.15	18(02,45)	10.46	South Dakota	1.76	0.28	26(01,50)	4.02
Kansas	1.80	0.15	23(04,47)	6.64	Tennessee	1.73	0.10	29(10,45)	2.53
Kentucky	2.16	0.13	06(01,26)	27.73 ^c	Texas	1.40	0.05	46(37,50)	-17.16 ^c
Louisiana	1.38	0.10	47(28,51)	-18.23 ^c	Utah	1.93	0.18	12(01,45)	14.28
Maine	1.89	0.21	14(01,48)	11.94	Vermont	1.85	0.29	19(01,50)	9.42
Maryland	1.59	0.10	37(16,49)	-5.78	Virginia	1.73	0.09	30(10,44)	2.13
Massachusetts	1.95	0.10	11(03,32)	15.36 ^c	Washington	1.83	0.10	20(06,39)	8.47
Michigan	1.56	0.07	39(23,48)	-7.73	West Virginia	1.99	0.19	10(01,43)	17.39 ^c
Minnesota	1.59	0.10	38(15,49)	-6.12	Wisconsin	1.54	0.10	41(20,49)	-8.89
Mississippi	1.35	0.13	48(26,51)	-20.07 ^c	Wyoming	2.13	0.38	07(01,50)	26.21 ^c
Missouri	1.82	0.10	21(06,41)	7.31					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.18
Myeloma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.34	0.01		
High Five States				
D.C.	4.20	0.38	01(01,42)	25.72 ^c
Alabama	3.91	0.12	02(01,19)	17.00 ^c
South Carolina	3.86	0.12	03(01,23)	15.69 ^c
Mississippi	3.82	0.16	04(01,30)	14.43
Louisiana	3.82	0.13	05(01,27)	14.24
Low Five States				
Arizona	2.90	0.09	47(36,51)	-13.07
Nevada	2.83	0.15	48(32,51)	-15.35 ^c
Rhode Island	2.81	0.21	49(21,51)	-15.75 ^c
New Mexico	2.75	0.16	50(34,51)	-17.60 ^c
Hawaii	2.45	0.17	51(45,51)	-26.69 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.91	0.12	02(01,19)	17.00 ^c	Montana	3.50	0.24	22(01,48)	4.70
Alaska	3.49	0.41	23(01,51)	4.51	Nebraska	3.29	0.18	36(04,50)	-1.58
Arizona	2.90	0.09	47(36,51)	-13.07	Nevada	2.83	0.15	48(32,51)	-15.35 ^c
Arkansas	3.41	0.14	29(04,45)	2.01	New Hampshire	3.30	0.21	34(02,50)	-1.32
California	3.12	0.04	40(33,47)	-6.72	New Jersey	3.17	0.08	39(26,47)	-5.22
Colorado	3.08	0.12	42(25,50)	-7.80	New Mexico	2.75	0.16	50(34,51)	-17.60 ^c
Connecticut	3.00	0.12	46(28,50)	-10.08	New York	3.06	0.05	43(34,48)	-8.26
Delaware	3.51	0.26	20(01,49)	5.07	North Carolina	3.66	0.09	11(03,29)	9.58
D.C.	4.20	0.38	01(01,42)	25.72 ^c	North Dakota	3.60	0.30	15(01,49)	7.70
Florida	3.02	0.05	45(37,49)	-9.60	Ohio	3.59	0.07	17(05,30)	7.39
Georgia	3.69	0.09	08(02,29)	10.46	Oklahoma	3.43	0.13	28(05,44)	2.66
Hawaii	2.45	0.17	51(45,51)	-26.69 ^c	Oregon	3.46	0.13	26(04,42)	3.53
Idaho	3.39	0.21	30(02,49)	1.56	Pennsylvania	3.29	0.06	35(21,42)	-1.39
Illinois	3.37	0.07	33(15,39)	0.95	Rhode Island	2.81	0.21	49(21,51)	-15.75 ^c
Indiana	3.65	0.10	12(02,31)	9.39	South Carolina	3.86	0.12	03(01,23)	15.69 ^c
Iowa	3.38	0.14	31(06,45)	1.34	South Dakota	3.59	0.27	16(01,48)	7.55
Kansas	3.70	0.15	07(01,37)	10.81	Tennessee	3.69	0.10	10(02,29)	10.35
Kentucky	3.46	0.12	25(05,41)	3.58	Texas	3.20	0.05	38(27,44)	-4.14
Louisiana	3.82	0.13	05(01,27)	14.24	Utah	3.48	0.18	24(01,46)	4.24
Maine	3.09	0.19	41(11,51)	-7.54	Vermont	3.02	0.28	44(03,51)	-9.47
Maryland	3.72	0.11	06(01,29)	11.26	Virginia	3.69	0.10	09(02,28)	10.45
Massachusetts	3.20	0.09	37(21,47)	-4.09	Washington	3.38	0.10	32(10,42)	1.14
Michigan	3.61	0.08	14(04,30)	8.00	West Virginia	3.53	0.17	19(01,44)	5.57
Minnesota	3.44	0.11	27(06,41)	2.87	Wisconsin	3.58	0.11	18(03,34)	7.32
Mississippi	3.82	0.16	04(01,30)	14.43	Wyoming	3.65	0.36	13(01,50)	9.25
Missouri	3.50	0.10	21(05,37)	4.92					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.19
Myeloma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.22	0.02		
High Five States				
D.C.	5.01	0.67	01(01,51)	18.90 ^c
Alabama	4.95	0.22	02(01,27)	17.46 ^c
Louisiana	4.95	0.23	03(01,30)	17.32 ^c
South Carolina	4.83	0.22	04(01,32)	14.63
Indiana	4.83	0.18	05(01,28)	14.49
Low Five States				
Rhode Island	3.65	0.38	47(08,51)	-13.42
Arizona	3.61	0.15	48(35,51)	-14.36
New Mexico	3.47	0.27	49(26,51)	-17.78 ^c
Nevada	3.44	0.25	50(28,51)	-18.49 ^c
Hawaii	3.34	0.30	51(25,51)	-20.81 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.95	0.22	02(01,27)	17.46 ^c	Montana	4.28	0.40	29(01,51)	1.58
Alaska	4.51	0.72	16(01,51)	7.01	Nebraska	4.23	0.31	33(02,50)	0.36
Arizona	3.61	0.15	48(35,51)	-14.36	Nevada	3.44	0.25	50(28,51)	-18.49 ^c
Arkansas	4.33	0.25	26(02,47)	2.63	New Hampshire	4.29	0.37	28(01,50)	1.89
California	4.00	0.07	40(27,45)	-5.16	New Jersey	4.09	0.14	37(17,47)	-2.97
Colorado	3.99	0.20	41(12,50)	-5.25	New Mexico	3.47	0.27	49(26,51)	-17.78 ^c
Connecticut	3.71	0.21	46(23,51)	-11.92	New York	3.79	0.09	43(34,49)	-10.14
Delaware	4.46	0.44	23(01,51)	5.73	North Carolina	4.43	0.15	24(05,40)	5.00
D.C.	5.01	0.67	01(01,51)	18.90 ^c	North Dakota	4.75	0.53	08(01,50)	12.68
Florida	3.78	0.08	44(34,49)	-10.25	Ohio	4.48	0.13	20(06,35)	6.21
Georgia	4.53	0.16	15(03,37)	7.49	Oklahoma	4.50	0.23	17(01,43)	6.68
Hawaii	3.34	0.30	51(25,51)	-20.81 ^c	Oregon	4.49	0.22	18(02,41)	6.61
Idaho	4.30	0.35	27(01,50)	2.12	Pennsylvania	4.10	0.11	36(21,45)	-2.69
Illinois	4.27	0.13	30(12,42)	1.27	Rhode Island	3.65	0.38	47(08,51)	-13.42
Indiana	4.83	0.18	05(01,28)	14.49	South Carolina	4.83	0.22	04(01,32)	14.63
Iowa	4.46	0.24	22(02,45)	5.78	South Dakota	4.22	0.45	35(01,51)	0.22
Kansas	4.81	0.27	06(01,36)	14.00	Tennessee	4.75	0.18	07(01,32)	12.79
Kentucky	4.48	0.22	19(02,43)	6.27	Texas	4.08	0.09	38(23,44)	-3.29
Louisiana	4.95	0.23	03(01,30)	17.32 ^c	Utah	4.35	0.31	25(01,49)	3.15
Maine	3.73	0.32	45(11,51)	-11.60	Vermont	4.01	0.49	39(01,51)	-4.95
Maryland	4.23	0.19	32(07,47)	0.38	Virginia	4.69	0.17	11(01,31)	11.37
Massachusetts	3.90	0.16	42(23,50)	-7.48	Washington	4.23	0.17	34(08,45)	0.33
Michigan	4.59	0.14	14(03,33)	8.96	West Virginia	4.24	0.29	31(02,50)	0.53
Minnesota	4.61	0.19	13(01,37)	9.27	Wisconsin	4.74	0.19	09(01,32)	12.43
Mississippi	4.71	0.28	10(01,41)	11.65	Wyoming	4.66	0.62	12(01,51)	10.43
Missouri	4.47	0.18	21(03,39)	6.05					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 18.20
Myeloma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.69	0.02		
High Five States				
D.C.	3.52	0.45	01(01,47)	30.56 ^c
Maryland	3.37	0.14	02(01,14)	25.07 ^c
Mississippi	3.24	0.19	03(01,30)	20.28 ^c
South Carolina	3.19	0.15	04(01,24)	18.47 ^c
Alabama	3.18	0.14	05(01,23)	18.02 ^c
Low Five States				
Nevada	2.29	0.18	47(17,51)	-15.16 ^c
Rhode Island	2.27	0.25	48(10,51)	-15.80 ^c
Vermont	2.24	0.32	49(03,51)	-16.80 ^c
New Mexico	2.17	0.19	50(22,51)	-19.41 ^c
Hawaii	1.78	0.20	51(41,51)	-34.04 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.18	0.14	05(01,23)	18.02 ^c	Montana	2.81	0.29	19(01,50)	4.16
Alaska	2.69	0.48	30(01,51)	-0.22	Nebraska	2.56	0.21	36(05,50)	-4.89
Arizona	2.30	0.11	46(30,51)	-14.63	Nevada	2.29	0.18	47(17,51)	-15.16 ^c
Arkansas	2.72	0.17	24(04,48)	1.07	New Hampshire	2.52	0.25	42(04,51)	-6.58
California	2.46	0.05	43(30,48)	-8.74	New Jersey	2.52	0.09	39(19,48)	-6.31
Colorado	2.41	0.14	45(19,50)	-10.50	New Mexico	2.17	0.19	50(22,51)	-19.41 ^c
Connecticut	2.52	0.14	41(13,50)	-6.49	New York	2.56	0.06	37(22,46)	-4.97
Delaware	2.84	0.31	17(01,50)	5.22	North Carolina	3.13	0.10	06(01,21)	16.16 ^c
D.C.	3.52	0.45	01(01,47)	30.56 ^c	North Dakota	2.78	0.36	20(01,51)	3.08
Florida	2.42	0.06	44(30,49)	-10.37	Ohio	2.98	0.09	11(03,26)	10.50
Georgia	3.08	0.11	07(01,25)	14.18	Oklahoma	2.58	0.15	35(10,50)	-4.37
Hawaii	1.78	0.20	51(41,51)	-34.04 ^c	Oregon	2.62	0.15	31(08,49)	-2.86
Idaho	2.61	0.25	32(02,51)	-3.00	Pennsylvania	2.72	0.08	26(12,40)	0.93
Illinois	2.72	0.08	25(11,41)	0.99	Rhode Island	2.27	0.25	48(10,51)	-15.80 ^c
Indiana	2.83	0.12	18(05,40)	4.87	South Carolina	3.19	0.15	04(01,24)	18.47 ^c
Iowa	2.59	0.16	33(08,49)	-3.95	South Dakota	3.06	0.33	08(01,49)	13.54
Kansas	2.90	0.18	15(01,45)	7.44	Tennessee	2.94	0.12	13(03,33)	9.20
Kentucky	2.72	0.14	27(05,47)	0.80	Texas	2.54	0.06	38(23,46)	-5.72
Louisiana	3.04	0.15	09(01,35)	13.01	Utah	2.78	0.22	21(01,49)	2.99
Maine	2.59	0.23	34(04,50)	-4.05	Vermont	2.24	0.32	49(03,51)	-16.80 ^c
Maryland	3.37	0.14	02(01,14)	25.07 ^c	Virginia	2.99	0.11	10(02,31)	10.88
Massachusetts	2.70	0.11	29(10,45)	0.39	Washington	2.71	0.12	28(08,45)	0.72
Michigan	2.87	0.10	16(06,34)	6.66	West Virginia	2.92	0.21	14(01,46)	8.49
Minnesota	2.52	0.13	40(15,49)	-6.39	Wisconsin	2.74	0.12	23(06,44)	1.82
Mississippi	3.24	0.19	03(01,30)	20.28 ^c	Wyoming	2.96	0.44	12(01,51)	9.89
Missouri	2.76	0.12	22(07,44)	2.40					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 19.22
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	6.01	0.02		
High Five States				
Minnesota	7.00	0.16	01(01,11)	16.48 ^c
Kentucky	6.89	0.17	02(01,17)	14.76
Michigan	6.86	0.11	03(01,11)	14.29
Iowa	6.73	0.19	04(01,22)	12.11
Oklahoma	6.69	0.18	05(01,22)	11.46
Low Five States				
Hawaii	5.40	0.26	47(19,51)	-10.10
Maryland	5.36	0.13	48(30,51)	-10.83
Colorado	5.28	0.15	49(32,51)	-12.17
New Mexico	4.95	0.21	50(37,51)	-17.53 ^c
D.C.	4.88	0.42	51(19,51)	-18.77 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	6.14	0.15	22(07,35)	2.24	Montana	5.77	0.31	28(05,51)	-3.91
Alaska	5.86	0.53	26(01,51)	-2.49	Nebraska	6.18	0.25	21(03,44)	2.94
Arizona	5.56	0.13	38(25,49)	-7.44	Nevada	5.49	0.21	43(21,51)	-8.55
Arkansas	6.59	0.20	08(01,25)	9.66	New Hampshire	5.43	0.27	46(19,51)	-9.57
California	5.69	0.06	33(26,43)	-5.26	New Jersey	5.61	0.11	37(25,48)	-6.55
Colorado	5.28	0.15	49(32,51)	-12.17	New Mexico	4.95	0.21	50(37,51)	-17.53 ^c
Connecticut	5.56	0.16	39(22,50)	-7.51	New York	5.72	0.07	31(24,43)	-4.79
Delaware	6.10	0.34	23(01,49)	1.49	North Carolina	5.76	0.11	29(22,44)	-4.06
D.C.	4.88	0.42	51(19,51)	-18.77 ^c	North Dakota	5.46	0.37	44(09,51)	-9.17
Florida	5.75	0.07	30(23,42)	-4.22	Ohio	6.58	0.10	09(02,19)	9.56
Georgia	5.44	0.11	45(29,50)	-9.40	Oklahoma	6.69	0.18	05(01,22)	11.46
Hawaii	5.40	0.26	47(19,51)	-10.10	Oregon	6.34	0.17	16(03,29)	5.60
Idaho	6.32	0.28	18(01,43)	5.26	Pennsylvania	6.49	0.09	11(04,21)	8.04
Illinois	6.21	0.10	20(11,28)	3.32	Rhode Island	5.52	0.30	42(13,51)	-8.06
Indiana	6.59	0.14	07(01,22)	9.68	South Carolina	5.68	0.15	36(20,48)	-5.37
Iowa	6.73	0.19	04(01,22)	12.11	South Dakota	5.69	0.34	35(04,51)	-5.34
Kansas	6.34	0.20	17(02,32)	5.47	Tennessee	6.46	0.14	14(02,24)	7.54
Kentucky	6.89	0.17	02(01,17)	14.76	Texas	5.85	0.07	27(22,39)	-2.65
Louisiana	6.56	0.17	10(01,24)	9.29	Utah	5.69	0.24	34(15,50)	-5.30
Maine	6.46	0.28	13(01,38)	7.55	Vermont	6.32	0.42	19(01,49)	5.21
Maryland	5.36	0.13	48(30,51)	-10.83	Virginia	5.92	0.12	25(17,40)	-1.52
Massachusetts	5.54	0.12	40(25,49)	-7.72	Washington	6.35	0.14	15(04,26)	5.65
Michigan	6.86	0.11	03(01,11)	14.29	West Virginia	6.59	0.24	06(01,27)	9.76
Minnesota	7.00	0.16	01(01,11)	16.48 ^c	Wisconsin	6.48	0.14	12(02,24)	7.93
Mississippi	5.70	0.19	32(18,49)	-5.13	Wyoming	5.54	0.44	41(03,51)	-7.72
Missouri	6.01	0.13	24(14,39)	0.08					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 19.23
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	7.75	0.03		
High Five States				
Michigan	9.05	0.20	01(01,12)	16.76 ^c
Minnesota	9.01	0.27	02(01,15)	16.32 ^c
Kentucky	8.94	0.31	03(01,19)	15.40 ^c
Iowa	8.78	0.33	04(01,22)	13.24
Louisiana	8.75	0.31	05(01,21)	12.87
Low Five States				
Wyoming	6.53	0.74	47(02,51)	-15.69 ^c
Rhode Island	6.44	0.50	48(17,51)	-16.85 ^c
North Dakota	6.32	0.60	49(15,51)	-18.39 ^c
New Mexico	6.13	0.35	50(35,51)	-20.95 ^c
D.C.	5.95	0.72	51(14,51)	-23.28 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	7.71	0.27	20(08,43)	-0.50	Montana	7.62	0.54	24(01,50)	-1.72
Alaska	7.18	0.86	37(01,51)	-7.29	Nebraska	7.53	0.42	27(04,49)	-2.78
Arizona	7.06	0.21	40(22,49)	-8.95	Nevada	6.97	0.35	44(17,50)	-10.09
Arkansas	8.23	0.34	15(01,39)	6.21	New Hampshire	6.90	0.47	45(12,51)	-10.96
California	7.27	0.10	35(25,44)	-6.16	New Jersey	7.33	0.19	33(19,46)	-5.38
Colorado	6.74	0.26	46(25,51)	-12.97	New Mexico	6.13	0.35	50(35,51)	-20.95 ^c
Connecticut	7.22	0.29	36(17,49)	-6.86	New York	7.38	0.13	32(21,43)	-4.73
Delaware	7.50	0.58	28(01,50)	-3.21	North Carolina	7.46	0.19	30(17,44)	-3.78
D.C.	5.95	0.72	51(14,51)	-23.28 ^c	North Dakota	6.32	0.60	49(15,51)	-18.39 ^c
Florida	7.41	0.11	31(21,42)	-4.35	Ohio	8.55	0.18	07(02,19)	10.27
Georgia	7.32	0.21	34(18,47)	-5.49	Oklahoma	8.63	0.32	06(01,25)	11.38
Hawaii	7.10	0.45	39(07,51)	-8.37	Oregon	8.34	0.30	12(01,29)	7.68
Idaho	7.67	0.47	23(02,49)	-0.96	Pennsylvania	8.45	0.16	10(03,20)	8.98
Illinois	8.04	0.17	19(08,29)	3.71	Rhode Island	6.44	0.50	48(17,51)	-16.85 ^c
Indiana	8.48	0.24	09(01,24)	9.47	South Carolina	7.58	0.27	25(11,45)	-2.20
Iowa	8.78	0.33	04(01,22)	13.24	South Dakota	7.04	0.57	41(06,51)	-9.09
Kansas	8.26	0.35	13(01,38)	6.56	Tennessee	8.53	0.25	08(01,23)	10.12
Kentucky	8.94	0.31	03(01,19)	15.40 ^c	Texas	7.55	0.13	26(18,39)	-2.54
Louisiana	8.75	0.31	05(01,21)	12.87	Utah	7.01	0.39	43(14,51)	-9.49
Maine	8.18	0.48	17(01,44)	5.55	Vermont	8.22	0.73	16(01,49)	6.02
Maryland	7.04	0.24	42(21,49)	-9.14	Virginia	7.69	0.22	22(11,41)	-0.80
Massachusetts	7.14	0.22	38(21,48)	-7.91	Washington	8.08	0.23	18(04,32)	4.27
Michigan	9.05	0.20	01(01,12)	16.76 ^c	West Virginia	8.25	0.41	14(01,40)	6.40
Minnesota	9.01	0.27	02(01,15)	16.32 ^c	Wisconsin	8.37	0.25	11(02,26)	7.97
Mississippi	7.49	0.35	29(08,48)	-3.29	Wyoming	6.53	0.74	47(02,51)	-15.69 ^c
Missouri	7.69	0.23	21(11,43)	-0.78					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 19.24
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.69	0.02		
High Five States				
Kentucky	5.38	0.20	01(01,23)	14.89
Minnesota	5.38	0.18	02(01,23)	14.89
West Virginia	5.35	0.29	03(01,33)	14.19
Arkansas	5.35	0.24	04(01,28)	14.17
Iowa	5.27	0.23	05(01,29)	12.53
Low Five States				
Georgia	4.11	0.13	47(32,51)	-12.22
D.C.	4.08	0.50	48(01,51)	-13.04
Colorado	4.07	0.18	49(28,51)	-13.25
Hawaii	4.04	0.30	50(17,51)	-13.80
New Mexico	3.97	0.26	51(24,51)	-15.26 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.99	0.18	15(02,37)	6.51	Montana	4.26	0.37	44(04,51)	-9.02
Alaska	4.56	0.64	31(01,51)	-2.81	Nebraska	5.11	0.30	11(01,43)	8.99
Arizona	4.34	0.15	41(23,50)	-7.50	Nevada	4.23	0.25	45(15,51)	-9.74
Arkansas	5.35	0.24	04(01,28)	14.17	New Hampshire	4.32	0.32	42(06,51)	-7.72
California	4.48	0.07	35(25,44)	-4.49	New Jersey	4.41	0.12	37(22,49)	-5.90
Colorado	4.07	0.18	49(28,51)	-13.25	New Mexico	3.97	0.26	51(24,51)	-15.26 ^c
Connecticut	4.35	0.19	40(16,51)	-7.22	New York	4.50	0.09	34(23,44)	-3.93
Delaware	4.97	0.42	17(01,51)	6.04	North Carolina	4.53	0.13	33(18,46)	-3.35
D.C.	4.08	0.50	48(01,51)	-13.04	North Dakota	4.78	0.46	24(01,51)	2.00
Florida	4.40	0.08	39(25,47)	-6.12	Ohio	5.13	0.12	10(02,25)	9.49
Georgia	4.11	0.13	47(32,51)	-12.22	Oklahoma	5.17	0.21	08(01,32)	10.37
Hawaii	4.04	0.30	50(17,51)	-13.80	Oregon	4.74	0.20	26(05,45)	1.20
Idaho	5.10	0.34	12(01,47)	8.75	Pennsylvania	5.04	0.11	14(04,27)	7.45
Illinois	4.89	0.11	19(07,32)	4.36	Rhode Island	4.73	0.36	27(01,51)	1.03
Indiana	5.23	0.16	06(01,26)	11.53	South Carolina	4.30	0.17	43(20,51)	-8.27
Iowa	5.27	0.23	05(01,29)	12.53	South Dakota	4.64	0.41	28(01,51)	-0.99
Kansas	4.84	0.23	22(02,46)	3.32	Tennessee	4.87	0.16	21(04,39)	3.91
Kentucky	5.38	0.20	01(01,23)	14.89	Texas	4.55	0.09	32(21,43)	-2.98
Louisiana	4.87	0.19	20(03,42)	3.92	Utah	4.62	0.28	29(03,51)	-1.47
Maine	5.15	0.32	09(01,45)	9.94	Vermont	4.81	0.48	23(01,51)	2.55
Maryland	4.15	0.16	46(27,51)	-11.45	Virginia	4.60	0.14	30(14,46)	-1.81
Massachusetts	4.41	0.14	38(21,49)	-5.92	Washington	4.99	0.16	16(02,33)	6.37
Michigan	5.21	0.13	07(01,24)	11.16	West Virginia	5.35	0.29	03(01,33)	14.19
Minnesota	5.38	0.18	02(01,23)	14.89	Wisconsin	5.06	0.17	13(02,30)	7.99
Mississippi	4.45	0.23	36(11,51)	-5.10	Wyoming	4.94	0.56	18(01,51)	5.30
Missouri	4.77	0.16	25(06,42)	1.80					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 20.24
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.44	0.01		
High Five States				
Mississippi	3.68	0.15	01(01,03)	50.80 ^c
D.C.	3.33	0.33	02(01,31)	36.18 ^c
Louisiana	3.27	0.12	03(01,07)	33.77 ^c
Arkansas	3.08	0.14	04(02,13)	26.19 ^c
Tennessee	3.05	0.09	05(02,11)	24.72 ^c
Low Five States				
Kansas	1.98	0.11	47(31,51)	-18.79 ^c
Colorado	1.96	0.09	48(35,51)	-19.74 ^c
Arizona	1.93	0.07	49(38,51)	-20.99 ^c
Connecticut	1.92	0.09	50(36,51)	-21.44 ^c
Utah	1.60	0.12	51(45,51)	-34.45 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	2.67	0.10	11(05,30)	9.40	Montana	2.07	0.18	45(15,51)	-15.43 ^c
Alaska	2.08	0.28	43(05,51)	-14.80	Nebraska	2.29	0.15	33(10,49)	-6.18
Arizona	1.93	0.07	49(38,51)	-20.99 ^c	Nevada	2.39	0.13	27(09,45)	-2.34
Arkansas	3.08	0.14	04(02,13)	26.19 ^c	New Hampshire	2.12	0.17	41(12,51)	-13.23
California	2.43	0.04	25(17,33)	-0.52	New Jersey	1.99	0.06	46(37,50)	-18.51 ^c
Colorado	1.96	0.09	48(35,51)	-19.74 ^c	New Mexico	2.24	0.14	36(12,50)	-8.47
Connecticut	1.92	0.09	50(36,51)	-21.44 ^c	New York	2.19	0.04	37(30,44)	-10.28
Delaware	2.63	0.22	14(02,46)	7.65	North Carolina	2.57	0.07	18(09,31)	5.11
D.C.	3.33	0.33	02(01,31)	36.18 ^c	North Dakota	2.15	0.24	38(08,51)	-11.93
Florida	2.63	0.05	13(09,23)	7.75	Ohio	2.48	0.06	22(12,34)	1.36
Georgia	2.60	0.08	16(08,29)	6.27	Oklahoma	2.66	0.11	12(05,33)	8.74
Hawaii	2.53	0.18	19(04,44)	3.61	Oregon	2.32	0.10	32(14,44)	-4.87
Idaho	2.43	0.17	26(06,48)	-0.63	Pennsylvania	2.12	0.05	40(33,47)	-13.12
Illinois	2.35	0.06	29(19,40)	-3.90	Rhode Island	2.07	0.18	44(13,51)	-15.18 ^c
Indiana	2.34	0.08	30(16,42)	-4.16	South Carolina	2.96	0.11	06(02,14)	21.30 ^c
Iowa	2.25	0.11	35(16,48)	-7.86	South Dakota	2.12	0.21	39(10,51)	-13.09
Kansas	1.98	0.11	47(31,51)	-18.79 ^c	Tennessee	3.05	0.09	05(02,11)	24.72 ^c
Kentucky	2.93	0.11	07(02,16)	19.82 ^c	Texas	2.59	0.05	17(10,25)	6.12
Louisiana	3.27	0.12	03(01,07)	33.77 ^c	Utah	1.60	0.12	51(45,51)	-34.45 ^c
Maine	2.75	0.18	10(02,37)	12.47	Vermont	2.62	0.26	15(02,49)	7.31
Maryland	2.36	0.09	28(14,43)	-3.31	Virginia	2.33	0.07	31(18,42)	-4.45
Massachusetts	2.47	0.08	23(10,37)	1.15	Washington	2.84	0.09	08(03,18)	16.40 ^c
Michigan	2.48	0.07	20(12,34)	1.63	West Virginia	2.79	0.15	09(02,31)	14.04
Minnesota	2.09	0.08	42(30,50)	-14.56	Wisconsin	2.44	0.09	24(11,39)	-0.09
Mississippi	3.68	0.15	01(01,03)	50.80 ^c	Wyoming	2.28	0.27	34(04,51)	-6.67
Missouri	2.48	0.08	21(10,37)	1.57					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 20.25
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.77	0.02		
High Five States				
Mississippi	6.19	0.30	01(01,03)	64.32 ^c
D.C.	5.72	0.67	02(01,22)	51.70 ^c
Louisiana	5.45	0.23	03(01,06)	44.59 ^c
Tennessee	4.86	0.18	04(02,11)	29.08 ^c
South Carolina	4.78	0.20	05(02,13)	26.82 ^c
Low Five States				
Minnesota	3.04	0.15	47(31,50)	-19.34 ^c
New Jersey	3.03	0.12	48(35,50)	-19.59 ^c
Kansas	2.97	0.20	49(29,50)	-21.24 ^c
Colorado	2.66	0.16	50(41,51)	-29.35 ^c
Utah	2.04	0.20	51(49,51)	-45.90 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.28	0.19	11(05,25)	13.53	Montana	3.09	0.33	45(12,51)	-17.89 ^c
Alaska	3.35	0.54	37(04,51)	-11.20	Nebraska	3.38	0.27	35(11,50)	-10.37
Arizona	3.06	0.14	46(32,50)	-18.70 ^c	Nevada	3.48	0.23	32(11,49)	-7.60
Arkansas	4.67	0.25	06(02,20)	23.91 ^c	New Hampshire	3.30	0.30	38(11,50)	-12.33
California	3.67	0.07	24(19,35)	-2.51	New Jersey	3.03	0.12	48(35,50)	-19.59 ^c
Colorado	2.66	0.16	50(41,51)	-29.35 ^c	New Mexico	3.26	0.25	41(16,50)	-13.44
Connecticut	3.15	0.18	44(24,50)	-16.45 ^c	New York	3.40	0.08	33(25,44)	-9.72
Delaware	4.32	0.43	10(02,43)	14.72	North Carolina	4.04	0.13	16(08,28)	7.16
D.C.	5.72	0.67	02(01,22)	51.70 ^c	North Dakota	3.24	0.43	42(06,51)	-13.95
Florida	4.08	0.08	12(09,23)	8.31	Ohio	3.84	0.11	19(11,34)	1.85
Georgia	4.07	0.14	15(08,29)	7.94	Oklahoma	4.07	0.21	14(06,34)	8.15
Hawaii	3.82	0.32	21(05,48)	1.36	Oregon	3.38	0.18	36(19,48)	-10.38
Idaho	3.64	0.31	27(07,49)	-3.47	Pennsylvania	3.28	0.10	40(28,47)	-12.84
Illinois	3.58	0.11	30(17,41)	-4.86	Rhode Island	3.21	0.35	43(10,50)	-14.83
Indiana	3.61	0.15	28(13,42)	-4.09	South Carolina	4.78	0.20	05(02,13)	26.82 ^c
Iowa	3.54	0.21	31(11,48)	-5.95	South Dakota	3.39	0.40	34(06,50)	-9.96
Kansas	2.97	0.20	49(29,50)	-21.24 ^c	Tennessee	4.86	0.18	04(02,11)	29.08 ^c
Kentucky	4.53	0.21	08(03,20)	20.13 ^c	Texas	4.08	0.09	13(08,23)	8.19
Louisiana	5.45	0.23	03(01,06)	44.59 ^c	Utah	2.04	0.20	51(49,51)	-45.90 ^c
Maine	4.51	0.34	09(02,32)	19.71 ^c	Vermont	3.67	0.46	23(03,50)	-2.49
Maryland	3.59	0.16	29(14,44)	-4.64	Virginia	3.65	0.14	25(13,41)	-3.11
Massachusetts	3.84	0.15	20(10,37)	1.85	Washington	3.99	0.16	17(08,32)	5.94
Michigan	3.89	0.12	18(10,33)	3.26	West Virginia	4.66	0.30	07(02,22)	23.72 ^c
Minnesota	3.04	0.15	47(31,50)	-19.34 ^c	Wisconsin	3.65	0.16	26(13,42)	-3.16
Mississippi	6.19	0.30	01(01,03)	64.32 ^c	Wyoming	3.29	0.48	39(05,51)	-12.55
Missouri	3.77	0.16	22(10,39)	-0.03					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 20.26
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	1.33	0.01		
High Five States				
Washington	1.86	0.10	01(01,07)	39.67 ^c
Arkansas	1.74	0.14	02(01,24)	30.55 ^c
Mississippi	1.71	0.14	03(01,30)	27.97 ^c
Vermont	1.65	0.27	04(01,49)	23.79 ^c
Kentucky	1.60	0.11	05(01,31)	20.03 ^c
Low Five States				
West Virginia	1.11	0.13	46(08,50)	-16.51 ^c
New Hampshire	1.08	0.17	47(06,50)	-18.83 ^c
South Dakota	1.03	0.20	48(04,50)	-23.05 ^c
Arizona	0.95	0.07	49(39,50)	-28.66 ^c
Connecticut	0.94	0.09	50(37,50)	-29.57 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	1.33	0.09	29(04,45)	-0.32	Montana	1.12	0.19	45(03,50)	-15.87 ^c
Alaska	-	-	-(-,-)	-	Nebraska	1.36	0.15	21(02,49)	1.79
Arizona	0.95	0.07	49(39,50)	-28.66 ^c	Nevada	1.39	0.14	15(02,47)	4.21
Arkansas	1.74	0.14	02(01,24)	30.55 ^c	New Hampshire	1.08	0.17	47(06,50)	-18.83 ^c
California	1.39	0.04	17(08,33)	3.98	New Jersey	1.18	0.07	40(18,48)	-11.69
Colorado	1.34	0.10	24(04,46)	0.57	New Mexico	1.35	0.15	22(02,49)	1.25
Connecticut	0.94	0.09	50(37,50)	-29.57 ^c	New York	1.23	0.04	37(21,45)	-7.99
Delaware	1.21	0.20	38(02,50)	-9.15	North Carolina	1.38	0.07	18(05,40)	3.88
D.C.	1.48	0.29	08(01,50)	10.71	North Dakota	1.25	0.25	35(01,50)	-6.20
Florida	1.39	0.05	16(07,35)	4.06	Ohio	1.33	0.06	28(07,42)	-0.17
Georgia	1.39	0.07	14(04,41)	4.35	Oklahoma	1.42	0.11	11(02,44)	6.61
Hawaii	1.40	0.18	12(01,49)	5.14	Oregon	1.42	0.11	10(02,44)	6.81
Idaho	1.31	0.18	30(01,50)	-1.37	Pennsylvania	1.17	0.05	42(25,47)	-12.15
Illinois	1.33	0.06	27(08,42)	-0.01	Rhode Island	1.17	0.19	41(02,50)	-11.94
Indiana	1.27	0.08	34(08,46)	-4.82	South Carolina	1.46	0.10	09(02,41)	9.92
Iowa	1.13	0.10	44(13,50)	-15.24 ^c	South Dakota	1.03	0.20	48(04,50)	-23.05 ^c
Kansas	1.15	0.12	43(09,50)	-13.57	Tennessee	1.55	0.09	06(02,30)	16.26 ^c
Kentucky	1.60	0.11	05(01,31)	20.03 ^c	Texas	1.35	0.05	23(08,38)	1.04
Louisiana	1.48	0.10	07(02,41)	10.72	Utah	1.20	0.15	39(05,50)	-9.57
Maine	1.24	0.16	36(02,50)	-6.67	Vermont	1.65	0.27	04(01,49)	23.79 ^c
Maryland	1.34	0.09	26(06,45)	0.35	Virginia	1.27	0.07	33(09,46)	-4.45
Massachusetts	1.34	0.08	25(06,44)	0.37	Washington	1.86	0.10	01(01,07)	39.67 ^c
Michigan	1.31	0.06	31(08,44)	-1.66	West Virginia	1.11	0.13	46(08,50)	-16.51 ^c
Minnesota	1.28	0.09	32(06,46)	-3.63	Wisconsin	1.39	0.09	13(04,43)	4.59
Mississippi	1.71	0.14	03(01,30)	27.97 ^c	Wyoming	1.36	0.30	20(01,50)	2.07
Missouri	1.37	0.08	19(04,43)	2.50					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 21.14
Cancer of the Ovary (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	7.55	0.03		
High Five States				
Oregon	8.90	0.27	01(01,10)	17.93 ^c
Montana	8.43	0.51	02(01,43)	11.73
Washington	8.40	0.21	03(01,18)	11.25
Oklahoma	8.33	0.27	04(01,29)	10.42
Alabama	8.23	0.23	05(01,27)	9.02
Low Five States				
Louisiana	6.77	0.23	47(29,50)	-10.35
Rhode Island	6.44	0.43	48(18,51)	-14.64
Alaska	6.42	0.70	49(02,51)	-14.97
North Dakota	6.15	0.55	50(12,51)	-18.47 ^c
Hawaii	5.40	0.36	51(47,51)	-28.42 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	8.23	0.23	05(01,27)	9.02	Montana	8.43	0.51	02(01,43)	11.73
Alaska	6.42	0.70	49(02,51)	-14.97	Nebraska	7.11	0.36	40(06,50)	-5.81
Arizona	7.36	0.20	35(12,46)	-2.48	Nevada	7.09	0.32	42(09,50)	-6.02
Arkansas	8.04	0.30	10(01,39)	6.54	New Hampshire	7.62	0.43	23(01,49)	1.04
California	7.48	0.09	31(18,39)	-0.94	New Jersey	7.84	0.17	15(04,35)	3.95
Colorado	8.13	0.25	07(01,33)	7.78	New Mexico	7.38	0.35	33(04,49)	-2.19
Connecticut	7.15	0.25	37(13,49)	-5.31	New York	7.64	0.11	22(11,36)	1.20
Delaware	7.50	0.50	28(01,50)	-0.60	North Carolina	7.39	0.16	32(13,44)	-2.13
D.C.	7.13	0.65	39(01,51)	-5.48	North Dakota	6.15	0.55	50(12,51)	-18.47 ^c
Florida	6.90	0.10	46(35,49)	-8.61	Ohio	7.86	0.15	14(04,32)	4.09
Georgia	7.71	0.17	18(06,38)	2.13	Oklahoma	8.33	0.27	04(01,29)	10.42
Hawaii	5.40	0.36	51(47,51)	-28.42 ^c	Oregon	8.90	0.27	01(01,10)	17.93 ^c
Idaho	7.69	0.42	21(01,48)	1.87	Pennsylvania	8.08	0.14	09(02,24)	7.04
Illinois	7.71	0.14	17(07,36)	2.22	Rhode Island	6.44	0.43	48(18,51)	-14.64
Indiana	7.87	0.20	13(03,37)	4.23	South Carolina	7.50	0.23	29(06,45)	-0.65
Iowa	8.10	0.29	08(01,36)	7.40	South Dakota	6.98	0.52	43(02,51)	-7.47
Kansas	7.60	0.30	25(03,47)	0.67	Tennessee	7.62	0.20	24(06,41)	0.96
Kentucky	7.36	0.24	34(07,47)	-2.46	Texas	6.98	0.11	45(33,49)	-7.55
Louisiana	6.77	0.23	47(29,50)	-10.35	Utah	7.17	0.35	36(06,50)	-5.03
Maine	7.13	0.39	38(04,50)	-5.45	Vermont	7.55	0.62	26(01,50)	0.08
Maryland	7.55	0.21	27(06,43)	0.01	Virginia	7.69	0.18	20(05,39)	1.96
Massachusetts	7.70	0.19	19(04,39)	2.08	Washington	8.40	0.21	03(01,18)	11.25
Michigan	7.98	0.16	11(03,30)	5.81	West Virginia	7.96	0.35	12(01,42)	5.51
Minnesota	7.50	0.22	30(07,45)	-0.68	Wisconsin	7.74	0.21	16(04,40)	2.57
Mississippi	6.98	0.28	44(15,50)	-7.54	Wyoming	8.17	0.73	06(01,50)	8.21
Missouri	7.10	0.19	41(22,49)	-5.92					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.18
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	10.90	0.03		
High Five States				
D.C.	13.53	0.69	01(01,09)	24.16 ^c
Louisiana	12.80	0.23	02(01,04)	17.45 ^c
Mississippi	12.48	0.28	03(01,07)	14.48
Michigan	11.70	0.14	04(03,18)	7.31
Vermont	11.68	0.56	05(01,45)	7.10
Low Five States				
Utah	9.82	0.30	47(31,51)	-9.88
Montana	9.68	0.40	48(28,51)	-11.20
Colorado	9.64	0.20	49(42,51)	-11.56
Wyoming	9.60	0.57	50(13,51)	-11.93
New Mexico	9.49	0.29	51(39,51)	-12.91

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	11.29	0.20	12(04,34)	3.61	Montana	9.68	0.40	48(28,51)	-11.20
Alaska	10.95	0.69	29(01,51)	0.47	Nebraska	10.79	0.33	32(04,48)	-1.04
Arizona	10.06	0.17	45(34,50)	-7.68	Nevada	10.56	0.28	37(11,49)	-3.13
Arkansas	11.15	0.26	22(04,40)	2.30	New Hampshire	11.29	0.39	13(03,44)	3.58
California	10.45	0.08	39(31,44)	-4.17	New Jersey	11.58	0.15	07(04,23)	6.24
Colorado	9.64	0.20	49(42,51)	-11.56	New Mexico	9.49	0.29	51(39,51)	-12.91
Connecticut	11.24	0.23	17(04,37)	3.10	New York	11.29	0.10	14(06,27)	3.55
Delaware	11.25	0.46	16(02,47)	3.17	North Carolina	10.61	0.14	34(24,44)	-2.69
D.C.	13.53	0.69	01(01,09)	24.16 ^c	North Dakota	9.85	0.50	46(16,51)	-9.66
Florida	10.31	0.09	42(33,47)	-5.46	Ohio	11.42	0.13	10(04,26)	4.72
Georgia	10.40	0.15	41(28,47)	-4.59	Oklahoma	11.02	0.23	28(06,41)	1.10
Hawaii	11.17	0.37	19(03,45)	2.49	Oregon	10.57	0.22	36(18,47)	-3.06
Idaho	10.90	0.37	30(04,48)	-0.04	Pennsylvania	11.49	0.12	08(04,23)	5.44
Illinois	11.20	0.13	18(07,32)	2.74	Rhode Island	10.59	0.41	35(05,51)	-2.87
Indiana	11.10	0.18	24(06,36)	1.85	South Carolina	11.33	0.21	11(04,34)	3.97
Iowa	10.68	0.24	33(11,46)	-2.00	South Dakota	10.41	0.47	40(04,51)	-4.48
Kansas	11.07	0.27	26(04,41)	1.57	Tennessee	11.04	0.18	27(07,37)	1.25
Kentucky	11.17	0.22	20(04,37)	2.42	Texas	10.22	0.10	43(35,48)	-6.28
Louisiana	12.80	0.23	02(01,04)	17.45 ^c	Utah	9.82	0.30	47(31,51)	-9.88
Maine	11.25	0.36	15(03,43)	3.22	Vermont	11.68	0.56	05(01,45)	7.10
Maryland	11.64	0.20	06(03,26)	6.76	Virginia	10.86	0.16	31(13,40)	-0.41
Massachusetts	11.16	0.17	21(06,34)	2.38	Washington	11.08	0.18	25(07,37)	1.65
Michigan	11.70	0.14	04(03,18)	7.31	West Virginia	10.19	0.29	44(20,51)	-6.50
Minnesota	10.47	0.19	38(24,47)	-3.96	Wisconsin	11.43	0.19	09(04,31)	4.89
Mississippi	12.48	0.28	03(01,07)	14.48	Wyoming	9.60	0.57	50(13,51)	-11.93
Missouri	11.11	0.18	23(06,36)	1.92					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.19
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	12.54	0.04		
High Five States				
Louisiana	14.66	0.38	01(01,08)	16.89 ^c
D.C.	14.65	1.12	02(01,47)	16.88 ^c
Mississippi	14.34	0.47	03(01,19)	14.41 ^c
Maryland	13.76	0.33	04(01,24)	9.73
New Hampshire	13.50	0.64	05(01,44)	7.68
Low Five States				
New Mexico	11.19	0.47	47(27,51)	-10.73 ^c
Montana	11.11	0.64	48(17,51)	-11.39 ^c
Wyoming	11.10	0.91	49(05,51)	-11.49 ^c
South Dakota	10.93	0.71	50(14,51)	-12.81 ^c
Colorado	10.43	0.32	51(44,51)	-16.78 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	13.29	0.34	11(02,34)	6.01	Montana	11.11	0.64	48(17,51)	-11.39 ^c
Alaska	12.57	1.11	30(01,51)	0.25	Nebraska	12.37	0.52	35(04,49)	-1.36
Arizona	11.34	0.26	45(35,50)	-9.51	Nevada	12.17	0.45	37(07,49)	-2.93
Arkansas	13.17	0.42	12(02,39)	5.07	New Hampshire	13.50	0.64	05(01,44)	7.68
California	11.78	0.12	43(34,47)	-6.03	New Jersey	13.30	0.25	09(03,30)	6.11
Colorado	10.43	0.32	51(44,51)	-16.78 ^c	New Mexico	11.19	0.47	47(27,51)	-10.73 ^c
Connecticut	12.69	0.38	22(05,45)	1.22	New York	13.08	0.17	16(05,29)	4.36
Delaware	13.44	0.76	06(01,47)	7.21	North Carolina	12.43	0.24	34(13,43)	-0.88
D.C.	14.65	1.12	02(01,47)	16.88 ^c	North Dakota	12.23	0.83	36(01,51)	-2.44
Florida	11.98	0.14	40(30,46)	-4.45	Ohio	13.10	0.21	13(05,32)	4.48
Georgia	12.13	0.26	38(19,47)	-3.25	Oklahoma	12.56	0.37	31(05,46)	0.20
Hawaii	12.66	0.59	26(02,49)	0.94	Oregon	12.07	0.35	39(13,49)	-3.77
Idaho	12.80	0.59	21(02,48)	2.08	Pennsylvania	13.30	0.20	10(03,25)	6.04
Illinois	12.89	0.21	19(07,36)	2.84	Rhode Island	12.89	0.70	20(01,49)	2.80
Indiana	13.06	0.30	17(04,37)	4.18	South Carolina	13.09	0.35	14(03,39)	4.44
Iowa	12.53	0.39	32(05,46)	-0.04	South Dakota	10.93	0.71	50(14,51)	-12.81 ^c
Kansas	12.60	0.43	29(04,47)	0.46	Tennessee	12.69	0.29	24(06,41)	1.20
Kentucky	13.03	0.36	18(03,39)	3.90	Texas	11.69	0.16	44(34,49)	-6.74
Louisiana	14.66	0.38	01(01,08)	16.89 ^c	Utah	11.27	0.48	46(22,51)	-10.12 ^c
Maine	11.84	0.56	42(07,51)	-5.60	Vermont	13.36	0.90	08(01,49)	6.56
Maryland	13.76	0.33	04(01,24)	9.73	Virginia	12.65	0.27	27(07,41)	0.90
Massachusetts	12.67	0.28	25(07,42)	1.08	Washington	12.63	0.29	28(07,42)	0.70
Michigan	13.44	0.23	07(03,25)	7.17	West Virginia	11.97	0.48	41(09,50)	-4.52
Minnesota	12.43	0.31	33(09,46)	-0.84	Wisconsin	13.09	0.30	15(03,36)	4.44
Mississippi	14.34	0.47	03(01,19)	14.41 ^c	Wyoming	11.10	0.91	49(05,51)	-11.49 ^c
Missouri	12.69	0.29	23(06,41)	1.21					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.20
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	9.54	0.03		
High Five States				
D.C.	12.38	0.86	01(01,21)	29.66 ^c
Louisiana	11.21	0.29	02(01,07)	17.44 ^c
Mississippi	10.88	0.35	03(01,18)	13.96
Maine	10.66	0.47	04(01,33)	11.65
Michigan	10.25	0.18	05(03,23)	7.41
Low Five States				
Utah	8.60	0.39	47(17,51)	-9.85
Montana	8.33	0.51	48(16,51)	-12.75
Wyoming	8.31	0.72	49(04,51)	-12.94
New Mexico	8.08	0.36	50(33,51)	-15.40 ^c
North Dakota	7.83	0.59	51(24,51)	-17.97 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	9.62	0.25	24(06,42)	0.81	Montana	8.33	0.51	48(16,51)	-12.75
Alaska	9.37	0.86	31(01,51)	-1.86	Nebraska	9.39	0.41	30(04,49)	-1.62
Arizona	8.91	0.21	42(24,50)	-6.60	Nevada	9.15	0.36	37(07,50)	-4.13
Arkansas	9.40	0.32	29(05,48)	-1.54	New Hampshire	9.35	0.47	32(04,50)	-2.06
California	9.33	0.10	33(22,40)	-2.27	New Jersey	10.21	0.19	06(03,24)	6.98
Colorado	8.94	0.26	41(19,50)	-6.29	New Mexico	8.08	0.36	50(33,51)	-15.40 ^c
Connecticut	10.12	0.29	08(03,33)	6.01	New York	9.89	0.13	15(06,29)	3.60
Delaware	9.46	0.56	27(02,50)	-0.89	North Carolina	9.16	0.18	36(21,47)	-4.01
D.C.	12.38	0.86	01(01,21)	29.66 ^c	North Dakota	7.83	0.59	51(24,51)	-17.97 ^c
Florida	8.88	0.11	43(32,48)	-6.95	Ohio	10.06	0.16	09(04,26)	5.37
Georgia	9.03	0.19	39(23,48)	-5.41	Oklahoma	9.74	0.29	22(04,43)	2.04
Hawaii	9.86	0.48	16(02,48)	3.30	Oregon	9.30	0.28	34(09,48)	-2.57
Idaho	9.22	0.47	35(04,51)	-3.41	Pennsylvania	10.02	0.15	11(04,26)	5.03
Illinois	9.83	0.16	18(06,32)	2.98	Rhode Island	8.82	0.50	44(05,51)	-7.54
Indiana	9.54	0.22	26(07,41)	-0.03	South Carolina	9.94	0.26	13(03,36)	4.18
Iowa	9.10	0.30	38(11,49)	-4.67	South Dakota	9.78	0.61	20(01,50)	2.43
Kansas	9.77	0.34	21(03,45)	2.39	Tennessee	9.73	0.22	23(05,38)	1.93
Kentucky	9.58	0.27	25(05,43)	0.32	Texas	8.96	0.12	40(30,48)	-6.12
Louisiana	11.21	0.29	02(01,07)	17.44 ^c	Utah	8.60	0.39	47(17,51)	-9.85
Maine	10.66	0.47	04(01,33)	11.65	Vermont	10.20	0.70	07(01,49)	6.82
Maryland	10.04	0.24	10(03,33)	5.21	Virginia	9.40	0.20	28(12,44)	-1.52
Massachusetts	9.92	0.21	14(04,33)	3.93	Washington	9.83	0.23	17(04,36)	2.98
Michigan	10.25	0.18	05(03,23)	7.41	West Virginia	8.72	0.36	46(18,51)	-8.61
Minnesota	8.77	0.23	45(26,50)	-8.08	Wisconsin	10.00	0.24	12(03,32)	4.79
Mississippi	10.88	0.35	03(01,18)	13.96	Wyoming	8.31	0.72	49(04,51)	-12.94
Missouri	9.78	0.23	19(04,38)	2.43					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 23.14
Cancer of the Prostate (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	20.65	0.06		
High Five States				
D.C.	35.66	1.84	01(01,01)	72.66 ^c
Mississippi	27.37	0.70	02(02,03)	32.53 ^c
Alabama	25.17	0.51	03(02,06)	21.87 ^c
South Carolina	24.27	0.52	04(03,09)	17.53 ^c
Georgia	23.87	0.40	05(03,09)	15.58 ^c
Low Five States				
Kansas	18.92	0.54	47(31,50)	-8.40
Wyoming	18.91	1.27	48(09,50)	-8.42
Arizona	18.69	0.35	49(37,50)	-9.50
Florida	18.11	0.18	50(46,50)	-12.32 ^c
Hawaii	13.94	0.63	51(51,51)	-32.52 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	25.17	0.51	03(02,06)	21.87 ^c	Montana	22.51	0.94	08(03,38)	8.99
Alaska	20.48	1.62	32(03,50)	-0.84	Nebraska	21.62	0.71	17(06,41)	4.68
Arizona	18.69	0.35	49(37,50)	-9.50	Nevada	21.29	0.65	23(07,44)	3.11
Arkansas	21.80	0.58	15(06,36)	5.57	New Hampshire	20.05	0.83	36(09,50)	-2.94
California	20.55	0.17	29(24,37)	-0.48	New Jersey	19.47	0.32	43(32,49)	-5.74
Colorado	22.03	0.50	14(06,32)	6.68	New Mexico	21.42	0.69	21(06,42)	3.72
Connecticut	19.22	0.47	44(30,50)	-6.96	New York	19.93	0.22	38(30,44)	-3.50
Delaware	20.05	0.99	35(07,50)	-2.91	North Carolina	22.21	0.35	11(06,25)	7.53
D.C.	35.66	1.84	01(01,01)	72.66 ^c	North Dakota	19.97	1.08	37(07,50)	-3.30
Florida	18.11	0.18	50(46,50)	-12.32 ^c	Ohio	20.92	0.28	28(16,37)	1.30
Georgia	23.87	0.40	05(03,09)	15.58 ^c	Oklahoma	21.63	0.52	16(07,35)	4.72
Hawaii	13.94	0.63	51(51,51)	-32.52 ^c	Oregon	22.05	0.49	13(06,31)	6.75
Idaho	23.65	0.85	06(03,25)	14.49 ^c	Pennsylvania	20.38	0.25	33(23,41)	-1.29
Illinois	21.52	0.29	19(10,31)	4.19	Rhode Island	20.54	0.90	30(07,50)	-0.55
Indiana	21.18	0.40	26(10,37)	2.58	South Carolina	24.27	0.52	04(03,09)	17.53 ^c
Iowa	19.82	0.51	40(22,49)	-4.04	South Dakota	20.27	0.99	34(07,50)	-1.87
Kansas	18.92	0.54	47(31,50)	-8.40	Tennessee	21.47	0.41	20(08,35)	3.96
Kentucky	20.52	0.49	31(14,45)	-0.66	Texas	19.19	0.22	45(36,49)	-7.08
Louisiana	23.14	0.52	07(04,20)	12.07 ^c	Utah	22.32	0.72	10(04,35)	8.06
Maine	21.10	0.78	27(06,47)	2.17	Vermont	21.27	1.19	25(03,50)	3.01
Maryland	21.29	0.43	24(09,36)	3.07	Virginia	22.19	0.38	12(06,27)	7.46
Massachusetts	19.88	0.36	39(26,48)	-3.73	Washington	21.33	0.39	22(09,36)	3.28
Michigan	19.68	0.30	41(30,47)	-4.73	West Virginia	19.56	0.65	42(20,50)	-5.29
Minnesota	21.56	0.43	18(08,35)	4.40	Wisconsin	22.41	0.41	09(06,25)	8.51
Mississippi	27.37	0.70	02(02,03)	32.53 ^c	Wyoming	18.91	1.27	48(09,50)	-8.42
Missouri	18.97	0.38	46(35,50)	-8.12					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.18
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.30	0.01		
High Five States				
Hawaii	4.99	0.25	01(01,04)	51.45 ^c
D.C.	4.83	0.42	02(01,13)	46.56 ^c
Alaska	4.59	0.45	03(01,18)	39.39 ^c
Louisiana	4.25	0.14	04(02,08)	28.98 ^c
California	4.18	0.05	05(02,07)	26.91 ^c
Low Five States				
Idaho	2.22	0.17	47(36,51)	-32.58 ^c
Iowa	2.17	0.11	48(41,51)	-34.03 ^c
Utah	2.09	0.14	49(41,51)	-36.61 ^c
Maine	2.08	0.15	50(40,51)	-36.81 ^c
Nebraska	2.02	0.14	51(42,51)	-38.56 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.32	0.11	18(11,26)	0.90	Montana	2.26	0.20	46(30,51)	-31.54 ^c
Alaska	4.59	0.45	03(01,18)	39.39 ^c	Nebraska	2.02	0.14	51(42,51)	-38.56 ^c
Arizona	2.82	0.09	32(22,42)	-14.42	Nevada	2.96	0.15	28(16,42)	-10.32
Arkansas	2.98	0.14	25(16,41)	-9.58	New Hampshire	2.34	0.18	44(30,51)	-28.84 ^c
California	4.18	0.05	05(02,07)	26.91 ^c	New Jersey	3.69	0.09	10(06,16)	12.12
Colorado	2.77	0.11	35(23,43)	-15.91 ^c	New Mexico	3.98	0.19	07(02,16)	20.70 ^c
Connecticut	3.52	0.13	16(07,22)	6.88	New York	3.85	0.06	09(05,13)	16.90 ^c
Delaware	3.56	0.26	14(03,34)	8.05	North Carolina	3.13	0.08	22(16,31)	-4.96
D.C.	4.83	0.42	02(01,13)	46.56 ^c	North Dakota	2.73	0.26	36(15,50)	-17.24 ^c
Florida	2.93	0.05	30(22,35)	-11.06	Ohio	2.80	0.07	34(25,41)	-15.06 ^c
Georgia	3.53	0.09	15(09,19)	7.20	Oklahoma	2.80	0.12	33(21,43)	-14.89
Hawaii	4.99	0.25	01(01,04)	51.45 ^c	Oregon	2.51	0.11	41(31,47)	-23.74 ^c
Idaho	2.22	0.17	47(36,51)	-32.58 ^c	Pennsylvania	2.96	0.06	27(21,35)	-10.06
Illinois	3.65	0.07	13(07,17)	10.65	Rhode Island	3.67	0.24	12(03,26)	11.32
Indiana	2.68	0.09	37(27,44)	-18.52 ^c	South Carolina	3.87	0.12	08(04,15)	17.57 ^c
Iowa	2.17	0.11	48(41,51)	-34.03 ^c	South Dakota	2.47	0.23	42(20,51)	-24.90 ^c
Kansas	2.42	0.12	43(33,50)	-26.59 ^c	Tennessee	3.23	0.10	19(14,29)	-2.10
Kentucky	2.97	0.11	26(17,40)	-9.99	Texas	3.67	0.06	11(08,16)	11.42
Louisiana	4.25	0.14	04(02,08)	28.98 ^c	Utah	2.09	0.14	49(41,51)	-36.61 ^c
Maine	2.08	0.15	50(40,51)	-36.81 ^c	Vermont	2.55	0.26	40(17,51)	-22.56 ^c
Maryland	3.34	0.11	17(12,25)	1.51	Virginia	3.04	0.09	24(17,35)	-7.83
Massachusetts	3.17	0.09	21(15,31)	-3.94	Washington	2.93	0.09	29(19,39)	-10.96
Michigan	3.05	0.07	23(18,33)	-7.49	West Virginia	3.20	0.17	20(11,36)	-2.79
Minnesota	2.65	0.10	39(28,45)	-19.54 ^c	Wisconsin	2.83	0.09	31(22,41)	-14.04
Mississippi	4.00	0.16	06(02,15)	21.42 ^c	Wyoming	2.27	0.28	45(21,51)	-31.01 ^c
Missouri	2.66	0.09	38(29,44)	-19.38 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.19
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.47	0.03		
High Five States				
D.C.	6.98	0.79	01(01,16)	56.01 ^c
Hawaii	6.65	0.44	02(01,07)	48.70 ^c
Alaska	5.93	0.75	03(01,34)	32.64 ^c
Louisiana	5.86	0.24	04(01,11)	31.04 ^c
South Carolina	5.61	0.23	05(02,14)	25.47 ^c
Low Five States				
South Dakota	2.90	0.37	47(27,51)	-35.20 ^c
Utah	2.89	0.25	48(37,51)	-35.49 ^c
Nebraska	2.85	0.25	49(38,51)	-36.38 ^c
Montana	2.84	0.33	50(31,51)	-36.52 ^c
Maine	2.66	0.26	51(40,51)	-40.60 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.56	0.20	18(09,30)	2.04	Montana	2.84	0.33	50(31,51)	-36.52 ^c
Alaska	5.93	0.75	03(01,34)	32.64 ^c	Nebraska	2.85	0.25	49(38,51)	-36.38 ^c
Arizona	3.75	0.15	37(23,43)	-16.22 ^c	Nevada	3.84	0.26	34(17,46)	-14.24
Arkansas	3.91	0.24	30(17,44)	-12.60	New Hampshire	3.29	0.32	43(21,51)	-26.49 ^c
California	5.50	0.08	06(03,11)	22.86 ^c	New Jersey	5.10	0.16	11(04,18)	14.03
Colorado	3.76	0.19	36(21,45)	-15.87 ^c	New Mexico	5.06	0.32	13(03,27)	13.23
Connecticut	4.98	0.24	14(04,23)	11.28	New York	5.33	0.11	07(04,14)	19.19 ^c
Delaware	5.08	0.48	12(02,37)	13.52	North Carolina	4.20	0.14	23(17,35)	-6.08
D.C.	6.98	0.79	01(01,16)	56.01 ^c	North Dakota	3.41	0.43	41(16,51)	-23.86 ^c
Florida	3.86	0.08	32(25,40)	-13.69	Ohio	3.78	0.12	35(25,42)	-15.40 ^c
Georgia	4.78	0.16	16(08,23)	6.83	Oklahoma	4.04	0.21	27(16,42)	-9.57
Hawaii	6.65	0.44	02(01,07)	48.70 ^c	Oregon	3.33	0.19	42(31,49)	-25.58 ^c
Idaho	3.13	0.30	45(27,51)	-29.97 ^c	Pennsylvania	4.20	0.11	22(18,33)	-6.00
Illinois	5.11	0.13	10(05,17)	14.19	Rhode Island	5.18	0.44	09(02,32)	15.90 ^c
Indiana	3.86	0.16	33(20,43)	-13.79	South Carolina	5.61	0.23	05(02,14)	25.47 ^c
Iowa	3.02	0.19	46(38,51)	-32.59 ^c	South Dakota	2.90	0.37	47(27,51)	-35.20 ^c
Kansas	3.42	0.22	40(26,50)	-23.59 ^c	Tennessee	4.56	0.18	19(10,28)	1.88
Kentucky	4.10	0.20	26(16,42)	-8.45	Texas	4.86	0.10	15(09,19)	8.70
Louisiana	5.86	0.24	04(01,11)	31.04 ^c	Utah	2.89	0.25	48(37,51)	-35.49 ^c
Maine	2.66	0.26	51(40,51)	-40.60 ^c	Vermont	3.87	0.49	31(07,51)	-13.52
Maryland	4.70	0.19	17(07,26)	5.17	Virginia	4.13	0.15	25(18,38)	-7.75
Massachusetts	4.20	0.16	24(16,37)	-6.09	Washington	4.04	0.16	28(18,40)	-9.71
Michigan	4.27	0.13	21(16,33)	-4.64	West Virginia	4.45	0.30	20(07,39)	-0.42
Minnesota	3.64	0.17	39(24,45)	-18.52 ^c	Wisconsin	3.94	0.17	29(19,41)	-11.86
Mississippi	5.23	0.29	08(02,20)	16.91 ^c	Wyoming	3.17	0.50	44(16,51)	-29.02 ^c
Missouri	3.72	0.16	38(24,44)	-16.93 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.20
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.38	0.02		
High Five States				
Hawaii	3.65	0.28	01(01,08)	53.20 ^c
D.C.	3.35	0.46	02(01,31)	40.85 ^c
Alaska	3.30	0.52	03(01,36)	38.43 ^c
California	3.17	0.06	04(01,07)	32.97 ^c
New Mexico	3.07	0.23	05(01,17)	28.78 ^c
Low Five States				
Iowa	1.55	0.12	47(36,51)	-35.02 ^c
Wyoming	1.49	0.31	48(14,51)	-37.31 ^c
Utah	1.43	0.16	49(35,51)	-40.12 ^c
Idaho	1.41	0.18	50(33,51)	-40.79 ^c
Nebraska	1.37	0.16	51(38,51)	-42.32 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	2.41	0.13	15(07,32)	1.11	Montana	1.76	0.24	42(14,51)	-26.06 ^c
Alaska	3.30	0.52	03(01,36)	38.43 ^c	Nebraska	1.37	0.16	51(38,51)	-42.32 ^c
Arizona	2.04	0.11	31(18,43)	-14.45	Nevada	2.24	0.18	22(08,43)	-5.77
Arkansas	2.28	0.16	21(08,40)	-4.44	New Hampshire	1.59	0.20	45(23,51)	-33.19 ^c
California	3.17	0.06	04(01,07)	32.97 ^c	New Jersey	2.66	0.10	10(05,18)	11.90
Colorado	1.97	0.12	36(18,46)	-17.14 ^c	New Mexico	3.07	0.23	05(01,17)	28.78 ^c
Connecticut	2.39	0.14	16(07,34)	0.21	New York	2.77	0.07	08(05,13)	16.54 ^c
Delaware	2.29	0.28	20(03,48)	-3.88	North Carolina	2.32	0.09	19(11,31)	-2.72
D.C.	3.35	0.46	02(01,31)	40.85 ^c	North Dakota	2.15	0.32	27(03,50)	-9.84
Florida	2.16	0.06	26(18,35)	-9.07	Ohio	2.03	0.07	32(21,41)	-14.64
Georgia	2.59	0.10	12(06,22)	8.96	Oklahoma	1.83	0.13	39(22,49)	-23.10 ^c
Hawaii	3.65	0.28	01(01,08)	53.20 ^c	Oregon	1.84	0.13	38(22,48)	-22.72 ^c
Idaho	1.41	0.18	50(33,51)	-40.79 ^c	Pennsylvania	2.02	0.07	34(22,41)	-15.22 ^c
Illinois	2.54	0.08	13(08,22)	6.70	Rhode Island	2.60	0.27	11(02,40)	9.40
Indiana	1.81	0.10	41(27,48)	-24.17 ^c	South Carolina	2.45	0.13	14(07,30)	3.03
Iowa	1.55	0.12	47(36,51)	-35.02 ^c	South Dakota	2.06	0.28	30(06,50)	-13.34
Kansas	1.66	0.14	43(27,51)	-30.36 ^c	Tennessee	2.20	0.11	25(13,38)	-7.45
Kentucky	2.07	0.13	29(15,44)	-13.17	Texas	2.74	0.07	09(05,14)	15.27 ^c
Louisiana	3.01	0.15	07(01,14)	26.47 ^c	Utah	1.43	0.16	49(35,51)	-40.12 ^c
Maine	1.56	0.18	46(26,51)	-34.36 ^c	Vermont	1.60	0.28	44(14,51)	-32.67 ^c
Maryland	2.33	0.12	18(10,34)	-2.24	Virginia	2.20	0.10	24(13,38)	-7.44
Massachusetts	2.38	0.11	17(10,30)	0.05	Washington	2.03	0.10	33(18,43)	-14.80
Michigan	2.13	0.08	28(16,38)	-10.43	West Virginia	2.23	0.19	23(07,44)	-6.29
Minnesota	1.90	0.11	37(21,47)	-20.38 ^c	Wisconsin	1.98	0.11	35(19,44)	-16.89 ^c
Mississippi	3.05	0.19	06(01,14)	27.92 ^c	Wyoming	1.49	0.31	48(14,51)	-37.31 ^c
Missouri	1.82	0.10	40(27,48)	-23.68 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 25.14
Cancer of the Testis (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.25	0.01		
High Five States				
West Virginia	0.39	0.10	01(01,30)	55.81 ^c
Oklahoma	0.39	0.07	02(01,22)	54.20 ^c
Arizona	0.38	0.05	03(01,15)	51.90 ^c
Missouri	0.36	0.05	04(01,21)	43.18 ^c
California	0.32	0.02	05(02,14)	27.25 ^c
Low Five States				
South Carolina	0.19	0.04	28(07,32)	-24.23 ^c
Maryland	0.19	0.04	29(07,32)	-24.54 ^c
Massachusetts	0.15	0.03	30(17,32)	-41.15 ^c
Virginia	0.15	0.03	31(20,32)	-42.08 ^c
Minnesota	0.13	0.03	32(20,32)	-47.33 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.24	0.05	18(02,32)	-4.24	Montana	-	-	-(-,-)	-
Alaska	-	-	-(-,-)	-	Nebraska	-	-	-(-,-)	-
Arizona	0.38	0.05	03(01,15)	51.90 ^c	Nevada	0.28	0.06	10(01,31)	12.39
Arkansas	-	-	-(-,-)	-	New Hampshire	-	-	-(-,-)	-
California	0.32	0.02	05(02,14)	27.25 ^c	New Jersey	0.23	0.03	20(05,31)	-7.18
Colorado	0.29	0.05	09(01,29)	14.39	New Mexico	0.31	0.08	07(01,32)	22.98 ^c
Connecticut	-	-	-(-,-)	-	New York	0.22	0.02	23(10,30)	-13.33
Delaware	-	-	-(-,-)	-	North Carolina	0.25	0.03	16(04,29)	-0.32
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.27	0.02	12(04,25)	7.28	Ohio	0.25	0.03	17(04,29)	-0.74
Georgia	0.21	0.03	26(08,32)	-16.74 ^c	Oklahoma	0.39	0.07	02(01,22)	54.20 ^c
Hawaii	-	-	-(-,-)	-	Oregon	-	-	-(-,-)	-
Idaho	-	-	-(-,-)	-	Pennsylvania	0.26	0.03	15(04,28)	2.00
Illinois	0.19	0.02	27(13,32)	-23.79 ^c	Rhode Island	-	-	-(-,-)	-
Indiana	0.30	0.05	08(01,27)	21.04 ^c	South Carolina	0.19	0.04	28(07,32)	-24.23 ^c
Iowa	0.26	0.06	13(01,32)	4.72	South Dakota	-	-	-(-,-)	-
Kansas	0.22	0.06	24(02,32)	-13.36	Tennessee	0.23	0.04	19(03,31)	-6.63
Kentucky	0.31	0.05	06(01,28)	23.24 ^c	Texas	0.28	0.02	11(04,23)	10.17
Louisiana	0.21	0.04	25(04,32)	-16.58 ^c	Utah	-	-	-(-,-)	-
Maine	-	-	-(-,-)	-	Vermont	-	-	-(-,-)	-
Maryland	0.19	0.04	29(07,32)	-24.54 ^c	Virginia	0.15	0.03	31(20,32)	-42.08 ^c
Massachusetts	0.15	0.03	30(17,32)	-41.15 ^c	Washington	0.22	0.04	22(05,32)	-11.48
Michigan	0.26	0.03	14(03,29)	2.54	West Virginia	0.39	0.10	01(01,30)	55.81 ^c
Minnesota	0.13	0.03	32(20,32)	-47.33 ^c	Wisconsin	0.23	0.04	21(03,32)	-7.98
Mississippi	-	-	-(-,-)	-	Wyoming	-	-	-(-,-)	-
Missouri	0.36	0.05	04(01,21)	43.18 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 26.18
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.51	0.01		
High Five States				
Hawaii	0.79	0.10	01(01,23)	55.15 ^c
D.C.	0.76	0.16	02(01,50)	49.62 ^c
Idaho	0.68	0.09	03(01,45)	33.71 ^c
Alaska	0.66	0.17	04(01,50)	29.24 ^c
Rhode Island	0.63	0.10	05(01,49)	24.42 ^c
Low Five States				
Mississippi	0.43	0.05	46(07,50)	-15.90 ^c
Maryland	0.43	0.04	47(14,50)	-15.97 ^c
Florida	0.43	0.02	48(29,50)	-16.10 ^c
Connecticut	0.42	0.05	49(10,50)	-16.62 ^c
Georgia	0.41	0.03	50(25,50)	-20.01 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.45	0.04	39(08,50)	-10.44	Montana	0.47	0.09	35(02,50)	-7.97
Alaska	0.66	0.17	04(01,50)	29.24 ^c	Nebraska	0.57	0.07	08(01,48)	13.20
Arizona	0.47	0.04	34(08,50)	-7.90	Nevada	0.53	0.06	14(02,49)	5.27
Arkansas	0.46	0.05	38(05,50)	-10.07	New Hampshire	0.45	0.08	41(03,50)	-10.54
California	0.59	0.02	06(03,18)	16.23 ^c	New Jersey	0.50	0.03	27(06,46)	-2.29
Colorado	0.49	0.05	30(05,49)	-3.03	New Mexico	0.57	0.07	10(01,49)	11.80
Connecticut	0.42	0.05	49(10,50)	-16.62 ^c	New York	0.55	0.02	12(05,30)	8.51
Delaware	0.58	0.11	07(01,50)	13.98	North Carolina	0.51	0.03	18(05,44)	0.78
D.C.	0.76	0.16	02(01,50)	49.62 ^c	North Dakota	0.46	0.10	37(01,50)	-8.38
Florida	0.43	0.02	48(29,50)	-16.10 ^c	Ohio	0.52	0.03	15(06,40)	2.70
Georgia	0.41	0.03	50(25,50)	-20.01 ^c	Oklahoma	0.47	0.05	36(05,50)	-8.11
Hawaii	0.79	0.10	01(01,23)	55.15 ^c	Oregon	0.52	0.05	16(03,49)	2.29
Idaho	0.68	0.09	03(01,45)	33.71 ^c	Pennsylvania	0.52	0.03	17(07,40)	1.53
Illinois	0.51	0.03	22(07,44)	-0.32	Rhode Island	0.63	0.10	05(01,49)	24.42 ^c
Indiana	0.47	0.04	31(08,49)	-6.71	South Carolina	0.43	0.04	44(10,50)	-14.69
Iowa	0.47	0.05	32(05,50)	-7.72	South Dakota	0.47	0.10	33(01,50)	-7.86
Kansas	0.50	0.06	25(03,50)	-1.20	Tennessee	0.44	0.04	42(10,50)	-12.52
Kentucky	0.51	0.05	23(04,49)	-0.44	Texas	0.56	0.02	11(04,29)	9.56
Louisiana	0.43	0.04	43(10,50)	-14.57	Utah	0.49	0.07	29(03,50)	-2.97
Maine	0.43	0.07	45(04,50)	-14.96	Vermont	0.45	0.11	40(01,50)	-10.49
Maryland	0.43	0.04	47(14,50)	-15.97 ^c	Virginia	0.50	0.03	26(06,47)	-1.63
Massachusetts	0.57	0.04	09(02,37)	13.02	Washington	0.50	0.04	24(05,47)	-0.54
Michigan	0.49	0.03	28(08,46)	-2.76	West Virginia	0.51	0.06	19(02,50)	0.26
Minnesota	0.51	0.04	21(04,48)	-0.13	Wisconsin	0.54	0.04	13(03,43)	6.78
Mississippi	0.43	0.05	46(07,50)	-15.90 ^c	Wyoming	-	-	-(-,-)	-
Missouri	0.51	0.04	20(04,47)	-0.07					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 26.19
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.51	0.01		
High Five States				
Rhode Island	0.93	0.19	01(01,37)	81.85 ^c
Idaho	0.70	0.14	02(01,43)	35.53 ^c
Nebraska	0.65	0.12	03(01,43)	25.87 ^c
Massachusetts	0.63	0.06	04(01,32)	23.08 ^c
Hawaii	0.62	0.13	05(01,43)	21.60 ^c
Low Five States				
Arkansas	0.41	0.08	39(05,43)	-19.23 ^c
Connecticut	0.41	0.07	40(06,43)	-19.47 ^c
Louisiana	0.41	0.07	41(08,43)	-19.87 ^c
Maryland	0.40	0.06	42(13,43)	-21.90 ^c
Mississippi	0.40	0.08	43(05,43)	-22.50 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.43	0.06	36(07,43)	-15.93 ^c	Montana	0.60	0.14	07(01,43)	17.00 ^c
Alaska	-	-	-(-,-)	-	Nebraska	0.65	0.12	03(01,43)	25.87 ^c
Arizona	0.54	0.06	18(02,40)	4.71	Nevada	0.47	0.08	33(03,43)	-8.49
Arkansas	0.41	0.08	39(05,43)	-19.23 ^c	New Hampshire	0.49	0.12	30(01,43)	-5.27
California	0.53	0.03	20(07,33)	4.00	New Jersey	0.52	0.05	22(04,41)	0.76
Colorado	0.54	0.07	16(02,42)	5.61	New Mexico	0.43	0.09	37(03,43)	-16.57 ^c
Connecticut	0.41	0.07	40(06,43)	-19.47 ^c	New York	0.56	0.04	14(04,33)	8.53
Delaware	-	-	-(-,-)	-	North Carolina	0.55	0.05	15(03,39)	6.43
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.44	0.03	35(20,42)	-14.78	Ohio	0.54	0.04	17(03,38)	5.57
Georgia	0.43	0.05	38(12,43)	-17.20 ^c	Oklahoma	0.57	0.08	13(01,42)	10.57
Hawaii	0.62	0.13	05(01,43)	21.60 ^c	Oregon	0.60	0.08	08(01,41)	16.82 ^c
Idaho	0.70	0.14	02(01,43)	35.53 ^c	Pennsylvania	0.50	0.04	25(06,40)	-2.56
Illinois	0.45	0.04	34(12,43)	-12.84	Rhode Island	0.93	0.19	01(01,37)	81.85 ^c
Indiana	0.50	0.06	27(04,43)	-3.48	South Carolina	0.47	0.07	32(04,43)	-7.62
Iowa	0.54	0.08	19(02,43)	4.36	South Dakota	-	-	-(-,-)	-
Kansas	0.61	0.10	06(01,42)	18.30 ^c	Tennessee	0.49	0.06	28(04,43)	-4.36
Kentucky	0.53	0.07	21(02,43)	3.11	Texas	0.59	0.04	10(03,28)	14.17
Louisiana	0.41	0.07	41(08,43)	-19.87 ^c	Utah	0.60	0.11	09(01,43)	16.76 ^c
Maine	-	-	-(-,-)	-	Vermont	-	-	-(-,-)	-
Maryland	0.40	0.06	42(13,43)	-21.90 ^c	Virginia	0.58	0.06	11(02,37)	12.23
Massachusetts	0.63	0.06	04(01,32)	23.08 ^c	Washington	0.49	0.06	29(04,43)	-4.79
Michigan	0.50	0.05	26(06,42)	-3.02	West Virginia	0.48	0.10	31(02,43)	-6.48
Minnesota	0.57	0.07	12(02,41)	11.53	Wisconsin	0.51	0.06	23(03,42)	-0.76
Mississippi	0.40	0.08	43(05,43)	-22.50 ^c	Wyoming	-	-	-(-,-)	-
Missouri	0.50	0.06	24(03,43)	-2.47					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 26.20
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.50	0.01		
High Five States				
Hawaii	0.92	0.15	01(01,16)	85.47 ^c
Idaho	0.67	0.13	02(01,43)	35.07 ^c
New Mexico	0.67	0.11	03(01,40)	33.92 ^c
California	0.63	0.02	04(02,11)	26.55 ^c
North Dakota	0.60	0.16	05(01,44)	21.54 ^c
Low Five States				
Tennessee	0.40	0.05	40(10,44)	-19.09 ^c
South Carolina	0.40	0.05	41(07,44)	-19.58 ^c
Oklahoma	0.40	0.06	42(07,44)	-20.23 ^c
Iowa	0.40	0.06	43(05,44)	-20.28 ^c
Georgia	0.38	0.04	44(18,44)	-23.87 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.47	0.06	24(04,44)	-5.89	Montana	-	-	-(-,-)	-
Alaska	-	-	-(-,-)	-	Nebraska	0.46	0.08	26(02,44)	-7.18
Arizona	0.41	0.05	39(11,44)	-18.43 ^c	Nevada	0.57	0.09	07(01,43)	14.09
Arkansas	0.48	0.07	23(02,44)	-3.63	New Hampshire	0.41	0.10	37(02,44)	-17.00 ^c
California	0.63	0.02	04(02,11)	26.55 ^c	New Jersey	0.48	0.04	21(05,41)	-2.81
Colorado	0.45	0.06	29(04,44)	-9.05	New Mexico	0.67	0.11	03(01,40)	33.92 ^c
Connecticut	0.42	0.06	35(06,44)	-15.67 ^c	New York	0.54	0.03	09(04,28)	8.62
Delaware	0.60	0.14	06(01,44)	20.59 ^c	North Carolina	0.49	0.04	19(04,41)	-1.23
D.C.	-	-	-(-,-)	-	North Dakota	0.60	0.16	05(01,44)	21.54 ^c
Florida	0.41	0.02	36(20,44)	-16.82 ^c	Ohio	0.50	0.04	18(05,38)	0.04
Georgia	0.38	0.04	44(18,44)	-23.87 ^c	Oklahoma	0.40	0.06	42(07,44)	-20.23 ^c
Hawaii	0.92	0.15	01(01,16)	85.47 ^c	Oregon	0.47	0.06	25(03,44)	-6.35
Idaho	0.67	0.13	02(01,43)	35.07 ^c	Pennsylvania	0.53	0.03	12(04,33)	5.73
Illinois	0.53	0.04	11(03,33)	7.34	Rhode Island	-	-	-(-,-)	-
Indiana	0.44	0.05	33(06,44)	-11.08	South Carolina	0.40	0.05	41(07,44)	-19.58 ^c
Iowa	0.40	0.06	43(05,44)	-20.28 ^c	South Dakota	-	-	-(-,-)	-
Kansas	0.42	0.07	34(04,44)	-14.48	Tennessee	0.40	0.05	40(10,44)	-19.09 ^c
Kentucky	0.48	0.06	22(03,44)	-3.55	Texas	0.52	0.03	13(05,31)	5.62
Louisiana	0.45	0.06	30(05,44)	-10.43	Utah	0.41	0.08	38(03,44)	-18.34 ^c
Maine	0.53	0.11	10(01,44)	7.47	Vermont	-	-	-(-,-)	-
Maryland	0.44	0.05	32(06,44)	-10.80	Virginia	0.44	0.04	31(07,44)	-10.58
Massachusetts	0.52	0.05	14(03,40)	4.73	Washington	0.51	0.05	15(03,42)	2.15
Michigan	0.48	0.04	20(05,40)	-2.52	West Virginia	0.50	0.08	16(02,44)	0.93
Minnesota	0.45	0.05	28(05,44)	-8.94	Wisconsin	0.57	0.06	08(02,35)	13.98
Mississippi	0.46	0.07	27(03,44)	-8.31	Wyoming	-	-	-(-,-)	-
Missouri	0.50	0.05	17(04,42)	0.18					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 27.18
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.41	0.02		
High Five States				
Maine	5.99	0.26	01(01,04)	35.96 ^c
Vermont	5.60	0.39	02(01,25)	27.09 ^c
Nevada	5.28	0.21	03(01,19)	19.84 ^c
New Hampshire	5.13	0.26	04(01,29)	16.35 ^c
Ohio	5.11	0.09	05(02,14)	15.85 ^c
Low Five States				
Mississippi	3.71	0.16	47(33,50)	-15.79 ^c
Texas	3.71	0.06	48(42,50)	-15.83 ^c
South Dakota	3.70	0.27	49(21,50)	-16.04 ^c
Alaska	3.28	0.40	50(21,51)	-25.65 ^c
Hawaii	2.68	0.18	51(50,51)	-39.18 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.91	0.12	45(31,50)	-11.30	Montana	4.91	0.28	10(02,38)	11.34
Alaska	3.28	0.40	50(21,51)	-25.65 ^c	Nebraska	4.03	0.20	40(20,50)	-8.65
Arizona	4.07	0.11	37(25,48)	-7.67	Nevada	5.28	0.21	03(01,19)	19.84 ^c
Arkansas	4.56	0.17	21(06,38)	3.49	New Hampshire	5.13	0.26	04(01,29)	16.35 ^c
California	3.96	0.05	42(35,47)	-10.03	New Jersey	4.85	0.10	13(04,22)	10.00
Colorado	3.93	0.13	44(28,50)	-10.91	New Mexico	3.99	0.19	41(20,50)	-9.37
Connecticut	4.92	0.15	09(02,24)	11.65	New York	4.38	0.06	27(20,36)	-0.54
Delaware	4.62	0.30	20(02,47)	4.85	North Carolina	4.27	0.09	30(21,42)	-3.08
D.C.	3.85	0.37	46(08,51)	-12.68	North Dakota	4.22	0.32	34(03,50)	-4.26
Florida	4.42	0.06	26(19,34)	0.27	Ohio	5.11	0.09	05(02,14)	15.85 ^c
Georgia	4.45	0.10	24(14,36)	0.98	Oklahoma	4.53	0.15	23(08,38)	2.77
Hawaii	2.68	0.18	51(50,51)	-39.18 ^c	Oregon	4.76	0.15	17(04,30)	8.06
Idaho	4.21	0.23	35(08,49)	-4.38	Pennsylvania	4.84	0.08	15(05,22)	9.91
Illinois	4.34	0.08	28(20,38)	-1.42	Rhode Island	4.94	0.28	08(01,36)	12.18
Indiana	4.86	0.12	12(03,23)	10.30	South Carolina	4.25	0.13	32(19,45)	-3.56
Iowa	4.43	0.15	25(11,42)	0.51	South Dakota	3.70	0.27	49(21,50)	-16.04 ^c
Kansas	4.06	0.16	38(21,49)	-7.75	Tennessee	4.25	0.11	31(20,44)	-3.48
Kentucky	4.80	0.14	16(03,27)	8.98	Texas	3.71	0.06	48(42,50)	-15.83 ^c
Louisiana	4.16	0.14	36(20,47)	-5.61	Utah	3.95	0.20	43(21,50)	-10.35
Maine	5.99	0.26	01(01,04)	35.96 ^c	Vermont	5.60	0.39	02(01,25)	27.09 ^c
Maryland	4.99	0.13	07(02,21)	13.23	Virginia	4.23	0.10	33(21,44)	-4.02
Massachusetts	5.02	0.11	06(02,20)	13.90	Washington	4.88	0.12	11(03,22)	10.80
Michigan	4.85	0.09	14(04,22)	9.98	West Virginia	4.62	0.20	19(03,40)	4.90
Minnesota	4.04	0.12	39(25,48)	-8.25	Wisconsin	4.69	0.12	18(06,30)	6.32
Mississippi	3.71	0.16	47(33,50)	-15.79 ^c	Wyoming	4.55	0.40	22(02,50)	3.25
Missouri	4.28	0.11	29(19,43)	-2.79					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 27.19
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	7.66	0.03		
High Five States				
Maine	10.52	0.54	01(01,05)	37.33 ^c
Vermont	9.41	0.79	02(01,39)	22.85 ^c
New Hampshire	9.24	0.55	03(01,25)	20.62 ^c
Ohio	8.93	0.18	04(02,15)	16.52 ^c
Maryland	8.92	0.28	05(02,19)	16.43 ^c
Low Five States				
South Dakota	6.50	0.56	47(16,51)	-15.16 ^c
Texas	6.43	0.12	48(41,49)	-16.02 ^c
D.C.	5.13	0.69	49(31,51)	-33.06 ^c
Hawaii	4.96	0.38	50(48,51)	-35.32 ^c
Alaska	4.77	0.72	51(34,51)	-37.71 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	7.12	0.27	36(20,48)	-7.10	Montana	8.77	0.58	07(01,37)	14.40
Alaska	4.77	0.72	51(34,51)	-37.71 ^c	Nebraska	6.98	0.40	42(17,49)	-8.96
Arizona	7.00	0.21	40(25,48)	-8.63	Nevada	8.78	0.41	06(01,27)	14.62
Arkansas	7.93	0.34	21(04,41)	3.45	New Hampshire	9.24	0.55	03(01,25)	20.62 ^c
California	6.81	0.10	44(34,47)	-11.16	New Jersey	8.49	0.21	13(03,23)	10.75
Colorado	6.72	0.27	45(27,49)	-12.26	New Mexico	6.98	0.39	41(17,49)	-8.85
Connecticut	8.53	0.32	11(02,27)	11.26	New York	7.75	0.13	22(17,33)	1.09
Delaware	7.21	0.59	35(04,49)	-5.90	North Carolina	7.45	0.20	29(19,42)	-2.83
D.C.	5.13	0.69	49(31,51)	-33.06 ^c	North Dakota	7.43	0.64	30(02,49)	-3.07
Florida	7.60	0.11	26(19,35)	-0.81	Ohio	8.93	0.18	04(02,15)	16.52 ^c
Georgia	7.93	0.23	20(08,34)	3.50	Oklahoma	7.73	0.31	23(09,43)	0.83
Hawaii	4.96	0.38	50(48,51)	-35.32 ^c	Oregon	8.32	0.30	18(03,30)	8.64
Idaho	7.27	0.46	34(08,48)	-5.09	Pennsylvania	8.64	0.16	08(03,19)	12.76
Illinois	7.53	0.17	27(18,39)	-1.69	Rhode Island	8.49	0.58	12(01,42)	10.86
Indiana	8.59	0.25	09(02,23)	12.10	South Carolina	7.29	0.28	31(17,47)	-4.85
Iowa	7.99	0.32	19(05,38)	4.33	South Dakota	6.50	0.56	47(16,51)	-15.16 ^c
Kansas	7.03	0.33	37(19,48)	-8.29	Tennessee	7.65	0.24	24(14,41)	-0.20
Kentucky	8.42	0.31	14(02,30)	9.88	Texas	6.43	0.12	48(41,49)	-16.02 ^c
Louisiana	7.02	0.28	39(21,48)	-8.37	Utah	6.84	0.40	43(19,49)	-10.79
Maine	10.52	0.54	01(01,05)	37.33 ^c	Vermont	9.41	0.79	02(01,39)	22.85 ^c
Maryland	8.92	0.28	05(02,19)	16.43 ^c	Virginia	7.28	0.22	32(20,45)	-4.98
Massachusetts	8.54	0.24	10(02,23)	11.48	Washington	8.37	0.24	16(03,26)	9.30
Michigan	8.36	0.19	17(04,24)	9.16	West Virginia	7.46	0.39	28(09,48)	-2.67
Minnesota	7.02	0.24	38(23,48)	-8.33	Wisconsin	8.40	0.25	15(03,27)	9.70
Mississippi	6.58	0.34	46(25,49)	-14.07	Wyoming	7.61	0.78	25(02,49)	-0.70
Missouri	7.28	0.23	33(20,46)	-5.02					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 27.20
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.17	0.01		
High Five States				
D.C.	3.07	0.43	01(01,40)	41.45 ^c
Delaware	2.96	0.32	02(01,34)	36.53 ^c
Vermont	2.93	0.38	03(01,42)	35.06 ^c
Maine	2.78	0.24	04(01,32)	27.90 ^c
Massachusetts	2.73	0.11	05(01,16)	25.72 ^c
Low Five States				
New Mexico	1.78	0.17	47(17,50)	-18.08 ^c
Utah	1.77	0.18	48(17,50)	-18.36 ^c
Alabama	1.77	0.11	49(31,50)	-18.56 ^c
South Dakota	1.76	0.25	50(07,51)	-19.15 ^c
Hawaii	0.98	0.14	51(50,51)	-54.68 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	1.77	0.11	49(31,50)	-18.56 ^c	Montana	2.00	0.25	36(04,50)	-7.81
Alaska	1.94	0.43	40(01,51)	-10.85	Nebraska	2.01	0.19	35(07,50)	-7.33
Arizona	1.81	0.10	44(31,50)	-16.59 ^c	Nevada	2.54	0.19	08(01,37)	16.84 ^c
Arkansas	2.21	0.15	23(05,45)	1.99	New Hampshire	2.18	0.23	28(02,50)	0.23
California	1.95	0.04	39(30,45)	-9.95	New Jersey	2.50	0.09	11(03,25)	15.03 ^c
Colorado	1.88	0.12	41(21,50)	-13.21	New Mexico	1.78	0.17	47(17,50)	-18.08 ^c
Connecticut	2.53	0.14	09(02,32)	16.76 ^c	New York	2.19	0.06	26(14,37)	1.07
Delaware	2.96	0.32	02(01,34)	36.53 ^c	North Carolina	2.20	0.09	25(11,40)	1.34
D.C.	3.07	0.43	01(01,40)	41.45 ^c	North Dakota	1.87	0.29	42(04,51)	-14.04
Florida	2.04	0.05	32(24,43)	-6.21	Ohio	2.52	0.08	10(03,22)	16.16 ^c
Georgia	2.20	0.09	24(10,40)	1.47	Oklahoma	2.27	0.14	18(05,43)	4.68
Hawaii	0.98	0.14	51(50,51)	-54.68 ^c	Oregon	2.14	0.13	30(09,47)	-1.47
Idaho	1.81	0.20	45(12,50)	-16.63 ^c	Pennsylvania	2.31	0.07	17(09,33)	6.33
Illinois	2.25	0.08	22(10,37)	3.46	Rhode Island	2.67	0.27	06(01,42)	22.92 ^c
Indiana	2.35	0.11	16(05,36)	8.45	South Carolina	2.25	0.13	21(06,42)	3.79
Iowa	2.03	0.14	33(12,49)	-6.71	South Dakota	1.76	0.25	50(07,51)	-19.15 ^c
Kansas	2.02	0.15	34(10,50)	-6.92	Tennessee	1.99	0.10	38(18,49)	-8.52
Kentucky	2.38	0.13	15(03,37)	9.57	Texas	1.81	0.05	46(37,50)	-16.82 ^c
Louisiana	2.27	0.13	19(05,42)	4.40	Utah	1.77	0.18	48(17,50)	-18.36 ^c
Maine	2.78	0.24	04(01,32)	27.90 ^c	Vermont	2.93	0.38	03(01,42)	35.06 ^c
Maryland	2.47	0.12	13(03,32)	13.79	Virginia	2.19	0.10	27(10,42)	0.93
Massachusetts	2.73	0.11	05(01,16)	25.72 ^c	Washington	2.38	0.11	14(04,34)	9.80
Michigan	2.48	0.09	12(04,25)	14.14	West Virginia	2.61	0.20	07(01,36)	20.21 ^c
Minnesota	1.99	0.11	37(17,49)	-8.18	Wisconsin	2.16	0.11	29(10,44)	-0.38
Mississippi	1.86	0.14	43(19,50)	-14.23	Wyoming	2.12	0.37	31(01,50)	-2.19
Missouri	2.26	0.11	20(07,39)	4.32					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.