

# SEER Incidence\* and US Death Rates,# 2000-2003 5-Year Relative Survival Rates,^ 1996-2002 All Cancer Combined, by Race and Sex




Figure 1-9


\* Incidence rates are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey) and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

# Death rates are from the NCHS public use data file for the total US and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

^ Survival rates are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2002. The remaining 13 SEER Areas contribute cases for the entire period 1996-2002. Relative survival rates are expressed as percents.

Figure I-11


## 5-Year Relative Survival Rates SEER Program, 1996-2002 Both Sexes, by Race



Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2002. The remaining 13 SEER Areas contribute cases for the entire period 1996-2002.

\* Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

# SEER Cancer Incidence and US Death Rates, 2000-2003 By Cancer Site and Race



Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey) and NCHS public use data file for the total US.

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

\* Incidence data for American Indians/Alaska Natives include cases from Connecticut, Detroit, Iowa, New Mexico, Seattle, Utah, Atlanta, and the Alaska Native Registry for the time period 1999-2002. Mortality data for American Indians/Alaska Natives include the total US for the time period 2000-2003.

^ Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Mortality data for Hispanics exclude cases from Minnesota, New Hampshire, and North Dakota.