

Table I-17
 US AND SEER DEATH RATES BY PRIMARY CANCER SITE AND RACE/ETHNICITY, 1992-2001

| Site | | Total United States ^a | | | | | | | SEER 12 Areas ^a | | | | | | |
|--------------------------------|------------|----------------------------------|-------|-------|--------------------|------------------|-------------------|----------------------|----------------------------|-------|-------|--------------------|------------------|-------------------|----------------------|
| | | Total | White | Black | AI/AN ^b | API ^c | Hisp ^d | W-NHisp ^d | Total | White | Black | AI/AN ^b | API ^c | Hisp ^d | W-NHisp ^d |
| All Sites | Both Sexes | 205.2 | 201.8 | 261.5 | 136.2 | 126.6 | 138.1 | 204.5 | 191.2 | 191.5 | 250.9 | 143.7 | 137.7 | 137.8 | 196.5 |
| | Male | 260.6 | 254.3 | 364.7 | 168.1 | 157.9 | 177.1 | 257.9 | 236.7 | 235.2 | 337.0 | 172.8 | 173.7 | 173.0 | 240.3 |
| | Female | 169.9 | 168.6 | 200.1 | 115.1 | 103.0 | 112.4 | 170.5 | 162.4 | 164.5 | 198.9 | 123.5 | 110.0 | 115.7 | 169.3 |
| Oral Cavity & Pharynx | Both Sexes | 3.0 | 2.8 | 4.8 | 2.3 | 2.6 | 1.9 | 2.9 | 2.9 | 2.8 | 4.1 | 3.4 | 3.0 | 1.7 | 2.9 |
| | Male | 4.6 | 4.2 | 8.4 | 3.7 | 3.9 | 3.3 | 4.3 | 4.3 | 4.0 | 7.0 | 5.6 | 4.5 | 2.8 | 4.2 |
| | Female | 1.7 | 1.7 | 2.2 | 1.2 | 1.4 | 0.9 | 1.7 | 1.8 | 1.8 | 2.0 | 1.7 | 1.7 | 0.8 | 1.9 |
| Esophagus | Both Sexes | 4.3 | 4.0 | 7.5 | 2.7 | 2.2 | 2.5 | 4.1 | 4.1 | 4.0 | 6.6 | 3.3 | 2.4 | 2.3 | 4.1 |
| | Male | 7.6 | 7.1 | 13.2 | 4.7 | 3.7 | 4.5 | 7.3 | 7.0 | 6.9 | 11.1 | 5.6 | 4.3 | 4.3 | 7.1 |
| | Female | 1.8 | 1.6 | 3.5 | 1.1 | 0.9 | 0.9 | 1.6 | 1.8 | 1.7 | 3.5 | 1.6 | 0.9 | 0.8 | 1.8 |
| Stomach | Both Sexes | 5.0 | 4.4 | 9.7 | 5.5 | 10.0 | 7.4 | 4.1 | 5.7 | 4.8 | 9.2 | 7.9 | 11.0 | 8.7 | 4.2 |
| | Male | 7.2 | 6.4 | 14.3 | 7.6 | 13.0 | 10.1 | 6.0 | 8.1 | 6.9 | 13.3 | 11.1 | 14.6 | 11.9 | 6.1 |
| | Female | 3.5 | 3.0 | 6.7 | 4.1 | 7.7 | 5.5 | 2.8 | 4.0 | 3.3 | 6.3 | 5.5 | 8.2 | 6.5 | 2.8 |
| Colon & Rectum | Both Sexes | 21.8 | 21.3 | 28.8 | 13.7 | 13.6 | 14.2 | 21.4 | 20.2 | 19.9 | 27.6 | 15.4 | 15.3 | 14.1 | 20.3 |
| | Male | 26.6 | 26.1 | 35.0 | 16.1 | 16.4 | 18.1 | 26.2 | 24.4 | 24.3 | 33.1 | 16.6 | 18.7 | 18.3 | 24.6 |
| | Female | 18.5 | 18.0 | 24.9 | 12.0 | 11.4 | 11.5 | 18.0 | 17.1 | 16.9 | 24.2 | 14.4 | 12.5 | 11.3 | 17.2 |
| Liver & Intrahepatic Bile Duct | Both Sexes | 4.4 | 4.0 | 6.0 | 5.6 | 10.8 | 7.1 | 3.8 | 4.9 | 4.1 | 6.1 | 6.7 | 10.9 | 7.1 | 3.8 |
| | Male | 6.4 | 5.8 | 9.1 | 7.7 | 15.9 | 10.1 | 5.5 | 7.3 | 6.0 | 9.4 | 8.9 | 16.2 | 10.1 | 5.4 |
| | Female | 2.9 | 2.7 | 3.7 | 4.1 | 6.6 | 4.8 | 2.6 | 3.1 | 2.7 | 3.7 | 5.2 | 6.6 | 4.8 | 2.5 |
| Pancreas | Both Sexes | 10.6 | 10.3 | 14.5 | 6.2 | 7.5 | 8.4 | 10.3 | 10.4 | 10.2 | 14.8 | 7.2 | 8.1 | 8.5 | 10.3 |
| | Male | 12.3 | 12.0 | 16.6 | 6.2 | 8.5 | 9.5 | 12.0 | 11.8 | 11.7 | 16.2 | 7.7 | 9.4 | 9.1 | 11.9 |
| | Female | 9.2 | 8.9 | 12.9 | 6.0 | 6.7 | 7.5 | 8.9 | 9.2 | 9.0 | 13.5 | 6.7 | 7.1 | 7.9 | 9.1 |
| Larynx | Both Sexes | 1.5 | 1.3 | 2.9 | 1.1 | 0.5 | 1.1 | 1.3 | 1.3 | 1.2 | 2.7 | 1.1 | 0.5 | 0.9 | 1.2 |
| | Male | 2.7 | 2.5 | 5.8 | 2.0 | 1.0 | 2.3 | 2.4 | 2.3 | 2.2 | 5.2 | 2.3 | 1.0 | 1.8 | 2.2 |
| | Female | 0.5 | 0.5 | 1.0 | 0.5 | 0.1 | 0.2 | 0.5 | 0.5 | 0.5 | 0.9 | 0.3 | 0.1 | 0.2 | 0.5 |
| Lung & Bronchus | Both Sexes | 57.3 | 57.1 | 67.4 | 36.6 | 29.0 | 25.7 | 59.2 | 49.4 | 49.9 | 65.6 | 32.8 | 31.9 | 23.8 | 52.8 |
| | Male | 81.6 | 80.0 | 110.7 | 51.0 | 41.6 | 40.9 | 82.7 | 67.0 | 66.3 | 101.5 | 45.7 | 46.3 | 36.1 | 69.3 |
| | Female | 40.2 | 41.0 | 39.2 | 26.2 | 19.2 | 14.8 | 42.5 | 37.0 | 38.5 | 41.6 | 23.2 | 20.7 | 15.3 | 41.3 |
| Melanoma of the Skin | Both Sexes | 2.7 | 3.0 | 0.5 | 0.6 | 0.4 | 0.8 | 3.2 | 2.4 | 2.9 | 0.4 | 0.4 | 0.4 | 0.8 | 3.2 |
| | Male | 3.9 | 4.4 | 0.5 | 0.8 | 0.5 | 1.1 | 4.6 | 3.5 | 4.1 | 0.4 | 0.6 | 0.6 | 0.9 | 4.6 |
| | Female | 1.8 | 2.0 | 0.5 | 0.4 | 0.3 | 0.6 | 2.1 | 1.6 | 1.9 | 0.4 | 0.3 | 0.3 | 0.6 | 2.1 |
| Breast | Female | 28.8 | 28.3 | 36.4 | 14.5 | 12.9 | 17.9 | 28.6 | 27.8 | 28.3 | 36.7 | 16.6 | 14.9 | 17.7 | 29.4 |

^a NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 U. S. standard population by 5-year age groups.

^b American Indian/Alaskan Native.

^c Asian/Pacific Islander.

^d Mortality data for Hispanics (Hisp) and White Non-Hispanics (W-NHisp) do not include cases from Connecticut, Maine, Maryland, New Hampshire, New York, North Dakota, Oklahoma and Vermont. Hispanic is not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives.

- Statistic could not be calculated.

Table I-17 - continued
 US AND SEER DEATH RATES BY PRIMARY CANCER SITE AND RACE/ETHNICITY, 1992-2001

| Site | | Total United States ^a | | | | | | | SEER 12 Areas ^a | | | | | | |
|------------------------|------------|----------------------------------|-------|-------|--------------------|------------------|-----------------------|--------------------------|----------------------------|-------|-------|--------------------|------------------|-----------------------|--------------------------|
| | | Total | White | Black | AI/AN ^b | API ^c | Hispanic ^d | W-NHispanic ^d | Total | White | Black | AI/AN ^b | API ^c | Hispanic ^d | W-NHispanic ^d |
| Cervix | Female | 3.1 | 2.7 | 6.3 | 3.2 | 3.0 | 3.9 | 2.6 | 2.7 | 2.5 | 4.8 | 2.7 | 2.8 | 4.0 | 2.3 |
| Corpus & Uterus, NOS | Female | 4.1 | 3.9 | 7.0 | 2.5 | 2.2 | 3.2 | 3.9 | 4.1 | 4.0 | 6.3 | 2.5 | 2.5 | 3.2 | 4.1 |
| Ovary | Female | 9.0 | 9.3 | 7.6 | 5.1 | 4.7 | 6.3 | 9.4 | 9.0 | 9.6 | 7.0 | 6.0 | 5.0 | 6.5 | 10.1 |
| Prostate | Male | 34.6 | 31.8 | 74.9 | 22.3 | 14.7 | 25.0 | 32.1 | 32.5 | 31.6 | 67.8 | 18.4 | 15.7 | 25.1 | 32.4 |
| Testis | Male | 0.3 | 0.3 | 0.1 | 0.2 | 0.1 | 0.3 | 0.3 | 0.3 | 0.3 | 0.1 | 0.4 | 0.1 | 0.3 | 0.3 |
| Urinary Bladder | Both Sexes | 4.4 | 4.5 | 4.0 | 1.6 | 1.8 | 2.4 | 4.5 | 4.1 | 4.3 | 4.0 | 1.6 | 2.0 | 2.3 | 4.4 |
| | Male | 7.7 | 8.0 | 5.9 | 2.5 | 2.9 | 4.2 | 8.1 | 7.0 | 7.6 | 5.9 | 2.6 | 3.1 | 3.6 | 7.9 |
| | Female | 2.3 | 2.3 | 3.0 | 1.0 | 1.1 | 1.3 | 2.3 | 2.2 | 2.3 | 2.9 | 1.0 | 1.1 | 1.4 | 2.3 |
| Kidney & Renal Pelvis | Both Sexes | 4.2 | 4.3 | 4.2 | 4.8 | 1.9 | 3.8 | 4.3 | 3.9 | 4.0 | 4.0 | 5.4 | 2.2 | 3.7 | 4.1 |
| | Male | 6.2 | 6.2 | 6.2 | 6.7 | 2.7 | 5.5 | 6.3 | 5.6 | 5.9 | 5.7 | 7.5 | 3.2 | 5.3 | 5.9 |
| | Female | 2.8 | 2.9 | 2.8 | 3.4 | 1.2 | 2.5 | 2.9 | 2.6 | 2.7 | 2.7 | 3.9 | 1.3 | 2.5 | 2.7 |
| Brain & Nervous System | Both Sexes | 4.7 | 5.0 | 2.7 | 2.0 | 1.9 | 2.9 | 5.2 | 4.5 | 5.0 | 2.9 | 2.0 | 2.2 | 3.0 | 5.3 |
| | Male | 5.7 | 6.1 | 3.3 | 2.5 | 2.3 | 3.5 | 6.3 | 5.6 | 6.2 | 3.4 | 2.5 | 2.6 | 3.5 | 6.6 |
| | Female | 3.8 | 4.1 | 2.3 | 1.6 | 1.6 | 2.5 | 4.3 | 3.6 | 4.0 | 2.5 | 1.7 | 1.8 | 2.6 | 4.3 |
| Thyroid | Both Sexes | 0.5 | 0.5 | 0.4 | 0.3 | 0.7 | 0.6 | 0.4 | 0.5 | 0.5 | 0.3 | 0.5 | 0.8 | 0.8 | 0.4 |
| | Male | 0.4 | 0.4 | 0.3 | 0.2 | 0.5 | 0.4 | 0.4 | 0.4 | 0.4 | 0.3 | 0.4 | 0.6 | 0.6 | 0.4 |
| | Female | 0.5 | 0.5 | 0.5 | 0.4 | 0.8 | 0.7 | 0.4 | 0.5 | 0.5 | 0.4 | 0.6 | 0.9 | 1.0 | 0.4 |
| Hodgkin Lymphoma | Both Sexes | 0.5 | 0.5 | 0.5 | 0.2 | 0.2 | 0.6 | 0.5 | 0.5 | 0.6 | 0.5 | 0.1 | 0.2 | 0.6 | 0.6 |
| | Male | 0.7 | 0.7 | 0.6 | 0.2 | 0.2 | 0.8 | 0.7 | 0.7 | 0.7 | 0.7 | 0.1 | 0.2 | 0.8 | 0.7 |
| | Female | 0.4 | 0.4 | 0.4 | 0.2 | 0.1 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.1 | 0.1 | 0.5 | 0.4 |
| Non-Hodgkin Lymphoma | Both Sexes | 8.4 | 8.8 | 5.8 | 4.6 | 5.3 | 6.5 | 8.9 | 8.2 | 8.7 | 5.7 | 4.2 | 5.7 | 6.5 | 8.9 |
| | Male | 10.5 | 10.9 | 7.5 | 5.4 | 6.8 | 8.1 | 11.0 | 10.2 | 10.8 | 7.2 | 5.5 | 7.6 | 8.2 | 11.1 |
| | Female | 6.9 | 7.2 | 4.5 | 4.0 | 4.2 | 5.3 | 7.3 | 6.7 | 7.2 | 4.7 | 3.2 | 4.3 | 5.3 | 7.3 |
| Myeloma | Both Sexes | 3.9 | 3.6 | 7.6 | 3.0 | 1.8 | 3.1 | 3.6 | 3.8 | 3.7 | 7.6 | 3.2 | 1.9 | 3.4 | 3.8 |
| | Male | 4.8 | 4.5 | 9.2 | 3.5 | 2.2 | 3.8 | 4.5 | 4.7 | 4.7 | 8.6 | 4.0 | 2.3 | 4.1 | 4.8 |
| | Female | 3.2 | 2.9 | 6.6 | 2.7 | 1.5 | 2.7 | 3.0 | 3.2 | 3.0 | 7.0 | 2.6 | 1.6 | 2.9 | 3.1 |
| Leukemia | Both Sexes | 7.8 | 7.9 | 7.0 | 4.1 | 4.3 | 5.3 | 8.0 | 7.5 | 7.8 | 7.1 | 3.6 | 4.8 | 5.4 | 8.0 |
| | Male | 10.4 | 10.7 | 9.4 | 5.2 | 5.5 | 6.7 | 10.8 | 10.0 | 10.5 | 9.2 | 4.6 | 6.2 | 6.6 | 10.9 |
| | Female | 6.0 | 6.1 | 5.5 | 3.4 | 3.4 | 4.4 | 6.1 | 5.8 | 6.0 | 5.7 | 3.0 | 3.8 | 4.5 | 6.1 |

^a NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 U. S. standard population by 5-year age groups.

^b American Indian/Alaskan Native.

^c Asian/Pacific Islander.

^d Mortality data for Hispanics (Hispanic) and White Non-Hispanics (W-NHispanic) do not include cases from Connecticut, Maine, Maryland, New Hampshire, New York, North Dakota, Oklahoma and Vermont. Hispanic is not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives.

- Statistic could not be calculated.