


Table 2.18  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	181.32	0.11		
High Five States				
Kentucky	213.58 <sup>b</sup>	0.99	(01)	17.79 <sup>c</sup>
Louisiana	208.41 <sup>b</sup>	0.98	(02)	14.94 <sup>c</sup>
West Virginia	207.83 <sup>b</sup>	1.37	(03)	14.62 <sup>c</sup>
Mississippi	206.82 <sup>b</sup>	1.19	(04)	14.06 <sup>c</sup>
Tennessee	202.83 <sup>b</sup>	0.80	(05)	11.86 <sup>c</sup>
Low Five States				
New Mexico	160.84 <sup>b</sup>	1.28	(47)	-11.29 <sup>c</sup>
Arizona	156.15 <sup>b</sup>	0.70	(48)	-13.88 <sup>c</sup>
Colorado	156.09 <sup>b</sup>	0.88	(49)	-13.91 <sup>c</sup>
Hawaii	149.15 <sup>b</sup>	1.44	(50)	-17.74 <sup>c</sup>
Utah	131.78 <sup>b</sup>	1.18	(51)	-27.32 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	199.86 <sup>b</sup>	0.90	(07)	10.23 <sup>c</sup>	Montana	175.70	1.81	(36)	-3.10
Alaska	181.15	3.11	(30)	-0.09	Nebraska	175.43 <sup>b</sup>	1.36	(37)	-3.25
Arizona	156.15 <sup>b</sup>	0.70	(48)	-13.88 <sup>c</sup>	Nevada	186.14 <sup>b</sup>	1.30	(21)	2.66
Arkansas	201.70 <sup>b</sup>	1.14	(06)	11.24 <sup>c</sup>	New Hampshire	184.17	1.64	(24)	1.57
California	165.13 <sup>b</sup>	0.32	(46)	-8.93 <sup>c</sup>	New Jersey	182.57	0.63	(28)	0.69
Colorado	156.09 <sup>b</sup>	0.88	(49)	-13.91 <sup>c</sup>	New Mexico	160.84 <sup>b</sup>	1.28	(47)	-11.29 <sup>c</sup>
Connecticut	176.91 <sup>b</sup>	0.96	(35)	-2.43	New York	169.64 <sup>b</sup>	0.40	(44)	-6.44 <sup>c</sup>
Delaware	196.62 <sup>b</sup>	2.06	(11)	8.44 <sup>c</sup>	North Carolina	189.55 <sup>b</sup>	0.65	(17)	4.54
D.C.	198.33 <sup>b</sup>	2.63	(08)	9.38 <sup>c</sup>	North Dakota	172.97 <sup>b</sup>	2.17	(39)	-4.60
Florida	172.49 <sup>b</sup>	0.39	(40)	-4.87	Ohio	197.23 <sup>b</sup>	0.56	(09)	8.77 <sup>c</sup>
Georgia	183.14	0.69	(25)	1.00	Oklahoma	195.87 <sup>b</sup>	1.01	(13)	8.03 <sup>c</sup>
Hawaii	149.15 <sup>b</sup>	1.44	(50)	-17.74 <sup>c</sup>	Oregon	183.02	0.96	(26)	0.94
Idaho	167.84 <sup>b</sup>	1.55	(45)	-7.43 <sup>c</sup>	Pennsylvania	190.25 <sup>b</sup>	0.50	(16)	4.92
Illinois	189.29 <sup>b</sup>	0.55	(19)	4.40	Rhode Island	184.95	1.75	(23)	2.00
Indiana	197.18 <sup>b</sup>	0.78	(10)	8.75 <sup>c</sup>	South Carolina	191.08 <sup>b</sup>	0.92	(15)	5.38 <sup>c</sup>
Iowa	180.52	1.02	(32)	-0.44	South Dakota	172.37 <sup>b</sup>	1.96	(41)	-4.93
Kansas	180.75	1.11	(31)	-0.32	Tennessee	202.83 <sup>b</sup>	0.80	(05)	11.86 <sup>c</sup>
Kentucky	213.58 <sup>b</sup>	0.99	(01)	17.79 <sup>c</sup>	Texas	174.75 <sup>b</sup>	0.42	(38)	-3.62
Louisiana	208.41 <sup>b</sup>	0.98	(02)	14.94 <sup>c</sup>	Utah	131.78 <sup>b</sup>	1.18	(51)	-27.32 <sup>c</sup>
Maine	195.96 <sup>b</sup>	1.58	(12)	8.08 <sup>c</sup>	Vermont	178.44	2.28	(34)	-1.59
Maryland	186.82 <sup>b</sup>	0.83	(20)	3.03	Virginia	185.65 <sup>b</sup>	0.71	(22)	2.39
Massachusetts	183.01	0.72	(27)	0.93	Washington	178.59 <sup>b</sup>	0.76	(33)	-1.51
Michigan	189.30 <sup>b</sup>	0.60	(18)	4.40	West Virginia	207.83 <sup>b</sup>	1.37	(03)	14.62 <sup>c</sup>
Minnesota	171.47 <sup>b</sup>	0.81	(42)	-5.43 <sup>c</sup>	Wisconsin	181.48	0.78	(29)	0.09
Mississippi	206.82 <sup>b</sup>	1.19	(04)	14.06 <sup>c</sup>	Wyoming	171.02 <sup>b</sup>	2.57	(43)	-5.68 <sup>c</sup>
Missouri	194.46 <sup>b</sup>	0.78	(14)	7.25 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 2.19  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	223.04	0.19		
High Five States				
Mississippi	276.12 <sup>b</sup>	2.18	(01)	23.80 <sup>c</sup>
Kentucky	271.17 <sup>b</sup>	1.75	(02)	21.58 <sup>c</sup>
Louisiana	268.14 <sup>b</sup>	1.76	(03)	20.22 <sup>c</sup>
Alabama	261.97 <sup>b</sup>	1.63	(04)	17.45 <sup>c</sup>
Tennessee	261.08 <sup>b</sup>	1.44	(05)	17.05 <sup>c</sup>
Low Five States				
New Mexico	193.02 <sup>b</sup>	2.13	(47)	-13.46 <sup>c</sup>
Colorado	187.26 <sup>b</sup>	1.51	(48)	-16.04 <sup>c</sup>
Arizona	186.67 <sup>b</sup>	1.15	(49)	-16.31 <sup>c</sup>
Hawaii	186.22 <sup>b</sup>	2.42	(50)	-16.51 <sup>c</sup>
Utah	158.27 <sup>b</sup>	1.97	(51)	-29.04 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	261.97 <sup>b</sup>	1.63	(04)	17.45 <sup>c</sup>	Montana	208.07 <sup>b</sup>	2.98	(42)	-6.71 <sup>c</sup>
Alaska	212.35	5.20	(38)	-4.79	Nebraska	217.13	2.33	(32)	-2.65
Arizona	186.67 <sup>b</sup>	1.15	(49)	-16.31 <sup>c</sup>	Nevada	214.66 <sup>b</sup>	2.08	(34)	-3.76
Arkansas	254.87 <sup>b</sup>	1.97	(08)	14.27 <sup>c</sup>	New Hampshire	223.41	2.82	(27)	0.17
California	197.45 <sup>b</sup>	0.54	(46)	-11.48 <sup>c</sup>	New Jersey	218.49 <sup>b</sup>	1.08	(29)	-2.04
Colorado	187.26 <sup>b</sup>	1.51	(48)	-16.04 <sup>c</sup>	New Mexico	193.02 <sup>b</sup>	2.13	(47)	-13.46 <sup>c</sup>
Connecticut	216.40 <sup>b</sup>	1.66	(33)	-2.98	New York	204.58 <sup>b</sup>	0.70	(43)	-8.28 <sup>c</sup>
Delaware	238.46 <sup>b</sup>	3.51	(16)	6.91 <sup>c</sup>	North Carolina	241.37 <sup>b</sup>	1.17	(15)	8.22 <sup>c</sup>
D.C.	260.44 <sup>b</sup>	4.89	(06)	16.77 <sup>c</sup>	North Dakota	212.84	3.63	(37)	-4.58
Florida	209.42 <sup>b</sup>	0.64	(40)	-6.11 <sup>c</sup>	Ohio	246.49 <sup>b</sup>	0.99	(10)	10.51 <sup>c</sup>
Georgia	237.08 <sup>b</sup>	1.28	(17)	6.29 <sup>c</sup>	Oklahoma	245.45 <sup>b</sup>	1.75	(12)	10.04 <sup>c</sup>
Hawaii	186.22 <sup>b</sup>	2.42	(50)	-16.51 <sup>c</sup>	Oregon	217.68	1.60	(31)	-2.41
Idaho	197.85 <sup>b</sup>	2.54	(45)	-11.29 <sup>c</sup>	Pennsylvania	235.61 <sup>b</sup>	0.87	(18)	5.63 <sup>c</sup>
Illinois	233.25 <sup>b</sup>	0.96	(20)	4.58	Rhode Island	234.39 <sup>b</sup>	3.10	(19)	5.09 <sup>c</sup>
Indiana	247.29 <sup>b</sup>	1.38	(09)	10.87 <sup>c</sup>	South Carolina	245.71 <sup>b</sup>	1.64	(11)	10.16 <sup>c</sup>
Iowa	224.69	1.75	(25)	0.74	South Dakota	214.17	3.30	(36)	-3.98
Kansas	224.66	1.91	(26)	0.72	Tennessee	261.08 <sup>b</sup>	1.44	(05)	17.05 <sup>c</sup>
Kentucky	271.17 <sup>b</sup>	1.75	(02)	21.58 <sup>c</sup>	Texas	217.82 <sup>b</sup>	0.74	(30)	-2.34
Louisiana	268.14 <sup>b</sup>	1.76	(03)	20.22 <sup>c</sup>	Utah	158.27 <sup>b</sup>	1.97	(51)	-29.04 <sup>c</sup>
Maine	243.44 <sup>b</sup>	2.74	(13)	9.15 <sup>c</sup>	Vermont	214.22	3.89	(35)	-3.95
Maryland	229.69 <sup>b</sup>	1.47	(23)	2.98	Virginia	232.72 <sup>b</sup>	1.27	(21)	4.34
Massachusetts	227.26	1.26	(24)	1.89	Washington	211.94 <sup>b</sup>	1.27	(39)	-4.98
Michigan	231.08 <sup>b</sup>	1.04	(22)	3.60	West Virginia	257.12 <sup>b</sup>	2.37	(07)	15.28 <sup>c</sup>
Minnesota	208.79 <sup>b</sup>	1.38	(41)	-6.39 <sup>c</sup>	Wisconsin	222.78	1.34	(28)	-0.12
Mississippi	276.12 <sup>b</sup>	2.18	(01)	23.80 <sup>c</sup>	Wyoming	199.36 <sup>b</sup>	4.22	(44)	-10.62 <sup>c</sup>
Missouri	242.03 <sup>b</sup>	1.36	(14)	8.51 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 2.20  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	153.25	0.13		
High Five States				
Kentucky	175.08 <sup>b</sup>	1.19	(01)	14.24 <sup>c</sup>
West Virginia	174.05 <sup>b</sup>	1.69	(02)	13.57 <sup>c</sup>
Louisiana	168.57 <sup>b</sup>	1.17	(03)	10.00 <sup>c</sup>
Delaware	167.54 <sup>b</sup>	2.54	(04)	9.33 <sup>c</sup>
Ohio	165.50 <sup>b</sup>	0.68	(05)	7.99 <sup>c</sup>
Low Five States				
New Mexico	136.85 <sup>b</sup>	1.59	(47)	-10.70 <sup>c</sup>
Colorado	135.65 <sup>b</sup>	1.09	(48)	-11.48 <sup>c</sup>
Arizona	132.43 <sup>b</sup>	0.88	(49)	-13.58 <sup>c</sup>
Hawaii	120.66 <sup>b</sup>	1.77	(50)	-21.27 <sup>c</sup>
Utah	112.39 <sup>b</sup>	1.47	(51)	-26.66 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	158.71 <sup>b</sup>	1.06	(22)	3.56	Montana	152.95	2.30	(32)	-0.19
Alaska	157.23	3.89	(23)	2.60	Nebraska	147.19 <sup>b</sup>	1.68	(40)	-3.95
Arizona	132.43 <sup>b</sup>	0.88	(49)	-13.58 <sup>c</sup>	Nevada	163.02 <sup>b</sup>	1.65	(10)	6.37 <sup>c</sup>
Arkansas	164.07 <sup>b</sup>	1.38	(08)	7.06 <sup>c</sup>	New Hampshire	159.10	2.04	(20)	3.82
California	143.44 <sup>b</sup>	0.40	(45)	-6.40 <sup>c</sup>	New Jersey	160.63 <sup>b</sup>	0.78	(18)	4.81
Colorado	135.65 <sup>b</sup>	1.09	(48)	-11.48 <sup>c</sup>	New Mexico	136.85 <sup>b</sup>	1.59	(47)	-10.70 <sup>c</sup>
Connecticut	152.49	1.18	(33)	-0.49	New York	147.95 <sup>b</sup>	0.50	(38)	-3.46
Delaware	167.54 <sup>b</sup>	2.54	(04)	9.33 <sup>c</sup>	North Carolina	155.47	0.78	(28)	1.45
D.C.	161.10	3.09	(16)	5.12 <sup>c</sup>	North Dakota	146.02	2.72	(41)	-4.72
Florida	143.87 <sup>b</sup>	0.48	(44)	-6.12 <sup>c</sup>	Ohio	165.50 <sup>b</sup>	0.68	(05)	7.99 <sup>c</sup>
Georgia	149.54 <sup>b</sup>	0.81	(37)	-2.42	Oklahoma	161.49 <sup>b</sup>	1.24	(14)	5.37 <sup>c</sup>
Hawaii	120.66 <sup>b</sup>	1.77	(50)	-21.27 <sup>c</sup>	Oregon	158.71 <sup>b</sup>	1.21	(21)	3.57
Idaho	145.67 <sup>b</sup>	1.97	(42)	-4.94	Pennsylvania	161.07 <sup>b</sup>	0.62	(17)	5.10 <sup>c</sup>
Illinois	162.03 <sup>b</sup>	0.67	(13)	5.73 <sup>c</sup>	Rhode Island	155.02	2.13	(29)	1.16
Indiana	164.82 <sup>b</sup>	0.95	(06)	7.55 <sup>c</sup>	South Carolina	153.86	1.09	(31)	0.40
Iowa	151.73	1.27	(34)	-0.99	South Dakota	142.67 <sup>b</sup>	2.43	(46)	-6.90 <sup>c</sup>
Kansas	151.32	1.37	(35)	-1.26	Tennessee	163.98 <sup>b</sup>	0.95	(09)	7.00 <sup>c</sup>
Kentucky	175.08 <sup>b</sup>	1.19	(01)	14.24 <sup>c</sup>	Texas	145.15 <sup>b</sup>	0.51	(43)	-5.29 <sup>c</sup>
Louisiana	168.57 <sup>b</sup>	1.17	(03)	10.00 <sup>c</sup>	Utah	112.39 <sup>b</sup>	1.47	(51)	-26.66 <sup>c</sup>
Maine	164.74 <sup>b</sup>	1.94	(07)	7.50 <sup>c</sup>	Vermont	155.49	2.85	(27)	1.46
Maryland	159.71 <sup>b</sup>	1.01	(19)	4.22	Virginia	155.50	0.86	(26)	1.47
Massachusetts	155.97	0.88	(24)	1.78	Washington	155.71	0.95	(25)	1.60
Michigan	162.05 <sup>b</sup>	0.74	(12)	5.74 <sup>c</sup>	West Virginia	174.05 <sup>b</sup>	1.69	(02)	13.57 <sup>c</sup>
Minnesota	147.63 <sup>b</sup>	1.01	(39)	-3.67	Wisconsin	154.34	0.97	(30)	0.71
Mississippi	161.37 <sup>b</sup>	1.39	(15)	5.30 <sup>c</sup>	Wyoming	150.71	3.28	(36)	-1.66
Missouri	162.66 <sup>b</sup>	0.96	(11)	6.14 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 3.18  
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.27	0.02		
High Five States				
Iowa	5.32 <sup>b</sup>	0.18	(01)	24.61 <sup>c</sup>
Washington	5.24 <sup>b</sup>	0.13	(02)	22.66 <sup>c</sup>
Idaho	5.22	0.27	(03)	22.29 <sup>c</sup>
Nebraska	5.15 <sup>b</sup>	0.24	(04)	20.62 <sup>c</sup>
Oregon	5.14 <sup>b</sup>	0.16	(05)	20.55 <sup>c</sup>
Low Five States				
Georgia	3.80 <sup>b</sup>	0.10	(47)	-10.97
New Jersey	3.74 <sup>b</sup>	0.09	(48)	-12.36
New York	3.67 <sup>b</sup>	0.06	(49)	-13.91
D.C.	3.28	0.34	(50)	-23.14 <sup>c</sup>
Hawaii	2.27 <sup>b</sup>	0.18	(51)	-46.90 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.44	0.14	(26)	4.10	Montana	4.74	0.30	(17)	11.01
Alaska	4.18	0.43	(36)	-2.15	Nebraska	5.15 <sup>b</sup>	0.24	(04)	20.62 <sup>c</sup>
Arizona	4.08	0.11	(40)	-4.45	Nevada	4.21	0.19	(34)	-1.30
Arkansas	4.92 <sup>b</sup>	0.18	(09)	15.28 <sup>c</sup>	New Hampshire	4.89	0.27	(10)	14.69
California	4.28	0.05	(33)	0.31	New Jersey	3.74 <sup>b</sup>	0.09	(48)	-12.36
Colorado	4.33	0.14	(32)	1.37	New Mexico	3.82	0.20	(46)	-10.54
Connecticut	4.11	0.15	(37)	-3.79	New York	3.67 <sup>b</sup>	0.06	(49)	-13.91
Delaware	4.45	0.31	(24)	4.21	North Carolina	4.10	0.10	(38)	-3.88
D.C.	3.28	0.34	(50)	-23.14 <sup>c</sup>	North Dakota	4.84	0.38	(12)	13.34
Florida	3.94 <sup>b</sup>	0.06	(42)	-7.62	Ohio	4.44	0.09	(25)	4.12
Georgia	3.80 <sup>b</sup>	0.10	(47)	-10.97	Oklahoma	4.74	0.16	(16)	11.12
Hawaii	2.27 <sup>b</sup>	0.18	(51)	-46.90 <sup>c</sup>	Oregon	5.14 <sup>b</sup>	0.16	(05)	20.55 <sup>c</sup>
Idaho	5.22	0.27	(03)	22.29 <sup>c</sup>	Pennsylvania	4.07	0.08	(41)	-4.60
Illinois	3.84 <sup>b</sup>	0.08	(44)	-9.94	Rhode Island	4.42	0.28	(29)	3.57
Indiana	4.65	0.12	(19)	8.90	South Carolina	4.43	0.14	(28)	3.72
Iowa	5.32 <sup>b</sup>	0.18	(01)	24.61 <sup>c</sup>	South Dakota	4.85	0.34	(11)	13.57
Kansas	5.13 <sup>b</sup>	0.19	(06)	20.24 <sup>c</sup>	Tennessee	5.04 <sup>b</sup>	0.13	(07)	18.11 <sup>c</sup>
Kentucky	4.38	0.14	(30)	2.56	Texas	4.18	0.06	(35)	-2.08
Louisiana	4.55	0.14	(22)	6.56	Utah	4.52	0.21	(23)	6.02
Maine	5.03	0.26	(08)	17.94 <sup>c</sup>	Vermont	4.59	0.37	(20)	7.48
Maryland	3.83 <sup>b</sup>	0.12	(45)	-10.24	Virginia	3.90 <sup>b</sup>	0.10	(43)	-8.68
Massachusetts	4.10	0.11	(39)	-4.01	Washington	5.24 <sup>b</sup>	0.13	(02)	22.66 <sup>c</sup>
Michigan	4.81 <sup>b</sup>	0.10	(13)	12.74	West Virginia	4.66	0.21	(18)	9.23
Minnesota	4.33	0.13	(31)	1.46	Wisconsin	4.77 <sup>b</sup>	0.13	(15)	11.74
Mississippi	4.78	0.18	(14)	11.93	Wyoming	4.44	0.41	(27)	3.92
Missouri	4.57	0.12	(21)	7.03					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 3.19  
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.22	0.03		
High Five States				
Iowa	6.76 <sup>b</sup>	0.30	(01)	29.66 <sup>c</sup>
Washington	6.46 <sup>b</sup>	0.21	(02)	23.76 <sup>c</sup>
Kansas	6.35 <sup>b</sup>	0.32	(03)	21.73 <sup>c</sup>
Oregon	6.29 <sup>b</sup>	0.26	(04)	20.52 <sup>c</sup>
New Hampshire	6.25	0.45	(05)	19.86 <sup>c</sup>
Low Five States				
Alaska	4.63	0.67	(47)	-11.33
Georgia	4.58 <sup>b</sup>	0.16	(48)	-12.20
Maryland	4.51 <sup>b</sup>	0.19	(49)	-13.65
D.C.	4.50	0.61	(50)	-13.71
Hawaii	2.92 <sup>b</sup>	0.30	(51)	-43.99 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.40	0.22	(25)	3.60	Montana	5.83	0.48	(12)	11.80
Alaska	4.63	0.67	(47)	-11.33	Nebraska	5.93	0.38	(09)	13.70
Arizona	4.78	0.18	(41)	-8.31	Nevada	5.18	0.30	(33)	-0.63
Arkansas	5.75	0.29	(16)	10.17	New Hampshire	6.25	0.45	(05)	19.86 <sup>c</sup>
California	5.29	0.08	(29)	1.31	New Jersey	4.65 <sup>b</sup>	0.15	(46)	-10.80
Colorado	5.15	0.23	(34)	-1.27	New Mexico	4.72	0.32	(43)	-9.49
Connecticut	4.79	0.24	(40)	-8.10	New York	4.70 <sup>b</sup>	0.10	(45)	-9.92
Delaware	5.49	0.52	(20)	5.23	North Carolina	5.12	0.16	(35)	-1.93
D.C.	4.50	0.61	(50)	-13.71	North Dakota	5.80	0.62	(13)	11.14
Florida	4.76 <sup>b</sup>	0.10	(42)	-8.70	Ohio	5.38	0.14	(27)	3.19
Georgia	4.58 <sup>b</sup>	0.16	(48)	-12.20	Oklahoma	5.75	0.26	(15)	10.23
Hawaii	2.92 <sup>b</sup>	0.30	(51)	-43.99 <sup>c</sup>	Oregon	6.29 <sup>b</sup>	0.26	(04)	20.52 <sup>c</sup>
Idaho	5.38	0.40	(28)	3.04	Pennsylvania	4.97	0.13	(38)	-4.76
Illinois	4.71 <sup>b</sup>	0.13	(44)	-9.66	Rhode Island	5.52	0.46	(18)	5.89
Indiana	5.48	0.20	(21)	4.97	South Carolina	5.47	0.24	(23)	4.77
Iowa	6.76 <sup>b</sup>	0.30	(01)	29.66 <sup>c</sup>	South Dakota	5.78	0.55	(14)	10.77
Kansas	6.35 <sup>b</sup>	0.32	(03)	21.73 <sup>c</sup>	Tennessee	6.17 <sup>b</sup>	0.21	(06)	18.25 <sup>c</sup>
Kentucky	5.39	0.24	(26)	3.29	Texas	4.94	0.10	(39)	-5.39
Louisiana	5.84	0.25	(10)	11.97	Utah	5.46	0.34	(24)	4.71
Maine	6.14	0.42	(07)	17.64 <sup>c</sup>	Vermont	5.28	0.58	(30)	1.16
Maryland	4.51 <sup>b</sup>	0.19	(49)	-13.65	Virginia	4.98	0.18	(37)	-4.57
Massachusetts	5.22	0.18	(31)	0.01	Washington	6.46 <sup>b</sup>	0.21	(02)	23.76 <sup>c</sup>
Michigan	6.01 <sup>b</sup>	0.16	(08)	15.15 <sup>c</sup>	West Virginia	5.50	0.34	(19)	5.50
Minnesota	5.20	0.21	(32)	-0.30	Wisconsin	5.84	0.21	(11)	11.90
Mississippi	5.73	0.30	(17)	9.77	Wyoming	5.11	0.64	(36)	-2.11
Missouri	5.47	0.20	(22)	4.81					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 3.20  
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.48	0.02		
High Five States				
Idaho	5.02 <sup>b</sup>	0.37	(01)	44.32 <sup>c</sup>
Nebraska	4.41	0.30	(02)	26.81 <sup>c</sup>
Arkansas	4.22	0.23	(03)	21.38 <sup>c</sup>
Washington	4.20 <sup>b</sup>	0.16	(04)	20.72 <sup>c</sup>
Oregon	4.18	0.20	(05)	20.35 <sup>c</sup>
Low Five States				
New Mexico	3.04	0.24	(47)	-12.60
New Jersey	2.98 <sup>b</sup>	0.11	(48)	-14.20
New York	2.86 <sup>b</sup>	0.07	(49)	-17.81 <sup>c</sup>
D.C.	2.26	0.38	(50)	-34.88 <sup>c</sup>
Hawaii	1.69 <sup>b</sup>	0.22	(51)	-51.50 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.63	0.16	(27)	4.48	Montana	3.77	0.37	(21)	8.47
Alaska	3.76	0.58	(22)	8.26	Nebraska	4.41	0.30	(02)	26.81 <sup>c</sup>
Arizona	3.47	0.15	(36)	-0.13	Nevada	3.30	0.23	(39)	-5.17
Arkansas	4.22	0.23	(03)	21.38 <sup>c</sup>	New Hampshire	3.72	0.32	(23)	7.08
California	3.43	0.06	(37)	-1.21	New Jersey	2.98 <sup>b</sup>	0.11	(48)	-14.20
Colorado	3.61	0.18	(29)	3.88	New Mexico	3.04	0.24	(47)	-12.60
Connecticut	3.50	0.19	(33)	0.69	New York	2.86 <sup>b</sup>	0.07	(49)	-17.81 <sup>c</sup>
Delaware	3.60	0.38	(30)	3.54	North Carolina	3.27	0.11	(40)	-5.99
D.C.	2.26	0.38	(50)	-34.88 <sup>c</sup>	North Dakota	3.95	0.47	(14)	13.57
Florida	3.23	0.08	(42)	-7.01	Ohio	3.65	0.10	(24)	5.13
Georgia	3.18	0.12	(44)	-8.57	Oklahoma	3.84	0.19	(16)	10.52
Hawaii	1.69 <sup>b</sup>	0.22	(51)	-51.50 <sup>c</sup>	Oregon	4.18	0.20	(05)	20.35 <sup>c</sup>
Idaho	5.02 <sup>b</sup>	0.37	(01)	44.32 <sup>c</sup>	Pennsylvania	3.35	0.09	(38)	-3.59
Illinois	3.16	0.10	(45)	-8.99	Rhode Island	3.56	0.34	(31)	2.49
Indiana	3.96	0.15	(12)	14.09	South Carolina	3.63	0.17	(26)	4.55
Iowa	4.10	0.22	(08)	18.12 <sup>c</sup>	South Dakota	4.08	0.43	(10)	17.53 <sup>c</sup>
Kansas	4.11	0.23	(07)	18.19 <sup>c</sup>	Tennessee	4.14 <sup>b</sup>	0.15	(06)	19.24 <sup>c</sup>
Kentucky	3.48	0.17	(35)	0.26	Texas	3.55	0.08	(32)	2.06
Louisiana	3.49	0.17	(34)	0.36	Utah	3.63	0.26	(25)	4.58
Maine	4.10	0.32	(09)	18.06 <sup>c</sup>	Vermont	3.89	0.47	(15)	11.89
Maryland	3.26	0.15	(41)	-6.22	Virginia	3.05	0.12	(46)	-12.16
Massachusetts	3.19	0.13	(43)	-8.24	Washington	4.20 <sup>b</sup>	0.16	(04)	20.72 <sup>c</sup>
Michigan	3.83	0.12	(18)	10.09	West Virginia	3.95	0.27	(13)	13.63
Minnesota	3.61	0.16	(28)	4.02	Wisconsin	3.81	0.16	(19)	9.73
Mississippi	4.01	0.22	(11)	15.48 <sup>c</sup>	Wyoming	3.80	0.53	(20)	9.47
Missouri	3.83	0.15	(17)	10.33					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 4.24  
Cancer of the Female Breast (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	23.47	0.05		
High Five States				
D.C.	27.64	1.28	(01)	17.79 <sup>c</sup>
Louisiana	26.81 <sup>b</sup>	0.47	(02)	14.23 <sup>c</sup>
New Jersey	26.45 <sup>b</sup>	0.32	(03)	12.71 <sup>c</sup>
Ohio	25.95 <sup>b</sup>	0.27	(04)	10.56 <sup>c</sup>
Maryland	25.57 <sup>b</sup>	0.40	(05)	8.96
Low Five States				
Idaho	21.19	0.75	(47)	-9.72
Arizona	20.97 <sup>b</sup>	0.35	(48)	-10.66 <sup>c</sup>
Montana	20.72	0.86	(49)	-11.71 <sup>c</sup>
Colorado	20.49 <sup>b</sup>	0.42	(50)	-12.69 <sup>c</sup>
Hawaii	17.80 <sup>b</sup>	0.69	(51)	-24.14 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	24.46	0.42	(11)	4.24	Montana	20.72	0.86	(49)	-11.71 <sup>c</sup>
Alaska	21.67	1.37	(43)	-7.65	Nebraska	21.98	0.66	(39)	-6.34
Arizona	20.97 <sup>b</sup>	0.35	(48)	-10.66 <sup>c</sup>	Nevada	23.54	0.62	(21)	0.29
Arkansas	23.96	0.53	(19)	2.11	New Hampshire	22.82	0.77	(27)	-2.76
California	22.53 <sup>b</sup>	0.16	(29)	-4.00	New Jersey	26.45 <sup>b</sup>	0.32	(03)	12.71 <sup>c</sup>
Colorado	20.49 <sup>b</sup>	0.42	(50)	-12.69 <sup>c</sup>	New Mexico	21.45	0.63	(46)	-8.60
Connecticut	23.25	0.46	(23)	-0.93	New York	23.08	0.20	(26)	-1.66
Delaware	24.26	0.98	(15)	3.37	North Carolina	24.35	0.31	(13)	3.77
D.C.	27.64	1.28	(01)	17.79 <sup>c</sup>	North Dakota	22.35	1.08	(32)	-4.78
Florida	21.89 <sup>b</sup>	0.19	(40)	-6.73	Ohio	25.95 <sup>b</sup>	0.27	(04)	10.56 <sup>c</sup>
Georgia	23.24	0.32	(24)	-0.99	Oklahoma	24.13	0.48	(17)	2.84
Hawaii	17.80 <sup>b</sup>	0.69	(51)	-24.14 <sup>c</sup>	Oregon	22.47	0.46	(30)	-4.24
Idaho	21.19	0.75	(47)	-9.72	Pennsylvania	24.85 <sup>b</sup>	0.25	(09)	5.88
Illinois	24.73 <sup>b</sup>	0.26	(10)	5.39	Rhode Island	22.18	0.81	(34)	-5.48
Indiana	24.04	0.36	(18)	2.42	South Carolina	24.26	0.44	(16)	3.36
Iowa	22.12	0.49	(35)	-5.73	South Dakota	21.78	0.96	(41)	-7.20
Kansas	23.10	0.54	(25)	-1.58	Tennessee	24.45	0.37	(12)	4.20
Kentucky	23.53	0.44	(22)	0.28	Texas	22.59 <sup>b</sup>	0.20	(28)	-3.74
Louisiana	26.81 <sup>b</sup>	0.47	(02)	14.23 <sup>c</sup>	Utah	22.09	0.65	(36)	-5.86
Maine	21.50	0.70	(45)	-8.37	Vermont	21.69	1.06	(42)	-7.59
Maryland	25.57 <sup>b</sup>	0.40	(05)	8.96	Virginia	25.06 <sup>b</sup>	0.34	(08)	6.77
Massachusetts	22.25 <sup>b</sup>	0.33	(33)	-5.18	Washington	22.43	0.36	(31)	-4.41
Michigan	24.35	0.29	(14)	3.77	West Virginia	23.93	0.63	(20)	1.99
Minnesota	21.56 <sup>b</sup>	0.39	(44)	-8.13	Wisconsin	22.07 <sup>b</sup>	0.37	(38)	-5.97
Mississippi	25.49 <sup>b</sup>	0.56	(06)	8.62	Wyoming	22.09	1.26	(37)	-5.88
Missouri	25.42 <sup>b</sup>	0.38	(07)	8.32					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 5.14  
Cancer of the Cervix Uteri (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.42	0.02		
High Five States				
Mississippi	4.03 <sup>b</sup>	0.23	(01)	66.95 <sup>c</sup>
Louisiana	3.35 <sup>b</sup>	0.17	(02)	38.55 <sup>c</sup>
Arkansas	3.24 <sup>b</sup>	0.21	(03)	34.34 <sup>c</sup>
D.C.	3.19	0.45	(04)	32.00 <sup>c</sup>
Alabama	3.07 <sup>b</sup>	0.15	(05)	27.06 <sup>c</sup>
Low Five States				
Rhode Island	1.61	0.23	(47)	-33.41 <sup>c</sup>
Minnesota	1.57 <sup>b</sup>	0.11	(48)	-34.83 <sup>c</sup>
Vermont	1.48	0.28	(49)	-38.65 <sup>c</sup>
Massachusetts	1.39 <sup>b</sup>	0.09	(50)	-42.60 <sup>c</sup>
Utah	1.31 <sup>b</sup>	0.16	(51)	-45.77 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.07 <sup>b</sup>	0.15	(05)	27.06 <sup>c</sup>	Montana	1.87	0.27	(39)	-22.41 <sup>c</sup>
Alaska	2.21	0.41	(27)	-8.35	Nebraska	1.80	0.20	(43)	-25.37 <sup>c</sup>
Arizona	1.98 <sup>b</sup>	0.11	(38)	-18.04 <sup>c</sup>	Nevada	2.71	0.21	(12)	12.16
Arkansas	3.24 <sup>b</sup>	0.21	(03)	34.34 <sup>c</sup>	New Hampshire	2.20	0.24	(28)	-8.96
California	2.36	0.05	(23)	-2.23	New Jersey	2.47	0.10	(19)	2.14
Colorado	1.84 <sup>b</sup>	0.12	(40)	-23.80 <sup>c</sup>	New Mexico	2.64	0.23	(15)	9.17
Connecticut	1.67 <sup>b</sup>	0.13	(45)	-30.66 <sup>c</sup>	New York	2.43	0.07	(21)	0.68
Delaware	2.24	0.31	(26)	-7.24	North Carolina	2.37	0.10	(22)	-1.86
D.C.	3.19	0.45	(04)	32.00 <sup>c</sup>	North Dakota	1.61	0.31	(46)	-33.30 <sup>c</sup>
Florida	2.66	0.07	(14)	10.25	Ohio	2.49	0.09	(18)	3.23
Georgia	2.73	0.11	(10)	12.87	Oklahoma	2.89	0.18	(07)	19.48 <sup>c</sup>
Hawaii	1.99	0.24	(36)	-17.44 <sup>c</sup>	Oregon	1.83 <sup>b</sup>	0.14	(41)	-24.34 <sup>c</sup>
Idaho	2.04	0.24	(33)	-15.35 <sup>c</sup>	Pennsylvania	2.26	0.08	(25)	-6.50
Illinois	2.70	0.09	(13)	11.74	Rhode Island	1.61	0.23	(47)	-33.41 <sup>c</sup>
Indiana	2.46	0.12	(20)	1.91	South Carolina	2.72	0.15	(11)	12.42
Iowa	2.11	0.16	(31)	-12.53	South Dakota	1.70	0.29	(44)	-29.42 <sup>c</sup>
Kansas	2.14	0.18	(30)	-11.43	Tennessee	2.86	0.13	(09)	18.32 <sup>c</sup>
Kentucky	2.64	0.15	(16)	9.12	Texas	3.05 <sup>b</sup>	0.07	(06)	26.43 <sup>c</sup>
Louisiana	3.35 <sup>b</sup>	0.17	(02)	38.55 <sup>c</sup>	Utah	1.31 <sup>b</sup>	0.16	(51)	-45.77 <sup>c</sup>
Maine	1.98	0.22	(37)	-18.01 <sup>c</sup>	Vermont	1.48	0.28	(49)	-38.65 <sup>c</sup>
Maryland	2.31	0.12	(24)	-4.30	Virginia	2.14	0.10	(29)	-11.21
Massachusetts	1.39 <sup>b</sup>	0.09	(50)	-42.60 <sup>c</sup>	Washington	2.00 <sup>b</sup>	0.11	(35)	-17.19 <sup>c</sup>
Michigan	2.03 <sup>b</sup>	0.09	(34)	-15.96 <sup>c</sup>	West Virginia	2.88	0.24	(08)	19.08 <sup>c</sup>
Minnesota	1.57 <sup>b</sup>	0.11	(48)	-34.83 <sup>c</sup>	Wisconsin	1.82 <sup>b</sup>	0.11	(42)	-24.60 <sup>c</sup>
Mississippi	4.03 <sup>b</sup>	0.23	(01)	66.95 <sup>c</sup>	Wyoming	2.11	0.41	(32)	-12.73
Missouri	2.62	0.13	(17)	8.51					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 6.25  
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	17.15	0.03		
High Five States				
Louisiana	20.28 <sup>b</sup>	0.31	(01)	18.27 <sup>c</sup>
Mississippi	20.22 <sup>b</sup>	0.37	(02)	17.88 <sup>c</sup>
Kentucky	20.11 <sup>b</sup>	0.30	(03)	17.29 <sup>c</sup>
D.C.	20.11 <sup>b</sup>	0.83	(04)	17.25 <sup>c</sup>
West Virginia	20.08 <sup>b</sup>	0.43	(05)	17.07 <sup>c</sup>
Low Five States				
Minnesota	15.20 <sup>b</sup>	0.24	(47)	-11.39 <sup>c</sup>
Idaho	14.78 <sup>b</sup>	0.46	(48)	-13.79 <sup>c</sup>
Arizona	14.46 <sup>b</sup>	0.21	(49)	-15.70 <sup>c</sup>
Hawaii	14.38 <sup>b</sup>	0.45	(50)	-16.12 <sup>c</sup>
Utah	12.20 <sup>b</sup>	0.36	(51)	-28.84 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	18.71 <sup>b</sup>	0.28	(11)	9.13	Montana	15.54	0.54	(43)	-9.38
Alaska	17.24	0.99	(26)	0.50	Nebraska	18.82 <sup>b</sup>	0.44	(09)	9.75
Arizona	14.46 <sup>b</sup>	0.21	(49)	-15.70 <sup>c</sup>	Nevada	18.72 <sup>b</sup>	0.42	(10)	9.17
Arkansas	18.91 <sup>b</sup>	0.35	(08)	10.30 <sup>c</sup>	New Hampshire	16.74	0.49	(35)	-2.38
California	15.50 <sup>b</sup>	0.10	(44)	-9.64	New Jersey	18.70 <sup>b</sup>	0.20	(12)	9.04
Colorado	15.50 <sup>b</sup>	0.28	(45)	-9.64	New Mexico	16.16	0.41	(39)	-5.78
Connecticut	15.59 <sup>b</sup>	0.28	(42)	-9.10	New York	16.83	0.13	(32)	-1.87
Delaware	17.70	0.62	(20)	3.19	North Carolina	16.82	0.19	(33)	-1.89
D.C.	20.11 <sup>b</sup>	0.83	(04)	17.25 <sup>c</sup>	North Dakota	17.64	0.68	(21)	2.89
Florida	15.78 <sup>b</sup>	0.12	(41)	-7.99	Ohio	18.99 <sup>b</sup>	0.17	(07)	10.73 <sup>c</sup>
Georgia	16.86	0.21	(30)	-1.67	Oklahoma	18.49 <sup>b</sup>	0.31	(16)	7.80
Hawaii	14.38 <sup>b</sup>	0.45	(50)	-16.12 <sup>c</sup>	Oregon	16.33	0.29	(38)	-4.77
Idaho	14.78 <sup>b</sup>	0.46	(48)	-13.79 <sup>c</sup>	Pennsylvania	18.61 <sup>b</sup>	0.16	(14)	8.54
Illinois	19.03 <sup>b</sup>	0.17	(06)	11.00 <sup>c</sup>	Rhode Island	16.35	0.52	(37)	-4.67
Indiana	18.69 <sup>b</sup>	0.24	(13)	8.98	South Carolina	17.32	0.28	(25)	0.99
Iowa	17.96	0.32	(19)	4.75	South Dakota	17.16	0.61	(29)	0.06
Kansas	17.63	0.35	(23)	2.83	Tennessee	18.59 <sup>b</sup>	0.24	(15)	8.41
Kentucky	20.11 <sup>b</sup>	0.30	(03)	17.29 <sup>c</sup>	Texas	16.60 <sup>b</sup>	0.13	(36)	-3.22
Louisiana	20.28 <sup>b</sup>	0.31	(01)	18.27 <sup>c</sup>	Utah	12.20 <sup>b</sup>	0.36	(51)	-28.84 <sup>c</sup>
Maine	17.64	0.47	(22)	2.89	Vermont	17.23	0.71	(27)	0.49
Maryland	18.19 <sup>b</sup>	0.26	(17)	6.07	Virginia	17.16	0.22	(28)	0.07
Massachusetts	16.75	0.22	(34)	-2.34	Washington	15.36 <sup>b</sup>	0.22	(46)	-10.41 <sup>c</sup>
Michigan	17.42	0.18	(24)	1.57	West Virginia	20.08 <sup>b</sup>	0.43	(05)	17.07 <sup>c</sup>
Minnesota	15.20 <sup>b</sup>	0.24	(47)	-11.39 <sup>c</sup>	Wisconsin	16.07 <sup>b</sup>	0.23	(40)	-6.30
Mississippi	20.22 <sup>b</sup>	0.37	(02)	17.88 <sup>c</sup>	Wyoming	16.85	0.81	(31)	-1.74
Missouri	18.04 <sup>b</sup>	0.24	(18)	5.19					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 6.26  
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	20.69	0.06		
High Five States				
Louisiana	25.82 <sup>b</sup>	0.55	(01)	24.82 <sup>c</sup>
Mississippi	25.19 <sup>b</sup>	0.66	(02)	21.80 <sup>c</sup>
Kentucky	24.42 <sup>b</sup>	0.53	(03)	18.04 <sup>c</sup>
West Virginia	24.35 <sup>b</sup>	0.73	(04)	17.72 <sup>c</sup>
Alabama	23.60 <sup>b</sup>	0.49	(05)	14.10 <sup>c</sup>
Low Five States				
Connecticut	18.09 <sup>b</sup>	0.48	(47)	-12.53 <sup>c</sup>
Montana	17.52 <sup>b</sup>	0.86	(48)	-15.31 <sup>c</sup>
Arizona	17.48 <sup>b</sup>	0.35	(49)	-15.49 <sup>c</sup>
Idaho	15.91 <sup>b</sup>	0.72	(50)	-23.10 <sup>c</sup>
Utah	14.60 <sup>b</sup>	0.60	(51)	-29.42 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	23.60 <sup>b</sup>	0.49	(05)	14.10 <sup>c</sup>	Montana	17.52 <sup>b</sup>	0.86	(48)	-15.31 <sup>c</sup>
Alaska	21.46	1.68	(20)	3.74	Nebraska	22.94	0.76	(12)	10.91 <sup>c</sup>
Arizona	17.48 <sup>b</sup>	0.35	(49)	-15.49 <sup>c</sup>	Nevada	21.32	0.66	(21)	3.09
Arkansas	23.19 <sup>b</sup>	0.60	(08)	12.13 <sup>c</sup>	New Hampshire	20.45	0.86	(32)	-1.14
California	18.43 <sup>b</sup>	0.16	(43)	-10.93 <sup>c</sup>	New Jersey	22.59 <sup>b</sup>	0.35	(15)	9.19
Colorado	18.26 <sup>b</sup>	0.47	(44)	-11.74 <sup>c</sup>	New Mexico	19.61	0.68	(38)	-5.21
Connecticut	18.09 <sup>b</sup>	0.48	(47)	-12.53 <sup>c</sup>	New York	20.15	0.22	(35)	-2.58
Delaware	20.82	1.04	(26)	0.64	North Carolina	20.39	0.34	(33)	-1.43
D.C.	23.02	1.45	(11)	11.31 <sup>c</sup>	North Dakota	22.16	1.17	(17)	7.12
Florida	18.70 <sup>b</sup>	0.19	(42)	-9.61	Ohio	23.25 <sup>b</sup>	0.30	(07)	12.42 <sup>c</sup>
Georgia	20.65	0.38	(28)	-0.16	Oklahoma	23.26 <sup>b</sup>	0.54	(06)	12.45 <sup>c</sup>
Hawaii	18.78	0.77	(41)	-9.19	Oregon	19.04	0.47	(40)	-7.96
Idaho	15.91 <sup>b</sup>	0.72	(50)	-23.10 <sup>c</sup>	Pennsylvania	22.71 <sup>b</sup>	0.27	(13)	9.80
Illinois	23.16 <sup>b</sup>	0.30	(09)	11.95 <sup>c</sup>	Rhode Island	20.56	0.92	(30)	-0.59
Indiana	23.09 <sup>b</sup>	0.42	(10)	11.64 <sup>c</sup>	South Carolina	20.92	0.48	(24)	1.15
Iowa	21.32	0.54	(22)	3.08	South Dakota	20.47	1.02	(31)	-1.02
Kansas	21.76	0.60	(19)	5.17	Tennessee	22.68 <sup>b</sup>	0.43	(14)	9.65
Kentucky	24.42 <sup>b</sup>	0.53	(03)	18.04 <sup>c</sup>	Texas	20.70	0.23	(27)	0.09
Louisiana	25.82 <sup>b</sup>	0.55	(01)	24.82 <sup>c</sup>	Utah	14.60 <sup>b</sup>	0.60	(51)	-29.42 <sup>c</sup>
Maine	20.85	0.81	(25)	0.81	Vermont	20.18	1.20	(34)	-2.42
Maryland	22.58 <sup>b</sup>	0.46	(16)	9.14	Virginia	20.97	0.38	(23)	1.37
Massachusetts	20.15	0.38	(36)	-2.60	Washington	18.20 <sup>b</sup>	0.37	(45)	-12.03 <sup>c</sup>
Michigan	20.62	0.31	(29)	-0.30	West Virginia	24.35 <sup>b</sup>	0.73	(04)	17.72 <sup>c</sup>
Minnesota	18.18 <sup>b</sup>	0.41	(46)	-12.12 <sup>c</sup>	Wisconsin	19.36	0.40	(39)	-6.40
Mississippi	25.19 <sup>b</sup>	0.66	(02)	21.80 <sup>c</sup>	Wyoming	19.89	1.34	(37)	-3.86
Missouri	22.11	0.41	(18)	6.89					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 6.27  
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	14.49	0.04		
High Five States				
D.C.	18.13 <sup>b</sup>	1.02	(01)	25.09 <sup>c</sup>
Kentucky	17.00 <sup>b</sup>	0.37	(02)	17.26 <sup>c</sup>
West Virginia	16.95 <sup>b</sup>	0.52	(03)	16.93 <sup>c</sup>
Mississippi	16.62 <sup>b</sup>	0.44	(04)	14.64 <sup>c</sup>
Nevada	16.37 <sup>b</sup>	0.53	(05)	12.97 <sup>c</sup>
Low Five States				
Washington	13.08 <sup>b</sup>	0.27	(47)	-9.79
Minnesota	12.96 <sup>b</sup>	0.29	(48)	-10.56 <sup>c</sup>
Arizona	11.94 <sup>b</sup>	0.26	(49)	-17.63 <sup>c</sup>
Hawaii	10.71 <sup>b</sup>	0.52	(50)	-26.13 <sup>c</sup>
Utah	10.18 <sup>b</sup>	0.44	(51)	-29.75 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	15.21	0.33	(17)	4.96	Montana	13.89	0.68	(36)	-4.13
Alaska	13.54	1.18	(40)	-6.55	Nebraska	15.59	0.53	(12)	7.59
Arizona	11.94 <sup>b</sup>	0.26	(49)	-17.63 <sup>c</sup>	Nevada	16.37 <sup>b</sup>	0.53	(05)	12.97 <sup>c</sup>
Arkansas	15.55	0.42	(14)	7.32	New Hampshire	13.94	0.59	(35)	-3.79
California	13.26 <sup>b</sup>	0.12	(46)	-8.54	New Jersey	15.97 <sup>b</sup>	0.24	(09)	10.16 <sup>c</sup>
Colorado	13.31	0.34	(45)	-8.19	New Mexico	13.36	0.50	(43)	-7.83
Connecticut	13.81	0.35	(37)	-4.71	New York	14.49	0.15	(26)	-0.01
Delaware	15.01	0.75	(22)	3.57	North Carolina	14.19	0.23	(33)	-2.08
D.C.	18.13 <sup>b</sup>	1.02	(01)	25.09 <sup>c</sup>	North Dakota	14.27	0.81	(32)	-1.54
Florida	13.34 <sup>b</sup>	0.14	(44)	-7.96	Ohio	15.98 <sup>b</sup>	0.21	(08)	10.29 <sup>c</sup>
Georgia	14.31	0.25	(31)	-1.27	Oklahoma	14.86	0.37	(23)	2.55
Hawaii	10.71 <sup>b</sup>	0.52	(50)	-26.13 <sup>c</sup>	Oregon	14.13	0.36	(34)	-2.52
Idaho	13.79	0.60	(38)	-4.88	Pennsylvania	15.77 <sup>b</sup>	0.19	(10)	8.82
Illinois	16.18 <sup>b</sup>	0.21	(07)	11.61 <sup>c</sup>	Rhode Island	13.52	0.61	(41)	-6.74
Indiana	15.60 <sup>b</sup>	0.29	(11)	7.62	South Carolina	14.56	0.33	(25)	0.46
Iowa	15.52	0.39	(15)	7.10	South Dakota	14.32	0.74	(30)	-1.21
Kansas	14.47	0.42	(27)	-0.13	Tennessee	15.58 <sup>b</sup>	0.29	(13)	7.51
Kentucky	17.00 <sup>b</sup>	0.37	(02)	17.26 <sup>c</sup>	Texas	13.44 <sup>b</sup>	0.15	(42)	-7.27
Louisiana	16.30 <sup>b</sup>	0.36	(06)	12.48 <sup>c</sup>	Utah	10.18 <sup>b</sup>	0.44	(51)	-29.75 <sup>c</sup>
Maine	15.37	0.58	(16)	6.03	Vermont	15.03	0.87	(19)	3.68
Maryland	15.03	0.31	(21)	3.67	Virginia	14.41	0.26	(28)	-0.55
Massachusetts	14.37	0.26	(29)	-0.84	Washington	13.08 <sup>b</sup>	0.27	(47)	-9.79
Michigan	15.06	0.22	(18)	3.89	West Virginia	16.95 <sup>b</sup>	0.52	(03)	16.93 <sup>c</sup>
Minnesota	12.96 <sup>b</sup>	0.29	(48)	-10.56 <sup>c</sup>	Wisconsin	13.61	0.28	(39)	-6.08
Mississippi	16.62 <sup>b</sup>	0.44	(04)	14.64 <sup>c</sup>	Wyoming	14.61	1.01	(24)	0.78
Missouri	15.03	0.28	(20)	3.67					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 7.14  
Cancer of the Corpus and Uterus, NOS (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.17	0.02		
High Five States				
D.C.	5.74	0.58	(01)	37.75 <sup>c</sup>
Delaware	5.35	0.45	(02)	28.24 <sup>c</sup>
New Jersey	5.14 <sup>b</sup>	0.14	(03)	23.34 <sup>c</sup>
New York	5.05 <sup>b</sup>	0.09	(04)	21.13 <sup>c</sup>
Nebraska	4.96	0.31	(05)	18.98 <sup>c</sup>
Low Five States				
Arkansas	3.04 <sup>b</sup>	0.19	(47)	-27.07 <sup>c</sup>
Nevada	3.03 <sup>b</sup>	0.22	(48)	-27.26 <sup>c</sup>
Arizona	3.01 <sup>b</sup>	0.13	(49)	-27.76 <sup>c</sup>
Utah	2.99 <sup>b</sup>	0.24	(50)	-28.22 <sup>c</sup>
Alaska	2.13 <sup>b</sup>	0.44	(51)	-48.85 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.39 <sup>b</sup>	0.15	(43)	-18.57 <sup>c</sup>	Montana	4.66	0.40	(12)	11.87
Alaska	2.13 <sup>b</sup>	0.44	(51)	-48.85 <sup>c</sup>	Nebraska	4.96	0.31	(05)	18.98 <sup>c</sup>
Arizona	3.01 <sup>b</sup>	0.13	(49)	-27.76 <sup>c</sup>	Nevada	3.03 <sup>b</sup>	0.22	(48)	-27.26 <sup>c</sup>
Arkansas	3.04 <sup>b</sup>	0.19	(47)	-27.07 <sup>c</sup>	New Hampshire	4.39	0.34	(20)	5.21
California	4.06	0.07	(27)	-2.52	New Jersey	5.14 <sup>b</sup>	0.14	(03)	23.34 <sup>c</sup>
Colorado	3.18 <sup>b</sup>	0.17	(46)	-23.79 <sup>c</sup>	New Mexico	3.37	0.25	(44)	-19.09 <sup>c</sup>
Connecticut	4.49	0.20	(18)	7.78	New York	5.05 <sup>b</sup>	0.09	(04)	21.13 <sup>c</sup>
Delaware	5.35	0.45	(02)	28.24 <sup>c</sup>	North Carolina	4.04	0.13	(29)	-2.99
D.C.	5.74	0.58	(01)	37.75 <sup>c</sup>	North Dakota	4.93	0.50	(06)	18.24 <sup>c</sup>
Florida	3.61 <sup>b</sup>	0.08	(38)	-13.45	Ohio	4.65 <sup>b</sup>	0.11	(13)	11.57
Georgia	3.63 <sup>b</sup>	0.13	(37)	-12.88	Oklahoma	3.93	0.19	(32)	-5.66
Hawaii	3.56	0.31	(40)	-14.52	Oregon	3.84	0.19	(33)	-7.81
Idaho	4.05	0.33	(28)	-2.79	Pennsylvania	4.88 <sup>b</sup>	0.11	(07)	17.03 <sup>c</sup>
Illinois	4.53	0.11	(16)	8.62	Rhode Island	3.95	0.34	(31)	-5.36
Indiana	4.50	0.16	(17)	8.06	South Carolina	4.16	0.18	(25)	-0.18
Iowa	4.75	0.22	(08)	13.99	South Dakota	3.98	0.41	(30)	-4.55
Kansas	3.45 <sup>b</sup>	0.20	(42)	-17.35 <sup>c</sup>	Tennessee	3.74	0.14	(36)	-10.21
Kentucky	3.54 <sup>b</sup>	0.17	(41)	-15.10 <sup>c</sup>	Texas	3.60 <sup>b</sup>	0.08	(39)	-13.75
Louisiana	3.75	0.17	(35)	-9.96	Utah	2.99 <sup>b</sup>	0.24	(50)	-28.22 <sup>c</sup>
Maine	4.49	0.32	(19)	7.76	Vermont	4.73	0.49	(10)	13.56
Maryland	4.67	0.17	(11)	11.96	Virginia	4.20	0.14	(22)	0.86
Massachusetts	4.17	0.14	(24)	0.12	Washington	4.13	0.15	(26)	-0.83
Michigan	4.74 <sup>b</sup>	0.13	(09)	13.81	West Virginia	4.54	0.27	(15)	9.02
Minnesota	4.19	0.17	(23)	0.59	Wisconsin	4.64	0.17	(14)	11.36
Mississippi	3.81	0.21	(34)	-8.50	Wyoming	3.21	0.48	(45)	-23.11 <sup>c</sup>
Missouri	4.29	0.15	(21)	2.82					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 8.18  
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.34	0.02		
High Five States				
Maine	5.94 <sup>b</sup>	0.27	(01)	36.84 <sup>c</sup>
New Hampshire	5.77 <sup>b</sup>	0.29	(02)	32.95 <sup>c</sup>
D.C.	5.74	0.45	(03)	32.06 <sup>c</sup>
Rhode Island	5.54 <sup>b</sup>	0.31	(04)	27.63 <sup>c</sup>
Massachusetts	5.29 <sup>b</sup>	0.12	(05)	21.79 <sup>c</sup>
Low Five States				
Texas	3.68 <sup>b</sup>	0.06	(47)	-15.27 <sup>c</sup>
California	3.61 <sup>b</sup>	0.05	(48)	-16.81 <sup>c</sup>
New Mexico	3.50 <sup>b</sup>	0.19	(49)	-19.49 <sup>c</sup>
Utah	3.12 <sup>b</sup>	0.18	(50)	-28.08 <sup>c</sup>
Hawaii	3.01 <sup>b</sup>	0.20	(51)	-30.67 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.01	0.13	(43)	-7.55	Montana	4.60	0.29	(20)	5.93
Alaska	4.63	0.47	(19)	6.64	Nebraska	4.29	0.21	(35)	-1.29
Arizona	3.93 <sup>b</sup>	0.11	(44)	-9.40	Nevada	4.38	0.20	(32)	0.75
Arkansas	3.87	0.16	(45)	-10.97	New Hampshire	5.77 <sup>b</sup>	0.29	(02)	32.95 <sup>c</sup>
California	3.61 <sup>b</sup>	0.05	(48)	-16.81 <sup>c</sup>	New Jersey	4.17	0.09	(36)	-3.97
Colorado	4.04	0.14	(40)	-7.02	New Mexico	3.50 <sup>b</sup>	0.19	(49)	-19.49 <sup>c</sup>
Connecticut	4.48	0.15	(25)	3.20	New York	4.32	0.06	(33)	-0.63
Delaware	4.78	0.32	(13)	10.06	North Carolina	4.03	0.09	(41)	-7.11
D.C.	5.74	0.45	(03)	32.06 <sup>c</sup>	North Dakota	4.02	0.33	(42)	-7.47
Florida	4.13 <sup>b</sup>	0.06	(37)	-4.83	Ohio	5.21 <sup>b</sup>	0.09	(06)	20.05 <sup>c</sup>
Georgia	4.08	0.10	(39)	-6.12	Oklahoma	4.52	0.15	(23)	4.06
Hawaii	3.01 <sup>b</sup>	0.20	(51)	-30.67 <sup>c</sup>	Oregon	5.01 <sup>b</sup>	0.16	(09)	15.36 <sup>c</sup>
Idaho	4.47	0.25	(26)	2.99	Pennsylvania	5.04 <sup>b</sup>	0.08	(08)	16.15 <sup>c</sup>
Illinois	4.65 <sup>b</sup>	0.09	(18)	7.06	Rhode Island	5.54 <sup>b</sup>	0.31	(04)	27.63 <sup>c</sup>
Indiana	4.79 <sup>b</sup>	0.12	(11)	10.24	South Carolina	4.59	0.14	(21)	5.76
Iowa	4.78	0.17	(12)	10.12	South Dakota	4.68	0.33	(15)	7.69
Kansas	3.86	0.16	(46)	-11.21	Tennessee	4.12	0.11	(38)	-5.12
Kentucky	4.58	0.14	(22)	5.43	Texas	3.68 <sup>b</sup>	0.06	(47)	-15.27 <sup>c</sup>
Louisiana	4.30	0.14	(34)	-0.96	Utah	3.12 <sup>b</sup>	0.18	(50)	-28.08 <sup>c</sup>
Maine	5.94 <sup>b</sup>	0.27	(01)	36.84 <sup>c</sup>	Vermont	4.67	0.37	(17)	7.47
Maryland	4.42	0.13	(29)	1.81	Virginia	4.39	0.11	(31)	1.12
Massachusetts	5.29 <sup>b</sup>	0.12	(05)	21.79 <sup>c</sup>	Washington	4.69	0.12	(14)	8.03
Michigan	4.87 <sup>b</sup>	0.10	(10)	12.13	West Virginia	4.51	0.20	(24)	3.79
Minnesota	4.43	0.13	(27)	1.96	Wisconsin	5.17 <sup>b</sup>	0.13	(07)	18.95 <sup>c</sup>
Mississippi	4.40	0.17	(30)	1.33	Wyoming	4.68	0.42	(16)	7.68
Missouri	4.43	0.12	(28)	1.94					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 8.19  
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.78	0.03		
High Five States				
Maine	11.19 <sup>b</sup>	0.57	(01)	43.86 <sup>c</sup>
D.C.	10.30	0.94	(02)	32.40 <sup>c</sup>
New Hampshire	10.21 <sup>b</sup>	0.59	(03)	31.28 <sup>c</sup>
Rhode Island	10.11 <sup>b</sup>	0.64	(04)	30.06 <sup>c</sup>
Ohio	9.65 <sup>b</sup>	0.19	(05)	24.09 <sup>c</sup>
Low Five States				
Texas	6.70 <sup>b</sup>	0.13	(47)	-13.78
California	6.28 <sup>b</sup>	0.09	(48)	-19.29 <sup>c</sup>
New Mexico	6.14 <sup>b</sup>	0.37	(49)	-21.04 <sup>c</sup>
Utah	5.76 <sup>b</sup>	0.36	(50)	-25.94 <sup>c</sup>
Hawaii	5.29 <sup>b</sup>	0.40	(51)	-32.03 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.64	0.27	(34)	-1.80	Montana	7.79	0.56	(33)	0.12
Alaska	7.81	0.91	(30)	0.48	Nebraska	7.93	0.44	(27)	1.92
Arizona	6.98 <sup>b</sup>	0.22	(45)	-10.20	Nevada	7.46	0.38	(36)	-4.02
Arkansas	6.85	0.31	(46)	-11.97	New Hampshire	10.21 <sup>b</sup>	0.59	(03)	31.28 <sup>c</sup>
California	6.28 <sup>b</sup>	0.09	(48)	-19.29 <sup>c</sup>	New Jersey	7.26	0.19	(40)	-6.66
Colorado	7.23	0.28	(41)	-6.97	New Mexico	6.14 <sup>b</sup>	0.37	(49)	-21.04 <sup>c</sup>
Connecticut	8.22	0.32	(20)	5.73	New York	7.57	0.13	(35)	-2.63
Delaware	8.88	0.67	(09)	14.18	North Carolina	7.18	0.19	(43)	-7.72
D.C.	10.30	0.94	(02)	32.40 <sup>c</sup>	North Dakota	7.06	0.66	(44)	-9.19
Florida	7.26 <sup>b</sup>	0.12	(39)	-6.61	Ohio	9.65 <sup>b</sup>	0.19	(05)	24.09 <sup>c</sup>
Georgia	7.29	0.21	(38)	-6.29	Oklahoma	8.28	0.31	(18)	6.43
Hawaii	5.29 <sup>b</sup>	0.40	(51)	-32.03 <sup>c</sup>	Oregon	8.82	0.32	(11)	13.46
Idaho	7.81	0.49	(31)	0.46	Pennsylvania	9.27 <sup>b</sup>	0.17	(07)	19.27 <sup>c</sup>
Illinois	8.47 <sup>b</sup>	0.18	(15)	8.88	Rhode Island	10.11 <sup>b</sup>	0.64	(04)	30.06 <sup>c</sup>
Indiana	8.77 <sup>b</sup>	0.25	(12)	12.80	South Carolina	8.29	0.29	(17)	6.58
Iowa	8.84	0.34	(10)	13.66	South Dakota	8.55	0.66	(14)	9.96
Kansas	7.21	0.34	(42)	-7.27	Tennessee	7.45	0.23	(37)	-4.17
Kentucky	8.24	0.29	(19)	5.93	Texas	6.70 <sup>b</sup>	0.13	(47)	-13.78
Louisiana	7.79	0.29	(32)	0.18	Utah	5.76 <sup>b</sup>	0.36	(50)	-25.94 <sup>c</sup>
Maine	11.19 <sup>b</sup>	0.57	(01)	43.86 <sup>c</sup>	Vermont	8.33	0.75	(16)	7.08
Maryland	8.00	0.26	(26)	2.87	Virginia	7.84	0.22	(29)	0.77
Massachusetts	9.57 <sup>b</sup>	0.25	(06)	23.09 <sup>c</sup>	Washington	8.20	0.25	(21)	5.41
Michigan	8.68 <sup>b</sup>	0.20	(13)	11.67	West Virginia	8.13	0.41	(22)	4.54
Minnesota	7.84	0.26	(28)	0.80	Wisconsin	9.20 <sup>b</sup>	0.27	(08)	18.36 <sup>c</sup>
Mississippi	8.03	0.36	(23)	3.27	Wyoming	8.02	0.82	(25)	3.18
Missouri	8.03	0.24	(24)	3.25					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 8.20  
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.64	0.01		
High Five States				
D.C.	2.46	0.38	(01)	50.01 <sup>c</sup>
New Hampshire	2.30	0.24	(02)	39.81 <sup>c</sup>
Rhode Island	2.20	0.25	(03)	33.74 <sup>c</sup>
Massachusetts	2.05 <sup>b</sup>	0.10	(04)	25.05 <sup>c</sup>
Oregon	1.94	0.13	(05)	18.36 <sup>c</sup>
Low Five States				
New Mexico	1.29	0.16	(47)	-21.43 <sup>c</sup>
Kansas	1.22	0.12	(48)	-25.76 <sup>c</sup>
Alabama	1.19 <sup>b</sup>	0.09	(49)	-27.22 <sup>c</sup>
Hawaii	1.07	0.17	(50)	-35.05 <sup>c</sup>
Utah	0.85 <sup>b</sup>	0.13	(51)	-48.35 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.19 <sup>b</sup>	0.09	(49)	-27.22 <sup>c</sup>	Montana	1.86	0.25	(11)	13.46
Alaska	1.71	0.41	(18)	4.44	Nebraska	1.36	0.16	(44)	-17.12 <sup>c</sup>
Arizona	1.33 <sup>b</sup>	0.09	(45)	-19.01 <sup>c</sup>	Nevada	1.61	0.16	(29)	-1.69
Arkansas	1.45	0.13	(41)	-11.77	New Hampshire	2.30	0.24	(02)	39.81 <sup>c</sup>
California	1.53	0.04	(36)	-6.97	New Jersey	1.78	0.08	(14)	8.65
Colorado	1.42	0.11	(43)	-13.77	New Mexico	1.29	0.16	(47)	-21.43 <sup>c</sup>
Connecticut	1.68	0.12	(21)	2.45	New York	1.89 <sup>b</sup>	0.06	(07)	14.94
Delaware	1.64	0.25	(26)	-0.13	North Carolina	1.60	0.08	(31)	-2.73
D.C.	2.46	0.38	(01)	50.01 <sup>c</sup>	North Dakota	1.63	0.29	(28)	-0.89
Florida	1.53	0.05	(37)	-7.00	Ohio	1.86	0.07	(09)	13.50
Georgia	1.67	0.09	(24)	1.50	Oklahoma	1.50	0.12	(38)	-8.39
Hawaii	1.07	0.17	(50)	-35.05 <sup>c</sup>	Oregon	1.94	0.13	(05)	18.36 <sup>c</sup>
Idaho	1.50	0.20	(39)	-8.80	Pennsylvania	1.80	0.06	(13)	9.75
Illinois	1.78	0.07	(15)	8.36	Rhode Island	2.20	0.25	(03)	33.74 <sup>c</sup>
Indiana	1.68	0.10	(23)	2.31	South Carolina	1.71	0.11	(19)	4.43
Iowa	1.58	0.13	(33)	-3.59	South Dakota	1.42	0.24	(42)	-13.72
Kansas	1.22	0.12	(48)	-25.76 <sup>c</sup>	Tennessee	1.48	0.09	(40)	-9.92
Kentucky	1.61	0.11	(30)	-1.74	Texas	1.31 <sup>b</sup>	0.05	(46)	-20.51 <sup>c</sup>
Louisiana	1.59	0.11	(32)	-3.38	Utah	0.85 <sup>b</sup>	0.13	(51)	-48.35 <sup>c</sup>
Maine	1.76	0.20	(16)	7.08	Vermont	1.86	0.31	(10)	13.48
Maryland	1.71	0.10	(20)	4.01	Virginia	1.66	0.09	(25)	1.22
Massachusetts	2.05 <sup>b</sup>	0.10	(04)	25.05 <sup>c</sup>	Washington	1.87	0.10	(08)	13.70
Michigan	1.85	0.08	(12)	12.64	West Virginia	1.54	0.16	(35)	-5.96
Minnesota	1.68	0.11	(22)	2.43	Wisconsin	1.90	0.11	(06)	15.66 <sup>c</sup>
Mississippi	1.57	0.14	(34)	-4.30	Wyoming	1.75	0.36	(17)	6.33
Missouri	1.63	0.10	(27)	-0.47					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 9.18  
Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.41	0.01		
High Five States				
Louisiana	0.54	0.05	(01)	29.28 <sup>c</sup>
Arkansas	0.49	0.06	(02)	17.85 <sup>c</sup>
New Jersey	0.48	0.03	(03)	16.25 <sup>c</sup>
Nebraska	0.48	0.07	(04)	15.87 <sup>c</sup>
Maine	0.47	0.08	(05)	13.64
Low Five States				
Utah	0.33	0.06	(41)	-20.29 <sup>c</sup>
New Mexico	0.33	0.06	(42)	-20.58 <sup>c</sup>
Montana	0.31	0.08	(43)	-24.83 <sup>c</sup>
New Hampshire	0.30	0.07	(44)	-27.67 <sup>c</sup>
Hawaii	0.23	0.06	(45)	-45.00 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.41	0.04	(27)	-1.91	Montana	0.31	0.08	(43)	-24.83 <sup>c</sup>
Alaska	-	-	(-)	-	Nebraska	0.48	0.07	(04)	15.87 <sup>c</sup>
Arizona	0.34	0.03	(39)	-18.50 <sup>c</sup>	Nevada	0.41	0.06	(22)	0.05
Arkansas	0.49	0.06	(02)	17.85 <sup>c</sup>	New Hampshire	0.30	0.07	(44)	-27.67 <sup>c</sup>
California	0.42	0.02	(16)	2.17	New Jersey	0.48	0.03	(03)	16.25 <sup>c</sup>
Colorado	0.37	0.04	(36)	-11.29	New Mexico	0.33	0.06	(42)	-20.58 <sup>c</sup>
Connecticut	0.44	0.05	(13)	5.37	New York	0.44	0.02	(11)	6.43
Delaware	0.41	0.10	(26)	-1.83	North Carolina	0.39	0.03	(34)	-6.27
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.40	0.02	(29)	-4.32	Ohio	0.47	0.03	(07)	13.50
Georgia	0.36	0.03	(37)	-12.35	Oklahoma	0.39	0.05	(32)	-4.82
Hawaii	0.23	0.06	(45)	-45.00 <sup>c</sup>	Oregon	0.44	0.05	(10)	7.20
Idaho	0.41	0.08	(24)	-0.53	Pennsylvania	0.47	0.03	(06)	13.51
Illinois	0.41	0.03	(25)	-0.67	Rhode Island	0.42	0.08	(17)	2.15
Indiana	0.35	0.03	(38)	-16.09 <sup>c</sup>	South Carolina	0.39	0.04	(33)	-6.15
Iowa	0.40	0.05	(31)	-4.65	South Dakota	-	-	(-)	-
Kansas	0.40	0.05	(28)	-3.34	Tennessee	0.46	0.04	(08)	10.63
Kentucky	0.45	0.05	(09)	8.15	Texas	0.42	0.02	(18)	1.99
Louisiana	0.54	0.05	(01)	29.28 <sup>c</sup>	Utah	0.33	0.06	(41)	-20.29 <sup>c</sup>
Maine	0.47	0.08	(05)	13.64	Vermont	-	-	(-)	-
Maryland	0.40	0.04	(30)	-4.58	Virginia	0.42	0.03	(19)	0.28
Massachusetts	0.33	0.03	(40)	-19.97 <sup>c</sup>	Washington	0.39	0.03	(35)	-7.04
Michigan	0.44	0.03	(12)	5.67	West Virginia	0.41	0.06	(23)	-0.05
Minnesota	0.41	0.04	(20)	0.07	Wisconsin	0.43	0.04	(14)	4.80
Mississippi	0.41	0.05	(21)	0.06	Wyoming	-	-	(-)	-
Missouri	0.43	0.04	(15)	4.35					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 9.19  
Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.51	0.01		
High Five States				
Rhode Island	0.70	0.17	(01)	37.26 <sup>c</sup>
Nebraska	0.69	0.13	(02)	33.59 <sup>c</sup>
Louisiana	0.64	0.08	(03)	24.71 <sup>c</sup>
Pennsylvania	0.62	0.04	(04)	21.36 <sup>c</sup>
Maine	0.62	0.14	(05)	21.03 <sup>c</sup>
Low Five States				
West Virginia	0.43	0.10	(36)	-17.10 <sup>c</sup>
Oklahoma	0.42	0.07	(37)	-17.38 <sup>c</sup>
South Carolina	0.42	0.07	(38)	-18.06 <sup>c</sup>
Massachusetts	0.41	0.05	(39)	-20.41 <sup>c</sup>
Iowa	0.35	0.07	(40)	-32.11 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.48	0.07	(27)	-7.13	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.69	0.13	(02)	33.59 <sup>c</sup>
Arizona	0.45	0.06	(30)	-11.37	Nevada	0.56	0.10	(11)	9.00
Arkansas	0.59	0.09	(08)	14.20	New Hampshire	-	-	(-)	-
California	0.52	0.03	(18)	2.09	New Jersey	0.60	0.06	(07)	16.77 <sup>c</sup>
Colorado	0.52	0.07	(20)	0.66	New Mexico	0.44	0.10	(33)	-14.00
Connecticut	0.54	0.08	(14)	5.53	New York	0.55	0.04	(12)	7.22
Delaware	-	-	(-)	-	North Carolina	0.52	0.05	(19)	1.91
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.51	0.03	(22)	-1.51	Ohio	0.60	0.05	(06)	17.64 <sup>c</sup>
Georgia	0.44	0.05	(35)	-15.03 <sup>c</sup>	Oklahoma	0.42	0.07	(37)	-17.38 <sup>c</sup>
Hawaii	-	-	(-)	-	Oregon	0.54	0.08	(15)	5.34
Idaho	-	-	(-)	-	Pennsylvania	0.62	0.04	(04)	21.36 <sup>c</sup>
Illinois	0.51	0.04	(21)	0.33	Rhode Island	0.70	0.17	(01)	37.26 <sup>c</sup>
Indiana	0.48	0.06	(26)	-6.85	South Carolina	0.42	0.07	(38)	-18.06 <sup>c</sup>
Iowa	0.35	0.07	(40)	-32.11 <sup>c</sup>	South Dakota	-	-	(-)	-
Kansas	0.46	0.09	(29)	-11.04	Tennessee	0.46	0.06	(28)	-10.34
Kentucky	0.59	0.08	(09)	13.97	Texas	0.53	0.03	(16)	2.92
Louisiana	0.64	0.08	(03)	24.71 <sup>c</sup>	Utah	0.44	0.10	(32)	-13.76
Maine	0.62	0.14	(05)	21.03 <sup>c</sup>	Vermont	-	-	(-)	-
Maryland	0.44	0.06	(31)	-13.32	Virginia	0.52	0.06	(17)	2.16
Massachusetts	0.41	0.05	(39)	-20.41 <sup>c</sup>	Washington	0.50	0.06	(23)	-3.12
Michigan	0.57	0.05	(10)	11.21	West Virginia	0.43	0.10	(36)	-17.10 <sup>c</sup>
Minnesota	0.50	0.07	(25)	-3.38	Wisconsin	0.54	0.07	(13)	5.92
Mississippi	0.50	0.09	(24)	-3.13	Wyoming	-	-	(-)	-
Missouri	0.44	0.06	(34)	-14.86					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 9.20  
Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.33	0.01		
High Five States				
Louisiana	0.46	0.06	(01)	37.89 <sup>c</sup>
Tennessee	0.45	0.05	(02)	35.97 <sup>c</sup>
Idaho	0.45	0.11	(03)	34.78 <sup>c</sup>
Iowa	0.42	0.07	(04)	27.08 <sup>c</sup>
Missouri	0.41	0.05	(05)	23.07 <sup>c</sup>
Low Five States				
North Carolina	0.28	0.03	(33)	-16.93 <sup>c</sup>
Nevada	0.28	0.07	(34)	-17.07 <sup>c</sup>
Indiana	0.25	0.04	(35)	-25.21 <sup>c</sup>
Arizona	0.23	0.04	(36)	-30.25 <sup>c</sup>
Colorado	0.23	0.05	(37)	-30.41 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.34	0.05	(20)	2.61	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.31	0.08	(28)	-7.22
Arizona	0.23	0.04	(36)	-30.25 <sup>c</sup>	Nevada	0.28	0.07	(34)	-17.07 <sup>c</sup>
Arkansas	0.40	0.07	(06)	19.75 <sup>c</sup>	New Hampshire	-	-	(-)	-
California	0.34	0.02	(19)	2.70	New Jersey	0.39	0.04	(08)	16.50 <sup>c</sup>
Colorado	0.23	0.05	(37)	-30.41 <sup>c</sup>	New Mexico	-	-	(-)	-
Connecticut	0.34	0.06	(21)	2.54	New York	0.35	0.03	(17)	5.74
Delaware	-	-	(-)	-	North Carolina	0.28	0.03	(33)	-16.93 <sup>c</sup>
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.30	0.02	(29)	-8.90	Ohio	0.36	0.03	(13)	8.50
Georgia	0.30	0.04	(30)	-10.17	Oklahoma	0.38	0.06	(10)	12.41
Hawaii	-	-	(-)	-	Oregon	0.37	0.06	(11)	11.13
Idaho	0.45	0.11	(03)	34.78 <sup>c</sup>	Pennsylvania	0.35	0.03	(16)	5.93
Illinois	0.33	0.03	(25)	-0.51	Rhode Island	-	-	(-)	-
Indiana	0.25	0.04	(35)	-25.21 <sup>c</sup>	South Carolina	0.36	0.06	(12)	9.06
Iowa	0.42	0.07	(04)	27.08 <sup>c</sup>	South Dakota	-	-	(-)	-
Kansas	0.38	0.07	(09)	15.23 <sup>c</sup>	Tennessee	0.45	0.05	(02)	35.97 <sup>c</sup>
Kentucky	0.33	0.05	(24)	0.08	Texas	0.34	0.02	(23)	0.90
Louisiana	0.46	0.06	(01)	37.89 <sup>c</sup>	Utah	-	-	(-)	-
Maine	-	-	(-)	-	Vermont	-	-	(-)	-
Maryland	0.35	0.05	(18)	4.03	Virginia	0.32	0.04	(27)	-4.02
Massachusetts	0.28	0.04	(32)	-16.64 <sup>c</sup>	Washington	0.29	0.04	(31)	-13.02
Michigan	0.33	0.03	(26)	-1.39	West Virginia	0.39	0.08	(07)	16.75 <sup>c</sup>
Minnesota	0.36	0.05	(14)	6.59	Wisconsin	0.34	0.05	(22)	1.21
Mississippi	0.36	0.07	(15)	6.37	Wyoming	-	-	(-)	-
Missouri	0.41	0.05	(05)	23.07 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 11.18  
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.05	0.02		
High Five States				
North Dakota	5.58 <sup>b</sup>	0.39	(01)	37.98 <sup>c</sup>
Louisiana	5.21 <sup>b</sup>	0.15	(02)	28.84 <sup>c</sup>
Oklahoma	5.10 <sup>b</sup>	0.16	(03)	26.11 <sup>c</sup>
Iowa	4.91 <sup>b</sup>	0.17	(04)	21.30 <sup>c</sup>
Kentucky	4.72 <sup>b</sup>	0.15	(05)	16.73 <sup>c</sup>
Low Five States				
Connecticut	3.48 <sup>b</sup>	0.13	(47)	-13.85
New York	3.42 <sup>b</sup>	0.06	(48)	-15.50 <sup>c</sup>
Utah	3.16 <sup>b</sup>	0.18	(49)	-21.94 <sup>c</sup>
D.C.	2.87 <sup>b</sup>	0.32	(50)	-29.06 <sup>c</sup>
Hawaii	2.54 <sup>b</sup>	0.19	(51)	-37.21 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.94	0.13	(32)	-2.52	Montana	3.95	0.27	(31)	-2.47
Alaska	3.89	0.45	(35)	-3.73	Nebraska	4.46	0.22	(14)	10.19
Arizona	3.76	0.11	(41)	-7.16	Nevada	3.92	0.19	(34)	-3.17
Arkansas	4.44	0.17	(15)	9.81	New Hampshire	4.10	0.24	(26)	1.36
California	3.63 <sup>b</sup>	0.05	(43)	-10.24	New Jersey	3.51 <sup>b</sup>	0.09	(46)	-13.15
Colorado	3.71	0.13	(42)	-8.27	New Mexico	4.51	0.21	(13)	11.39
Connecticut	3.48 <sup>b</sup>	0.13	(47)	-13.85	New York	3.42 <sup>b</sup>	0.06	(48)	-15.50 <sup>c</sup>
Delaware	4.15	0.30	(24)	2.51	North Carolina	4.24	0.10	(22)	4.70
D.C.	2.87 <sup>b</sup>	0.32	(50)	-29.06 <sup>c</sup>	North Dakota	5.58 <sup>b</sup>	0.39	(01)	37.98 <sup>c</sup>
Florida	3.52 <sup>b</sup>	0.06	(45)	-13.03	Ohio	4.57 <sup>b</sup>	0.09	(11)	13.01
Georgia	3.88	0.10	(36)	-3.99	Oklahoma	5.10 <sup>b</sup>	0.16	(03)	26.11 <sup>c</sup>
Hawaii	2.54 <sup>b</sup>	0.19	(51)	-37.21 <sup>c</sup>	Oregon	3.78	0.14	(40)	-6.48
Idaho	3.97	0.24	(30)	-1.91	Pennsylvania	4.06	0.07	(28)	0.27
Illinois	4.27	0.08	(21)	5.68	Rhode Island	3.83	0.25	(38)	-5.28
Indiana	4.68 <sup>b</sup>	0.12	(07)	15.62 <sup>c</sup>	South Carolina	4.35	0.14	(18)	7.63
Iowa	4.91 <sup>b</sup>	0.17	(04)	21.30 <sup>c</sup>	South Dakota	4.66	0.32	(08)	15.30 <sup>c</sup>
Kansas	4.61	0.18	(10)	13.95	Tennessee	4.31	0.12	(19)	6.60
Kentucky	4.72 <sup>b</sup>	0.15	(05)	16.73 <sup>c</sup>	Texas	4.61 <sup>b</sup>	0.07	(09)	14.02
Louisiana	5.21 <sup>b</sup>	0.15	(02)	28.84 <sup>c</sup>	Utah	3.16 <sup>b</sup>	0.18	(49)	-21.94 <sup>c</sup>
Maine	4.14	0.23	(25)	2.37	Vermont	4.05	0.34	(29)	0.19
Maryland	3.84	0.12	(37)	-5.18	Virginia	3.92	0.10	(33)	-3.10
Massachusetts	3.81	0.10	(39)	-5.74	Washington	4.08	0.11	(27)	0.81
Michigan	4.20	0.09	(23)	3.93	West Virginia	4.68	0.21	(06)	15.77 <sup>c</sup>
Minnesota	4.31	0.13	(20)	6.59	Wisconsin	4.42	0.12	(16)	9.24
Mississippi	4.55	0.18	(12)	12.38	Wyoming	3.62	0.37	(44)	-10.59
Missouri	4.37	0.12	(17)	7.95					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 11.19  
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.87	0.03		
High Five States				
North Dakota	7.85	0.70	(01)	33.74 <sup>c</sup>
Louisiana	7.44 <sup>b</sup>	0.29	(02)	26.78 <sup>c</sup>
Oklahoma	7.30 <sup>b</sup>	0.30	(03)	24.30 <sup>c</sup>
Iowa	7.09 <sup>b</sup>	0.31	(04)	20.77 <sup>c</sup>
Kansas	6.88	0.33	(05)	17.12 <sup>c</sup>
Low Five States				
Vermont	5.04	0.58	(47)	-14.13
Connecticut	4.98 <sup>b</sup>	0.25	(48)	-15.25 <sup>c</sup>
Utah	4.29 <sup>b</sup>	0.32	(49)	-26.92 <sup>c</sup>
D.C.	4.00	0.57	(50)	-31.92 <sup>c</sup>
Hawaii	3.57 <sup>b</sup>	0.33	(51)	-39.25 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.42	0.23	(39)	-7.71	Montana	5.60	0.49	(35)	-4.59
Alaska	5.54	0.81	(37)	-5.66	Nebraska	5.89	0.38	(25)	0.28
Arizona	5.23	0.19	(42)	-10.95	Nevada	5.48	0.33	(38)	-6.68
Arkansas	6.43	0.31	(17)	9.50	New Hampshire	5.80	0.45	(30)	-1.23
California	5.32 <sup>b</sup>	0.09	(41)	-9.43	New Jersey	5.22 <sup>b</sup>	0.17	(43)	-11.16
Colorado	5.60	0.26	(34)	-4.57	New Mexico	6.17	0.38	(22)	4.99
Connecticut	4.98 <sup>b</sup>	0.25	(48)	-15.25 <sup>c</sup>	New York	5.10 <sup>b</sup>	0.11	(46)	-13.21
Delaware	6.43	0.57	(15)	9.54	North Carolina	6.52	0.19	(13)	11.11
D.C.	4.00	0.57	(50)	-31.92 <sup>c</sup>	North Dakota	7.85	0.70	(01)	33.74 <sup>c</sup>
Florida	5.18 <sup>b</sup>	0.10	(45)	-11.75	Ohio	6.59 <sup>b</sup>	0.16	(09)	12.24
Georgia	5.74	0.20	(32)	-2.31	Oklahoma	7.30 <sup>b</sup>	0.30	(03)	24.30 <sup>c</sup>
Hawaii	3.57 <sup>b</sup>	0.33	(51)	-39.25 <sup>c</sup>	Oregon	5.55	0.25	(36)	-5.52
Idaho	5.19	0.40	(44)	-11.53	Pennsylvania	5.79	0.14	(31)	-1.41
Illinois	6.24	0.15	(20)	6.24	Rhode Island	5.34	0.46	(40)	-9.03
Indiana	6.57	0.22	(11)	11.89	South Carolina	6.43	0.26	(16)	9.53
Iowa	7.09 <sup>b</sup>	0.31	(04)	20.77 <sup>c</sup>	South Dakota	6.76	0.58	(07)	15.13 <sup>c</sup>
Kansas	6.88	0.33	(05)	17.12 <sup>c</sup>	Tennessee	6.48	0.22	(14)	10.41
Kentucky	6.82	0.28	(06)	16.11 <sup>c</sup>	Texas	6.68 <sup>b</sup>	0.13	(08)	13.75
Louisiana	7.44 <sup>b</sup>	0.29	(02)	26.78 <sup>c</sup>	Utah	4.29 <sup>b</sup>	0.32	(49)	-26.92 <sup>c</sup>
Maine	5.91	0.42	(24)	0.62	Vermont	5.04	0.58	(47)	-14.13
Maryland	5.80	0.23	(29)	-1.22	Virginia	6.00	0.20	(23)	2.22
Massachusetts	5.88	0.20	(26)	0.21	Washington	5.71	0.21	(33)	-2.81
Michigan	5.80	0.16	(28)	-1.16	West Virginia	6.29	0.36	(18)	7.10
Minnesota	6.19	0.23	(21)	5.46	Wisconsin	6.28	0.22	(19)	6.88
Mississippi	6.58	0.33	(10)	12.05	Wyoming	5.83	0.70	(27)	-0.75
Missouri	6.54	0.22	(12)	11.36					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 11.20  
Cancer of the Kidney And Renal Pelvis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.65	0.02		
High Five States				
North Dakota	3.66	0.43	(01)	38.15 <sup>c</sup>
Louisiana	3.48 <sup>b</sup>	0.17	(02)	31.34 <sup>c</sup>
Oklahoma	3.37 <sup>b</sup>	0.18	(03)	27.29 <sup>c</sup>
West Virginia	3.37	0.23	(04)	27.26 <sup>c</sup>
Nebraska	3.28	0.25	(05)	23.89 <sup>c</sup>
Low Five States				
Utah	2.18	0.20	(47)	-17.96 <sup>c</sup>
Florida	2.16 <sup>b</sup>	0.06	(48)	-18.50 <sup>c</sup>
D.C.	1.87	0.34	(49)	-29.56 <sup>c</sup>
Hawaii	1.68 <sup>b</sup>	0.21	(50)	-36.61 <sup>c</sup>
Wyoming	1.67	0.34	(51)	-37.09 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.78	0.14	(26)	4.76	Montana	2.65	0.30	(31)	-0.08
Alaska	2.41	0.49	(39)	-9.03	Nebraska	3.28	0.25	(05)	23.89 <sup>c</sup>
Arizona	2.52	0.12	(34)	-5.09	Nevada	2.50	0.21	(35)	-5.53
Arkansas	2.85	0.18	(21)	7.40	New Hampshire	2.86	0.27	(19)	7.81
California	2.34 <sup>b</sup>	0.05	(43)	-11.90	New Jersey	2.28 <sup>b</sup>	0.09	(45)	-13.91
Colorado	2.29	0.14	(44)	-13.45	New Mexico	3.14	0.24	(09)	18.39 <sup>c</sup>
Connecticut	2.43	0.15	(37)	-8.48	New York	2.22 <sup>b</sup>	0.06	(46)	-16.12 <sup>c</sup>
Delaware	2.48	0.31	(36)	-6.50	North Carolina	2.54	0.10	(33)	-4.36
D.C.	1.87	0.34	(49)	-29.56 <sup>c</sup>	North Dakota	3.66	0.43	(01)	38.15 <sup>c</sup>
Florida	2.16 <sup>b</sup>	0.06	(48)	-18.50 <sup>c</sup>	Ohio	3.10 <sup>b</sup>	0.09	(11)	16.80 <sup>c</sup>
Georgia	2.58	0.11	(32)	-2.54	Oklahoma	3.37 <sup>b</sup>	0.18	(03)	27.29 <sup>c</sup>
Hawaii	1.68 <sup>b</sup>	0.21	(50)	-36.61 <sup>c</sup>	Oregon	2.39	0.15	(40)	-9.92
Idaho	2.91	0.28	(17)	9.69	Pennsylvania	2.76	0.08	(28)	4.24
Illinois	2.83	0.09	(22)	6.64	Rhode Island	2.73	0.28	(29)	2.93
Indiana	3.25 <sup>b</sup>	0.13	(08)	22.55 <sup>c</sup>	South Carolina	2.80	0.15	(24)	5.49
Iowa	3.27	0.18	(06)	23.47 <sup>c</sup>	South Dakota	2.96	0.35	(15)	11.75
Kansas	2.86	0.19	(20)	7.78	Tennessee	2.71	0.12	(30)	2.17
Kentucky	3.12	0.16	(10)	17.81 <sup>c</sup>	Texas	3.00 <sup>b</sup>	0.07	(14)	12.98
Louisiana	3.48 <sup>b</sup>	0.17	(02)	31.34 <sup>c</sup>	Utah	2.18	0.20	(47)	-17.96 <sup>c</sup>
Maine	2.87	0.26	(18)	8.26	Vermont	3.26	0.41	(07)	22.90 <sup>c</sup>
Maryland	2.42	0.12	(38)	-8.90	Virginia	2.38	0.11	(42)	-10.40
Massachusetts	2.39	0.11	(41)	-9.95	Washington	2.78	0.13	(27)	4.69
Michigan	3.00 <sup>b</sup>	0.10	(13)	13.22	West Virginia	3.37	0.23	(04)	27.26 <sup>c</sup>
Minnesota	2.80	0.14	(23)	5.62	Wisconsin	2.95	0.13	(16)	11.10
Mississippi	3.04	0.19	(12)	14.49	Wyoming	1.67	0.34	(51)	-37.09 <sup>c</sup>
Missouri	2.79	0.12	(25)	5.28					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.18  
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.20	0.01		
High Five States				
D.C.	1.93	0.26	(01)	61.12 <sup>c</sup>
West Virginia	1.89 <sup>b</sup>	0.13	(02)	57.36 <sup>c</sup>
Louisiana	1.66 <sup>b</sup>	0.09	(03)	38.88 <sup>c</sup>
Delaware	1.64	0.19	(04)	36.50 <sup>c</sup>
Kentucky	1.57 <sup>b</sup>	0.08	(05)	30.84 <sup>c</sup>
Low Five States				
South Dakota	0.81	0.14	(46)	-32.06 <sup>c</sup>
Oregon	0.78 <sup>b</sup>	0.06	(47)	-35.05 <sup>c</sup>
Colorado	0.73 <sup>b</sup>	0.06	(48)	-39.39 <sup>c</sup>
North Dakota	0.68 <sup>b</sup>	0.14	(49)	-43.10 <sup>c</sup>
Utah	0.38 <sup>b</sup>	0.06	(50)	-68.39 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.44	0.08	(07)	19.85 <sup>c</sup>	Montana	0.87	0.13	(43)	-27.70 <sup>c</sup>
Alaska	-	-	(-)	-	Nebraska	1.08	0.11	(32)	-10.11
Arizona	0.87 <sup>b</sup>	0.05	(42)	-27.70 <sup>c</sup>	Nevada	1.29	0.11	(19)	7.64
Arkansas	1.33	0.09	(15)	10.98	New Hampshire	1.08	0.13	(31)	-9.96
California	0.92 <sup>b</sup>	0.02	(38)	-23.28 <sup>c</sup>	New Jersey	1.30	0.05	(18)	8.64
Colorado	0.73 <sup>b</sup>	0.06	(48)	-39.39 <sup>c</sup>	New Mexico	1.06	0.10	(33)	-11.66
Connecticut	1.02	0.07	(35)	-14.60	New York	1.23	0.03	(21)	2.49
Delaware	1.64	0.19	(04)	36.50 <sup>c</sup>	North Carolina	1.39	0.06	(11)	15.81 <sup>c</sup>
D.C.	1.93	0.26	(01)	61.12 <sup>c</sup>	North Dakota	0.68 <sup>b</sup>	0.14	(49)	-43.10 <sup>c</sup>
Florida	1.40 <sup>b</sup>	0.04	(10)	16.69 <sup>c</sup>	Ohio	1.41 <sup>b</sup>	0.05	(08)	17.58 <sup>c</sup>
Georgia	1.16	0.05	(27)	-3.28	Oklahoma	1.19	0.08	(26)	-0.25
Hawaii	0.88	0.11	(41)	-26.80 <sup>c</sup>	Oregon	0.78 <sup>b</sup>	0.06	(47)	-35.05 <sup>c</sup>
Idaho	0.82 <sup>b</sup>	0.11	(45)	-31.94 <sup>c</sup>	Pennsylvania	1.32	0.04	(16)	9.89
Illinois	1.22	0.04	(22)	2.10	Rhode Island	1.25	0.15	(20)	4.56
Indiana	1.21	0.06	(23)	1.35	South Carolina	1.40	0.08	(09)	17.13 <sup>c</sup>
Iowa	1.05	0.08	(34)	-12.73	South Dakota	0.81	0.14	(46)	-32.06 <sup>c</sup>
Kansas	0.97	0.08	(36)	-18.65 <sup>c</sup>	Tennessee	1.50 <sup>b</sup>	0.07	(06)	25.46 <sup>c</sup>
Kentucky	1.57 <sup>b</sup>	0.08	(05)	30.84 <sup>c</sup>	Texas	1.21	0.03	(24)	1.21
Louisiana	1.66 <sup>b</sup>	0.09	(03)	38.88 <sup>c</sup>	Utah	0.38 <sup>b</sup>	0.06	(50)	-68.39 <sup>c</sup>
Maine	1.31	0.13	(17)	9.02	Vermont	0.89	0.16	(40)	-26.06 <sup>c</sup>
Maryland	1.37	0.07	(13)	14.75	Virginia	1.14	0.06	(29)	-4.45
Massachusetts	1.20	0.06	(25)	-0.24	Washington	0.92 <sup>b</sup>	0.05	(39)	-23.57 <sup>c</sup>
Michigan	1.38 <sup>b</sup>	0.05	(12)	14.87	West Virginia	1.89 <sup>b</sup>	0.13	(02)	57.36 <sup>c</sup>
Minnesota	0.96 <sup>b</sup>	0.06	(37)	-19.51 <sup>c</sup>	Wisconsin	1.09	0.06	(30)	-8.81
Mississippi	1.34	0.09	(14)	11.84	Wyoming	0.86	0.18	(44)	-28.25 <sup>c</sup>
Missouri	1.15	0.06	(28)	-3.84					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.19  
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	2.16	0.02		
High Five States				
D.C.	3.90	0.59	(01)	80.46 <sup>c</sup>
West Virginia	3.22 <sup>b</sup>	0.25	(02)	49.08 <sup>c</sup>
Louisiana	3.01 <sup>b</sup>	0.18	(03)	39.30 <sup>c</sup>
Delaware	2.94	0.37	(04)	36.09 <sup>c</sup>
Tennessee	2.82 <sup>b</sup>	0.14	(05)	30.34 <sup>c</sup>
Low Five States				
Wyoming	1.36	0.34	(46)	-37.01 <sup>c</sup>
North Dakota	1.29	0.28	(47)	-40.47 <sup>c</sup>
Colorado	1.21 <sup>b</sup>	0.12	(48)	-43.89 <sup>c</sup>
Oregon	1.17 <sup>b</sup>	0.11	(49)	-46.10 <sup>c</sup>
Utah	0.63 <sup>b</sup>	0.12	(50)	-70.80 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.58	0.15	(09)	19.29 <sup>c</sup>	Montana	1.45	0.24	(42)	-33.15 <sup>c</sup>
Alaska	-	-	(-)	-	Nebraska	1.92	0.21	(32)	-11.30
Arizona	1.45 <sup>b</sup>	0.10	(41)	-32.84 <sup>c</sup>	Nevada	2.01	0.20	(29)	-6.88
Arkansas	2.45	0.19	(13)	13.46	New Hampshire	1.83	0.25	(35)	-15.30 <sup>c</sup>
California	1.71 <sup>b</sup>	0.05	(38)	-20.98 <sup>c</sup>	New Jersey	2.38	0.11	(18)	9.98
Colorado	1.21 <sup>b</sup>	0.12	(48)	-43.89 <sup>c</sup>	New Mexico	1.85	0.20	(34)	-14.36
Connecticut	1.94	0.15	(31)	-10.35	New York	2.28	0.07	(20)	5.53
Delaware	2.94	0.37	(04)	36.09 <sup>c</sup>	North Carolina	2.61 <sup>b</sup>	0.12	(08)	20.75 <sup>c</sup>
D.C.	3.90	0.59	(01)	80.46 <sup>c</sup>	North Dakota	1.29	0.28	(47)	-40.47 <sup>c</sup>
Florida	2.44 <sup>b</sup>	0.07	(14)	12.63	Ohio	2.54 <sup>b</sup>	0.10	(11)	17.41 <sup>c</sup>
Georgia	2.17	0.12	(24)	0.44	Oklahoma	2.03	0.16	(28)	-6.10
Hawaii	1.69	0.23	(39)	-22.04 <sup>c</sup>	Oregon	1.17 <sup>b</sup>	0.11	(49)	-46.10 <sup>c</sup>
Idaho	1.39 <sup>b</sup>	0.21	(45)	-35.65 <sup>c</sup>	Pennsylvania	2.41	0.09	(16)	11.31
Illinois	2.24	0.09	(22)	3.72	Rhode Island	2.41	0.31	(15)	11.62
Indiana	2.13	0.12	(27)	-1.69	South Carolina	2.55	0.16	(10)	17.75 <sup>c</sup>
Iowa	1.85	0.16	(33)	-14.31	South Dakota	1.43	0.27	(44)	-33.97 <sup>c</sup>
Kansas	1.73	0.16	(36)	-20.09 <sup>c</sup>	Tennessee	2.82 <sup>b</sup>	0.14	(05)	30.34 <sup>c</sup>
Kentucky	2.80 <sup>b</sup>	0.17	(06)	29.54 <sup>c</sup>	Texas	2.29	0.07	(19)	5.73
Louisiana	3.01 <sup>b</sup>	0.18	(03)	39.30 <sup>c</sup>	Utah	0.63 <sup>b</sup>	0.12	(50)	-70.80 <sup>c</sup>
Maine	2.28	0.26	(21)	5.47	Vermont	1.44	0.30	(43)	-33.22 <sup>c</sup>
Maryland	2.52	0.15	(12)	16.55 <sup>c</sup>	Virginia	2.17	0.12	(23)	0.53
Massachusetts	2.14	0.12	(26)	-1.03	Washington	1.53 <sup>b</sup>	0.11	(40)	-29.47 <sup>c</sup>
Michigan	2.41	0.10	(17)	11.22	West Virginia	3.22 <sup>b</sup>	0.25	(02)	49.08 <sup>c</sup>
Minnesota	1.73 <sup>b</sup>	0.12	(37)	-20.17 <sup>c</sup>	Wisconsin	1.96	0.12	(30)	-9.54
Mississippi	2.71	0.21	(07)	25.23 <sup>c</sup>	Wyoming	1.36	0.34	(46)	-37.01 <sup>c</sup>
Missouri	2.14	0.13	(25)	-0.96					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.20  
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.46	0.01		
High Five States				
West Virginia	0.78	0.12	(01)	71.88 <sup>c</sup>
Nevada	0.68	0.11	(02)	48.42 <sup>c</sup>
Louisiana	0.64	0.07	(03)	40.37 <sup>c</sup>
Kentucky	0.64	0.07	(04)	39.46 <sup>c</sup>
Michigan	0.57	0.04	(05)	23.90 <sup>c</sup>
Low Five States				
Minnesota	0.37	0.05	(35)	-19.02 <sup>c</sup>
Connecticut	0.36	0.06	(36)	-21.68 <sup>c</sup>
Colorado	0.33	0.05	(37)	-27.19 <sup>c</sup>
California	0.33 <sup>b</sup>	0.02	(38)	-28.76 <sup>c</sup>
Mississippi	0.31	0.06	(39)	-31.27 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.55	0.06	(07)	19.86 <sup>c</sup>	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.42	0.09	(27)	-8.92
Arizona	0.37	0.05	(34)	-17.86 <sup>c</sup>	Nevada	0.68	0.11	(02)	48.42 <sup>c</sup>
Arkansas	0.45	0.07	(23)	-0.70	New Hampshire	0.51	0.12	(18)	10.83
California	0.33 <sup>b</sup>	0.02	(38)	-28.76 <sup>c</sup>	New Jersey	0.52	0.04	(14)	14.08
Colorado	0.33	0.05	(37)	-27.19 <sup>c</sup>	New Mexico	0.39	0.09	(30)	-14.57
Connecticut	0.36	0.06	(36)	-21.68 <sup>c</sup>	New York	0.45	0.03	(24)	-1.85
Delaware	-	-	(-)	-	North Carolina	0.51	0.04	(17)	11.03
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.54	0.03	(09)	17.44 <sup>c</sup>	Ohio	0.55	0.04	(08)	19.38 <sup>c</sup>
Georgia	0.41	0.04	(28)	-9.93	Oklahoma	0.53	0.07	(10)	15.74 <sup>c</sup>
Hawaii	-	-	(-)	-	Oregon	0.47	0.07	(20)	3.77
Idaho	-	-	(-)	-	Pennsylvania	0.51	0.04	(15)	12.54
Illinois	0.47	0.04	(21)	1.92	Rhode Island	-	-	(-)	-
Indiana	0.52	0.05	(12)	14.72	South Carolina	0.52	0.06	(13)	14.37
Iowa	0.46	0.07	(22)	0.42	South Dakota	-	-	(-)	-
Kansas	0.38	0.07	(33)	-16.72 <sup>c</sup>	Tennessee	0.50	0.05	(19)	8.51
Kentucky	0.64	0.07	(04)	39.46 <sup>c</sup>	Texas	0.39	0.03	(31)	-15.03 <sup>c</sup>
Louisiana	0.64	0.07	(03)	40.37 <sup>c</sup>	Utah	-	-	(-)	-
Maine	0.55	0.11	(06)	20.09 <sup>c</sup>	Vermont	-	-	(-)	-
Maryland	0.52	0.06	(11)	14.79	Virginia	0.39	0.04	(32)	-15.07 <sup>c</sup>
Massachusetts	0.51	0.05	(16)	11.51	Washington	0.44	0.05	(25)	-2.53
Michigan	0.57	0.04	(05)	23.90 <sup>c</sup>	West Virginia	0.78	0.12	(01)	71.88 <sup>c</sup>
Minnesota	0.37	0.05	(35)	-19.02 <sup>c</sup>	Wisconsin	0.43	0.05	(26)	-6.85
Mississippi	0.31	0.06	(39)	-31.27 <sup>c</sup>	Wyoming	-	-	(-)	-
Missouri	0.41	0.05	(29)	-11.12					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 13.26  
Leukemia

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	7.15	0.02		
High Five States				
Wisconsin	8.19 <sup>b</sup>	0.17	(01)	14.58
Indiana	7.90 <sup>b</sup>	0.16	(02)	10.53
Missouri	7.90 <sup>b</sup>	0.16	(03)	10.49
Idaho	7.85	0.34	(04)	9.81
Michigan	7.84 <sup>b</sup>	0.12	(05)	9.76
Low Five States				
D.C.	6.60	0.48	(47)	-7.67
New Mexico	6.38	0.26	(48)	-10.73
Nevada	6.38	0.24	(49)	-10.73
Arizona	6.18 <sup>b</sup>	0.14	(50)	-13.46
Hawaii	4.99 <sup>b</sup>	0.27	(51)	-30.22 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.49	0.18	(17)	4.87	Montana	7.82	0.39	(07)	9.39
Alaska	7.18	0.63	(28)	0.46	Nebraska	7.26	0.28	(27)	1.65
Arizona	6.18 <sup>b</sup>	0.14	(50)	-13.46	Nevada	6.38	0.24	(49)	-10.73
Arkansas	7.76	0.22	(08)	8.63	New Hampshire	6.86	0.32	(40)	-4.03
California	6.73 <sup>b</sup>	0.06	(45)	-5.82	New Jersey	6.96	0.12	(33)	-2.64
Colorado	6.66	0.18	(46)	-6.74	New Mexico	6.38	0.26	(48)	-10.73
Connecticut	6.77	0.19	(43)	-5.19	New York	6.80 <sup>b</sup>	0.08	(41)	-4.81
Delaware	6.88	0.39	(39)	-3.71	North Carolina	6.96	0.13	(34)	-2.65
D.C.	6.60	0.48	(47)	-7.67	North Dakota	7.37	0.44	(22)	3.14
Florida	6.75 <sup>b</sup>	0.08	(44)	-5.58	Ohio	7.49	0.11	(18)	4.86
Georgia	7.02	0.14	(32)	-1.73	Oklahoma	7.71	0.20	(12)	7.89
Hawaii	4.99 <sup>b</sup>	0.27	(51)	-30.22 <sup>c</sup>	Oregon	7.27	0.19	(26)	1.69
Idaho	7.85	0.34	(04)	9.81	Pennsylvania	7.56 <sup>b</sup>	0.10	(16)	5.84
Illinois	7.45	0.11	(20)	4.31	Rhode Island	7.73	0.36	(10)	8.13
Indiana	7.90 <sup>b</sup>	0.16	(02)	10.53	South Carolina	6.90	0.18	(38)	-3.42
Iowa	7.83	0.21	(06)	9.63	South Dakota	7.31	0.40	(24)	2.24
Kansas	7.72	0.23	(11)	8.07	Tennessee	7.59	0.16	(15)	6.27
Kentucky	7.35	0.19	(23)	2.88	Texas	6.92	0.08	(37)	-3.21
Louisiana	7.27	0.18	(25)	1.75	Utah	7.11	0.27	(31)	-0.49
Maine	7.13	0.30	(30)	-0.15	Vermont	7.14	0.46	(29)	-0.15
Maryland	6.94	0.16	(36)	-2.90	Virginia	6.95	0.14	(35)	-2.71
Massachusetts	6.80	0.14	(42)	-4.82	Washington	7.41	0.16	(21)	3.68
Michigan	7.84 <sup>b</sup>	0.12	(05)	9.76	West Virginia	7.61	0.27	(14)	6.45
Minnesota	7.74	0.17	(09)	8.26	Wisconsin	8.19 <sup>b</sup>	0.17	(01)	14.58
Mississippi	7.64	0.23	(13)	6.90	Wyoming	7.49	0.54	(19)	4.85
Missouri	7.90 <sup>b</sup>	0.16	(03)	10.49					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 13.27  
Leukemia

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	9.66	0.04		
High Five States				
Wisconsin	11.55 <sup>b</sup>	0.31	(01)	19.58 <sup>c</sup>
Missouri	11.04 <sup>b</sup>	0.29	(02)	14.32
Indiana	11.04 <sup>b</sup>	0.29	(03)	14.31
Iowa	10.81	0.39	(04)	11.88
Minnesota	10.80 <sup>b</sup>	0.32	(05)	11.76
Low Five States				
D.C.	8.85	0.91	(47)	-8.33
Nevada	8.67	0.43	(48)	-10.25
New Mexico	8.20	0.44	(49)	-15.06 <sup>c</sup>
Arizona	8.04 <sup>b</sup>	0.24	(50)	-16.71 <sup>c</sup>
Hawaii	6.67 <sup>b</sup>	0.46	(51)	-30.94 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	10.43	0.33	(10)	7.94	Montana	10.74	0.69	(06)	11.16
Alaska	8.92	1.08	(44)	-7.70	Nebraska	10.15	0.51	(17)	5.04
Arizona	8.04 <sup>b</sup>	0.24	(50)	-16.71 <sup>c</sup>	Nevada	8.67	0.43	(48)	-10.25
Arkansas	10.07	0.40	(19)	4.29	New Hampshire	8.96	0.57	(43)	-7.24
California	8.87 <sup>b</sup>	0.11	(46)	-8.15	New Jersey	9.37	0.23	(36)	-3.04
Colorado	9.06	0.34	(42)	-6.16	New Mexico	8.20	0.44	(49)	-15.06 <sup>c</sup>
Connecticut	9.12	0.34	(41)	-5.54	New York	9.26	0.15	(39)	-4.13
Delaware	9.22	0.70	(40)	-4.54	North Carolina	9.82	0.24	(25)	1.66
D.C.	8.85	0.91	(47)	-8.33	North Dakota	10.30	0.80	(13)	6.62
Florida	8.89 <sup>b</sup>	0.13	(45)	-7.92	Ohio	9.99	0.20	(22)	3.47
Georgia	9.66	0.26	(30)	0.01	Oklahoma	10.71	0.37	(07)	10.89
Hawaii	6.67 <sup>b</sup>	0.46	(51)	-30.94 <sup>c</sup>	Oregon	9.68	0.34	(29)	0.26
Idaho	9.63	0.56	(31)	-0.27	Pennsylvania	10.17	0.18	(16)	5.27
Illinois	10.23	0.20	(15)	5.87	Rhode Island	10.07	0.64	(20)	4.22
Indiana	11.04 <sup>b</sup>	0.29	(03)	14.31	South Carolina	9.57	0.33	(33)	-0.88
Iowa	10.81	0.39	(04)	11.88	South Dakota	9.61	0.70	(32)	-0.52
Kansas	10.32	0.41	(12)	6.81	Tennessee	10.70	0.30	(08)	10.78
Kentucky	9.80	0.34	(26)	1.43	Texas	9.35	0.15	(37)	-3.21
Louisiana	10.03	0.35	(21)	3.88	Utah	9.70	0.49	(28)	0.40
Maine	10.42	0.58	(11)	7.85	Vermont	9.96	0.85	(24)	3.09
Maryland	9.49	0.30	(35)	-1.75	Virginia	9.54	0.26	(34)	-1.23
Massachusetts	9.27	0.26	(38)	-4.06	Washington	10.24	0.28	(14)	6.00
Michigan	10.64 <sup>b</sup>	0.23	(09)	10.15	West Virginia	10.08	0.48	(18)	4.35
Minnesota	10.80 <sup>b</sup>	0.32	(05)	11.76	Wisconsin	11.55 <sup>b</sup>	0.31	(01)	19.58 <sup>c</sup>
Mississippi	9.97	0.42	(23)	3.23	Wyoming	9.74	0.93	(27)	0.86
Missouri	11.04 <sup>b</sup>	0.29	(02)	14.32					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 13.28  
Leukemia

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.37	0.02		
High Five States				
Idaho	6.40	0.41	(01)	19.07 <sup>c</sup>
Rhode Island	6.24	0.43	(02)	16.13 <sup>c</sup>
Arkansas	6.16	0.27	(03)	14.72
Mississippi	6.06	0.27	(04)	12.82
Kansas	5.93	0.27	(05)	10.29
Low Five States				
Colorado	4.99	0.21	(47)	-7.15
Maine	4.96	0.34	(48)	-7.65
Arizona	4.65 <sup>b</sup>	0.17	(49)	-13.49
Nevada	4.54	0.28	(50)	-15.60 <sup>c</sup>
Hawaii	3.59 <sup>b</sup>	0.31	(51)	-33.17 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.58	0.20	(21)	3.86	Montana	5.60	0.44	(19)	4.28
Alaska	5.81	0.76	(10)	8.20	Nebraska	5.13	0.31	(39)	-4.52
Arizona	4.65 <sup>b</sup>	0.17	(49)	-13.49	Nevada	4.54	0.28	(50)	-15.60 <sup>c</sup>
Arkansas	6.16	0.27	(03)	14.72	New Hampshire	5.46	0.38	(24)	1.68
California	5.16	0.07	(37)	-3.99	New Jersey	5.36	0.14	(28)	-0.17
Colorado	4.99	0.21	(47)	-7.15	New Mexico	4.99	0.31	(46)	-7.06
Connecticut	5.22	0.22	(33)	-2.80	New York	5.14	0.09	(38)	-4.27
Delaware	5.26	0.45	(32)	-2.12	North Carolina	5.01	0.14	(45)	-6.81
D.C.	5.01	0.54	(44)	-6.70	North Dakota	5.41	0.52	(26)	0.74
Florida	5.04 <sup>b</sup>	0.09	(41)	-6.26	Ohio	5.79	0.13	(11)	7.72
Georgia	5.29	0.15	(30)	-1.50	Oklahoma	5.60	0.23	(20)	4.19
Hawaii	3.59 <sup>b</sup>	0.31	(51)	-33.17 <sup>c</sup>	Oregon	5.45	0.22	(25)	1.39
Idaho	6.40	0.41	(01)	19.07 <sup>c</sup>	Pennsylvania	5.76	0.12	(12)	7.24
Illinois	5.63	0.12	(17)	4.83	Rhode Island	6.24	0.43	(02)	16.13 <sup>c</sup>
Indiana	5.84	0.18	(09)	8.71	South Carolina	5.03	0.20	(42)	-6.30
Iowa	5.87	0.25	(07)	9.18	South Dakota	5.51	0.47	(23)	2.59
Kansas	5.93	0.27	(05)	10.29	Tennessee	5.58	0.18	(22)	3.81
Kentucky	5.65	0.21	(16)	5.19	Texas	5.18	0.10	(35)	-3.51
Louisiana	5.41	0.21	(27)	0.60	Utah	5.13	0.31	(40)	-4.58
Maine	4.96	0.34	(48)	-7.65	Vermont	5.03	0.52	(43)	-6.33
Maryland	5.20	0.18	(34)	-3.16	Virginia	5.17	0.16	(36)	-3.87
Massachusetts	5.27	0.16	(31)	-1.84	Washington	5.34	0.18	(29)	-0.58
Michigan	5.86 <sup>b</sup>	0.14	(08)	9.11	West Virginia	5.87	0.31	(06)	9.33
Minnesota	5.69	0.20	(15)	5.82	Wisconsin	5.76	0.19	(13)	7.21
Mississippi	6.06	0.27	(04)	12.82	Wyoming	5.61	0.63	(18)	4.49
Missouri	5.71	0.18	(14)	6.29					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.18  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.33	0.02		
High Five States				
Hawaii	7.49 <sup>b</sup>	0.33	(01)	40.41 <sup>c</sup>
Louisiana	7.11 <sup>b</sup>	0.18	(02)	33.37 <sup>c</sup>
Texas	7.11 <sup>b</sup>	0.08	(03)	33.25 <sup>c</sup>
D.C.	6.95	0.49	(04)	30.38 <sup>c</sup>
California	6.59 <sup>b</sup>	0.06	(05)	23.64 <sup>c</sup>
Low Five States				
Idaho	3.96 <sup>b</sup>	0.24	(47)	-25.70 <sup>c</sup>
Vermont	3.78 <sup>b</sup>	0.33	(48)	-29.05 <sup>c</sup>
Nebraska	3.78 <sup>b</sup>	0.20	(49)	-29.07 <sup>c</sup>
Montana	3.77 <sup>b</sup>	0.26	(50)	-29.35 <sup>c</sup>
Utah	3.44 <sup>b</sup>	0.19	(51)	-35.48 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.40	0.15	(14)	1.21	Montana	3.77 <sup>b</sup>	0.26	(50)	-29.35 <sup>c</sup>
Alaska	6.02	0.54	(08)	12.94	Nebraska	3.78 <sup>b</sup>	0.20	(49)	-29.07 <sup>c</sup>
Arizona	5.05	0.13	(21)	-5.40	Nevada	5.95	0.23	(09)	11.66
Arkansas	5.75	0.19	(10)	7.77	New Hampshire	4.44	0.26	(40)	-16.68 <sup>c</sup>
California	6.59 <sup>b</sup>	0.06	(05)	23.64 <sup>c</sup>	New Jersey	5.07	0.10	(20)	-4.88
Colorado	4.59 <sup>b</sup>	0.15	(37)	-13.95	New Mexico	6.46 <sup>b</sup>	0.25	(06)	21.06 <sup>c</sup>
Connecticut	4.90	0.16	(27)	-8.15	New York	5.45	0.07	(13)	2.14
Delaware	5.04	0.33	(22)	-5.42	North Carolina	4.76 <sup>b</sup>	0.10	(31)	-10.81
D.C.	6.95	0.49	(04)	30.38 <sup>c</sup>	North Dakota	4.12 <sup>b</sup>	0.33	(44)	-22.82 <sup>c</sup>
Florida	5.12	0.07	(18)	-3.98	Ohio	4.59 <sup>b</sup>	0.08	(38)	-13.99
Georgia	4.60 <sup>b</sup>	0.11	(35)	-13.68	Oklahoma	5.03	0.16	(23)	-5.65
Hawaii	7.49 <sup>b</sup>	0.33	(01)	40.41 <sup>c</sup>	Oregon	4.76 <sup>b</sup>	0.15	(32)	-10.83
Idaho	3.96 <sup>b</sup>	0.24	(47)	-25.70 <sup>c</sup>	Pennsylvania	4.91 <sup>b</sup>	0.08	(26)	-7.90
Illinois	5.15	0.09	(16)	-3.38	Rhode Island	5.28	0.30	(15)	-1.00
Indiana	4.72 <sup>b</sup>	0.12	(33)	-11.52	South Carolina	4.87	0.15	(28)	-8.61
Iowa	4.11 <sup>b</sup>	0.15	(45)	-22.90 <sup>c</sup>	South Dakota	4.04 <sup>b</sup>	0.30	(46)	-24.18 <sup>c</sup>
Kansas	4.18 <sup>b</sup>	0.17	(43)	-21.67 <sup>c</sup>	Tennessee	4.96	0.12	(25)	-7.08
Kentucky	4.85	0.15	(29)	-9.08	Texas	7.11 <sup>b</sup>	0.08	(03)	33.25 <sup>c</sup>
Louisiana	7.11 <sup>b</sup>	0.18	(02)	33.37 <sup>c</sup>	Utah	3.44 <sup>b</sup>	0.19	(51)	-35.48 <sup>c</sup>
Maine	4.39 <sup>b</sup>	0.23	(41)	-17.61 <sup>c</sup>	Vermont	3.78 <sup>b</sup>	0.33	(48)	-29.05 <sup>c</sup>
Maryland	5.12	0.14	(17)	-3.98	Virginia	4.71 <sup>b</sup>	0.11	(34)	-11.69
Massachusetts	5.49	0.12	(12)	2.90	Washington	5.61	0.13	(11)	5.27
Michigan	5.09	0.10	(19)	-4.56	West Virginia	4.54 <sup>b</sup>	0.20	(39)	-14.92
Minnesota	4.60 <sup>b</sup>	0.13	(36)	-13.84	Wisconsin	4.76 <sup>b</sup>	0.13	(30)	-10.73
Mississippi	6.43 <sup>b</sup>	0.21	(07)	20.54 <sup>c</sup>	Wyoming	4.39	0.41	(42)	-17.64 <sup>c</sup>
Missouri	5.01	0.13	(24)	-6.12					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.19  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.88	0.03		
High Five States				
D.C.	11.38 <sup>b</sup>	0.96	(01)	44.47 <sup>c</sup>
Louisiana	10.79 <sup>b</sup>	0.34	(02)	37.01 <sup>c</sup>
Hawaii	10.61 <sup>b</sup>	0.57	(03)	34.63 <sup>c</sup>
Texas	10.56 <sup>b</sup>	0.16	(04)	34.03 <sup>c</sup>
Mississippi	9.90 <sup>b</sup>	0.41	(05)	25.70 <sup>c</sup>
Low Five States				
South Dakota	5.35 <sup>b</sup>	0.52	(47)	-32.14 <sup>c</sup>
North Dakota	5.25 <sup>b</sup>	0.57	(48)	-33.32 <sup>c</sup>
Vermont	5.07 <sup>b</sup>	0.57	(49)	-35.68 <sup>c</sup>
Montana	4.75 <sup>b</sup>	0.44	(50)	-39.72 <sup>c</sup>
Utah	4.69 <sup>b</sup>	0.33	(51)	-40.47 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	8.05	0.28	(16)	2.20	Montana	4.75 <sup>b</sup>	0.44	(50)	-39.72 <sup>c</sup>
Alaska	8.64	0.98	(09)	9.70	Nebraska	5.65 <sup>b</sup>	0.37	(46)	-28.30 <sup>c</sup>
Arizona	7.19	0.22	(29)	-8.79	Nevada	8.22	0.39	(14)	4.41
Arkansas	8.29	0.35	(12)	5.29	New Hampshire	6.39	0.46	(38)	-18.88 <sup>c</sup>
California	9.52 <sup>b</sup>	0.11	(06)	20.79 <sup>c</sup>	New Jersey	7.49	0.19	(20)	-4.94
Colorado	6.29 <sup>b</sup>	0.26	(40)	-20.09 <sup>c</sup>	New Mexico	9.37	0.45	(07)	18.96 <sup>c</sup>
Connecticut	7.55	0.30	(19)	-4.11	New York	8.25	0.14	(13)	4.73
Delaware	8.35	0.63	(11)	6.06	North Carolina	7.12 <sup>b</sup>	0.19	(30)	-9.67
D.C.	11.38 <sup>b</sup>	0.96	(01)	44.47 <sup>c</sup>	North Dakota	5.25 <sup>b</sup>	0.57	(48)	-33.32 <sup>c</sup>
Florida	7.85	0.12	(17)	-0.35	Ohio	6.85 <sup>b</sup>	0.16	(34)	-13.06
Georgia	6.83 <sup>b</sup>	0.20	(35)	-13.31	Oklahoma	7.31	0.29	(26)	-7.14
Hawaii	10.61 <sup>b</sup>	0.57	(03)	34.63 <sup>c</sup>	Oregon	6.70 <sup>b</sup>	0.27	(36)	-14.95
Idaho	5.85 <sup>b</sup>	0.42	(44)	-25.78 <sup>c</sup>	Pennsylvania	7.41	0.15	(24)	-5.98
Illinois	7.44	0.17	(22)	-5.50	Rhode Island	8.45	0.58	(10)	7.24
Indiana	7.01 <sup>b</sup>	0.23	(31)	-11.06	South Carolina	7.43	0.27	(23)	-5.68
Iowa	5.83 <sup>b</sup>	0.28	(45)	-26.00 <sup>c</sup>	South Dakota	5.35 <sup>b</sup>	0.52	(47)	-32.14 <sup>c</sup>
Kansas	6.15 <sup>b</sup>	0.31	(42)	-21.92 <sup>c</sup>	Tennessee	7.46	0.23	(21)	-5.33
Kentucky	7.00	0.28	(32)	-11.17	Texas	10.56 <sup>b</sup>	0.16	(04)	34.03 <sup>c</sup>
Louisiana	10.79 <sup>b</sup>	0.34	(02)	37.01 <sup>c</sup>	Utah	4.69 <sup>b</sup>	0.33	(51)	-40.47 <sup>c</sup>
Maine	6.39	0.43	(39)	-18.93 <sup>c</sup>	Vermont	5.07 <sup>b</sup>	0.57	(49)	-35.68 <sup>c</sup>
Maryland	7.81	0.26	(18)	-0.91	Virginia	7.29	0.22	(27)	-7.41
Massachusetts	8.87 <sup>b</sup>	0.24	(08)	12.57	Washington	8.14	0.24	(15)	3.36
Michigan	7.26 <sup>b</sup>	0.18	(28)	-7.88	West Virginia	6.18 <sup>b</sup>	0.36	(41)	-21.53 <sup>c</sup>
Minnesota	6.56 <sup>b</sup>	0.24	(37)	-16.67 <sup>c</sup>	Wisconsin	6.93 <sup>b</sup>	0.23	(33)	-12.02
Mississippi	9.90 <sup>b</sup>	0.41	(05)	25.70 <sup>c</sup>	Wyoming	5.99	0.72	(43)	-24.01 <sup>c</sup>
Missouri	7.34	0.23	(25)	-6.77					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.20  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.23	0.02		
High Five States				
Hawaii	4.65 <sup>b</sup>	0.34	(01)	43.78 <sup>c</sup>
Texas	4.26 <sup>b</sup>	0.09	(02)	31.74 <sup>c</sup>
Louisiana	4.22 <sup>b</sup>	0.18	(03)	30.39 <sup>c</sup>
California	4.11 <sup>b</sup>	0.07	(04)	27.19 <sup>c</sup>
New Mexico	3.95	0.27	(05)	22.06 <sup>c</sup>
Low Five States				
Kansas	2.49 <sup>b</sup>	0.17	(47)	-23.02 <sup>c</sup>
Utah	2.36 <sup>b</sup>	0.21	(48)	-26.85 <sup>c</sup>
Idaho	2.30 <sup>b</sup>	0.25	(49)	-28.94 <sup>c</sup>
Delaware	2.27	0.30	(50)	-29.91 <sup>c</sup>
Nebraska	2.24 <sup>b</sup>	0.21	(51)	-30.66 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.32	0.15	(12)	2.57	Montana	2.86	0.32	(33)	-11.39
Alaska	3.62	0.58	(09)	12.08	Nebraska	2.24 <sup>b</sup>	0.21	(51)	-30.66 <sup>c</sup>
Arizona	3.17	0.14	(16)	-2.01	Nevada	3.83	0.25	(06)	18.35 <sup>c</sup>
Arkansas	3.65	0.20	(08)	12.76	New Hampshire	2.76	0.27	(43)	-14.77
California	4.11 <sup>b</sup>	0.07	(04)	27.19 <sup>c</sup>	New Jersey	3.14	0.11	(18)	-2.90
Colorado	3.13	0.17	(19)	-3.21	New Mexico	3.95	0.27	(05)	22.06 <sup>c</sup>
Connecticut	2.81	0.16	(39)	-13.11	New York	3.22	0.07	(15)	-0.38
Delaware	2.27	0.30	(50)	-29.91 <sup>c</sup>	North Carolina	2.84 <sup>b</sup>	0.11	(36)	-12.07
D.C.	3.53	0.46	(10)	9.16	North Dakota	3.12	0.39	(20)	-3.56
Florida	2.77 <sup>b</sup>	0.07	(42)	-14.46	Ohio	2.79 <sup>b</sup>	0.09	(40)	-13.76
Georgia	2.86	0.11	(34)	-11.45	Oklahoma	3.10	0.17	(22)	-4.22
Hawaii	4.65 <sup>b</sup>	0.34	(01)	43.78 <sup>c</sup>	Oregon	3.04	0.17	(24)	-5.91
Idaho	2.30 <sup>b</sup>	0.25	(49)	-28.94 <sup>c</sup>	Pennsylvania	2.95 <sup>b</sup>	0.08	(28)	-8.88
Illinois	3.31	0.10	(13)	2.36	Rhode Island	2.84	0.28	(35)	-12.02
Indiana	2.90	0.13	(31)	-10.20	South Carolina	2.78	0.15	(41)	-14.04
Iowa	2.67	0.16	(46)	-17.29 <sup>c</sup>	South Dakota	2.87	0.34	(32)	-11.21
Kansas	2.49 <sup>b</sup>	0.17	(47)	-23.02 <sup>c</sup>	Tennessee	2.93	0.13	(30)	-9.41
Kentucky	3.17	0.16	(17)	-2.10	Texas	4.26 <sup>b</sup>	0.09	(02)	31.74 <sup>c</sup>
Louisiana	4.22 <sup>b</sup>	0.18	(03)	30.39 <sup>c</sup>	Utah	2.36 <sup>b</sup>	0.21	(48)	-26.85 <sup>c</sup>
Maine	2.81	0.25	(38)	-13.09	Vermont	2.72	0.39	(44)	-15.95 <sup>c</sup>
Maryland	2.99	0.14	(26)	-7.56	Virginia	2.70 <sup>b</sup>	0.11	(45)	-16.47 <sup>c</sup>
Massachusetts	2.84	0.12	(37)	-12.20	Washington	3.40	0.14	(11)	5.08
Michigan	3.30	0.11	(14)	2.04	West Virginia	3.09	0.22	(23)	-4.58
Minnesota	2.97	0.14	(27)	-8.26	Wisconsin	2.93	0.13	(29)	-9.37
Mississippi	3.82	0.21	(07)	18.02 <sup>c</sup>	Wyoming	3.04	0.46	(25)	-5.98
Missouri	3.10	0.13	(21)	-4.10					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 15.24  
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	51.61	0.06		
High Five States				
Kentucky	75.70 <sup>b</sup>	0.59	(01)	46.67 <sup>c</sup>
Arkansas	67.00 <sup>b</sup>	0.65	(02)	29.80 <sup>c</sup>
West Virginia	66.88 <sup>b</sup>	0.77	(03)	29.58 <sup>c</sup>
Tennessee	66.68 <sup>b</sup>	0.46	(04)	29.20 <sup>c</sup>
Mississippi	66.29 <sup>b</sup>	0.67	(05)	28.44 <sup>c</sup>
Low Five States				
California	40.82 <sup>b</sup>	0.16	(47)	-20.91 <sup>c</sup>
Hawaii	38.24 <sup>b</sup>	0.73	(48)	-25.91 <sup>c</sup>
Colorado	37.93 <sup>b</sup>	0.44	(49)	-26.50 <sup>c</sup>
New Mexico	36.58 <sup>b</sup>	0.61	(50)	-29.12 <sup>c</sup>
Utah	22.55 <sup>b</sup>	0.49	(51)	-56.32 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	61.52 <sup>b</sup>	0.50	(10)	19.20 <sup>c</sup>	Montana	49.79	0.97	(33)	-3.52
Alaska	53.51	1.68	(21)	3.67	Nebraska	47.84 <sup>b</sup>	0.72	(38)	-7.30 <sup>c</sup>
Arizona	42.06 <sup>b</sup>	0.37	(46)	-18.50 <sup>c</sup>	Nevada	55.89 <sup>b</sup>	0.71	(16)	8.29 <sup>c</sup>
Arkansas	67.00 <sup>b</sup>	0.65	(02)	29.80 <sup>c</sup>	New Hampshire	51.98	0.87	(28)	0.72
California	40.82 <sup>b</sup>	0.16	(47)	-20.91 <sup>c</sup>	New Jersey	47.37 <sup>b</sup>	0.32	(39)	-8.22 <sup>c</sup>
Colorado	37.93 <sup>b</sup>	0.44	(49)	-26.50 <sup>c</sup>	New Mexico	36.58 <sup>b</sup>	0.61	(50)	-29.12 <sup>c</sup>
Connecticut	46.92 <sup>b</sup>	0.49	(40)	-9.10 <sup>c</sup>	New York	44.57 <sup>b</sup>	0.21	(43)	-13.65 <sup>c</sup>
Delaware	60.22 <sup>b</sup>	1.14	(11)	16.67 <sup>c</sup>	North Carolina	58.14 <sup>b</sup>	0.36	(14)	12.64 <sup>c</sup>
D.C.	48.51	1.30	(36)	-6.02 <sup>c</sup>	North Dakota	45.42 <sup>b</sup>	1.13	(42)	-12.01 <sup>c</sup>
Florida	51.27	0.21	(29)	-0.65	Ohio	58.82 <sup>b</sup>	0.31	(13)	13.96 <sup>c</sup>
Georgia	55.25 <sup>b</sup>	0.38	(18)	7.05 <sup>c</sup>	Oklahoma	62.64 <sup>b</sup>	0.57	(07)	21.36 <sup>c</sup>
Hawaii	38.24 <sup>b</sup>	0.73	(48)	-25.91 <sup>c</sup>	Oregon	52.33	0.52	(26)	1.39
Idaho	42.45 <sup>b</sup>	0.78	(45)	-17.75 <sup>c</sup>	Pennsylvania	52.51	0.26	(24)	1.73
Illinois	53.36 <sup>b</sup>	0.29	(23)	3.39	Rhode Island	53.37	0.95	(22)	3.41
Indiana	61.90 <sup>b</sup>	0.44	(08)	19.93 <sup>c</sup>	South Carolina	57.57 <sup>b</sup>	0.50	(15)	11.55 <sup>c</sup>
Iowa	52.15	0.55	(27)	1.04	South Dakota	48.74	1.05	(35)	-5.56 <sup>c</sup>
Kansas	53.92 <sup>b</sup>	0.61	(20)	4.48	Tennessee	66.68 <sup>b</sup>	0.46	(04)	29.20 <sup>c</sup>
Kentucky	75.70 <sup>b</sup>	0.59	(01)	46.67 <sup>c</sup>	Texas	49.13 <sup>b</sup>	0.23	(34)	-4.81
Louisiana	62.80 <sup>b</sup>	0.54	(06)	21.68 <sup>c</sup>	Utah	22.55 <sup>b</sup>	0.49	(51)	-56.32 <sup>c</sup>
Maine	59.06 <sup>b</sup>	0.86	(12)	14.43 <sup>c</sup>	Vermont	51.21	1.22	(30)	-0.79
Maryland	52.47	0.44	(25)	1.67	Virginia	54.41 <sup>b</sup>	0.39	(19)	5.42 <sup>c</sup>
Massachusetts	51.19	0.38	(31)	-0.83	Washington	50.15 <sup>b</sup>	0.41	(32)	-2.83
Michigan	55.41 <sup>b</sup>	0.33	(17)	7.35 <sup>c</sup>	West Virginia	66.88 <sup>b</sup>	0.77	(03)	29.58 <sup>c</sup>
Minnesota	45.44 <sup>b</sup>	0.42	(41)	-11.95 <sup>c</sup>	Wisconsin	48.44 <sup>b</sup>	0.41	(37)	-6.15 <sup>c</sup>
Mississippi	66.29 <sup>b</sup>	0.67	(05)	28.44 <sup>c</sup>	Wyoming	44.40 <sup>b</sup>	1.31	(44)	-13.98 <sup>c</sup>
Missouri	61.78 <sup>b</sup>	0.44	(09)	19.69 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 15.25  
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	67.38	0.10		
High Five States				
Kentucky	102.97 <sup>b</sup>	1.06	(01)	52.81 <sup>c</sup>
Mississippi	98.93 <sup>b</sup>	1.29	(02)	46.82 <sup>c</sup>
Tennessee	93.95 <sup>b</sup>	0.85	(03)	39.43 <sup>c</sup>
Arkansas	93.20 <sup>b</sup>	1.18	(04)	38.32 <sup>c</sup>
Alabama	90.25 <sup>b</sup>	0.94	(05)	33.94 <sup>c</sup>
Low Five States				
Hawaii	51.83 <sup>b</sup>	1.28	(47)	-23.08 <sup>c</sup>
California	50.35 <sup>b</sup>	0.27	(48)	-25.28 <sup>c</sup>
Colorado	46.07 <sup>b</sup>	0.75	(49)	-31.64 <sup>c</sup>
New Mexico	45.50 <sup>b</sup>	1.03	(50)	-32.47 <sup>c</sup>
Utah	29.47 <sup>b</sup>	0.84	(51)	-56.27 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	90.25 <sup>b</sup>	0.94	(05)	33.94 <sup>c</sup>	Montana	59.51 <sup>b</sup>	1.58	(39)	-11.68 <sup>c</sup>
Alaska	62.29	2.74	(35)	-7.56 <sup>c</sup>	Nebraska	64.12	1.26	(29)	-4.85
Arizona	52.10 <sup>b</sup>	0.61	(45)	-22.68 <sup>c</sup>	Nevada	62.75 <sup>b</sup>	1.11	(33)	-6.88 <sup>c</sup>
Arkansas	93.20 <sup>b</sup>	1.18	(04)	38.32 <sup>c</sup>	New Hampshire	63.43	1.48	(31)	-5.87 <sup>c</sup>
California	50.35 <sup>b</sup>	0.27	(48)	-25.28 <sup>c</sup>	New Jersey	59.72 <sup>b</sup>	0.56	(37)	-11.37 <sup>c</sup>
Colorado	46.07 <sup>b</sup>	0.75	(49)	-31.64 <sup>c</sup>	New Mexico	45.50 <sup>b</sup>	1.03	(50)	-32.47 <sup>c</sup>
Connecticut	58.47 <sup>b</sup>	0.86	(41)	-13.22 <sup>c</sup>	New York	56.59 <sup>b</sup>	0.37	(43)	-16.01 <sup>c</sup>
Delaware	73.70	1.93	(16)	9.37 <sup>c</sup>	North Carolina	81.14 <sup>b</sup>	0.67	(12)	20.41 <sup>c</sup>
D.C.	68.56	2.49	(24)	1.75	North Dakota	59.26 <sup>b</sup>	1.92	(40)	-12.06 <sup>c</sup>
Florida	65.08 <sup>b</sup>	0.35	(28)	-3.42	Ohio	78.54 <sup>b</sup>	0.55	(14)	16.56 <sup>c</sup>
Georgia	78.90 <sup>b</sup>	0.72	(13)	17.10 <sup>c</sup>	Oklahoma	83.97 <sup>b</sup>	1.01	(08)	24.61 <sup>c</sup>
Hawaii	51.83 <sup>b</sup>	1.28	(47)	-23.08 <sup>c</sup>	Oregon	62.90 <sup>b</sup>	0.86	(32)	-6.66 <sup>c</sup>
Idaho	52.02 <sup>b</sup>	1.30	(46)	-22.80 <sup>c</sup>	Pennsylvania	69.94 <sup>b</sup>	0.47	(21)	3.79
Illinois	69.87 <sup>b</sup>	0.52	(22)	3.69	Rhode Island	69.00	1.68	(23)	2.41
Indiana	82.78 <sup>b</sup>	0.79	(10)	22.85 <sup>c</sup>	South Carolina	81.70 <sup>b</sup>	0.93	(11)	21.24 <sup>c</sup>
Iowa	69.97	0.97	(20)	3.85	South Dakota	65.38	1.82	(27)	-2.97
Kansas	71.79 <sup>b</sup>	1.08	(18)	6.54 <sup>c</sup>	Tennessee	93.95 <sup>b</sup>	0.85	(03)	39.43 <sup>c</sup>
Kentucky	102.97 <sup>b</sup>	1.06	(01)	52.81 <sup>c</sup>	Texas	65.74 <sup>b</sup>	0.40	(26)	-2.44
Louisiana	87.76 <sup>b</sup>	0.99	(07)	30.24 <sup>c</sup>	Utah	29.47 <sup>b</sup>	0.84	(51)	-56.27 <sup>c</sup>
Maine	75.61 <sup>b</sup>	1.50	(15)	12.22 <sup>c</sup>	Vermont	62.49	2.07	(34)	-7.26 <sup>c</sup>
Maryland	67.35	0.78	(25)	-0.04	Virginia	73.02 <sup>b</sup>	0.70	(17)	8.37 <sup>c</sup>
Massachusetts	64.04 <sup>b</sup>	0.67	(30)	-4.95	Washington	59.69 <sup>b</sup>	0.67	(38)	-11.41 <sup>c</sup>
Michigan	71.52 <sup>b</sup>	0.57	(19)	6.14 <sup>c</sup>	West Virginia	89.08 <sup>b</sup>	1.37	(06)	32.20 <sup>c</sup>
Minnesota	57.00 <sup>b</sup>	0.72	(42)	-15.41 <sup>c</sup>	Wisconsin	61.35 <sup>b</sup>	0.70	(36)	-8.94 <sup>c</sup>
Mississippi	98.93 <sup>b</sup>	1.29	(02)	46.82 <sup>c</sup>	Wyoming	52.48 <sup>b</sup>	2.15	(44)	-22.11 <sup>c</sup>
Missouri	83.07 <sup>b</sup>	0.79	(09)	23.28 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 15.26  
Cancer of the Lung And Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	40.12	0.07		
High Five States				
Kentucky	56.15 <sup>b</sup>	0.67	(01)	39.95 <sup>c</sup>
West Virginia	50.80 <sup>b</sup>	0.91	(02)	26.62 <sup>c</sup>
Delaware	50.26 <sup>b</sup>	1.39	(03)	25.28 <sup>c</sup>
Nevada	50.05 <sup>b</sup>	0.91	(04)	24.74 <sup>c</sup>
Arkansas	47.37 <sup>b</sup>	0.74	(05)	18.07 <sup>c</sup>
Low Five States				
California	33.87 <sup>b</sup>	0.19	(47)	-15.58 <sup>c</sup>
Colorado	32.30 <sup>b</sup>	0.54	(48)	-19.50 <sup>c</sup>
New Mexico	29.48 <sup>b</sup>	0.74	(49)	-26.52 <sup>c</sup>
Hawaii	27.45 <sup>b</sup>	0.84	(50)	-31.58 <sup>c</sup>
Utah	16.92 <sup>b</sup>	0.58	(51)	-57.83 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	41.02	0.54	(27)	2.24	Montana	42.39	1.22	(22)	5.67 <sup>c</sup>
Alaska	46.33	2.13	(11)	15.47 <sup>c</sup>	Nebraska	35.86 <sup>b</sup>	0.84	(42)	-10.61 <sup>c</sup>
Arizona	33.92 <sup>b</sup>	0.45	(46)	-15.46 <sup>c</sup>	Nevada	50.05 <sup>b</sup>	0.91	(04)	24.74 <sup>c</sup>
Arkansas	47.37 <sup>b</sup>	0.74	(05)	18.07 <sup>c</sup>	New Hampshire	43.75	1.08	(16)	9.05 <sup>c</sup>
California	33.87 <sup>b</sup>	0.19	(47)	-15.58 <sup>c</sup>	New Jersey	39.06	0.39	(35)	-2.63
Colorado	32.30 <sup>b</sup>	0.54	(48)	-19.50 <sup>c</sup>	New Mexico	29.48 <sup>b</sup>	0.74	(49)	-26.52 <sup>c</sup>
Connecticut	39.10	0.60	(34)	-2.54	New York	36.40 <sup>b</sup>	0.25	(40)	-9.28 <sup>c</sup>
Delaware	50.26 <sup>b</sup>	1.39	(03)	25.28 <sup>c</sup>	North Carolina	41.85 <sup>b</sup>	0.40	(25)	4.32
D.C.	35.14	1.44	(44)	-12.42 <sup>c</sup>	North Dakota	35.41	1.37	(43)	-11.75 <sup>c</sup>
Florida	40.09	0.25	(30)	-0.08	Ohio	44.96 <sup>b</sup>	0.36	(13)	12.07 <sup>c</sup>
Georgia	38.87	0.42	(36)	-3.10	Oklahoma	46.83 <sup>b</sup>	0.67	(09)	16.74 <sup>c</sup>
Hawaii	27.45 <sup>b</sup>	0.84	(50)	-31.58 <sup>c</sup>	Oregon	44.34 <sup>b</sup>	0.64	(14)	10.51 <sup>c</sup>
Idaho	34.93 <sup>b</sup>	0.97	(45)	-12.94 <sup>c</sup>	Pennsylvania	40.34	0.31	(29)	0.55
Illinois	41.99 <sup>b</sup>	0.34	(24)	4.67	Rhode Island	43.35	1.14	(17)	8.06 <sup>c</sup>
Indiana	47.23 <sup>b</sup>	0.51	(07)	17.73 <sup>c</sup>	South Carolina	39.93	0.56	(31)	-0.46
Iowa	39.29	0.65	(32)	-2.07	South Dakota	36.31	1.24	(41)	-9.50 <sup>c</sup>
Kansas	40.93	0.72	(28)	2.03	Tennessee	47.16 <sup>b</sup>	0.51	(08)	17.55 <sup>c</sup>
Kentucky	56.15 <sup>b</sup>	0.67	(01)	39.95 <sup>c</sup>	Texas	36.87 <sup>b</sup>	0.26	(39)	-8.09 <sup>c</sup>
Louisiana	44.99 <sup>b</sup>	0.61	(12)	12.13 <sup>c</sup>	Utah	16.92 <sup>b</sup>	0.58	(51)	-57.83 <sup>c</sup>
Maine	47.33 <sup>b</sup>	1.04	(06)	17.96 <sup>c</sup>	Vermont	43.15	1.51	(20)	7.55 <sup>c</sup>
Maryland	42.15 <sup>b</sup>	0.52	(23)	5.07 <sup>c</sup>	Virginia	41.29	0.44	(26)	2.91
Massachusetts	42.72 <sup>b</sup>	0.46	(21)	6.47 <sup>c</sup>	Washington	43.16 <sup>b</sup>	0.51	(19)	7.58 <sup>c</sup>
Michigan	43.93 <sup>b</sup>	0.39	(15)	9.50 <sup>c</sup>	West Virginia	50.80 <sup>b</sup>	0.91	(02)	26.62 <sup>c</sup>
Minnesota	37.25 <sup>b</sup>	0.52	(38)	-7.15 <sup>c</sup>	Wisconsin	39.17	0.49	(33)	-2.37
Mississippi	43.27 <sup>b</sup>	0.72	(18)	7.85 <sup>c</sup>	Wyoming	38.17	1.66	(37)	-4.86
Missouri	46.37 <sup>b</sup>	0.51	(10)	15.57 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 16.18  
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.70	0.01		
High Five States				
Oklahoma	3.45 <sup>b</sup>	0.14	(01)	27.66 <sup>c</sup>
Wyoming	3.39	0.36	(02)	25.33 <sup>c</sup>
Kentucky	3.32 <sup>b</sup>	0.12	(03)	23.01 <sup>c</sup>
Utah	3.22	0.18	(04)	19.21 <sup>c</sup>
Colorado	3.21 <sup>b</sup>	0.12	(05)	18.73 <sup>c</sup>
Low Five States				
Alaska	2.15	0.31	(47)	-20.47 <sup>c</sup>
Louisiana	2.12 <sup>b</sup>	0.10	(48)	-21.58 <sup>c</sup>
North Dakota	1.93	0.23	(49)	-28.72 <sup>c</sup>
Hawaii	1.53 <sup>b</sup>	0.15	(50)	-43.22 <sup>c</sup>
D.C.	1.38 <sup>b</sup>	0.21	(51)	-48.95 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.87	0.11	(24)	6.30	Montana	2.65	0.22	(33)	-1.81
Alaska	2.15	0.31	(47)	-20.47 <sup>c</sup>	Nebraska	2.99	0.18	(14)	10.58
Arizona	2.74	0.09	(30)	1.47	Nevada	2.83	0.16	(26)	4.89
Arkansas	2.58	0.13	(36)	-4.45	New Hampshire	3.02	0.21	(12)	11.72
California	2.58	0.04	(37)	-4.66	New Jersey	2.75	0.08	(28)	1.89
Colorado	3.21 <sup>b</sup>	0.12	(05)	18.73 <sup>c</sup>	New Mexico	2.65	0.17	(31)	-1.78
Connecticut	2.81	0.12	(27)	3.98	New York	2.17 <sup>b</sup>	0.05	(46)	-19.62 <sup>c</sup>
Delaware	2.91	0.25	(20)	7.53	North Carolina	2.98	0.08	(16)	10.18
D.C.	1.38 <sup>b</sup>	0.21	(51)	-48.95 <sup>c</sup>	North Dakota	1.93	0.23	(49)	-28.72 <sup>c</sup>
Florida	2.92 <sup>b</sup>	0.05	(19)	7.90	Ohio	2.65	0.07	(32)	-1.80
Georgia	2.46	0.08	(42)	-9.01	Oklahoma	3.45 <sup>b</sup>	0.14	(01)	27.66 <sup>c</sup>
Hawaii	1.53 <sup>b</sup>	0.15	(50)	-43.22 <sup>c</sup>	Oregon	3.01	0.12	(13)	11.51
Idaho	3.19	0.21	(06)	17.98 <sup>c</sup>	Pennsylvania	2.87	0.06	(25)	6.06
Illinois	2.55	0.06	(38)	-5.58	Rhode Island	2.89	0.22	(23)	6.78
Indiana	2.89	0.09	(22)	6.88	South Carolina	2.52	0.11	(41)	-6.82
Iowa	2.59	0.12	(35)	-4.02	South Dakota	2.54	0.24	(40)	-6.09
Kansas	3.11	0.15	(10)	15.03 <sup>c</sup>	Tennessee	2.99	0.10	(15)	10.46
Kentucky	3.32 <sup>b</sup>	0.12	(03)	23.01 <sup>c</sup>	Texas	2.54	0.05	(39)	-6.04
Louisiana	2.12 <sup>b</sup>	0.10	(48)	-21.58 <sup>c</sup>	Utah	3.22	0.18	(04)	19.21 <sup>c</sup>
Maine	2.92	0.19	(18)	7.97	Vermont	3.17	0.31	(07)	17.32 <sup>c</sup>
Maryland	2.75	0.10	(29)	1.65	Virginia	2.89	0.09	(21)	7.10
Massachusetts	2.96	0.09	(17)	9.58	Washington	3.02	0.10	(11)	11.83
Michigan	2.42 <sup>b</sup>	0.07	(43)	-10.54	West Virginia	3.13	0.17	(09)	15.80 <sup>c</sup>
Minnesota	2.30 <sup>b</sup>	0.09	(45)	-14.89	Wisconsin	2.62	0.09	(34)	-3.20
Mississippi	2.32	0.13	(44)	-14.04	Wyoming	3.39	0.36	(02)	25.33 <sup>c</sup>
Missouri	3.16 <sup>b</sup>	0.10	(08)	16.97 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 16.19  
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	4.04	0.02		
High Five States				
Oklahoma	5.16 <sup>b</sup>	0.25	(01)	27.84 <sup>c</sup>
Kentucky	4.99 <sup>b</sup>	0.23	(02)	23.71 <sup>c</sup>
Utah	4.89	0.34	(03)	21.07 <sup>c</sup>
Wyoming	4.86	0.66	(04)	20.33 <sup>c</sup>
Delaware	4.83	0.49	(05)	19.64 <sup>c</sup>
Low Five States				
Minnesota	3.21 <sup>b</sup>	0.17	(47)	-20.56 <sup>c</sup>
Alaska	2.90	0.53	(48)	-28.12 <sup>c</sup>
North Dakota	2.80	0.42	(49)	-30.63 <sup>c</sup>
D.C.	2.24 <sup>b</sup>	0.46	(50)	-44.40 <sup>c</sup>
Hawaii	2.23 <sup>b</sup>	0.26	(51)	-44.78 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.36	0.21	(20)	8.08	Montana	3.34	0.37	(44)	-17.25 <sup>c</sup>
Alaska	2.90	0.53	(48)	-28.12 <sup>c</sup>	Nebraska	4.16	0.32	(28)	3.06
Arizona	4.20	0.17	(27)	4.12	Nevada	4.41	0.29	(14)	9.26
Arkansas	3.91	0.24	(33)	-3.14	New Hampshire	4.39	0.38	(17)	8.86
California	3.90	0.07	(34)	-3.25	New Jersey	4.07	0.15	(29)	0.94
Colorado	4.75	0.23	(07)	17.70 <sup>c</sup>	New Mexico	3.60	0.29	(42)	-10.81
Connecticut	4.33	0.23	(21)	7.19	New York	3.24 <sup>b</sup>	0.09	(46)	-19.66 <sup>c</sup>
Delaware	4.83	0.49	(05)	19.64 <sup>c</sup>	North Carolina	4.41	0.16	(16)	9.17
D.C.	2.24 <sup>b</sup>	0.46	(50)	-44.40 <sup>c</sup>	North Dakota	2.80	0.42	(49)	-30.63 <sup>c</sup>
Florida	4.41 <sup>b</sup>	0.09	(15)	9.18	Ohio	3.98	0.12	(31)	-1.29
Georgia	3.77	0.16	(37)	-6.67	Oklahoma	5.16 <sup>b</sup>	0.25	(01)	27.84 <sup>c</sup>
Hawaii	2.23 <sup>b</sup>	0.26	(51)	-44.78 <sup>c</sup>	Oregon	4.28	0.22	(22)	6.14
Idaho	4.75	0.39	(06)	17.79 <sup>c</sup>	Pennsylvania	4.28	0.12	(23)	5.99
Illinois	3.75	0.12	(38)	-7.06	Rhode Island	4.45	0.42	(13)	10.25
Indiana	4.37	0.18	(19)	8.22	South Carolina	3.96	0.21	(32)	-1.83
Iowa	3.61	0.22	(41)	-10.54	South Dakota	3.53	0.43	(43)	-12.61
Kansas	4.52	0.27	(12)	11.97	Tennessee	4.61	0.19	(09)	14.10
Kentucky	4.99 <sup>b</sup>	0.23	(02)	23.71 <sup>c</sup>	Texas	3.85	0.10	(35)	-4.69
Louisiana	3.26 <sup>b</sup>	0.19	(45)	-19.29 <sup>c</sup>	Utah	4.89	0.34	(03)	21.07 <sup>c</sup>
Maine	4.06	0.35	(30)	0.51	Vermont	4.73	0.57	(08)	17.19 <sup>c</sup>
Maryland	4.23	0.20	(26)	4.93	Virginia	4.25	0.17	(24)	5.32
Massachusetts	4.24	0.17	(25)	5.06	Washington	4.38	0.18	(18)	8.43
Michigan	3.64	0.13	(40)	-9.87	West Virginia	4.53	0.31	(11)	12.22
Minnesota	3.21 <sup>b</sup>	0.17	(47)	-20.56 <sup>c</sup>	Wisconsin	3.77	0.17	(36)	-6.52
Mississippi	3.74	0.26	(39)	-7.22	Wyoming	4.86	0.66	(04)	20.33 <sup>c</sup>
Missouri	4.58	0.18	(10)	13.44					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 16.20  
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.71	0.01		
High Five States				
Wyoming	2.28	0.41	(01)	33.03 <sup>c</sup>
Oklahoma	2.14	0.15	(02)	25.14 <sup>c</sup>
Montana	2.14	0.28	(03)	25.06 <sup>c</sup>
Nebraska	2.11	0.21	(04)	22.98 <sup>c</sup>
Missouri	2.09	0.11	(05)	21.97 <sup>c</sup>
Low Five States				
Mississippi	1.36	0.13	(47)	-20.63 <sup>c</sup>
Louisiana	1.29 <sup>b</sup>	0.10	(48)	-24.44 <sup>c</sup>
North Dakota	1.15	0.24	(49)	-32.77 <sup>c</sup>
Hawaii	0.92 <sup>b</sup>	0.16	(50)	-46.07 <sup>c</sup>
D.C.	0.87 <sup>b</sup>	0.22	(51)	-49.11 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.80	0.11	(26)	5.20	Montana	2.14	0.28	(03)	25.06 <sup>c</sup>
Alaska	1.38	0.34	(46)	-19.48 <sup>c</sup>	Nebraska	2.11	0.21	(04)	22.98 <sup>c</sup>
Arizona	1.52	0.09	(39)	-11.37	Nevada	1.45	0.15	(44)	-15.50 <sup>c</sup>
Arkansas	1.52	0.13	(40)	-11.46	New Hampshire	1.97	0.23	(16)	15.01 <sup>c</sup>
California	1.57 <sup>b</sup>	0.04	(36)	-8.35	New Jersey	1.82	0.08	(21)	6.33
Colorado	2.03	0.13	(12)	18.36 <sup>c</sup>	New Mexico	1.92	0.19	(18)	12.01
Connecticut	1.76	0.13	(28)	2.64	New York	1.43 <sup>b</sup>	0.05	(45)	-16.28 <sup>c</sup>
Delaware	1.48	0.24	(42)	-13.63	North Carolina	1.97	0.09	(14)	15.26 <sup>c</sup>
D.C.	0.87 <sup>b</sup>	0.22	(51)	-49.11 <sup>c</sup>	North Dakota	1.15	0.24	(49)	-32.77 <sup>c</sup>
Florida	1.72	0.05	(30)	0.54	Ohio	1.71	0.07	(31)	-0.21
Georgia	1.57	0.08	(37)	-8.61	Oklahoma	2.14	0.15	(02)	25.14 <sup>c</sup>
Hawaii	0.92 <sup>b</sup>	0.16	(50)	-46.07 <sup>c</sup>	Oregon	2.03	0.14	(11)	18.51 <sup>c</sup>
Idaho	1.81	0.22	(25)	5.92	Pennsylvania	1.85	0.07	(20)	8.06
Illinois	1.69	0.07	(34)	-1.25	Rhode Island	1.82	0.23	(23)	6.04
Indiana	1.80	0.10	(27)	4.89	South Carolina	1.46	0.11	(43)	-15.04 <sup>c</sup>
Iowa	1.81	0.14	(24)	5.95	South Dakota	1.76	0.28	(29)	2.56
Kansas	2.06	0.16	(08)	20.40 <sup>c</sup>	Tennessee	1.82	0.10	(22)	6.22
Kentucky	2.09	0.13	(06)	21.81 <sup>c</sup>	Texas	1.56	0.05	(38)	-8.98
Louisiana	1.29 <sup>b</sup>	0.10	(48)	-24.44 <sup>c</sup>	Utah	1.86	0.19	(19)	8.68
Maine	2.06	0.22	(09)	20.26 <sup>c</sup>	Vermont	1.97	0.32	(15)	15.04 <sup>c</sup>
Maryland	1.71	0.10	(32)	-0.24	Virginia	1.95	0.10	(17)	14.04
Massachusetts	2.08 <sup>b</sup>	0.10	(07)	21.57 <sup>c</sup>	Washington	1.98	0.11	(13)	15.41 <sup>c</sup>
Michigan	1.52	0.07	(41)	-11.46	West Virginia	2.06	0.19	(10)	20.26 <sup>c</sup>
Minnesota	1.65	0.11	(35)	-3.96	Wisconsin	1.70	0.10	(33)	-0.63
Mississippi	1.36	0.13	(47)	-20.63 <sup>c</sup>	Wyoming	2.28	0.41	(01)	33.03 <sup>c</sup>
Missouri	2.09	0.11	(05)	21.97 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.18  
Myeloma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.48	0.02		
High Five States				
D.C.	5.48 <sup>b</sup>	0.44	(01)	57.50 <sup>c</sup>
Alabama	4.22 <sup>b</sup>	0.13	(02)	21.37 <sup>c</sup>
North Carolina	4.12 <sup>b</sup>	0.10	(03)	18.41 <sup>c</sup>
South Carolina	4.04 <sup>b</sup>	0.13	(04)	16.14 <sup>c</sup>
Tennessee	4.03 <sup>b</sup>	0.11	(05)	15.83 <sup>c</sup>
Low Five States				
Florida	3.03 <sup>b</sup>	0.05	(47)	-12.84
Rhode Island	3.00	0.22	(48)	-13.74
Alaska	2.83	0.40	(49)	-18.79 <sup>c</sup>
Nevada	2.71 <sup>b</sup>	0.16	(50)	-22.11 <sup>c</sup>
Hawaii	2.46 <sup>b</sup>	0.19	(51)	-29.18 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.22 <sup>b</sup>	0.13	(02)	21.37 <sup>c</sup>	Montana	3.49	0.26	(25)	0.20
Alaska	2.83	0.40	(49)	-18.79 <sup>c</sup>	Nebraska	3.31	0.19	(37)	-4.81
Arizona	3.09 <sup>b</sup>	0.10	(46)	-11.12	Nevada	2.71 <sup>b</sup>	0.16	(50)	-22.11 <sup>c</sup>
Arkansas	3.50	0.15	(24)	0.61	New Hampshire	3.42	0.22	(33)	-1.67
California	3.22 <sup>b</sup>	0.04	(43)	-7.62	New Jersey	3.41	0.09	(35)	-2.07
Colorado	3.26	0.13	(39)	-6.27	New Mexico	3.19	0.18	(44)	-8.42
Connecticut	3.25	0.13	(41)	-6.76	New York	3.10 <sup>b</sup>	0.05	(45)	-10.86
Delaware	3.41	0.27	(34)	-2.06	North Carolina	4.12 <sup>b</sup>	0.10	(03)	18.41 <sup>c</sup>
D.C.	5.48 <sup>b</sup>	0.44	(01)	57.50 <sup>c</sup>	North Dakota	3.71	0.31	(14)	6.45
Florida	3.03 <sup>b</sup>	0.05	(47)	-12.84	Ohio	3.76 <sup>b</sup>	0.08	(12)	8.03
Georgia	3.66	0.10	(17)	5.26	Oklahoma	3.44	0.13	(30)	-1.13
Hawaii	2.46 <sup>b</sup>	0.19	(51)	-29.18 <sup>c</sup>	Oregon	3.70	0.14	(15)	6.31
Idaho	3.58	0.23	(20)	2.94	Pennsylvania	3.47	0.07	(28)	-0.32
Illinois	3.61	0.08	(19)	3.67	Rhode Island	3.00	0.22	(48)	-13.74
Indiana	3.54	0.10	(22)	1.63	South Carolina	4.04 <sup>b</sup>	0.13	(04)	16.14 <sup>c</sup>
Iowa	3.43	0.14	(32)	-1.52	South Dakota	3.43	0.28	(31)	-1.47
Kansas	3.87	0.16	(09)	11.32	Tennessee	4.03 <sup>b</sup>	0.11	(05)	15.83 <sup>c</sup>
Kentucky	3.54	0.13	(23)	1.63	Texas	3.48	0.06	(26)	-0.09
Louisiana	3.91	0.14	(07)	12.32	Utah	3.48	0.19	(27)	-0.16
Maine	3.26	0.20	(40)	-6.42	Vermont	3.30	0.31	(38)	-5.24
Maryland	3.98 <sup>b</sup>	0.12	(06)	14.25	Virginia	3.89 <sup>b</sup>	0.10	(08)	11.71
Massachusetts	3.24	0.10	(42)	-6.86	Washington	3.32	0.10	(36)	-4.49
Michigan	3.78	0.09	(11)	8.46	West Virginia	3.57	0.18	(21)	2.49
Minnesota	3.72	0.12	(13)	6.92	Wisconsin	3.65	0.11	(18)	4.93
Mississippi	3.69	0.16	(16)	5.97	Wyoming	3.45	0.37	(29)	-0.78
Missouri	3.78	0.11	(10)	8.58					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.19  
Myeloma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	4.41	0.03		
High Five States				
D.C.	7.18	0.83	(01)	62.90 <sup>c</sup>
Tennessee	5.24 <sup>b</sup>	0.21	(02)	18.82 <sup>c</sup>
Alabama	5.16	0.23	(03)	17.15 <sup>c</sup>
Maryland	5.10	0.22	(04)	15.75 <sup>c</sup>
North Carolina	5.08 <sup>b</sup>	0.17	(05)	15.13 <sup>c</sup>
Low Five States				
Florida	3.82 <sup>b</sup>	0.09	(47)	-13.25
Nevada	3.48	0.27	(48)	-20.98 <sup>c</sup>
Rhode Island	3.34	0.37	(49)	-24.33 <sup>c</sup>
Hawaii	3.09 <sup>b</sup>	0.31	(50)	-30.03 <sup>c</sup>
Alaska	3.08	0.59	(51)	-30.06 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.16	0.23	(03)	17.15 <sup>c</sup>	Montana	4.68	0.45	(18)	6.20
Alaska	3.08	0.59	(51)	-30.06 <sup>c</sup>	Nebraska	4.15	0.32	(41)	-5.80
Arizona	3.89	0.17	(44)	-11.68	Nevada	3.48	0.27	(48)	-20.98 <sup>c</sup>
Arkansas	4.39	0.26	(29)	-0.38	New Hampshire	4.24	0.39	(37)	-3.92
California	4.09 <sup>b</sup>	0.08	(42)	-7.33	New Jersey	4.18	0.15	(38)	-5.22
Colorado	4.26	0.23	(35)	-3.45	New Mexico	3.89	0.30	(45)	-11.85
Connecticut	4.17	0.23	(40)	-5.50	New York	3.93 <sup>b</sup>	0.10	(43)	-10.85
Delaware	4.42	0.48	(28)	0.30	North Carolina	5.08 <sup>b</sup>	0.17	(05)	15.13 <sup>c</sup>
D.C.	7.18	0.83	(01)	62.90 <sup>c</sup>	North Dakota	4.74	0.54	(16)	7.54
Florida	3.82 <sup>b</sup>	0.09	(47)	-13.25	Ohio	4.66	0.14	(20)	5.59
Georgia	4.53	0.18	(24)	2.75	Oklahoma	4.63	0.24	(21)	5.10
Hawaii	3.09 <sup>b</sup>	0.31	(50)	-30.03 <sup>c</sup>	Oregon	4.89	0.24	(11)	10.98
Idaho	4.77	0.40	(15)	8.19	Pennsylvania	4.30	0.12	(34)	-2.42
Illinois	4.56	0.14	(23)	3.35	Rhode Island	3.34	0.37	(49)	-24.33 <sup>c</sup>
Indiana	4.73	0.19	(17)	7.30	South Carolina	5.04	0.24	(06)	14.41
Iowa	4.48	0.25	(27)	1.70	South Dakota	4.31	0.47	(33)	-2.25
Kansas	4.96	0.29	(08)	12.52	Tennessee	5.24 <sup>b</sup>	0.21	(02)	18.82 <sup>c</sup>
Kentucky	4.81	0.24	(14)	9.10	Texas	4.35	0.10	(32)	-1.23
Louisiana	4.95	0.24	(09)	12.18	Utah	4.37	0.33	(31)	-0.94
Maine	4.25	0.36	(36)	-3.55	Vermont	3.85	0.53	(46)	-12.66
Maryland	5.10	0.22	(04)	15.75 <sup>c</sup>	Virginia	4.89	0.19	(12)	10.89
Massachusetts	4.37	0.18	(30)	-0.83	Washington	4.52	0.19	(26)	2.57
Michigan	5.00 <sup>b</sup>	0.15	(07)	13.39	West Virginia	4.90	0.33	(10)	11.19
Minnesota	4.62	0.21	(22)	4.88	Wisconsin	4.67	0.19	(19)	5.84
Mississippi	4.17	0.27	(39)	-5.42	Wyoming	4.53	0.62	(25)	2.66
Missouri	4.86	0.19	(13)	10.13					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.20  
Myeloma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.82	0.02		
High Five States				
D.C.	4.59	0.53	(01)	62.48 <sup>c</sup>
Alabama	3.54 <sup>b</sup>	0.16	(02)	25.45 <sup>c</sup>
North Carolina	3.46 <sup>b</sup>	0.12	(03)	22.55 <sup>c</sup>
South Carolina	3.40 <sup>b</sup>	0.16	(04)	20.44 <sup>c</sup>
Mississippi	3.32	0.20	(05)	17.64 <sup>c</sup>
Low Five States				
Massachusetts	2.46	0.11	(47)	-12.90
Washington	2.44	0.12	(48)	-13.68
Florida	2.41 <sup>b</sup>	0.06	(49)	-14.58
Nevada	2.07 <sup>b</sup>	0.19	(50)	-26.77 <sup>c</sup>
Hawaii	1.99 <sup>b</sup>	0.22	(51)	-29.52 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.54 <sup>b</sup>	0.16	(02)	25.45 <sup>c</sup>	Montana	2.56	0.29	(41)	-9.21
Alaska	2.47	0.50	(46)	-12.61	Nebraska	2.73	0.22	(30)	-3.26
Arizona	2.47	0.12	(45)	-12.37	Nevada	2.07 <sup>b</sup>	0.19	(50)	-26.77 <sup>c</sup>
Arkansas	2.83	0.18	(24)	0.04	New Hampshire	2.86	0.27	(22)	1.29
California	2.59 <sup>b</sup>	0.05	(39)	-8.36	New Jersey	2.88	0.10	(21)	1.91
Colorado	2.55	0.15	(42)	-9.83	New Mexico	2.65	0.22	(34)	-6.25
Connecticut	2.58	0.15	(40)	-8.77	New York	2.54 <sup>b</sup>	0.07	(43)	-9.97
Delaware	2.65	0.32	(35)	-6.31	North Carolina	3.46 <sup>b</sup>	0.12	(03)	22.55 <sup>c</sup>
D.C.	4.59	0.53	(01)	62.48 <sup>c</sup>	North Dakota	2.91	0.37	(20)	2.89
Florida	2.41 <sup>b</sup>	0.06	(49)	-14.58	Ohio	3.15 <sup>b</sup>	0.09	(10)	11.43
Georgia	3.11	0.12	(12)	10.03	Oklahoma	2.59	0.15	(38)	-8.31
Hawaii	1.99 <sup>b</sup>	0.22	(51)	-29.52 <sup>c</sup>	Oregon	2.81	0.16	(25)	-0.39
Idaho	2.73	0.27	(29)	-3.17	Pennsylvania	2.91	0.08	(18)	3.09
Illinois	2.99	0.09	(15)	6.05	Rhode Island	2.79	0.28	(26)	-1.17
Indiana	2.74	0.12	(28)	-2.87	South Carolina	3.40 <sup>b</sup>	0.16	(04)	20.44 <sup>c</sup>
Iowa	2.65	0.16	(33)	-6.24	South Dakota	2.70	0.33	(31)	-4.45
Kansas	3.14	0.20	(11)	11.30	Tennessee	3.20	0.13	(08)	13.25
Kentucky	2.68	0.14	(32)	-5.05	Texas	2.84	0.07	(23)	0.54
Louisiana	3.19	0.16	(09)	13.01	Utah	2.78	0.23	(27)	-1.54
Maine	2.59	0.24	(37)	-8.31	Vermont	2.95	0.39	(17)	4.43
Maryland	3.25	0.14	(06)	15.19 <sup>c</sup>	Virginia	3.25	0.12	(07)	14.92
Massachusetts	2.46	0.11	(47)	-12.90	Washington	2.44	0.12	(48)	-13.68
Michigan	2.91	0.10	(19)	2.95	West Virginia	2.63	0.20	(36)	-6.74
Minnesota	3.06	0.15	(13)	8.27	Wisconsin	2.95	0.13	(16)	4.46
Mississippi	3.32	0.20	(05)	17.64 <sup>c</sup>	Wyoming	2.48	0.43	(44)	-12.11
Missouri	3.03	0.13	(14)	7.18					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 19.22  
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	6.71	0.02		
High Five States				
West Virginia	7.83 <sup>b</sup>	0.27	(01)	16.66 <sup>c</sup>
Indiana	7.49 <sup>b</sup>	0.15	(02)	11.58
Michigan	7.43 <sup>b</sup>	0.12	(03)	10.75
Wisconsin	7.42 <sup>b</sup>	0.16	(04)	10.51
Kentucky	7.39 <sup>b</sup>	0.19	(05)	10.16
Low Five States				
Georgia	6.11 <sup>b</sup>	0.13	(47)	-8.91
Nevada	5.86 <sup>b</sup>	0.23	(48)	-12.66
New Mexico	5.61 <sup>b</sup>	0.24	(49)	-16.40 <sup>c</sup>
Hawaii	5.60 <sup>b</sup>	0.28	(50)	-16.62 <sup>c</sup>
D.C.	5.54	0.44	(51)	-17.45 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.71	0.17	(26)	0.02	Montana	6.88	0.36	(20)	2.43
Alaska	6.13	0.59	(46)	-8.68	Nebraska	7.19	0.27	(12)	7.16
Arizona	6.17 <sup>b</sup>	0.14	(44)	-8.04	Nevada	5.86 <sup>b</sup>	0.23	(48)	-12.66
Arkansas	6.67	0.21	(28)	-0.68	New Hampshire	6.38	0.31	(37)	-5.01
California	6.43 <sup>b</sup>	0.06	(33)	-4.22	New Jersey	6.85	0.12	(21)	2.09
Colorado	6.20	0.18	(43)	-7.64	New Mexico	5.61 <sup>b</sup>	0.24	(49)	-16.40 <sup>c</sup>
Connecticut	6.57	0.18	(30)	-2.14	New York	6.29 <sup>b</sup>	0.08	(40)	-6.34
Delaware	6.74	0.38	(25)	0.45	North Carolina	6.41	0.12	(35)	-4.50
D.C.	5.54	0.44	(51)	-17.45 <sup>c</sup>	North Dakota	6.42	0.41	(34)	-4.30
Florida	6.38 <sup>b</sup>	0.07	(36)	-5.00	Ohio	7.21 <sup>b</sup>	0.11	(10)	7.34
Georgia	6.11 <sup>b</sup>	0.13	(47)	-8.91	Oklahoma	7.20	0.19	(11)	7.32
Hawaii	5.60 <sup>b</sup>	0.28	(50)	-16.62 <sup>c</sup>	Oregon	7.28	0.19	(08)	8.44
Idaho	6.85	0.31	(22)	2.08	Pennsylvania	7.34 <sup>b</sup>	0.10	(07)	9.28
Illinois	6.97	0.11	(19)	3.83	Rhode Island	6.57	0.33	(29)	-2.09
Indiana	7.49 <sup>b</sup>	0.15	(02)	11.58	South Carolina	6.22	0.17	(42)	-7.37
Iowa	7.13	0.20	(13)	6.25	South Dakota	6.84	0.39	(23)	1.90
Kansas	7.23	0.22	(09)	7.74	Tennessee	7.10	0.15	(17)	5.78
Kentucky	7.39 <sup>b</sup>	0.19	(05)	10.16	Texas	6.47	0.08	(31)	-3.66
Louisiana	7.10	0.18	(16)	5.80	Utah	6.31	0.26	(38)	-5.94
Maine	7.38	0.31	(06)	9.95	Vermont	6.15	0.42	(45)	-8.39
Maryland	6.30	0.15	(39)	-6.09	Virginia	6.44	0.13	(32)	-4.00
Massachusetts	6.71	0.14	(27)	-0.01	Washington	7.07	0.15	(18)	5.37
Michigan	7.43 <sup>b</sup>	0.12	(03)	10.75	West Virginia	7.83 <sup>b</sup>	0.27	(01)	16.66 <sup>c</sup>
Minnesota	7.13	0.17	(14)	6.17	Wisconsin	7.42 <sup>b</sup>	0.16	(04)	10.51
Mississippi	6.27	0.21	(41)	-6.53	Wyoming	7.13	0.53	(15)	6.16
Missouri	6.75	0.15	(24)	0.59					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 19.23  
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	8.55	0.04		
High Five States				
Indiana	9.94 <sup>b</sup>	0.28	(01)	16.26 <sup>c</sup>
Kansas	9.69	0.40	(02)	13.34
West Virginia	9.58	0.46	(03)	12.01
Minnesota	9.54	0.30	(04)	11.52
Wisconsin	9.46	0.28	(05)	10.57
Low Five States				
Vermont	7.72	0.75	(47)	-9.78
Alaska	7.69	1.05	(48)	-10.05
Hawaii	7.18	0.48	(49)	-16.11 <sup>c</sup>
Nevada	6.77 <sup>b</sup>	0.36	(50)	-20.83 <sup>c</sup>
New Mexico	6.63 <sup>b</sup>	0.39	(51)	-22.46 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	8.52	0.30	(26)	-0.43	Montana	8.51	0.61	(27)	-0.55
Alaska	7.69	1.05	(48)	-10.05	Nebraska	9.02	0.48	(19)	5.44
Arizona	7.74	0.24	(46)	-9.56	Nevada	6.77 <sup>b</sup>	0.36	(50)	-20.83 <sup>c</sup>
Arkansas	8.56	0.36	(24)	0.09	New Hampshire	8.30	0.55	(31)	-2.95
California	8.22	0.11	(34)	-3.87	New Jersey	8.47	0.21	(29)	-1.01
Colorado	8.25	0.32	(32)	-3.56	New Mexico	6.63 <sup>b</sup>	0.39	(51)	-22.46 <sup>c</sup>
Connecticut	8.18	0.32	(36)	-4.36	New York	7.98 <sup>b</sup>	0.14	(41)	-6.70
Delaware	9.02	0.69	(18)	5.46	North Carolina	7.98	0.21	(40)	-6.66
D.C.	8.80	0.88	(21)	2.83	North Dakota	7.96	0.70	(43)	-6.97
Florida	8.03 <sup>b</sup>	0.13	(39)	-6.15	Ohio	9.45 <sup>b</sup>	0.19	(06)	10.54
Georgia	7.97	0.24	(42)	-6.83	Oklahoma	9.17	0.34	(14)	7.17
Hawaii	7.18	0.48	(49)	-16.11 <sup>c</sup>	Oregon	9.11	0.33	(16)	6.51
Idaho	8.24	0.52	(33)	-3.68	Pennsylvania	9.38 <sup>b</sup>	0.17	(07)	9.69
Illinois	9.05	0.19	(17)	5.85	Rhode Island	9.14	0.61	(15)	6.81
Indiana	9.94 <sup>b</sup>	0.28	(01)	16.26 <sup>c</sup>	South Carolina	7.85	0.30	(44)	-8.24
Iowa	9.23	0.36	(13)	7.85	South Dakota	8.72	0.67	(22)	1.98
Kansas	9.69	0.40	(02)	13.34	Tennessee	9.26	0.27	(11)	8.25
Kentucky	9.35	0.33	(08)	9.26	Texas	8.21	0.14	(35)	-4.04
Louisiana	9.34	0.33	(09)	9.22	Utah	7.84	0.44	(45)	-8.32
Maine	9.32	0.54	(10)	9.01	Vermont	7.72	0.75	(47)	-9.78
Maryland	8.09	0.28	(38)	-5.45	Virginia	8.33	0.24	(30)	-2.59
Massachusetts	8.66	0.25	(23)	1.21	Washington	8.95	0.26	(20)	4.58
Michigan	9.23	0.21	(12)	7.96	West Virginia	9.58	0.46	(03)	12.01
Minnesota	9.54	0.30	(04)	11.52	Wisconsin	9.46	0.28	(05)	10.57
Mississippi	8.53	0.38	(25)	-0.23	Wyoming	8.12	0.86	(37)	-5.11
Missouri	8.49	0.26	(28)	-0.71					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 19.24  
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.36	0.02		
High Five States				
West Virginia	6.52 <sup>b</sup>	0.32	(01)	21.61 <sup>c</sup>
Wyoming	6.34	0.67	(02)	18.34 <sup>c</sup>
Michigan	6.20 <sup>b</sup>	0.14	(03)	15.57 <sup>c</sup>
Kentucky	6.00	0.22	(04)	11.85
Maine	5.99	0.37	(05)	11.67
Low Five States				
New Mexico	4.77	0.30	(47)	-11.03
Colorado	4.66	0.20	(48)	-13.04
Mississippi	4.65	0.23	(49)	-13.30
Hawaii	4.36	0.33	(50)	-18.60 <sup>c</sup>
D.C.	3.25 <sup>b</sup>	0.43	(51)	-39.40 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.49	0.20	(22)	2.36	Montana	5.55	0.43	(18)	3.58
Alaska	5.10	0.72	(34)	-4.92	Nebraska	5.92	0.33	(08)	10.37
Arizona	4.86	0.17	(44)	-9.26	Nevada	4.95	0.29	(43)	-7.74
Arkansas	5.25	0.24	(29)	-2.09	New Hampshire	5.10	0.36	(31)	-4.78
California	5.08 <sup>b</sup>	0.07	(36)	-5.20	New Jersey	5.72	0.15	(12)	6.67
Colorado	4.66	0.20	(48)	-13.04	New Mexico	4.77	0.30	(47)	-11.03
Connecticut	5.44	0.22	(25)	1.42	New York	5.09	0.09	(35)	-5.08
Delaware	5.07	0.44	(37)	-5.37	North Carolina	5.28	0.14	(27)	-1.52
D.C.	3.25 <sup>b</sup>	0.43	(51)	-39.40 <sup>c</sup>	North Dakota	5.10	0.49	(32)	-4.82
Florida	5.01 <sup>b</sup>	0.09	(40)	-6.58	Ohio	5.62	0.12	(16)	4.76
Georgia	4.83 <sup>b</sup>	0.15	(45)	-9.87	Oklahoma	5.71	0.23	(13)	6.61
Hawaii	4.36	0.33	(50)	-18.60 <sup>c</sup>	Oregon	5.93	0.23	(07)	10.56
Idaho	5.77	0.39	(11)	7.55	Pennsylvania	5.87 <sup>b</sup>	0.12	(09)	9.52
Illinois	5.57	0.12	(17)	3.86	Rhode Island	4.78	0.36	(46)	-10.80
Indiana	5.79	0.18	(10)	8.00	South Carolina	5.06	0.20	(39)	-5.58
Iowa	5.62	0.24	(15)	4.87	South Dakota	5.26	0.45	(28)	-1.94
Kansas	5.53	0.26	(20)	3.11	Tennessee	5.54	0.17	(19)	3.25
Kentucky	6.00	0.22	(04)	11.85	Texas	5.19	0.10	(30)	-3.24
Louisiana	5.49	0.21	(23)	2.35	Utah	4.97	0.31	(42)	-7.31
Maine	5.99	0.37	(05)	11.67	Vermont	5.10	0.51	(33)	-4.91
Maryland	4.99	0.18	(41)	-6.82	Virginia	5.07	0.15	(38)	-5.42
Massachusetts	5.44	0.16	(24)	1.47	Washington	5.65	0.18	(14)	5.42
Michigan	6.20 <sup>b</sup>	0.14	(03)	15.57 <sup>c</sup>	West Virginia	6.52 <sup>b</sup>	0.32	(01)	21.61 <sup>c</sup>
Minnesota	5.40	0.19	(26)	0.81	Wisconsin	5.94	0.19	(06)	10.74
Mississippi	4.65	0.23	(49)	-13.30	Wyoming	6.34	0.67	(02)	18.34 <sup>c</sup>
Missouri	5.50	0.17	(21)	2.67					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 20.24  
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.52	0.01		
High Five States				
D.C.	3.82 <sup>b</sup>	0.36	(01)	51.61 <sup>c</sup>
Mississippi	3.35 <sup>b</sup>	0.15	(02)	33.09 <sup>c</sup>
Louisiana	3.22 <sup>b</sup>	0.12	(03)	28.00 <sup>c</sup>
Alabama	3.02 <sup>b</sup>	0.11	(04)	20.14 <sup>c</sup>
South Carolina	3.02 <sup>b</sup>	0.11	(05)	19.98 <sup>c</sup>
Low Five States				
Colorado	2.04 <sup>b</sup>	0.10	(47)	-19.00 <sup>c</sup>
Minnesota	2.03 <sup>b</sup>	0.09	(48)	-19.31 <sup>c</sup>
Nebraska	1.97 <sup>b</sup>	0.15	(49)	-21.87 <sup>c</sup>
South Dakota	1.84	0.20	(50)	-27.09 <sup>c</sup>
Utah	1.42 <sup>b</sup>	0.12	(51)	-43.42 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.02 <sup>b</sup>	0.11	(04)	20.14 <sup>c</sup>	Montana	2.10	0.20	(43)	-16.47 <sup>c</sup>
Alaska	2.38	0.34	(33)	-5.63	Nebraska	1.97 <sup>b</sup>	0.15	(49)	-21.87 <sup>c</sup>
Arizona	2.13 <sup>b</sup>	0.08	(41)	-15.29 <sup>c</sup>	Nevada	2.52	0.15	(26)	0.05
Arkansas	3.00	0.14	(07)	19.33 <sup>c</sup>	New Hampshire	2.76	0.20	(12)	9.67
California	2.54	0.04	(25)	0.74	New Jersey	2.23 <sup>b</sup>	0.07	(38)	-11.55
Colorado	2.04 <sup>b</sup>	0.10	(47)	-19.00 <sup>c</sup>	New Mexico	2.17	0.15	(40)	-13.69
Connecticut	2.06 <sup>b</sup>	0.10	(46)	-18.35 <sup>c</sup>	New York	2.22 <sup>b</sup>	0.05	(39)	-11.65
Delaware	2.67	0.24	(16)	6.10	North Carolina	2.61	0.08	(20)	3.58
D.C.	3.82 <sup>b</sup>	0.36	(01)	51.61 <sup>c</sup>	North Dakota	2.10	0.24	(44)	-16.73 <sup>c</sup>
Florida	2.82 <sup>b</sup>	0.05	(11)	11.86	Ohio	2.67	0.06	(15)	6.25
Georgia	2.69	0.08	(14)	6.93	Oklahoma	2.66	0.12	(17)	5.76
Hawaii	3.02	0.21	(06)	19.86 <sup>c</sup>	Oregon	2.46	0.11	(31)	-2.44
Idaho	2.59	0.19	(23)	2.87	Pennsylvania	2.30 <sup>b</sup>	0.06	(35)	-8.62
Illinois	2.49	0.06	(27)	-0.91	Rhode Island	2.12	0.19	(42)	-15.63 <sup>c</sup>
Indiana	2.28	0.08	(36)	-9.40	South Carolina	3.02 <sup>b</sup>	0.11	(05)	19.98 <sup>c</sup>
Iowa	2.48	0.12	(29)	-1.58	South Dakota	1.84	0.20	(50)	-27.09 <sup>c</sup>
Kansas	2.08 <sup>b</sup>	0.12	(45)	-17.39 <sup>c</sup>	Tennessee	2.95 <sup>b</sup>	0.10	(08)	17.03 <sup>c</sup>
Kentucky	2.74	0.11	(13)	8.79	Texas	2.57	0.05	(24)	2.17
Louisiana	3.22 <sup>b</sup>	0.12	(03)	28.00 <sup>c</sup>	Utah	1.42 <sup>b</sup>	0.12	(51)	-43.42 <sup>c</sup>
Maine	2.59	0.18	(22)	3.04	Vermont	2.63	0.27	(18)	4.38
Maryland	2.59	0.10	(21)	3.04	Virginia	2.26	0.08	(37)	-10.18
Massachusetts	2.49	0.08	(28)	-1.05	Washington	2.94 <sup>b</sup>	0.10	(09)	16.81 <sup>c</sup>
Michigan	2.61	0.07	(19)	3.61	West Virginia	2.87	0.16	(10)	14.01
Minnesota	2.03 <sup>b</sup>	0.09	(48)	-19.31 <sup>c</sup>	Wisconsin	2.47	0.09	(30)	-1.69
Mississippi	3.35 <sup>b</sup>	0.15	(02)	33.09 <sup>c</sup>	Wyoming	2.41	0.30	(32)	-4.23
Missouri	2.33	0.09	(34)	-7.35					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 20.25  
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	3.85	0.02		
High Five States				
D.C.	6.41	0.74	(01)	66.17 <sup>c</sup>
Mississippi	5.60 <sup>b</sup>	0.30	(02)	45.40 <sup>c</sup>
Louisiana	5.33 <sup>b</sup>	0.24	(03)	38.22 <sup>c</sup>
South Carolina	4.89 <sup>b</sup>	0.22	(04)	26.83 <sup>c</sup>
Arkansas	4.86 <sup>b</sup>	0.27	(05)	26.17 <sup>c</sup>
Low Five States				
Minnesota	2.91 <sup>b</sup>	0.16	(47)	-24.47 <sup>c</sup>
Colorado	2.81 <sup>b</sup>	0.18	(48)	-27.02 <sup>c</sup>
Nebraska	2.70 <sup>b</sup>	0.26	(49)	-29.86 <sup>c</sup>
South Dakota	2.62	0.36	(50)	-32.03 <sup>c</sup>
Utah	2.00 <sup>b</sup>	0.21	(51)	-48.15 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.80 <sup>b</sup>	0.21	(06)	24.56 <sup>c</sup>	Montana	3.06	0.35	(45)	-20.63 <sup>c</sup>
Alaska	2.93	0.55	(46)	-23.90 <sup>c</sup>	Nebraska	2.70 <sup>b</sup>	0.26	(49)	-29.86 <sup>c</sup>
Arizona	3.15 <sup>b</sup>	0.15	(42)	-18.31 <sup>c</sup>	Nevada	3.56	0.26	(31)	-7.60
Arkansas	4.86 <sup>b</sup>	0.27	(05)	26.17 <sup>c</sup>	New Hampshire	4.01	0.37	(21)	4.09
California	3.79	0.07	(25)	-1.61	New Jersey	3.33 <sup>b</sup>	0.13	(38)	-13.59
Colorado	2.81 <sup>b</sup>	0.18	(48)	-27.02 <sup>c</sup>	New Mexico	3.54	0.28	(32)	-8.12
Connecticut	3.19	0.20	(40)	-17.23 <sup>c</sup>	New York	3.51 <sup>b</sup>	0.09	(35)	-9.07
Delaware	4.22	0.45	(14)	9.40	North Carolina	3.94	0.14	(22)	2.25
D.C.	6.41	0.74	(01)	66.17 <sup>c</sup>	North Dakota	3.13	0.44	(43)	-18.68 <sup>c</sup>
Florida	4.33 <sup>b</sup>	0.09	(12)	12.23	Ohio	4.09	0.12	(17)	6.18
Georgia	4.35	0.17	(10)	12.97	Oklahoma	4.06	0.22	(19)	5.33
Hawaii	4.56	0.38	(08)	18.39 <sup>c</sup>	Oregon	3.76	0.21	(26)	-2.46
Idaho	3.58	0.33	(29)	-7.01	Pennsylvania	3.57	0.11	(30)	-7.26
Illinois	3.74	0.12	(27)	-2.86	Rhode Island	3.11	0.35	(44)	-19.38 <sup>c</sup>
Indiana	3.43	0.16	(37)	-11.12	South Carolina	4.89 <sup>b</sup>	0.22	(04)	26.83 <sup>c</sup>
Iowa	3.71	0.22	(28)	-3.78	South Dakota	2.62	0.36	(50)	-32.03 <sup>c</sup>
Kansas	3.16	0.22	(41)	-18.09 <sup>c</sup>	Tennessee	4.68 <sup>b</sup>	0.18	(07)	21.28 <sup>c</sup>
Kentucky	4.31	0.21	(13)	11.80	Texas	4.08	0.10	(18)	5.85
Louisiana	5.33 <sup>b</sup>	0.24	(03)	38.22 <sup>c</sup>	Utah	2.00 <sup>b</sup>	0.21	(51)	-48.15 <sup>c</sup>
Maine	3.88	0.34	(24)	0.73	Vermont	4.39	0.54	(09)	13.89
Maryland	4.16	0.19	(16)	7.95	Virginia	3.53	0.15	(33)	-8.47
Massachusetts	3.93	0.16	(23)	1.92	Washington	4.18	0.17	(15)	8.40
Michigan	4.04	0.13	(20)	4.71	West Virginia	4.35	0.29	(11)	12.81
Minnesota	2.91 <sup>b</sup>	0.16	(47)	-24.47 <sup>c</sup>	Wisconsin	3.52	0.16	(34)	-8.61
Mississippi	5.60 <sup>b</sup>	0.30	(02)	45.40 <sup>c</sup>	Wyoming	3.20	0.52	(39)	-16.94 <sup>c</sup>
Missouri	3.46	0.16	(36)	-10.37					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 20.26  
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.42	0.01		
High Five States				
D.C.	1.97	0.34	(01)	39.10 <sup>c</sup>
Washington	1.88 <sup>b</sup>	0.11	(02)	32.56 <sup>c</sup>
Alaska	1.85	0.43	(03)	30.41 <sup>c</sup>
New Hampshire	1.72	0.22	(04)	21.60 <sup>c</sup>
Wyoming	1.72	0.35	(05)	21.52 <sup>c</sup>
Low Five States				
Kansas	1.21	0.12	(47)	-14.98
Connecticut	1.18	0.10	(48)	-16.81 <sup>c</sup>
South Dakota	1.14	0.21	(49)	-19.31 <sup>c</sup>
New Mexico	0.96	0.13	(50)	-32.55 <sup>c</sup>
Utah	0.87 <sup>b</sup>	0.13	(51)	-38.72 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.62	0.11	(08)	14.34	Montana	1.28	0.21	(41)	-9.97
Alaska	1.85	0.43	(03)	30.41 <sup>c</sup>	Nebraska	1.36	0.16	(34)	-4.29
Arizona	1.24	0.09	(44)	-12.38	Nevada	1.56	0.16	(10)	9.95
Arkansas	1.47	0.13	(23)	3.34	New Hampshire	1.72	0.22	(04)	21.60 <sup>c</sup>
California	1.51	0.04	(17)	6.35	New Jersey	1.35	0.07	(35)	-4.51
Colorado	1.42	0.11	(28)	0.34	New Mexico	0.96	0.13	(50)	-32.55 <sup>c</sup>
Connecticut	1.18	0.10	(48)	-16.81 <sup>c</sup>	New York	1.22 <sup>b</sup>	0.05	(46)	-13.99
Delaware	1.40	0.23	(29)	-0.93	North Carolina	1.49	0.08	(20)	4.89
D.C.	1.97	0.34	(01)	39.10 <sup>c</sup>	North Dakota	1.26	0.26	(42)	-10.88
Florida	1.51	0.05	(15)	6.71	Ohio	1.51	0.06	(18)	6.19
Georgia	1.45	0.08	(26)	1.94	Oklahoma	1.51	0.12	(16)	6.49
Hawaii	1.65	0.20	(06)	16.63 <sup>c</sup>	Oregon	1.35	0.11	(36)	-4.77
Idaho	1.63	0.21	(07)	15.12 <sup>c</sup>	Pennsylvania	1.30	0.06	(40)	-8.36
Illinois	1.48	0.06	(21)	4.35	Rhode Island	1.39	0.21	(31)	-1.95
Indiana	1.32	0.08	(39)	-6.60	South Carolina	1.49	0.11	(19)	5.05
Iowa	1.46	0.12	(24)	2.77	South Dakota	1.14	0.21	(49)	-19.31 <sup>c</sup>
Kansas	1.21	0.12	(47)	-14.98	Tennessee	1.51	0.09	(14)	6.76
Kentucky	1.46	0.11	(25)	2.70	Texas	1.34	0.05	(37)	-5.34
Louisiana	1.52	0.11	(13)	7.54	Utah	0.87 <sup>b</sup>	0.13	(51)	-38.72 <sup>c</sup>
Maine	1.53	0.18	(12)	7.99	Vermont	1.23	0.24	(45)	-13.52
Maryland	1.37	0.09	(33)	-3.67	Virginia	1.25	0.08	(43)	-11.87
Massachusetts	1.37	0.08	(32)	-3.31	Washington	1.88 <sup>b</sup>	0.11	(02)	32.56 <sup>c</sup>
Michigan	1.44	0.07	(27)	1.84	West Virginia	1.54	0.16	(11)	8.52
Minnesota	1.33	0.09	(38)	-6.05	Wisconsin	1.60	0.10	(09)	12.70
Mississippi	1.47	0.13	(22)	3.67	Wyoming	1.72	0.35	(05)	21.52 <sup>c</sup>
Missouri	1.39	0.09	(30)	-1.65					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 21.14  
Cancer of the Ovary (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	8.42	0.03		
High Five States				
Washington	9.98 <sup>b</sup>	0.24	(01)	18.56 <sup>c</sup>
Oregon	9.77 <sup>b</sup>	0.30	(02)	16.00 <sup>c</sup>
Wisconsin	9.51 <sup>b</sup>	0.24	(03)	12.98
Alabama	9.39 <sup>b</sup>	0.26	(04)	11.53
Pennsylvania	9.10 <sup>b</sup>	0.15	(05)	8.09
Low Five States				
Maine	7.65	0.42	(47)	-9.13
Missouri	7.61 <sup>b</sup>	0.21	(48)	-9.62
Vermont	7.57	0.63	(49)	-10.07
Alaska	6.95	0.80	(50)	-17.42 <sup>c</sup>
Hawaii	6.41 <sup>b</sup>	0.41	(51)	-23.92 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.39 <sup>b</sup>	0.26	(04)	11.53	Montana	8.93	0.55	(08)	6.09
Alaska	6.95	0.80	(50)	-17.42 <sup>c</sup>	Nebraska	7.90	0.39	(42)	-6.16
Arizona	8.21	0.22	(33)	-2.45	Nevada	7.79	0.36	(43)	-7.56
Arkansas	8.41	0.31	(26)	-0.17	New Hampshire	8.13	0.46	(36)	-3.52
California	8.45	0.10	(23)	0.37	New Jersey	8.82	0.18	(11)	4.79
Colorado	8.72	0.28	(15)	3.56	New Mexico	8.02	0.38	(40)	-4.82
Connecticut	8.35	0.28	(29)	-0.86	New York	8.41	0.12	(25)	-0.16
Delaware	8.49	0.57	(22)	0.83	North Carolina	8.06	0.18	(39)	-4.26
D.C.	7.98	0.69	(41)	-5.27	North Dakota	8.50	0.67	(21)	0.91
Florida	7.71 <sup>b</sup>	0.11	(45)	-8.44	Ohio	8.25	0.15	(31)	-2.01
Georgia	8.57	0.20	(18)	1.79	Oklahoma	8.22	0.28	(32)	-2.42
Hawaii	6.41 <sup>b</sup>	0.41	(51)	-23.92 <sup>c</sup>	Oregon	9.77 <sup>b</sup>	0.30	(02)	16.00 <sup>c</sup>
Idaho	8.67	0.48	(16)	2.94	Pennsylvania	9.10 <sup>b</sup>	0.15	(05)	8.09
Illinois	8.56	0.15	(20)	1.69	Rhode Island	8.65	0.51	(17)	2.70
Indiana	8.74	0.22	(14)	3.74	South Carolina	8.14	0.25	(35)	-3.39
Iowa	9.05	0.31	(06)	7.52	South Dakota	8.11	0.59	(38)	-3.69
Kansas	8.99	0.34	(07)	6.78	Tennessee	8.38	0.22	(28)	-0.49
Kentucky	8.17	0.26	(34)	-2.98	Texas	7.74 <sup>b</sup>	0.12	(44)	-8.08
Louisiana	8.12	0.26	(37)	-3.63	Utah	8.42	0.40	(24)	0.00
Maine	7.65	0.42	(47)	-9.13	Vermont	7.57	0.63	(49)	-10.07
Maryland	8.26	0.23	(30)	-1.94	Virginia	8.85	0.21	(10)	5.09
Massachusetts	8.57	0.21	(19)	1.78	Washington	9.98 <sup>b</sup>	0.24	(01)	18.56 <sup>c</sup>
Michigan	8.79	0.17	(13)	4.40	West Virginia	8.86	0.38	(09)	5.25
Minnesota	8.39	0.24	(27)	-0.42	Wisconsin	9.51 <sup>b</sup>	0.24	(03)	12.98
Mississippi	7.66	0.30	(46)	-9.06	Wyoming	8.81	0.79	(12)	4.63
Missouri	7.61 <sup>b</sup>	0.21	(48)	-9.62					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.18  
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	10.81	0.03		
High Five States				
D.C.	12.50	0.66	(01)	15.61 <sup>c</sup>
Louisiana	12.26 <sup>b</sup>	0.24	(02)	13.44
Connecticut	11.99 <sup>b</sup>	0.25	(03)	10.95
New Hampshire	11.72	0.41	(04)	8.43
Massachusetts	11.55 <sup>b</sup>	0.18	(05)	6.88
Low Five States				
South Dakota	9.97	0.47	(47)	-7.76
Colorado	9.89 <sup>b</sup>	0.22	(48)	-8.48
West Virginia	9.46 <sup>b</sup>	0.29	(49)	-12.48
Arizona	9.45 <sup>b</sup>	0.17	(50)	-12.57
Utah	8.73 <sup>b</sup>	0.30	(51)	-19.26 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	10.96	0.21	(26)	1.36	Montana	10.67	0.44	(34)	-1.28
Alaska	11.19	0.78	(17)	3.56	Nebraska	10.20	0.33	(42)	-5.66
Arizona	9.45 <sup>b</sup>	0.17	(50)	-12.57	Nevada	10.98	0.31	(22)	1.60
Arkansas	10.99	0.27	(21)	1.66	New Hampshire	11.72	0.41	(04)	8.43
California	10.42 <sup>b</sup>	0.08	(39)	-3.59	New Jersey	11.40 <sup>b</sup>	0.16	(10)	5.50
Colorado	9.89 <sup>b</sup>	0.22	(48)	-8.48	New Mexico	10.28	0.32	(40)	-4.91
Connecticut	11.99 <sup>b</sup>	0.25	(03)	10.95	New York	10.97	0.10	(23)	1.46
Delaware	10.86	0.48	(32)	0.45	North Carolina	10.95	0.16	(27)	1.31
D.C.	12.50	0.66	(01)	15.61 <sup>c</sup>	North Dakota	10.96	0.55	(24)	1.43
Florida	10.12 <sup>b</sup>	0.09	(44)	-6.39	Ohio	11.23	0.13	(16)	3.89
Georgia	10.46	0.17	(37)	-3.23	Oklahoma	10.09	0.23	(45)	-6.69
Hawaii	11.02	0.39	(20)	1.92	Oregon	11.28	0.24	(15)	4.34
Idaho	10.95	0.39	(28)	1.29	Pennsylvania	11.49 <sup>b</sup>	0.12	(08)	6.30
Illinois	11.42 <sup>b</sup>	0.13	(09)	5.67	Rhode Island	10.16	0.41	(43)	-6.01
Indiana	10.96	0.18	(25)	1.41	South Carolina	10.84	0.22	(33)	0.24
Iowa	10.25	0.24	(41)	-5.21	South Dakota	9.97	0.47	(47)	-7.76
Kansas	10.92	0.27	(30)	1.06	Tennessee	10.93	0.19	(29)	1.13
Kentucky	10.65	0.22	(35)	-1.45	Texas	10.05 <sup>b</sup>	0.10	(46)	-7.05
Louisiana	12.26 <sup>b</sup>	0.24	(02)	13.44	Utah	8.73 <sup>b</sup>	0.30	(51)	-19.26 <sup>c</sup>
Maine	11.32	0.38	(13)	4.71	Vermont	10.49	0.55	(36)	-2.93
Maryland	11.54 <sup>b</sup>	0.21	(06)	6.73	Virginia	11.32	0.18	(14)	4.68
Massachusetts	11.55 <sup>b</sup>	0.18	(05)	6.88	Washington	10.88	0.19	(31)	0.63
Michigan	11.51 <sup>b</sup>	0.15	(07)	6.45	West Virginia	9.46 <sup>b</sup>	0.29	(49)	-12.48
Minnesota	10.45	0.20	(38)	-3.34	Wisconsin	11.10	0.19	(18)	2.73
Mississippi	11.38	0.28	(11)	5.30	Wyoming	11.37	0.66	(12)	5.13
Missouri	11.05	0.19	(19)	2.23					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.19  
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	12.49	0.04		
High Five States				
D.C.	16.07	1.21	(01)	28.63 <sup>c</sup>
Connecticut	14.40 <sup>b</sup>	0.42	(02)	15.28 <sup>c</sup>
Louisiana	13.99 <sup>b</sup>	0.40	(03)	11.95 <sup>c</sup>
Mississippi	13.64	0.48	(04)	9.19
Michigan	13.58 <sup>b</sup>	0.25	(05)	8.66
Low Five States				
Vermont	11.47	0.87	(47)	-8.23
Arizona	11.35 <sup>b</sup>	0.28	(48)	-9.14
Colorado	11.24	0.36	(49)	-10.04 <sup>c</sup>
South Dakota	11.23	0.75	(50)	-10.15 <sup>c</sup>
Utah	9.66 <sup>b</sup>	0.48	(51)	-22.68 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	12.93	0.36	(12)	3.47	Montana	12.28	0.71	(32)	-1.75
Alaska	11.91	1.19	(38)	-4.65	Nebraska	12.18	0.55	(33)	-2.51
Arizona	11.35 <sup>b</sup>	0.28	(48)	-9.14	Nevada	12.11	0.49	(35)	-3.07
Arkansas	12.74	0.44	(21)	2.00	New Hampshire	12.84	0.67	(16)	2.74
California	11.81 <sup>b</sup>	0.13	(40)	-5.49	New Jersey	13.30	0.26	(07)	6.48
Colorado	11.24	0.36	(49)	-10.04 <sup>c</sup>	New Mexico	11.49	0.51	(46)	-8.06
Connecticut	14.40 <sup>b</sup>	0.42	(02)	15.28 <sup>c</sup>	New York	12.56	0.17	(25)	0.51
Delaware	12.07	0.78	(36)	-3.39	North Carolina	12.46	0.26	(26)	-0.25
D.C.	16.07	1.21	(01)	28.63 <sup>c</sup>	North Dakota	12.40	0.87	(28)	-0.75
Florida	11.87 <sup>b</sup>	0.15	(39)	-5.01	Ohio	13.13	0.22	(10)	5.08
Georgia	12.80	0.29	(19)	2.41	Oklahoma	11.76	0.38	(43)	-5.89
Hawaii	12.89	0.64	(14)	3.18	Oregon	12.33	0.38	(29)	-1.30
Idaho	11.62	0.61	(45)	-6.98	Pennsylvania	13.53 <sup>b</sup>	0.21	(06)	8.30
Illinois	13.16	0.23	(08)	5.31	Rhode Island	12.30	0.71	(30)	-1.56
Indiana	12.88	0.31	(15)	3.11	South Carolina	12.58	0.37	(24)	0.67
Iowa	12.05	0.40	(37)	-3.55	South Dakota	11.23	0.75	(50)	-10.15 <sup>c</sup>
Kansas	12.72	0.45	(22)	1.79	Tennessee	12.83	0.32	(17)	2.65
Kentucky	12.30	0.37	(31)	-1.57	Texas	11.80 <sup>b</sup>	0.17	(41)	-5.54
Louisiana	13.99 <sup>b</sup>	0.40	(03)	11.95 <sup>c</sup>	Utah	9.66 <sup>b</sup>	0.48	(51)	-22.68 <sup>c</sup>
Maine	12.71	0.61	(23)	1.72	Vermont	11.47	0.87	(47)	-8.23
Maryland	12.81	0.34	(18)	2.53	Virginia	13.07	0.30	(11)	4.62
Massachusetts	13.15	0.30	(09)	5.26	Washington	12.14	0.30	(34)	-2.80
Michigan	13.58 <sup>b</sup>	0.25	(05)	8.66	West Virginia	11.75	0.50	(44)	-5.97
Minnesota	11.79	0.33	(42)	-5.64	Wisconsin	12.76	0.32	(20)	2.11
Mississippi	13.64	0.48	(04)	9.19	Wyoming	12.43	1.04	(27)	-0.48
Missouri	12.92	0.31	(13)	3.43					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 22.20  
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	9.44	0.03		
High Five States				
New Hampshire	10.98	0.53	(01)	16.30 <sup>c</sup>
Louisiana	10.89 <sup>b</sup>	0.30	(02)	15.29 <sup>c</sup>
Maryland	10.55 <sup>b</sup>	0.26	(03)	11.69
Wyoming	10.41	0.85	(04)	10.25
Alaska	10.37	1.03	(05)	9.80
Low Five States				
Texas	8.64 <sup>b</sup>	0.12	(47)	-8.45
Florida	8.63 <sup>b</sup>	0.12	(48)	-8.62
Utah	7.93 <sup>b</sup>	0.39	(49)	-16.06 <sup>c</sup>
Arizona	7.79 <sup>b</sup>	0.21	(50)	-17.46 <sup>c</sup>
West Virginia	7.62 <sup>b</sup>	0.35	(51)	-19.31 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.39	0.25	(34)	-0.51	Montana	9.30	0.56	(37)	-1.55
Alaska	10.37	1.03	(05)	9.80	Nebraska	8.66	0.40	(46)	-8.34
Arizona	7.79 <sup>b</sup>	0.21	(50)	-17.46 <sup>c</sup>	Nevada	9.98	0.41	(13)	5.72
Arkansas	9.53	0.33	(26)	0.95	New Hampshire	10.98	0.53	(01)	16.30 <sup>c</sup>
California	9.28	0.10	(39)	-1.71	New Jersey	9.95	0.19	(14)	5.36
Colorado	8.78	0.28	(42)	-7.04	New Mexico	9.29	0.41	(38)	-1.58
Connecticut	10.14	0.30	(09)	7.38	New York	9.76	0.13	(20)	3.39
Delaware	9.83	0.61	(18)	4.08	North Carolina	9.67	0.19	(23)	2.44
D.C.	10.08	0.77	(10)	6.78	North Dakota	9.49	0.68	(29)	0.55
Florida	8.63 <sup>b</sup>	0.12	(48)	-8.62	Ohio	9.70	0.16	(22)	2.68
Georgia	8.76	0.20	(43)	-7.22	Oklahoma	8.70	0.29	(44)	-7.88
Hawaii	9.41	0.48	(33)	-0.37	Oregon	10.34	0.31	(06)	9.50
Idaho	10.21	0.52	(08)	8.15	Pennsylvania	9.83	0.15	(17)	4.08
Illinois	10.06 <sup>b</sup>	0.17	(11)	6.50	Rhode Island	8.66	0.50	(45)	-8.24
Indiana	9.50	0.23	(28)	0.62	South Carolina	9.45	0.27	(30)	0.08
Iowa	8.79	0.30	(41)	-6.89	South Dakota	9.16	0.61	(40)	-2.96
Kansas	9.43	0.34	(31)	-0.09	Tennessee	9.43	0.23	(32)	-0.11
Kentucky	9.30	0.27	(36)	-1.55	Texas	8.64 <sup>b</sup>	0.12	(47)	-8.45
Louisiana	10.89 <sup>b</sup>	0.30	(02)	15.29 <sup>c</sup>	Utah	7.93 <sup>b</sup>	0.39	(49)	-16.06 <sup>c</sup>
Maine	10.05	0.47	(12)	6.43	Vermont	9.57	0.71	(25)	1.37
Maryland	10.55 <sup>b</sup>	0.26	(03)	11.69	Virginia	9.91	0.22	(15)	4.99
Massachusetts	10.30 <sup>b</sup>	0.22	(07)	9.12	Washington	9.78	0.24	(19)	3.60
Michigan	9.85	0.18	(16)	4.35	West Virginia	7.62 <sup>b</sup>	0.35	(51)	-19.31 <sup>c</sup>
Minnesota	9.34	0.25	(35)	-1.12	Wisconsin	9.71	0.24	(21)	2.87
Mississippi	9.61	0.34	(24)	1.79	Wyoming	10.41	0.85	(04)	10.25
Missouri	9.53	0.23	(27)	0.95					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 23.14  
Cancer of the Prostate (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	24.38	0.06		
High Five States				
D.C.	41.74 <sup>b</sup>	2.08	(01)	71.21 <sup>c</sup>
Mississippi	31.67 <sup>b</sup>	0.79	(02)	29.88 <sup>c</sup>
Alabama	29.88 <sup>b</sup>	0.59	(03)	22.56 <sup>c</sup>
Georgia	28.62 <sup>b</sup>	0.49	(04)	17.39 <sup>c</sup>
Louisiana	28.61 <sup>b</sup>	0.62	(05)	17.34 <sup>c</sup>
Low Five States				
Kansas	22.16 <sup>b</sup>	0.62	(47)	-9.10
West Virginia	21.58 <sup>b</sup>	0.72	(48)	-11.49 <sup>c</sup>
Arizona	20.57 <sup>b</sup>	0.39	(49)	-15.64 <sup>c</sup>
Florida	20.29 <sup>b</sup>	0.20	(50)	-16.76 <sup>c</sup>
Hawaii	16.81 <sup>b</sup>	0.73	(51)	-31.06 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	29.88 <sup>b</sup>	0.59	(03)	22.56 <sup>c</sup>	Montana	28.00	1.13	(07)	14.83 <sup>c</sup>
Alaska	22.52	1.90	(46)	-7.65	Nebraska	24.87	0.81	(29)	2.02
Arizona	20.57 <sup>b</sup>	0.39	(49)	-15.64 <sup>c</sup>	Nevada	24.51	0.76	(32)	0.55
Arkansas	26.24	0.66	(16)	7.64	New Hampshire	25.08	1.00	(27)	2.88
California	23.60 <sup>b</sup>	0.19	(40)	-3.20	New Jersey	23.40	0.37	(41)	-4.00
Colorado	24.34	0.58	(35)	-0.17	New Mexico	24.62	0.80	(30)	1.00
Connecticut	25.74	0.59	(20)	5.59	New York	22.97 <sup>b</sup>	0.24	(43)	-5.79
Delaware	26.66	1.25	(12)	9.34	North Carolina	27.04 <sup>b</sup>	0.42	(10)	10.91 <sup>c</sup>
D.C.	41.74 <sup>b</sup>	2.08	(01)	71.21 <sup>c</sup>	North Dakota	25.87	1.26	(19)	6.11
Florida	20.29 <sup>b</sup>	0.20	(50)	-16.76 <sup>c</sup>	Ohio	26.26 <sup>b</sup>	0.34	(15)	7.71
Georgia	28.62 <sup>b</sup>	0.49	(04)	17.39 <sup>c</sup>	Oklahoma	23.89	0.58	(37)	-2.03
Hawaii	16.81 <sup>b</sup>	0.73	(51)	-31.06 <sup>c</sup>	Oregon	26.00	0.57	(18)	6.63
Idaho	27.28	0.99	(09)	11.88 <sup>c</sup>	Pennsylvania	24.52	0.29	(31)	0.59
Illinois	26.09 <sup>b</sup>	0.34	(17)	7.02	Rhode Island	23.81	1.01	(38)	-2.34
Indiana	25.21	0.47	(23)	3.42	South Carolina	28.46 <sup>b</sup>	0.60	(06)	16.75 <sup>c</sup>
Iowa	25.10	0.59	(26)	2.95	South Dakota	24.37	1.12	(33)	-0.06
Kansas	22.16 <sup>b</sup>	0.62	(47)	-9.10	Tennessee	26.35 <sup>b</sup>	0.49	(13)	8.06
Kentucky	25.56	0.58	(22)	4.84	Texas	22.57 <sup>b</sup>	0.25	(45)	-7.42
Louisiana	28.61 <sup>b</sup>	0.62	(05)	17.34 <sup>c</sup>	Utah	25.62	0.84	(21)	5.07
Maine	24.96	0.92	(28)	2.39	Vermont	24.34	1.38	(34)	-0.17
Maryland	27.51 <sup>b</sup>	0.54	(08)	12.85 <sup>c</sup>	Virginia	26.28 <sup>b</sup>	0.46	(14)	7.80
Massachusetts	24.10	0.43	(36)	-1.15	Washington	25.17	0.46	(24)	3.24
Michigan	23.63	0.35	(39)	-3.07	West Virginia	21.58 <sup>b</sup>	0.72	(48)	-11.49 <sup>c</sup>
Minnesota	25.12	0.50	(25)	3.04	Wisconsin	26.72 <sup>b</sup>	0.48	(11)	9.58
Mississippi	31.67 <sup>b</sup>	0.79	(02)	29.88 <sup>c</sup>	Wyoming	22.71	1.50	(44)	-6.87
Missouri	23.08	0.44	(42)	-5.35					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.18  
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.72	0.02		
High Five States				
Hawaii	6.58 <sup>b</sup>	0.30	(01)	76.86 <sup>c</sup>
D.C.	5.07	0.42	(02)	36.13 <sup>c</sup>
Louisiana	4.89 <sup>b</sup>	0.15	(03)	31.28 <sup>c</sup>
California	4.57 <sup>b</sup>	0.05	(04)	22.87 <sup>c</sup>
New York	4.53 <sup>b</sup>	0.07	(05)	21.59 <sup>c</sup>
Low Five States				
Nebraska	2.68 <sup>b</sup>	0.17	(47)	-28.03 <sup>c</sup>
Wyoming	2.67	0.32	(48)	-28.18 <sup>c</sup>
Iowa	2.65 <sup>b</sup>	0.12	(49)	-28.89 <sup>c</sup>
Utah	2.25 <sup>b</sup>	0.15	(50)	-39.47 <sup>c</sup>
Vermont	2.10 <sup>b</sup>	0.25	(51)	-43.57 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.83	0.13	(17)	2.90	Montana	2.74 <sup>b</sup>	0.23	(45)	-26.47 <sup>c</sup>
Alaska	4.00	0.45	(14)	7.44	Nebraska	2.68 <sup>b</sup>	0.17	(47)	-28.03 <sup>c</sup>
Arizona	3.10 <sup>b</sup>	0.10	(36)	-16.67 <sup>c</sup>	Nevada	3.50	0.18	(22)	-5.97
Arkansas	3.48	0.15	(24)	-6.58	New Hampshire	2.87 <sup>b</sup>	0.20	(43)	-22.92 <sup>c</sup>
California	4.57 <sup>b</sup>	0.05	(04)	22.87 <sup>c</sup>	New Jersey	4.52 <sup>b</sup>	0.10	(06)	21.52 <sup>c</sup>
Colorado	2.99 <sup>b</sup>	0.12	(41)	-19.70 <sup>c</sup>	New Mexico	4.51 <sup>b</sup>	0.22	(07)	21.09 <sup>c</sup>
Connecticut	4.22	0.15	(09)	13.38	New York	4.53 <sup>b</sup>	0.07	(05)	21.59 <sup>c</sup>
Delaware	3.92	0.29	(15)	5.42	North Carolina	3.44	0.09	(25)	-7.54
D.C.	5.07	0.42	(02)	36.13 <sup>c</sup>	North Dakota	3.62	0.31	(20)	-2.74
Florida	3.31 <sup>b</sup>	0.05	(30)	-10.98	Ohio	3.35 <sup>b</sup>	0.07	(29)	-10.04
Georgia	3.68	0.10	(19)	-1.19	Oklahoma	3.07 <sup>b</sup>	0.13	(37)	-17.43 <sup>c</sup>
Hawaii	6.58 <sup>b</sup>	0.30	(01)	76.86 <sup>c</sup>	Oregon	2.84 <sup>b</sup>	0.12	(44)	-23.68 <sup>c</sup>
Idaho	3.11	0.21	(35)	-16.53 <sup>c</sup>	Pennsylvania	3.51	0.07	(21)	-5.77
Illinois	4.08 <sup>b</sup>	0.08	(12)	9.46	Rhode Island	4.15	0.26	(10)	11.39
Indiana	2.93 <sup>b</sup>	0.10	(42)	-21.37 <sup>c</sup>	South Carolina	4.12	0.13	(11)	10.68
Iowa	2.65 <sup>b</sup>	0.12	(49)	-28.89 <sup>c</sup>	South Dakota	3.41	0.28	(27)	-8.53
Kansas	2.69 <sup>b</sup>	0.14	(46)	-27.81 <sup>c</sup>	Tennessee	3.37	0.10	(28)	-9.45
Kentucky	3.20 <sup>b</sup>	0.12	(32)	-13.94	Texas	4.02 <sup>b</sup>	0.06	(13)	7.89
Louisiana	4.89 <sup>b</sup>	0.15	(03)	31.28 <sup>c</sup>	Utah	2.25 <sup>b</sup>	0.15	(50)	-39.47 <sup>c</sup>
Maine	3.15	0.20	(33)	-15.41 <sup>c</sup>	Vermont	2.10 <sup>b</sup>	0.25	(51)	-43.57 <sup>c</sup>
Maryland	3.92	0.12	(16)	5.23	Virginia	3.42	0.10	(26)	-8.17
Massachusetts	3.76	0.10	(18)	1.08	Washington	3.23 <sup>b</sup>	0.10	(31)	-13.28
Michigan	3.48	0.08	(23)	-6.57	West Virginia	3.02 <sup>b</sup>	0.17	(38)	-18.80 <sup>c</sup>
Minnesota	3.01 <sup>b</sup>	0.11	(39)	-19.20 <sup>c</sup>	Wisconsin	3.12 <sup>b</sup>	0.10	(34)	-16.26 <sup>c</sup>
Mississippi	4.33 <sup>b</sup>	0.17	(08)	16.39 <sup>c</sup>	Wyoming	2.67	0.32	(48)	-28.18 <sup>c</sup>
Missouri	3.01 <sup>b</sup>	0.10	(40)	-19.23 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.19  
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.17	0.03		
High Five States				
Hawaii	9.43 <sup>b</sup>	0.55	(01)	82.50 <sup>c</sup>
D.C.	7.15	0.80	(02)	38.41 <sup>c</sup>
Louisiana	7.06 <sup>b</sup>	0.28	(03)	36.68 <sup>c</sup>
Rhode Island	6.59	0.52	(04)	27.52 <sup>c</sup>
New Mexico	6.40	0.39	(05)	23.85 <sup>c</sup>
Low Five States				
Iowa	3.80 <sup>b</sup>	0.23	(47)	-26.44 <sup>c</sup>
Nebraska	3.79 <sup>b</sup>	0.31	(48)	-26.57 <sup>c</sup>
Wyoming	3.06 <sup>b</sup>	0.51	(49)	-40.69 <sup>c</sup>
Vermont	2.97 <sup>b</sup>	0.46	(50)	-42.56 <sup>c</sup>
Utah	2.74 <sup>b</sup>	0.26	(51)	-47.03 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.38	0.24	(16)	4.17	Montana	3.82	0.40	(46)	-26.12 <sup>c</sup>
Alaska	4.96	0.75	(21)	-4.07	Nebraska	3.79 <sup>b</sup>	0.31	(48)	-26.57 <sup>c</sup>
Arizona	4.02 <sup>b</sup>	0.17	(43)	-22.26 <sup>c</sup>	Nevada	4.48	0.30	(33)	-13.32
Arkansas	4.54	0.26	(30)	-12.10	New Hampshire	4.16	0.38	(41)	-19.41 <sup>c</sup>
California	6.10 <sup>b</sup>	0.09	(09)	18.14 <sup>c</sup>	New Jersey	6.35 <sup>b</sup>	0.18	(06)	22.94 <sup>c</sup>
Colorado	4.13 <sup>b</sup>	0.22	(42)	-20.04 <sup>c</sup>	New Mexico	6.40	0.39	(05)	23.85 <sup>c</sup>
Connecticut	6.18 <sup>b</sup>	0.28	(08)	19.58 <sup>c</sup>	New York	6.30 <sup>b</sup>	0.12	(07)	21.82 <sup>c</sup>
Delaware	5.52	0.53	(14)	6.81	North Carolina	4.94	0.17	(23)	-4.44
D.C.	7.15	0.80	(02)	38.41 <sup>c</sup>	North Dakota	4.95	0.56	(22)	-4.30
Florida	4.51 <sup>b</sup>	0.09	(32)	-12.80	Ohio	4.79	0.14	(25)	-7.40
Georgia	5.09	0.19	(18)	-1.51	Oklahoma	4.25 <sup>b</sup>	0.23	(37)	-17.71 <sup>c</sup>
Hawaii	9.43 <sup>b</sup>	0.55	(01)	82.50 <sup>c</sup>	Oregon	3.84 <sup>b</sup>	0.21	(45)	-25.65 <sup>c</sup>
Idaho	4.25	0.37	(38)	-17.78 <sup>c</sup>	Pennsylvania	5.04	0.13	(19)	-2.49
Illinois	5.67	0.15	(12)	9.69	Rhode Island	6.59	0.52	(04)	27.52 <sup>c</sup>
Indiana	4.22 <sup>b</sup>	0.18	(40)	-18.42 <sup>c</sup>	South Carolina	5.98	0.25	(10)	15.79 <sup>c</sup>
Iowa	3.80 <sup>b</sup>	0.23	(47)	-26.44 <sup>c</sup>	South Dakota	4.67	0.49	(27)	-9.55
Kansas	3.87 <sup>b</sup>	0.25	(44)	-25.09 <sup>c</sup>	Tennessee	4.79	0.20	(24)	-7.34
Kentucky	4.61	0.23	(29)	-10.70	Texas	5.46	0.12	(15)	5.74
Louisiana	7.06 <sup>b</sup>	0.28	(03)	36.68 <sup>c</sup>	Utah	2.74 <sup>b</sup>	0.26	(51)	-47.03 <sup>c</sup>
Maine	4.65	0.38	(28)	-9.99	Vermont	2.97 <sup>b</sup>	0.46	(50)	-42.56 <sup>c</sup>
Maryland	5.28	0.22	(17)	2.23	Virginia	4.76	0.18	(26)	-7.84
Massachusetts	5.66	0.20	(13)	9.56	Washington	4.27 <sup>b</sup>	0.18	(36)	-17.42 <sup>c</sup>
Michigan	4.98	0.15	(20)	-3.72	West Virginia	4.23	0.31	(39)	-18.12 <sup>c</sup>
Minnesota	4.35 <sup>b</sup>	0.20	(34)	-15.74 <sup>c</sup>	Wisconsin	4.53	0.19	(31)	-12.27
Mississippi	5.93	0.32	(11)	14.71	Wyoming	3.06 <sup>b</sup>	0.51	(49)	-40.69 <sup>c</sup>
Missouri	4.32 <sup>b</sup>	0.18	(35)	-16.37 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.



Table 24.20  
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.66	0.02		
High Five States				
Hawaii	4.45 <sup>b</sup>	0.34	(01)	67.28 <sup>c</sup>
D.C.	3.59	0.46	(02)	35.13 <sup>c</sup>
California	3.45 <sup>b</sup>	0.06	(03)	29.66 <sup>c</sup>
New York	3.30 <sup>b</sup>	0.07	(04)	24.35 <sup>c</sup>
Louisiana	3.26 <sup>b</sup>	0.16	(05)	22.57 <sup>c</sup>
Low Five States				
Montana	1.83	0.25	(47)	-31.05 <sup>c</sup>
Nebraska	1.80 <sup>b</sup>	0.18	(48)	-32.14 <sup>c</sup>
Kansas	1.79 <sup>b</sup>	0.15	(49)	-32.51 <sup>c</sup>
Iowa	1.76 <sup>b</sup>	0.14	(50)	-33.80 <sup>c</sup>
Vermont	1.47 <sup>b</sup>	0.28	(51)	-44.51 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.73	0.14	(14)	2.80	Montana	1.83	0.25	(47)	-31.05 <sup>c</sup>
Alaska	3.13	0.54	(08)	17.72 <sup>c</sup>	Nebraska	1.80 <sup>b</sup>	0.18	(48)	-32.14 <sup>c</sup>
Arizona	2.38	0.12	(27)	-10.43	Nevada	2.65	0.21	(18)	-0.44
Arkansas	2.67	0.18	(17)	0.33	New Hampshire	1.84 <sup>b</sup>	0.21	(46)	-30.71 <sup>c</sup>
California	3.45 <sup>b</sup>	0.06	(03)	29.66 <sup>c</sup>	New Jersey	3.24 <sup>b</sup>	0.11	(06)	22.10 <sup>c</sup>
Colorado	2.14 <sup>b</sup>	0.14	(38)	-19.49 <sup>c</sup>	New Mexico	3.00	0.24	(10)	12.71
Connecticut	2.78	0.16	(13)	4.72	New York	3.30 <sup>b</sup>	0.07	(04)	24.35 <sup>c</sup>
Delaware	2.64	0.32	(20)	-0.72	North Carolina	2.35	0.10	(30)	-11.56
D.C.	3.59	0.46	(02)	35.13 <sup>c</sup>	North Dakota	2.64	0.36	(19)	-0.53
Florida	2.36 <sup>b</sup>	0.06	(29)	-11.24	Ohio	2.33 <sup>b</sup>	0.08	(32)	-12.43
Georgia	2.72	0.11	(15)	2.24	Oklahoma	2.16	0.14	(36)	-18.85 <sup>c</sup>
Hawaii	4.45 <sup>b</sup>	0.34	(01)	67.28 <sup>c</sup>	Oregon	2.09 <sup>b</sup>	0.14	(39)	-21.48 <sup>c</sup>
Idaho	2.14	0.24	(37)	-19.31 <sup>c</sup>	Pennsylvania	2.39 <sup>b</sup>	0.07	(26)	-10.06
Illinois	2.96	0.09	(11)	11.31	Rhode Island	2.62	0.27	(21)	-1.50
Indiana	2.02 <sup>b</sup>	0.11	(43)	-23.93 <sup>c</sup>	South Carolina	2.70	0.14	(16)	1.73
Iowa	1.76 <sup>b</sup>	0.14	(50)	-33.80 <sup>c</sup>	South Dakota	2.48	0.33	(22)	-6.79
Kansas	1.79 <sup>b</sup>	0.15	(49)	-32.51 <sup>c</sup>	Tennessee	2.37	0.11	(28)	-10.78
Kentucky	2.17 <sup>b</sup>	0.13	(35)	-18.32 <sup>c</sup>	Texas	2.90	0.07	(12)	9.21
Louisiana	3.26 <sup>b</sup>	0.16	(05)	22.57 <sup>c</sup>	Utah	1.86 <sup>b</sup>	0.19	(45)	-29.83 <sup>c</sup>
Maine	1.99	0.21	(44)	-25.06 <sup>c</sup>	Vermont	1.47 <sup>b</sup>	0.28	(51)	-44.51 <sup>c</sup>
Maryland	3.04	0.14	(09)	14.35	Virginia	2.45	0.11	(24)	-7.92
Massachusetts	2.47	0.11	(23)	-7.19	Washington	2.44	0.12	(25)	-8.31
Michigan	2.34 <sup>b</sup>	0.09	(31)	-11.89	West Virginia	2.18	0.19	(34)	-18.08 <sup>c</sup>
Minnesota	2.04 <sup>b</sup>	0.12	(41)	-23.17 <sup>c</sup>	Wisconsin	2.07 <sup>b</sup>	0.11	(40)	-22.04 <sup>c</sup>
Mississippi	3.17	0.19	(07)	19.20 <sup>c</sup>	Wyoming	2.31	0.40	(33)	-13.08
Missouri	2.03 <sup>b</sup>	0.11	(42)	-23.48 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 25.14  
Cancer of the Testis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.24	0.01		
High Five States				
California	0.33 <sup>b</sup>	0.02	(01)	35.49 <sup>c</sup>
Pennsylvania	0.30	0.03	(02)	24.33 <sup>c</sup>
Nevada	0.30	0.07	(03)	23.78 <sup>c</sup>
North Carolina	0.30	0.04	(04)	22.69 <sup>c</sup>
Alabama	0.29	0.05	(05)	21.08 <sup>c</sup>
Low Five States				
Wisconsin	0.18	0.04	(26)	-26.19 <sup>c</sup>
Minnesota	0.16	0.04	(27)	-32.12 <sup>c</sup>
Indiana	0.16	0.03	(28)	-34.63 <sup>c</sup>
Virginia	0.15	0.03	(29)	-36.19 <sup>c</sup>
Georgia	0.15	0.03	(30)	-37.58 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.29	0.05	(05)	21.08 <sup>c</sup>	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	-	-	(-)	-
Arizona	0.25	0.04	(12)	4.65	Nevada	0.30	0.07	(03)	23.78 <sup>c</sup>
Arkansas	0.26	0.06	(11)	6.78	New Hampshire	-	-	(-)	-
California	0.33 <sup>b</sup>	0.02	(01)	35.49 <sup>c</sup>	New Jersey	0.19	0.03	(24)	-20.97 <sup>c</sup>
Colorado	0.24	0.05	(14)	1.25	New Mexico	-	-	(-)	-
Connecticut	0.22	0.05	(20)	-8.82	New York	0.19	0.02	(23)	-20.54 <sup>c</sup>
Delaware	-	-	(-)	-	North Carolina	0.30	0.04	(04)	22.69 <sup>c</sup>
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.21	0.02	(22)	-13.74	Ohio	0.23	0.03	(17)	-3.57
Georgia	0.15	0.03	(30)	-37.58 <sup>c</sup>	Oklahoma	0.29	0.06	(06)	20.31 <sup>c</sup>
Hawaii	-	-	(-)	-	Oregon	0.24	0.05	(15)	-0.36
Idaho	-	-	(-)	-	Pennsylvania	0.30	0.03	(02)	24.33 <sup>c</sup>
Illinois	0.27	0.03	(09)	10.89	Rhode Island	-	-	(-)	-
Indiana	0.16	0.03	(28)	-34.63 <sup>c</sup>	South Carolina	-	-	(-)	-
Iowa	-	-	(-)	-	South Dakota	-	-	(-)	-
Kansas	-	-	(-)	-	Tennessee	0.23	0.04	(18)	-4.09
Kentucky	0.22	0.05	(21)	-10.56	Texas	0.28	0.02	(07)	17.42 <sup>c</sup>
Louisiana	0.24	0.05	(16)	-0.99	Utah	-	-	(-)	-
Maine	-	-	(-)	-	Vermont	-	-	(-)	-
Maryland	0.26	0.04	(10)	8.03	Virginia	0.15	0.03	(29)	-36.19 <sup>c</sup>
Massachusetts	0.18	0.03	(25)	-25.39 <sup>c</sup>	Washington	0.23	0.04	(19)	-4.63
Michigan	0.25	0.03	(13)	1.93	West Virginia	-	-	(-)	-
Minnesota	0.16	0.04	(27)	-32.12 <sup>c</sup>	Wisconsin	0.18	0.04	(26)	-26.19 <sup>c</sup>
Mississippi	-	-	(-)	-	Wyoming	-	-	(-)	-
Missouri	0.28	0.04	(08)	15.82 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 26.18  
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.49	0.01		
High Five States				
Wyoming	0.61	0.15	(01)	24.24 <sup>c</sup>
Colorado	0.59	0.05	(02)	19.39 <sup>c</sup>
California	0.59 <sup>b</sup>	0.02	(03)	19.14 <sup>c</sup>
Utah	0.56	0.08	(04)	13.73
Kansas	0.56	0.06	(05)	13.66
Low Five States				
Virginia	0.40	0.03	(45)	-17.98 <sup>c</sup>
Nebraska	0.38	0.06	(46)	-23.20 <sup>c</sup>
South Carolina	0.37	0.04	(47)	-24.02 <sup>c</sup>
Georgia	0.36 <sup>b</sup>	0.03	(48)	-26.53 <sup>c</sup>
Montana	0.35	0.08	(49)	-28.04 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.41	0.04	(43)	-16.04 <sup>c</sup>	Montana	0.35	0.08	(49)	-28.04 <sup>c</sup>
Alaska	-	-	(-)	-	Nebraska	0.38	0.06	(46)	-23.20 <sup>c</sup>
Arizona	0.53	0.04	(12)	8.71	Nevada	0.41	0.06	(41)	-15.64 <sup>c</sup>
Arkansas	0.43	0.05	(39)	-12.47	New Hampshire	0.47	0.08	(31)	-5.41
California	0.59 <sup>b</sup>	0.02	(03)	19.14 <sup>c</sup>	New Jersey	0.53	0.03	(11)	8.71
Colorado	0.59	0.05	(02)	19.39 <sup>c</sup>	New Mexico	0.50	0.07	(21)	2.43
Connecticut	0.49	0.05	(24)	0.24	New York	0.51	0.02	(20)	2.70
Delaware	0.52	0.11	(17)	5.66	North Carolina	0.43	0.03	(40)	-12.86
D.C.	-	-	(-)	-	North Dakota	0.46	0.11	(36)	-6.84
Florida	0.41 <sup>b</sup>	0.02	(42)	-15.65 <sup>c</sup>	Ohio	0.51	0.03	(19)	3.68
Georgia	0.36 <sup>b</sup>	0.03	(48)	-26.53 <sup>c</sup>	Oklahoma	0.46	0.05	(33)	-5.92
Hawaii	0.55	0.09	(07)	12.56	Oregon	0.44	0.05	(37)	-10.99
Idaho	0.54	0.09	(09)	9.34	Pennsylvania	0.49	0.03	(26)	-0.43
Illinois	0.49	0.03	(23)	0.58	Rhode Island	0.47	0.08	(30)	-4.88
Indiana	0.46	0.04	(34)	-6.33	South Carolina	0.37	0.04	(47)	-24.02 <sup>c</sup>
Iowa	0.53	0.05	(13)	7.19	South Dakota	0.44	0.10	(38)	-11.44
Kansas	0.56	0.06	(05)	13.66	Tennessee	0.52	0.04	(16)	5.87
Kentucky	0.52	0.05	(15)	6.25	Texas	0.52	0.02	(18)	5.02
Louisiana	0.46	0.05	(35)	-6.51	Utah	0.56	0.08	(04)	13.73
Maine	0.49	0.08	(25)	0.02	Vermont	0.54	0.13	(10)	8.92
Maryland	0.54	0.04	(08)	9.74	Virginia	0.40	0.03	(45)	-17.98 <sup>c</sup>
Massachusetts	0.46	0.04	(32)	-5.66	Washington	0.56	0.04	(06)	13.26
Michigan	0.47	0.03	(28)	-3.45	West Virginia	0.41	0.06	(44)	-17.20 <sup>c</sup>
Minnesota	0.50	0.04	(22)	1.81	Wisconsin	0.53	0.04	(14)	6.77
Mississippi	0.47	0.06	(29)	-4.63	Wyoming	0.61	0.15	(01)	24.24 <sup>c</sup>
Missouri	0.48	0.04	(27)	-2.53					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 26.19  
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.49	0.01		
High Five States				
Utah	0.70	0.13	(01)	44.11 <sup>c</sup>
Colorado	0.69	0.09	(02)	42.85 <sup>c</sup>
Kansas	0.66	0.11	(03)	34.78 <sup>c</sup>
Iowa	0.60	0.09	(04)	23.90 <sup>c</sup>
New Hampshire	0.59	0.15	(05)	21.29 <sup>c</sup>
Low Five States				
South Carolina	0.39	0.07	(34)	-19.24 <sup>c</sup>
Arkansas	0.37	0.07	(35)	-22.87 <sup>c</sup>
Georgia	0.37	0.05	(36)	-22.96 <sup>c</sup>
Louisiana	0.37	0.06	(37)	-24.39 <sup>c</sup>
Nevada	0.37	0.09	(38)	-24.71 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.45	0.07	(28)	-7.60	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	-	-	(-)	-
Arizona	0.50	0.06	(16)	2.92	Nevada	0.37	0.09	(38)	-24.71 <sup>c</sup>
Arkansas	0.37	0.07	(35)	-22.87 <sup>c</sup>	New Hampshire	0.59	0.15	(05)	21.29 <sup>c</sup>
California	0.55	0.03	(08)	12.58	New Jersey	0.52	0.05	(10)	7.29
Colorado	0.69	0.09	(02)	42.85 <sup>c</sup>	New Mexico	0.52	0.11	(09)	7.63
Connecticut	0.50	0.08	(17)	2.78	New York	0.52	0.04	(11)	6.98
Delaware	-	-	(-)	-	North Carolina	0.44	0.05	(29)	-9.68
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.43	0.03	(32)	-12.23	Ohio	0.56	0.05	(06)	15.04 <sup>c</sup>
Georgia	0.37	0.05	(36)	-22.96 <sup>c</sup>	Oklahoma	0.43	0.07	(31)	-12.19
Hawaii	-	-	(-)	-	Oregon	0.47	0.07	(23)	-3.19
Idaho	-	-	(-)	-	Pennsylvania	0.47	0.04	(25)	-3.73
Illinois	0.40	0.04	(33)	-17.23 <sup>c</sup>	Rhode Island	-	-	(-)	-
Indiana	0.49	0.06	(20)	1.69	South Carolina	0.39	0.07	(34)	-19.24 <sup>c</sup>
Iowa	0.60	0.09	(04)	23.90 <sup>c</sup>	South Dakota	-	-	(-)	-
Kansas	0.66	0.11	(03)	34.78 <sup>c</sup>	Tennessee	0.50	0.06	(15)	3.26
Kentucky	0.50	0.07	(19)	2.15	Texas	0.51	0.04	(14)	4.58
Louisiana	0.37	0.06	(37)	-24.39 <sup>c</sup>	Utah	0.70	0.13	(01)	44.11 <sup>c</sup>
Maine	0.52	0.12	(13)	6.36	Vermont	-	-	(-)	-
Maryland	0.55	0.07	(07)	13.84	Virginia	0.47	0.06	(26)	-3.91
Massachusetts	0.52	0.06	(12)	6.87	Washington	0.50	0.06	(18)	2.30
Michigan	0.47	0.05	(24)	-3.58	West Virginia	-	-	(-)	-
Minnesota	0.43	0.06	(30)	-11.38	Wisconsin	0.45	0.06	(27)	-6.66
Mississippi	0.49	0.09	(21)	0.21	Wyoming	-	-	(-)	-
Missouri	0.48	0.06	(22)	-1.52					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 26.20  
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.49	0.01		
High Five States				
Idaho	0.68	0.13	(01)	37.23 <sup>c</sup>
Hawaii	0.65	0.13	(02)	31.17 <sup>c</sup>
California	0.62 <sup>b</sup>	0.03	(03)	25.14 <sup>c</sup>
Washington	0.60	0.06	(04)	21.87 <sup>c</sup>
Wisconsin	0.57	0.06	(05)	15.71 <sup>c</sup>
Low Five States				
New Hampshire	0.39	0.10	(39)	-20.66 <sup>c</sup>
Alabama	0.38	0.05	(40)	-23.60 <sup>c</sup>
South Carolina	0.36	0.05	(41)	-26.14 <sup>c</sup>
Virginia	0.36	0.04	(42)	-26.78 <sup>c</sup>
Georgia	0.35 <sup>b</sup>	0.04	(43)	-28.64 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.38	0.05	(40)	-23.60 <sup>c</sup>	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.41	0.09	(35)	-16.02 <sup>c</sup>
Arizona	0.56	0.06	(07)	13.85	Nevada	0.47	0.09	(27)	-5.39
Arkansas	0.46	0.07	(29)	-6.58	New Hampshire	0.39	0.10	(39)	-20.66 <sup>c</sup>
California	0.62 <sup>b</sup>	0.03	(03)	25.14 <sup>c</sup>	New Jersey	0.54	0.05	(08)	10.66
Colorado	0.48	0.06	(21)	-1.57	New Mexico	0.49	0.09	(18)	-0.59
Connecticut	0.48	0.06	(22)	-2.63	New York	0.50	0.03	(17)	1.21
Delaware	-	-	(-)	-	North Carolina	0.41	0.04	(37)	-17.60 <sup>c</sup>
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.40 <sup>b</sup>	0.02	(38)	-18.35 <sup>c</sup>	Ohio	0.47	0.04	(26)	-5.10
Georgia	0.35 <sup>b</sup>	0.04	(43)	-28.64 <sup>c</sup>	Oklahoma	0.49	0.07	(20)	-1.22
Hawaii	0.65	0.13	(02)	31.17 <sup>c</sup>	Oregon	0.41	0.06	(36)	-16.57 <sup>c</sup>
Idaho	0.68	0.13	(01)	37.23 <sup>c</sup>	Pennsylvania	0.50	0.03	(16)	1.83
Illinois	0.56	0.04	(06)	14.07	Rhode Island	0.48	0.11	(23)	-2.95
Indiana	0.42	0.05	(34)	-14.50	South Carolina	0.36	0.05	(41)	-26.14 <sup>c</sup>
Iowa	0.47	0.07	(25)	-4.36	South Dakota	-	-	(-)	-
Kansas	0.50	0.08	(15)	2.46	Tennessee	0.53	0.05	(11)	7.50
Kentucky	0.53	0.07	(09)	8.03	Texas	0.52	0.03	(13)	5.38
Louisiana	0.52	0.07	(12)	6.22	Utah	0.44	0.09	(32)	-11.36
Maine	0.46	0.10	(30)	-7.45	Vermont	-	-	(-)	-
Maryland	0.52	0.06	(14)	4.66	Virginia	0.36	0.04	(42)	-26.78 <sup>c</sup>
Massachusetts	0.43	0.05	(33)	-13.42	Washington	0.60	0.06	(04)	21.87 <sup>c</sup>
Michigan	0.46	0.04	(28)	-5.80	West Virginia	0.49	0.09	(19)	-0.92
Minnesota	0.53	0.06	(10)	7.58	Wisconsin	0.57	0.06	(05)	15.71 <sup>c</sup>
Mississippi	0.45	0.07	(31)	-7.79	Wyoming	-	-	(-)	-
Missouri	0.47	0.05	(24)	-3.65					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 27.18  
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.38	0.02		
High Five States				
Maine	6.09 <sup>b</sup>	0.28	(01)	39.16 <sup>c</sup>
Vermont	5.64	0.40	(02)	28.82 <sup>c</sup>
Delaware	5.49	0.35	(03)	25.33 <sup>c</sup>
Rhode Island	5.38	0.30	(04)	22.80 <sup>c</sup>
Nevada	5.24 <sup>b</sup>	0.22	(05)	19.74 <sup>c</sup>
Low Five States				
Texas	3.82 <sup>b</sup>	0.06	(47)	-12.72
Mississippi	3.57 <sup>b</sup>	0.16	(48)	-18.40 <sup>c</sup>
New Mexico	3.52 <sup>b</sup>	0.19	(49)	-19.48 <sup>c</sup>
Utah	3.38 <sup>b</sup>	0.19	(50)	-22.83 <sup>c</sup>
Hawaii	2.66 <sup>b</sup>	0.19	(51)	-39.13 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.83 <sup>b</sup>	0.13	(46)	-12.55	Montana	4.42	0.28	(22)	1.00
Alaska	4.00	0.49	(39)	-8.65	Nebraska	3.96	0.20	(41)	-9.55
Arizona	3.96 <sup>b</sup>	0.11	(40)	-9.51	Nevada	5.24 <sup>b</sup>	0.22	(05)	19.74 <sup>c</sup>
Arkansas	4.10	0.16	(36)	-6.25	New Hampshire	5.07	0.27	(07)	15.84 <sup>c</sup>
California	4.15 <sup>b</sup>	0.05	(34)	-5.30	New Jersey	5.02 <sup>b</sup>	0.10	(08)	14.57
Colorado	3.95	0.14	(42)	-9.68	New Mexico	3.52 <sup>b</sup>	0.19	(49)	-19.48 <sup>c</sup>
Connecticut	4.86	0.16	(11)	11.07	New York	4.41	0.06	(25)	0.69
Delaware	5.49	0.35	(03)	25.33 <sup>c</sup>	North Carolina	4.10	0.10	(37)	-6.29
D.C.	4.39	0.39	(27)	0.28	North Dakota	4.41	0.33	(24)	0.70
Florida	4.41	0.06	(23)	0.74	Ohio	4.99 <sup>b</sup>	0.09	(09)	14.04
Georgia	3.89 <sup>b</sup>	0.10	(45)	-11.16	Oklahoma	4.13	0.15	(35)	-5.68
Hawaii	2.66 <sup>b</sup>	0.19	(51)	-39.13 <sup>c</sup>	Oregon	4.75	0.15	(13)	8.54
Idaho	4.39	0.25	(26)	0.32	Pennsylvania	4.68 <sup>b</sup>	0.08	(16)	7.02
Illinois	4.52	0.08	(20)	3.18	Rhode Island	5.38	0.30	(04)	22.80 <sup>c</sup>
Indiana	4.56	0.12	(17)	4.17	South Carolina	3.94	0.13	(43)	-9.97
Iowa	4.49	0.16	(21)	2.52	South Dakota	4.31	0.30	(29)	-1.51
Kansas	3.91	0.16	(44)	-10.75	Tennessee	4.17	0.12	(32)	-4.64
Kentucky	4.55	0.15	(18)	4.00	Texas	3.82 <sup>b</sup>	0.06	(47)	-12.72
Louisiana	4.21	0.14	(31)	-3.92	Utah	3.38 <sup>b</sup>	0.19	(50)	-22.83 <sup>c</sup>
Maine	6.09 <sup>b</sup>	0.28	(01)	39.16 <sup>c</sup>	Vermont	5.64	0.40	(02)	28.82 <sup>c</sup>
Maryland	4.54	0.13	(19)	3.79	Virginia	4.17	0.11	(33)	-4.82
Massachusetts	5.09 <sup>b</sup>	0.12	(06)	16.33 <sup>c</sup>	Washington	4.71	0.12	(15)	7.61
Michigan	4.73 <sup>b</sup>	0.10	(14)	7.97	West Virginia	4.86	0.21	(12)	10.94
Minnesota	4.01	0.12	(38)	-8.49	Wisconsin	4.90 <sup>b</sup>	0.13	(10)	11.90
Mississippi	3.57 <sup>b</sup>	0.16	(48)	-18.40 <sup>c</sup>	Wyoming	4.37	0.41	(28)	-0.05
Missouri	4.28	0.12	(30)	-2.15					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 27.19  
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.67	0.04		
High Five States				
Maine	10.48 <sup>b</sup>	0.58	(01)	36.63 <sup>c</sup>
Vermont	10.37	0.88	(02)	35.25 <sup>c</sup>
Rhode Island	10.08 <sup>b</sup>	0.65	(03)	31.43 <sup>c</sup>
Delaware	9.65	0.73	(04)	25.85 <sup>c</sup>
Massachusetts	9.31 <sup>b</sup>	0.26	(05)	21.46 <sup>c</sup>
Low Five States				
Mississippi	6.63	0.35	(47)	-13.48
South Carolina	6.59 <sup>b</sup>	0.28	(48)	-14.03
Utah	6.15 <sup>b</sup>	0.40	(49)	-19.76 <sup>c</sup>
New Mexico	5.55 <sup>b</sup>	0.37	(50)	-27.68 <sup>c</sup>
Hawaii	4.36 <sup>b</sup>	0.37	(51)	-43.09 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.74	0.27	(43)	-12.06	Montana	7.70	0.59	(26)	0.48
Alaska	7.35	1.03	(32)	-4.09	Nebraska	7.03	0.43	(39)	-8.29
Arizona	6.66 <sup>b</sup>	0.22	(46)	-13.09	Nevada	8.38	0.43	(12)	9.23
Arkansas	6.88	0.33	(41)	-10.30	New Hampshire	8.98	0.58	(07)	17.08 <sup>c</sup>
California	7.17 <sup>b</sup>	0.11	(36)	-6.47	New Jersey	8.95 <sup>b</sup>	0.23	(08)	16.76 <sup>c</sup>
Colorado	6.89	0.30	(40)	-10.14	New Mexico	5.55 <sup>b</sup>	0.37	(50)	-27.68 <sup>c</sup>
Connecticut	8.58	0.34	(09)	11.92	New York	7.77	0.14	(24)	1.28
Delaware	9.65	0.73	(04)	25.85 <sup>c</sup>	North Carolina	7.08	0.21	(38)	-7.62
D.C.	6.67	0.82	(44)	-12.97	North Dakota	7.96	0.70	(20)	3.83
Florida	7.42	0.12	(30)	-3.28	Ohio	9.14 <sup>b</sup>	0.20	(06)	19.16 <sup>c</sup>
Georgia	6.87	0.23	(42)	-10.43	Oklahoma	7.32	0.31	(33)	-4.52
Hawaii	4.36 <sup>b</sup>	0.37	(51)	-43.09 <sup>c</sup>	Oregon	8.17	0.32	(17)	6.53
Idaho	7.52	0.51	(27)	-1.88	Pennsylvania	8.34 <sup>b</sup>	0.17	(14)	8.75
Illinois	7.94	0.18	(22)	3.59	Rhode Island	10.08 <sup>b</sup>	0.65	(03)	31.43 <sup>c</sup>
Indiana	7.94	0.25	(21)	3.60	South Carolina	6.59 <sup>b</sup>	0.28	(48)	-14.03
Iowa	8.37	0.34	(13)	9.21	South Dakota	8.32	0.66	(15)	8.45
Kansas	7.20	0.35	(35)	-6.06	Tennessee	7.49	0.26	(28)	-2.26
Kentucky	7.97	0.31	(19)	3.97	Texas	6.67 <sup>b</sup>	0.14	(45)	-13.01
Louisiana	7.11	0.30	(37)	-7.27	Utah	6.15 <sup>b</sup>	0.40	(49)	-19.76 <sup>c</sup>
Maine	10.48 <sup>b</sup>	0.58	(01)	36.63 <sup>c</sup>	Vermont	10.37	0.88	(02)	35.25 <sup>c</sup>
Maryland	7.80	0.28	(23)	1.67	Virginia	7.49	0.24	(29)	-2.26
Massachusetts	9.31 <sup>b</sup>	0.26	(05)	21.46 <sup>c</sup>	Washington	8.21	0.26	(16)	7.04
Michigan	8.55 <sup>b</sup>	0.21	(11)	11.47	West Virginia	8.04	0.43	(18)	4.82
Minnesota	7.27	0.26	(34)	-5.15	Wisconsin	8.56	0.27	(10)	11.68
Mississippi	6.63	0.35	(47)	-13.48	Wyoming	7.36	0.83	(31)	-4.06
Missouri	7.75	0.25	(25)	1.03					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 27.20  
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2004-2008

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.22	0.02		
High Five States				
D.C.	3.11	0.42	(01)	40.07 <sup>c</sup>
Maine	3.08	0.26	(02)	38.62 <sup>c</sup>
West Virginia	2.72	0.20	(03)	22.55 <sup>c</sup>
Nevada	2.70	0.22	(04)	21.60 <sup>c</sup>
Connecticut	2.60	0.15	(05)	17.22 <sup>c</sup>
Low Five States				
Kansas	1.82	0.14	(46)	-17.86 <sup>c</sup>
Mississippi	1.73 <sup>b</sup>	0.14	(47)	-22.32 <sup>c</sup>
South Dakota	1.52	0.24	(48)	-31.53 <sup>c</sup>
Hawaii	1.42 <sup>b</sup>	0.19	(49)	-35.96 <sup>c</sup>
Utah	1.27 <sup>b</sup>	0.15	(50)	-42.84 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.01	0.12	(38)	-9.38	Montana	2.13	0.26	(29)	-4.23
Alaska	-	-	(-)	-	Nebraska	2.02	0.19	(35)	-8.92
Arizona	1.89	0.10	(43)	-14.99	Nevada	2.70	0.22	(04)	21.60 <sup>c</sup>
Arkansas	2.24	0.15	(25)	0.78	New Hampshire	2.48	0.25	(11)	11.80
California	2.10	0.05	(31)	-5.46	New Jersey	2.60 <sup>b</sup>	0.10	(06)	16.87 <sup>c</sup>
Colorado	1.95	0.13	(41)	-12.15	New Mexico	1.97	0.19	(40)	-11.34
Connecticut	2.60	0.15	(05)	17.22 <sup>c</sup>	New York	2.32	0.06	(21)	4.41
Delaware	2.54	0.31	(09)	14.55	North Carolina	2.29	0.09	(23)	3.14
D.C.	3.11	0.42	(01)	40.07 <sup>c</sup>	North Dakota	1.89	0.28	(44)	-15.13 <sup>c</sup>
Florida	2.15	0.06	(28)	-3.34	Ohio	2.43	0.08	(13)	9.62
Georgia	2.12	0.10	(30)	-4.61	Oklahoma	1.97	0.13	(39)	-11.12
Hawaii	1.42 <sup>b</sup>	0.19	(49)	-35.96 <sup>c</sup>	Oregon	2.35	0.14	(20)	5.68
Idaho	2.02	0.23	(37)	-9.19	Pennsylvania	2.40	0.07	(16)	8.04
Illinois	2.41	0.08	(14)	8.59	Rhode Island	2.45	0.26	(12)	10.35
Indiana	2.38	0.11	(18)	7.14	South Carolina	2.19	0.13	(26)	-1.18
Iowa	2.06	0.14	(33)	-7.04	South Dakota	1.52	0.24	(48)	-31.53 <sup>c</sup>
Kansas	1.82	0.14	(46)	-17.86 <sup>c</sup>	Tennessee	2.05	0.10	(34)	-7.62
Kentucky	2.36	0.13	(19)	6.40	Texas	1.94 <sup>b</sup>	0.06	(42)	-12.62
Louisiana	2.39	0.14	(17)	7.83	Utah	1.27 <sup>b</sup>	0.15	(50)	-42.84 <sup>c</sup>
Maine	3.08	0.26	(02)	38.62 <sup>c</sup>	Vermont	2.40	0.34	(15)	8.13
Maryland	2.57	0.13	(08)	15.60 <sup>c</sup>	Virginia	2.06	0.10	(32)	-7.03
Massachusetts	2.53	0.11	(10)	14.02	Washington	2.31	0.11	(22)	3.80
Michigan	2.28	0.09	(24)	2.48	West Virginia	2.72	0.20	(03)	22.55 <sup>c</sup>
Minnesota	1.86	0.11	(45)	-16.08 <sup>c</sup>	Wisconsin	2.57	0.12	(07)	15.86 <sup>c</sup>
Mississippi	1.73 <sup>b</sup>	0.14	(47)	-22.32 <sup>c</sup>	Wyoming	2.18	0.39	(27)	-1.85
Missouri	2.02	0.10	(36)	-8.99					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.