


Table 2.18  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	176.44	0.11		
High Five States				
Kentucky	209.49	0.96	(01)	18.73 <sup>c</sup>
Mississippi	203.54	1.17	(02)	15.36 <sup>c</sup>
Louisiana	201.30	0.96	(03)	14.09 <sup>c</sup>
West Virginia	201.16	1.33	(04)	14.01 <sup>c</sup>
Tennessee	199.07	0.78	(05)	12.82 <sup>c</sup>
Low Five States				
New Mexico	155.46	1.23	(47)	-11.89 <sup>c</sup>
Arizona	155.19	0.69	(48)	-12.04 <sup>c</sup>
Colorado	152.84	0.85	(49)	-13.37 <sup>c</sup>
Hawaii	143.68	1.38	(50)	-18.57 <sup>c</sup>
Utah	131.33	1.16	(51)	-25.57 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	196.45	0.88	(08)	11.34 <sup>c</sup>	Montana	166.34	1.72	(43)	-5.72 <sup>c</sup>
Alaska	182.47	3.04	(20)	3.42	Nebraska	170.69	1.32	(35)	-3.26
Arizona	155.19	0.69	(48)	-12.04 <sup>c</sup>	Nevada	179.00	1.24	(23)	1.45
Arkansas	198.04	1.11	(07)	12.24 <sup>c</sup>	New Hampshire	177.31	1.59	(28)	0.49
California	160.33	0.31	(46)	-9.13 <sup>c</sup>	New Jersey	175.00	0.61	(32)	-0.82
Colorado	152.84	0.85	(49)	-13.37 <sup>c</sup>	New Mexico	155.46	1.23	(47)	-11.89 <sup>c</sup>
Connecticut	166.35	0.91	(42)	-5.72 <sup>c</sup>	New York	166.60	0.40	(41)	-5.58 <sup>c</sup>
Delaware	187.71	1.96	(14)	6.38 <sup>c</sup>	North Carolina	182.81	0.62	(19)	3.61
D.C.	198.95	2.68	(06)	12.76 <sup>c</sup>	North Dakota	164.80	2.07	(44)	-6.60 <sup>c</sup>
Florida	168.35	0.38	(39)	-4.58	Ohio	192.36	0.55	(11)	9.02 <sup>c</sup>
Georgia	179.18	0.67	(22)	1.55	Oklahoma	193.26	0.99	(09)	9.53 <sup>c</sup>
Hawaii	143.68	1.38	(50)	-18.57 <sup>c</sup>	Oregon	176.29	0.93	(29)	-0.09
Idaho	162.81	1.49	(45)	-7.73 <sup>c</sup>	Pennsylvania	184.53	0.49	(17)	4.58
Illinois	183.92	0.53	(18)	4.24	Rhode Island	177.45	1.71	(27)	0.57
Indiana	192.59	0.76	(10)	9.15 <sup>c</sup>	South Carolina	187.62	0.89	(15)	6.33 <sup>c</sup>
Iowa	174.32	0.99	(34)	-1.20	South Dakota	168.38	1.92	(38)	-4.57
Kansas	174.68	1.08	(33)	-1.00	Tennessee	199.07	0.78	(05)	12.82 <sup>c</sup>
Kentucky	209.49	0.96	(01)	18.73 <sup>c</sup>	Texas	169.69	0.41	(36)	-3.83
Louisiana	201.30	0.96	(03)	14.09 <sup>c</sup>	Utah	131.33	1.16	(51)	-25.57 <sup>c</sup>
Maine	187.73	1.51	(13)	6.40 <sup>c</sup>	Vermont	177.88	2.24	(25)	0.82
Maryland	178.99	0.80	(24)	1.45	Virginia	179.31	0.69	(21)	1.62
Massachusetts	177.59	0.70	(26)	0.65	Washington	175.02	0.74	(31)	-0.81
Michigan	185.09	0.59	(16)	4.90	West Virginia	201.16	1.33	(04)	14.01 <sup>c</sup>
Minnesota	168.86	0.79	(37)	-4.30	Wisconsin	175.58	0.75	(30)	-0.49
Mississippi	203.54	1.17	(02)	15.36 <sup>c</sup>	Wyoming	167.80	2.49	(40)	-4.90
Missouri	189.17	0.76	(12)	7.21 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 2.19  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	215.27	0.18		
High Five States				
Mississippi	270.64	2.15	(01)	25.73 <sup>c</sup>
Kentucky	262.69	1.69	(02)	22.03 <sup>c</sup>
Tennessee	256.44	1.40	(03)	19.13 <sup>c</sup>
Alabama	255.38	1.59	(04)	18.64 <sup>c</sup>
Louisiana	254.70	1.69	(05)	18.32 <sup>c</sup>
Low Five States				
New Mexico	187.39	2.05	(47)	-12.95 <sup>c</sup>
Arizona	185.97	1.14	(48)	-13.61 <sup>c</sup>
Colorado	181.37	1.43	(49)	-15.75 <sup>c</sup>
Hawaii	178.37	2.31	(50)	-17.14 <sup>c</sup>
Utah	157.14	1.93	(51)	-27.00 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	255.38	1.59	(04)	18.64 <sup>c</sup>	Montana	195.26	2.80	(44)	-9.29 <sup>c</sup>
Alaska	210.49	5.02	(31)	-2.22	Nebraska	208.40	2.23	(34)	-3.19
Arizona	185.97	1.14	(48)	-13.61 <sup>c</sup>	Nevada	211.41	2.04	(30)	-1.79
Arkansas	250.59	1.93	(06)	16.41 <sup>c</sup>	New Hampshire	211.65	2.67	(29)	-1.68
California	189.83	0.51	(46)	-11.81 <sup>c</sup>	New Jersey	207.79	1.04	(35)	-3.47
Colorado	181.37	1.43	(49)	-15.75 <sup>c</sup>	New Mexico	187.39	2.05	(47)	-12.95 <sup>c</sup>
Connecticut	199.53	1.54	(42)	-7.31 <sup>c</sup>	New York	199.72	0.68	(41)	-7.22 <sup>c</sup>
Delaware	228.02	3.34	(17)	5.92 <sup>c</sup>	North Carolina	232.19	1.11	(13)	7.86 <sup>c</sup>
D.C.	248.47	4.75	(07)	15.42 <sup>c</sup>	North Dakota	203.58	3.48	(40)	-5.43 <sup>c</sup>
Florida	204.50	0.62	(38)	-5.00 <sup>c</sup>	Ohio	236.51	0.94	(12)	9.87 <sup>c</sup>
Georgia	228.32	1.22	(16)	6.06 <sup>c</sup>	Oklahoma	238.80	1.70	(11)	10.93 <sup>c</sup>
Hawaii	178.37	2.31	(50)	-17.14 <sup>c</sup>	Oregon	208.47	1.53	(33)	-3.16
Idaho	192.88	2.44	(45)	-10.40 <sup>c</sup>	Pennsylvania	225.63	0.84	(18)	4.82
Illinois	223.28	0.92	(19)	3.72	Rhode Island	221.79	2.97	(21)	3.03
Indiana	238.83	1.32	(10)	10.95 <sup>c</sup>	South Carolina	240.12	1.59	(09)	11.54 <sup>c</sup>
Iowa	212.61	1.67	(27)	-1.23	South Dakota	206.80	3.22	(37)	-3.93
Kansas	214.41	1.82	(25)	-0.40	Tennessee	256.44	1.40	(03)	19.13 <sup>c</sup>
Kentucky	262.69	1.69	(02)	22.03 <sup>c</sup>	Texas	209.32	0.71	(32)	-2.76
Louisiana	254.70	1.69	(05)	18.32 <sup>c</sup>	Utah	157.14	1.93	(51)	-27.00 <sup>c</sup>
Maine	231.11	2.59	(14)	7.36 <sup>c</sup>	Vermont	212.02	3.76	(28)	-1.51
Maryland	216.93	1.38	(23)	0.78	Virginia	221.11	1.20	(22)	2.71
Massachusetts	216.58	1.21	(24)	0.61	Washington	207.21	1.22	(36)	-3.74
Michigan	223.00	0.99	(20)	3.59	West Virginia	246.92	2.27	(08)	14.70 <sup>c</sup>
Minnesota	203.86	1.33	(39)	-5.30 <sup>c</sup>	Wisconsin	213.68	1.28	(26)	-0.74
Mississippi	270.64	2.15	(01)	25.73 <sup>c</sup>	Wyoming	196.05	4.06	(43)	-8.93 <sup>c</sup>
Missouri	231.09	1.30	(15)	7.35 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 2.20  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	149.67	0.13		
High Five States				
Kentucky	173.29	1.17	(01)	15.78 <sup>c</sup>
West Virginia	168.59	1.64	(02)	12.64 <sup>c</sup>
D.C.	167.63	3.23	(03)	12.00 <sup>c</sup>
Louisiana	164.42	1.15	(04)	9.86 <sup>c</sup>
Ohio	162.60	0.67	(05)	8.64 <sup>c</sup>
Low Five States				
Colorado	133.17	1.05	(47)	-11.02 <sup>c</sup>
Arizona	131.68	0.86	(48)	-12.02 <sup>c</sup>
New Mexico	131.56	1.52	(49)	-12.10 <sup>c</sup>
Hawaii	117.11	1.70	(50)	-21.75 <sup>c</sup>
Utah	112.34	1.45	(51)	-24.94 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	156.42	1.04	(18)	4.51	Montana	145.07	2.20	(37)	-3.07
Alaska	160.98	3.85	(07)	7.56 <sup>c</sup>	Nebraska	143.80	1.63	(40)	-3.92
Arizona	131.68	0.86	(48)	-12.02 <sup>c</sup>	Nevada	153.58	1.56	(23)	2.61
Arkansas	160.86	1.35	(09)	7.48 <sup>c</sup>	New Hampshire	154.18	1.99	(21)	3.02
California	139.81	0.38	(45)	-6.58 <sup>c</sup>	New Jersey	154.26	0.76	(19)	3.07
Colorado	133.17	1.05	(47)	-11.02 <sup>c</sup>	New Mexico	131.56	1.52	(49)	-12.10 <sup>c</sup>
Connecticut	144.75	1.13	(39)	-3.29	New York	145.37	0.49	(36)	-2.87
Delaware	159.29	2.42	(13)	6.43 <sup>c</sup>	North Carolina	149.93	0.75	(29)	0.18
D.C.	167.63	3.23	(03)	12.00 <sup>c</sup>	North Dakota	137.18	2.58	(46)	-8.34 <sup>c</sup>
Florida	140.73	0.47	(43)	-5.97 <sup>c</sup>	Ohio	162.60	0.67	(05)	8.64 <sup>c</sup>
Georgia	147.83	0.80	(33)	-1.23	Oklahoma	160.98	1.22	(08)	7.56 <sup>c</sup>
Hawaii	117.11	1.70	(50)	-21.75 <sup>c</sup>	Oregon	153.41	1.17	(24)	2.50
Idaho	140.25	1.88	(44)	-6.29 <sup>c</sup>	Pennsylvania	157.02	0.61	(17)	4.91
Illinois	158.52	0.66	(15)	5.92 <sup>c</sup>	Rhode Island	149.57	2.09	(31)	-0.06
Indiana	161.53	0.93	(06)	7.93 <sup>c</sup>	South Carolina	151.69	1.07	(27)	1.35
Iowa	148.39	1.24	(32)	-0.85	South Dakota	141.26	2.39	(42)	-5.61 <sup>c</sup>
Kansas	146.47	1.33	(35)	-2.13	Tennessee	160.35	0.93	(10)	7.14 <sup>c</sup>
Kentucky	173.29	1.17	(01)	15.78 <sup>c</sup>	Texas	141.81	0.50	(41)	-5.25 <sup>c</sup>
Louisiana	164.42	1.15	(04)	9.86 <sup>c</sup>	Utah	112.34	1.45	(51)	-24.94 <sup>c</sup>
Maine	158.31	1.87	(16)	5.77 <sup>c</sup>	Vermont	154.25	2.81	(20)	3.07
Maryland	153.99	0.98	(22)	2.89	Virginia	151.38	0.84	(28)	1.14
Massachusetts	152.48	0.86	(25)	1.88	Washington	152.16	0.93	(26)	1.67
Michigan	159.46	0.73	(12)	6.54 <sup>c</sup>	West Virginia	168.59	1.64	(02)	12.64 <sup>c</sup>
Minnesota	145.04	0.99	(38)	-3.09	Wisconsin	149.72	0.94	(30)	0.03
Mississippi	159.09	1.37	(14)	6.30 <sup>c</sup>	Wyoming	146.75	3.17	(34)	-1.95
Missouri	160.20	0.94	(11)	7.04 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 3.18  
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.25	0.02		
High Five States				
Iowa	5.37	0.18	(01)	26.24 <sup>c</sup>
South Dakota	5.36	0.36	(02)	25.94 <sup>c</sup>
Washington	5.10	0.12	(03)	19.95 <sup>c</sup>
Maine	5.03	0.25	(04)	18.20 <sup>c</sup>
Nebraska	4.95	0.23	(05)	16.41 <sup>c</sup>
Low Five States				
New York	3.73	0.06	(47)	-12.30
New Mexico	3.68	0.19	(48)	-13.55
New Jersey	3.66	0.09	(49)	-13.91
D.C.	3.62	0.36	(50)	-14.84
Hawaii	2.29	0.18	(51)	-46.16 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.78	0.14	(14)	12.39	Montana	4.44	0.29	(25)	4.35
Alaska	4.29	0.44	(31)	0.96	Nebraska	4.95	0.23	(05)	16.41 <sup>c</sup>
Arizona	4.05	0.11	(39)	-4.73	Nevada	4.03	0.18	(40)	-5.11
Arkansas	4.93	0.18	(06)	15.84 <sup>c</sup>	New Hampshire	4.86	0.26	(10)	14.33
California	4.24	0.05	(32)	-0.18	New Jersey	3.66	0.09	(49)	-13.91
Colorado	4.48	0.14	(22)	5.24	New Mexico	3.68	0.19	(48)	-13.55
Connecticut	4.24	0.15	(33)	-0.38	New York	3.73	0.06	(47)	-12.30
Delaware	4.20	0.30	(35)	-1.22	North Carolina	4.09	0.09	(37)	-3.90
D.C.	3.62	0.36	(50)	-14.84	North Dakota	4.47	0.36	(23)	5.16
Florida	3.91	0.06	(44)	-8.04	Ohio	4.48	0.08	(21)	5.41
Georgia	3.96	0.10	(43)	-6.98	Oklahoma	4.61	0.15	(17)	8.32
Hawaii	2.29	0.18	(51)	-46.16 <sup>c</sup>	Oregon	4.91	0.16	(08)	15.38 <sup>c</sup>
Idaho	4.92	0.26	(07)	15.62 <sup>c</sup>	Pennsylvania	4.12	0.08	(36)	-3.09
Illinois	3.99	0.08	(42)	-6.14	Rhode Island	4.01	0.26	(41)	-5.79
Indiana	4.51	0.12	(20)	6.02	South Carolina	4.32	0.14	(30)	1.66
Iowa	5.37	0.18	(01)	26.24 <sup>c</sup>	South Dakota	5.36	0.36	(02)	25.94 <sup>c</sup>
Kansas	4.84	0.18	(11)	13.76	Tennessee	4.83	0.12	(12)	13.53
Kentucky	4.61	0.14	(16)	8.48	Texas	4.08	0.06	(38)	-3.99
Louisiana	4.33	0.14	(29)	1.90	Utah	4.37	0.20	(27)	2.72
Maine	5.03	0.25	(04)	18.20 <sup>c</sup>	Vermont	4.77	0.37	(15)	12.10
Maryland	3.86	0.12	(46)	-9.24	Virginia	3.87	0.10	(45)	-9.09
Massachusetts	4.22	0.11	(34)	-0.75	Washington	5.10	0.12	(03)	19.95 <sup>c</sup>
Michigan	4.86	0.10	(09)	14.41	West Virginia	4.35	0.20	(28)	2.40
Minnesota	4.45	0.13	(24)	4.71	Wisconsin	4.82	0.13	(13)	13.46
Mississippi	4.42	0.17	(26)	3.98	Wyoming	4.55	0.41	(18)	6.93
Missouri	4.52	0.12	(19)	6.25					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 3.19  
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.19	0.03		
High Five States				
Iowa	6.71	0.30	(01)	29.17 <sup>c</sup>
Washington	6.38	0.20	(02)	22.89 <sup>c</sup>
New Hampshire	6.31	0.45	(03)	21.45 <sup>c</sup>
South Dakota	6.28	0.56	(04)	20.84 <sup>c</sup>
Maine	6.20	0.42	(05)	19.39 <sup>c</sup>
Low Five States				
New York	4.73	0.10	(47)	-8.96
New Jersey	4.62	0.15	(48)	-10.99
D.C.	4.49	0.62	(49)	-13.59
New Mexico	4.36	0.30	(50)	-15.95 <sup>c</sup>
Hawaii	2.86	0.29	(51)	-44.92 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.69	0.23	(14)	9.57	Montana	5.37	0.45	(24)	3.49
Alaska	5.09	0.74	(35)	-1.97	Nebraska	5.79	0.37	(13)	11.45
Arizona	4.83	0.18	(42)	-7.03	Nevada	5.47	0.31	(17)	5.42
Arkansas	5.96	0.29	(10)	14.74	New Hampshire	6.31	0.45	(03)	21.45 <sup>c</sup>
California	5.17	0.08	(30)	-0.54	New Jersey	4.62	0.15	(48)	-10.99
Colorado	5.49	0.23	(16)	5.69	New Mexico	4.36	0.30	(50)	-15.95 <sup>c</sup>
Connecticut	5.20	0.24	(28)	0.08	New York	4.73	0.10	(47)	-8.96
Delaware	5.46	0.51	(19)	5.19	North Carolina	5.06	0.16	(36)	-2.57
D.C.	4.49	0.62	(49)	-13.59	North Dakota	5.12	0.56	(34)	-1.35
Florida	4.82	0.10	(43)	-7.27	Ohio	5.41	0.14	(21)	4.13
Georgia	4.79	0.16	(44)	-7.78	Oklahoma	5.47	0.25	(18)	5.37
Hawaii	2.86	0.29	(51)	-44.92 <sup>c</sup>	Oregon	6.07	0.25	(06)	16.96 <sup>c</sup>
Idaho	5.81	0.40	(12)	11.79	Pennsylvania	4.95	0.12	(37)	-4.65
Illinois	4.88	0.13	(40)	-5.99	Rhode Island	4.91	0.43	(39)	-5.55
Indiana	5.40	0.19	(23)	3.95	South Carolina	5.66	0.24	(15)	8.98
Iowa	6.71	0.30	(01)	29.17 <sup>c</sup>	South Dakota	6.28	0.56	(04)	20.84 <sup>c</sup>
Kansas	5.99	0.30	(08)	15.39 <sup>c</sup>	Tennessee	5.84	0.20	(11)	12.49
Kentucky	5.46	0.23	(20)	5.05	Texas	4.84	0.10	(41)	-6.78
Louisiana	5.18	0.23	(29)	-0.18	Utah	5.15	0.32	(31)	-0.84
Maine	6.20	0.42	(05)	19.39 <sup>c</sup>	Vermont	5.27	0.57	(27)	1.49
Maryland	4.73	0.19	(46)	-8.92	Virginia	4.74	0.17	(45)	-8.82
Massachusetts	5.30	0.18	(26)	2.08	Washington	6.38	0.20	(02)	22.89 <sup>c</sup>
Michigan	6.02	0.16	(07)	15.88 <sup>c</sup>	West Virginia	4.92	0.31	(38)	-5.27
Minnesota	5.34	0.21	(25)	2.78	Wisconsin	5.99	0.21	(09)	15.36 <sup>c</sup>
Mississippi	5.14	0.29	(32)	-1.01	Wyoming	5.14	0.62	(33)	-1.08
Missouri	5.41	0.19	(22)	4.10					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 3.20  
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.46	0.02		
High Five States				
South Dakota	4.53	0.46	(01)	30.95 <sup>c</sup>
Iowa	4.24	0.22	(02)	22.57 <sup>c</sup>
Vermont	4.16	0.47	(03)	20.06 <sup>c</sup>
Nebraska	4.13	0.29	(04)	19.44 <sup>c</sup>
Arkansas	4.07	0.22	(05)	17.67 <sup>c</sup>
Low Five States				
New York	2.93	0.07	(47)	-15.25 <sup>c</sup>
D.C.	2.93	0.44	(48)	-15.48 <sup>c</sup>
New Jersey	2.88	0.11	(49)	-16.87 <sup>c</sup>
Nevada	2.76	0.21	(50)	-20.17 <sup>c</sup>
Hawaii	1.83	0.22	(51)	-47.26 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.99	0.17	(09)	15.25 <sup>c</sup>	Montana	3.60	0.36	(29)	3.97
Alaska	3.70	0.56	(23)	6.97	Nebraska	4.13	0.29	(04)	19.44 <sup>c</sup>
Arizona	3.36	0.14	(35)	-2.86	Nevada	2.76	0.21	(50)	-20.17 <sup>c</sup>
Arkansas	4.07	0.22	(05)	17.67 <sup>c</sup>	New Hampshire	3.63	0.31	(26)	4.99
California	3.46	0.06	(32)	-0.17	New Jersey	2.88	0.11	(49)	-16.87 <sup>c</sup>
Colorado	3.61	0.17	(28)	4.42	New Mexico	3.06	0.24	(46)	-11.48
Connecticut	3.44	0.18	(33)	-0.73	New York	2.93	0.07	(47)	-15.25 <sup>c</sup>
Delaware	3.20	0.35	(42)	-7.42	North Carolina	3.30	0.11	(38)	-4.78
D.C.	2.93	0.44	(48)	-15.48 <sup>c</sup>	North Dakota	3.79	0.44	(20)	9.55
Florida	3.13	0.07	(45)	-9.69	Ohio	3.68	0.10	(25)	6.37
Georgia	3.31	0.12	(37)	-4.45	Oklahoma	3.83	0.19	(17)	10.53
Hawaii	1.83	0.22	(51)	-47.26 <sup>c</sup>	Oregon	3.90	0.19	(14)	12.53
Idaho	4.06	0.32	(07)	17.16 <sup>c</sup>	Pennsylvania	3.43	0.09	(34)	-0.98
Illinois	3.27	0.10	(41)	-5.65	Rhode Island	3.28	0.32	(40)	-5.31
Indiana	3.77	0.15	(21)	9.02	South Carolina	3.29	0.16	(39)	-5.06
Iowa	4.24	0.22	(02)	22.57 <sup>c</sup>	South Dakota	4.53	0.46	(01)	30.95 <sup>c</sup>
Kansas	3.79	0.22	(19)	9.56	Tennessee	4.01	0.15	(08)	15.82 <sup>c</sup>
Kentucky	3.88	0.18	(15)	12.08	Texas	3.46	0.08	(31)	0.02
Louisiana	3.58	0.17	(30)	3.44	Utah	3.63	0.26	(27)	4.88
Maine	4.06	0.31	(06)	17.42 <sup>c</sup>	Vermont	4.16	0.47	(03)	20.06 <sup>c</sup>
Maryland	3.17	0.14	(43)	-8.54	Virginia	3.15	0.12	(44)	-9.09
Massachusetts	3.34	0.13	(36)	-3.37	Washington	3.96	0.15	(10)	14.41
Michigan	3.91	0.12	(12)	12.88	West Virginia	3.90	0.26	(13)	12.80
Minnesota	3.69	0.16	(24)	6.52	Wisconsin	3.82	0.16	(18)	10.47
Mississippi	3.88	0.22	(16)	12.04	Wyoming	3.93	0.52	(11)	13.46
Missouri	3.77	0.15	(22)	8.98					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 4.24  
Cancer of the Female Breast (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	22.64	0.05		
High Five States				
D.C.	29.79	1.37	(01)	31.63 <sup>c</sup>
Louisiana	25.44	0.45	(02)	12.39 <sup>c</sup>
New Jersey	25.16	0.31	(03)	11.15 <sup>c</sup>
Ohio	24.81	0.26	(04)	9.59
Mississippi	24.72	0.54	(05)	9.18
Low Five States				
South Dakota	20.29	0.91	(47)	-10.35 <sup>c</sup>
Nebraska	20.14	0.62	(48)	-11.02 <sup>c</sup>
Montana	19.83	0.82	(49)	-12.38 <sup>c</sup>
Colorado	19.57	0.40	(50)	-13.56 <sup>c</sup>
Hawaii	16.41	0.65	(51)	-27.50 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	23.33	0.41	(19)	3.05	Montana	19.83	0.82	(49)	-12.38 <sup>c</sup>
Alaska	24.23	1.44	(07)	7.05	Nebraska	20.14	0.62	(48)	-11.02 <sup>c</sup>
Arizona	20.62	0.34	(45)	-8.90	Nevada	22.99	0.60	(22)	1.59
Arkansas	23.41	0.52	(17)	3.44	New Hampshire	21.33	0.74	(36)	-5.76
California	21.85	0.15	(27)	-3.46	New Jersey	25.16	0.31	(03)	11.15 <sup>c</sup>
Colorado	19.57	0.40	(50)	-13.56 <sup>c</sup>	New Mexico	20.91	0.61	(41)	-7.64
Connecticut	21.57	0.44	(33)	-4.72	New York	22.30	0.19	(24)	-1.48
Delaware	22.79	0.93	(23)	0.69	North Carolina	23.09	0.29	(21)	2.01
D.C.	29.79	1.37	(01)	31.63 <sup>c</sup>	North Dakota	21.61	1.04	(32)	-4.52
Florida	21.26	0.19	(39)	-6.07	Ohio	24.81	0.26	(04)	9.59
Georgia	23.44	0.31	(16)	3.55	Oklahoma	23.85	0.47	(12)	5.37
Hawaii	16.41	0.65	(51)	-27.50 <sup>c</sup>	Oregon	21.62	0.44	(31)	-4.48
Idaho	21.63	0.74	(30)	-4.44	Pennsylvania	23.84	0.24	(13)	5.31
Illinois	23.59	0.25	(14)	4.21	Rhode Island	20.78	0.79	(44)	-8.20
Indiana	23.85	0.36	(11)	5.38	South Carolina	23.46	0.42	(15)	3.65
Iowa	21.34	0.48	(35)	-5.72	South Dakota	20.29	0.91	(47)	-10.35 <sup>c</sup>
Kansas	22.18	0.52	(25)	-2.01	Tennessee	23.35	0.36	(18)	3.15
Kentucky	23.12	0.43	(20)	2.12	Texas	21.82	0.19	(29)	-3.60
Louisiana	25.44	0.45	(02)	12.39 <sup>c</sup>	Utah	21.82	0.64	(28)	-3.60
Maine	20.87	0.69	(43)	-7.81	Vermont	20.39	1.02	(46)	-9.91
Maryland	24.47	0.39	(06)	8.10	Virginia	24.04	0.33	(09)	6.22
Massachusetts	21.25	0.33	(40)	-6.10	Washington	21.52	0.35	(34)	-4.95
Michigan	23.97	0.28	(10)	5.89	West Virginia	22.06	0.60	(26)	-2.53
Minnesota	20.87	0.37	(42)	-7.79	Wisconsin	21.28	0.36	(38)	-5.98
Mississippi	24.72	0.54	(05)	9.18	Wyoming	21.32	1.20	(37)	-5.82
Missouri	24.17	0.37	(08)	6.79					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 5.14  
Cancer of the Cervix Uteri (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.35	0.02		
High Five States				
Mississippi	3.83	0.22	(01)	63.02 <sup>c</sup>
Arkansas	3.38	0.21	(02)	43.65 <sup>c</sup>
Louisiana	3.23	0.17	(03)	37.26 <sup>c</sup>
Oklahoma	2.93	0.17	(04)	24.42 <sup>c</sup>
Alabama	2.92	0.15	(05)	24.05 <sup>c</sup>
Low Five States				
Minnesota	1.47	0.10	(47)	-37.32 <sup>c</sup>
North Dakota	1.40	0.28	(48)	-40.61 <sup>c</sup>
Massachusetts	1.34	0.08	(49)	-43.10 <sup>c</sup>
Vermont	1.21	0.26	(50)	-48.36 <sup>c</sup>
Utah	1.20	0.15	(51)	-48.82 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.92	0.15	(05)	24.05 <sup>c</sup>	Montana	1.71	0.25	(43)	-27.38 <sup>c</sup>
Alaska	2.20	0.40	(27)	-6.60	Nebraska	1.77	0.20	(41)	-24.75 <sup>c</sup>
Arizona	1.90	0.11	(36)	-19.23 <sup>c</sup>	Nevada	2.44	0.19	(19)	3.63
Arkansas	3.38	0.21	(02)	43.65 <sup>c</sup>	New Hampshire	2.03	0.23	(32)	-13.64
California	2.36	0.05	(22)	0.45	New Jersey	2.38	0.10	(21)	1.12
Colorado	1.76	0.12	(42)	-25.32 <sup>c</sup>	New Mexico	2.40	0.21	(20)	1.91
Connecticut	1.59	0.12	(45)	-32.22 <sup>c</sup>	New York	2.33	0.06	(24)	-0.89
Delaware	2.57	0.33	(17)	9.22	North Carolina	2.16	0.09	(28)	-8.29
D.C.	2.58	0.42	(16)	9.73	North Dakota	1.40	0.28	(48)	-40.61 <sup>c</sup>
Florida	2.64	0.07	(13)	12.23	Ohio	2.67	0.09	(12)	13.59
Georgia	2.73	0.11	(11)	16.32 <sup>c</sup>	Oklahoma	2.93	0.17	(04)	24.42 <sup>c</sup>
Hawaii	1.79	0.22	(39)	-23.85 <sup>c</sup>	Oregon	1.85	0.14	(38)	-21.15 <sup>c</sup>
Idaho	2.07	0.24	(31)	-11.92	Pennsylvania	2.21	0.08	(26)	-6.20
Illinois	2.60	0.09	(15)	10.47	Rhode Island	1.61	0.23	(44)	-31.66 <sup>c</sup>
Indiana	2.36	0.12	(23)	0.42	South Carolina	2.88	0.15	(06)	22.37 <sup>c</sup>
Iowa	2.08	0.16	(30)	-11.43	South Dakota	1.92	0.30	(35)	-18.27 <sup>c</sup>
Kansas	1.94	0.17	(34)	-17.56 <sup>c</sup>	Tennessee	2.62	0.12	(14)	11.64
Kentucky	2.81	0.15	(09)	19.71 <sup>c</sup>	Texas	2.85	0.07	(07)	21.09 <sup>c</sup>
Louisiana	3.23	0.17	(03)	37.26 <sup>c</sup>	Utah	1.20	0.15	(51)	-48.82 <sup>c</sup>
Maine	1.78	0.21	(40)	-24.46 <sup>c</sup>	Vermont	1.21	0.26	(50)	-48.36 <sup>c</sup>
Maryland	2.26	0.12	(25)	-3.80	Virginia	2.10	0.10	(29)	-10.77
Massachusetts	1.34	0.08	(49)	-43.10 <sup>c</sup>	Washington	1.89	0.10	(37)	-19.70 <sup>c</sup>
Michigan	2.02	0.09	(33)	-13.92	West Virginia	2.77	0.24	(10)	17.98 <sup>c</sup>
Minnesota	1.47	0.10	(47)	-37.32 <sup>c</sup>	Wisconsin	1.57	0.10	(46)	-33.19 <sup>c</sup>
Mississippi	3.83	0.22	(01)	63.02 <sup>c</sup>	Wyoming	2.82	0.46	(08)	19.98 <sup>c</sup>
Missouri	2.54	0.13	(18)	7.96					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 6.25  
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	16.37	0.03		
High Five States				
D.C.	20.29	0.85	(01)	23.92 <sup>c</sup>
Mississippi	20.07	0.37	(02)	22.58 <sup>c</sup>
Louisiana	19.32	0.30	(03)	18.02 <sup>c</sup>
Kentucky	19.30	0.29	(04)	17.87 <sup>c</sup>
West Virginia	19.25	0.41	(05)	17.58 <sup>c</sup>
Low Five States				
Arizona	14.24	0.21	(47)	-13.02 <sup>c</sup>
Idaho	14.08	0.44	(48)	-14.00 <sup>c</sup>
Connecticut	13.86	0.26	(49)	-15.38 <sup>c</sup>
Hawaii	13.80	0.43	(50)	-15.72 <sup>c</sup>
Utah	12.40	0.36	(51)	-24.29 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	18.14	0.27	(07)	10.76 <sup>c</sup>	Montana	15.30	0.52	(39)	-6.57
Alaska	17.37	0.95	(18)	6.07	Nebraska	17.89	0.42	(11)	9.26
Arizona	14.24	0.21	(47)	-13.02 <sup>c</sup>	Nevada	17.71	0.39	(14)	8.15
Arkansas	18.67	0.34	(06)	14.00 <sup>c</sup>	New Hampshire	15.23	0.47	(41)	-7.01
California	14.83	0.09	(43)	-9.40	New Jersey	17.46	0.19	(16)	6.66
Colorado	14.30	0.26	(46)	-12.66 <sup>c</sup>	New Mexico	15.41	0.39	(38)	-5.88
Connecticut	13.86	0.26	(49)	-15.38 <sup>c</sup>	New York	16.03	0.12	(32)	-2.09
Delaware	16.58	0.59	(24)	1.23	North Carolina	15.79	0.18	(34)	-3.58
D.C.	20.29	0.85	(01)	23.92 <sup>c</sup>	North Dakota	17.09	0.66	(20)	4.38
Florida	15.24	0.11	(40)	-6.90	Ohio	18.05	0.17	(10)	10.25 <sup>c</sup>
Georgia	16.39	0.20	(29)	0.11	Oklahoma	17.80	0.30	(12)	8.74
Hawaii	13.80	0.43	(50)	-15.72 <sup>c</sup>	Oregon	15.47	0.27	(36)	-5.49
Idaho	14.08	0.44	(48)	-14.00 <sup>c</sup>	Pennsylvania	17.74	0.15	(13)	8.37
Illinois	18.06	0.17	(09)	10.31 <sup>c</sup>	Rhode Island	15.44	0.50	(37)	-5.68
Indiana	17.40	0.23	(17)	6.27	South Carolina	16.90	0.27	(21)	3.23
Iowa	17.28	0.31	(19)	5.54	South Dakota	16.61	0.60	(23)	1.42
Kansas	16.47	0.33	(27)	0.56	Tennessee	18.09	0.24	(08)	10.45 <sup>c</sup>
Kentucky	19.30	0.29	(04)	17.87 <sup>c</sup>	Texas	16.16	0.13	(31)	-1.33
Louisiana	19.32	0.30	(03)	18.02 <sup>c</sup>	Utah	12.40	0.36	(51)	-24.29 <sup>c</sup>
Maine	16.53	0.45	(25)	0.98	Vermont	15.92	0.67	(33)	-2.75
Maryland	16.76	0.24	(22)	2.35	Virginia	16.22	0.21	(30)	-0.93
Massachusetts	15.64	0.21	(35)	-4.48	Washington	14.61	0.21	(45)	-10.75 <sup>c</sup>
Michigan	16.46	0.17	(28)	0.51	West Virginia	19.25	0.41	(05)	17.58 <sup>c</sup>
Minnesota	14.76	0.23	(44)	-9.87	Wisconsin	15.00	0.22	(42)	-8.40
Mississippi	20.07	0.37	(02)	22.58 <sup>c</sup>	Wyoming	16.49	0.78	(26)	0.70
Missouri	17.48	0.23	(15)	6.78					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 6.26  
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	19.65	0.05		
High Five States				
Mississippi	24.82	0.65	(01)	26.30 <sup>c</sup>
Louisiana	24.05	0.52	(02)	22.38 <sup>c</sup>
West Virginia	23.43	0.70	(03)	19.27 <sup>c</sup>
Kentucky	23.43	0.51	(04)	19.23 <sup>c</sup>
Arkansas	22.70	0.58	(05)	15.54 <sup>c</sup>
Low Five States				
Colorado	16.74	0.43	(47)	-14.78 <sup>c</sup>
Montana	16.73	0.82	(48)	-14.85 <sup>c</sup>
Connecticut	16.15	0.44	(49)	-17.79 <sup>c</sup>
Idaho	16.13	0.70	(50)	-17.91 <sup>c</sup>
Utah	14.35	0.58	(51)	-26.97 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	22.47	0.47	(06)	14.34 <sup>c</sup>	Montana	16.73	0.82	(48)	-14.85 <sup>c</sup>
Alaska	19.62	1.51	(28)	-0.17	Nebraska	21.41	0.71	(13)	8.94
Arizona	17.08	0.35	(46)	-13.07 <sup>c</sup>	Nevada	21.00	0.65	(18)	6.88
Arkansas	22.70	0.58	(05)	15.54 <sup>c</sup>	New Hampshire	17.93	0.78	(40)	-8.75
California	17.53	0.16	(42)	-10.78 <sup>c</sup>	New Jersey	21.02	0.33	(17)	6.96
Colorado	16.74	0.43	(47)	-14.78 <sup>c</sup>	New Mexico	18.80	0.65	(33)	-4.34
Connecticut	16.15	0.44	(49)	-17.79 <sup>c</sup>	New York	19.16	0.21	(32)	-2.47
Delaware	20.43	1.00	(21)	3.98	North Carolina	19.41	0.32	(30)	-1.24
D.C.	22.20	1.41	(07)	12.98 <sup>c</sup>	North Dakota	21.22	1.12	(15)	8.00
Florida	18.14	0.19	(37)	-7.70	Ohio	21.85	0.29	(09)	11.20 <sup>c</sup>
Georgia	20.17	0.36	(24)	2.66	Oklahoma	22.09	0.52	(08)	12.40 <sup>c</sup>
Hawaii	17.27	0.71	(44)	-12.11 <sup>c</sup>	Oregon	18.11	0.45	(38)	-7.81
Idaho	16.13	0.70	(50)	-17.91 <sup>c</sup>	Pennsylvania	21.45	0.26	(12)	9.19
Illinois	21.69	0.29	(11)	10.40 <sup>c</sup>	Rhode Island	18.60	0.86	(36)	-5.36
Indiana	21.32	0.40	(14)	8.51	South Carolina	20.22	0.46	(22)	2.92
Iowa	19.98	0.51	(26)	1.69	South Dakota	19.83	0.99	(27)	0.90
Kansas	20.56	0.57	(19)	4.64	Tennessee	21.80	0.41	(10)	10.95 <sup>c</sup>
Kentucky	23.43	0.51	(04)	19.23 <sup>c</sup>	Texas	20.03	0.22	(25)	1.93
Louisiana	24.05	0.52	(02)	22.38 <sup>c</sup>	Utah	14.35	0.58	(51)	-26.97 <sup>c</sup>
Maine	20.19	0.77	(23)	2.77	Vermont	17.37	1.09	(43)	-11.62 <sup>c</sup>
Maryland	20.51	0.42	(20)	4.38	Virginia	19.18	0.35	(31)	-2.39
Massachusetts	18.72	0.36	(34)	-4.74	Washington	17.12	0.35	(45)	-12.87 <sup>c</sup>
Michigan	19.41	0.29	(29)	-1.21	West Virginia	23.43	0.70	(03)	19.27 <sup>c</sup>
Minnesota	17.58	0.39	(41)	-10.52 <sup>c</sup>	Wisconsin	17.96	0.37	(39)	-8.58
Mississippi	24.82	0.65	(01)	26.30 <sup>c</sup>	Wyoming	18.70	1.24	(35)	-4.85
Missouri	21.16	0.39	(16)	7.70					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 6.27  
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	13.85	0.04		
High Five States				
D.C.	18.93	1.08	(01)	36.66 <sup>c</sup>
Mississippi	16.51	0.44	(02)	19.21 <sup>c</sup>
Kentucky	16.21	0.35	(03)	17.07 <sup>c</sup>
West Virginia	16.13	0.50	(04)	16.44 <sup>c</sup>
Louisiana	15.70	0.35	(05)	13.34 <sup>c</sup>
Low Five States				
Colorado	12.40	0.32	(47)	-10.48 <sup>c</sup>
Connecticut	12.09	0.32	(48)	-12.73 <sup>c</sup>
Arizona	11.86	0.26	(49)	-14.35 <sup>c</sup>
Hawaii	10.90	0.51	(50)	-21.30 <sup>c</sup>
Utah	10.69	0.45	(51)	-22.81 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	15.00	0.32	(12)	8.27	Montana	14.02	0.67	(24)	1.21
Alaska	15.08	1.20	(11)	8.87	Nebraska	14.96	0.51	(14)	7.99
Arizona	11.86	0.26	(49)	-14.35 <sup>c</sup>	Nevada	14.88	0.49	(16)	7.42
Arkansas	15.56	0.42	(06)	12.32 <sup>c</sup>	New Hampshire	13.12	0.57	(38)	-5.29
California	12.71	0.11	(42)	-8.21	New Jersey	14.90	0.23	(15)	7.59
Colorado	12.40	0.32	(47)	-10.48 <sup>c</sup>	New Mexico	12.70	0.47	(43)	-8.29
Connecticut	12.09	0.32	(48)	-12.73 <sup>c</sup>	New York	13.78	0.15	(30)	-0.49
Delaware	13.34	0.69	(33)	-3.66	North Carolina	13.12	0.22	(39)	-5.30
D.C.	18.93	1.08	(01)	36.66 <sup>c</sup>	North Dakota	13.85	0.79	(29)	0.01
Florida	12.85	0.14	(40)	-7.24	Ohio	15.24	0.20	(09)	10.05 <sup>c</sup>
Georgia	13.76	0.24	(31)	-0.67	Oklahoma	14.51	0.36	(21)	4.73
Hawaii	10.90	0.51	(50)	-21.30 <sup>c</sup>	Oregon	13.26	0.34	(34)	-4.23
Idaho	12.43	0.56	(46)	-10.24 <sup>c</sup>	Pennsylvania	15.00	0.18	(13)	8.26
Illinois	15.43	0.20	(07)	11.43 <sup>c</sup>	Rhode Island	13.25	0.61	(35)	-4.36
Indiana	14.51	0.28	(20)	4.77	South Carolina	14.22	0.32	(23)	2.65
Iowa	15.21	0.38	(10)	9.78	South Dakota	13.86	0.72	(28)	0.05
Kansas	13.14	0.39	(36)	-5.11	Tennessee	15.28	0.29	(08)	10.32 <sup>c</sup>
Kentucky	16.21	0.35	(03)	17.07 <sup>c</sup>	Texas	13.12	0.15	(37)	-5.28
Louisiana	15.70	0.35	(05)	13.34 <sup>c</sup>	Utah	10.69	0.45	(51)	-22.81 <sup>c</sup>
Maine	13.91	0.55	(27)	0.40	Vermont	14.76	0.85	(17)	6.55
Maryland	13.95	0.29	(26)	0.74	Virginia	13.98	0.25	(25)	0.92
Massachusetts	13.42	0.25	(32)	-3.09	Washington	12.50	0.26	(45)	-9.74
Michigan	14.24	0.22	(22)	2.85	West Virginia	16.13	0.50	(04)	16.44 <sup>c</sup>
Minnesota	12.56	0.28	(44)	-9.31	Wisconsin	12.74	0.27	(41)	-8.04
Mississippi	16.51	0.44	(02)	19.21 <sup>c</sup>	Wyoming	14.67	1.00	(18)	5.94
Missouri	14.65	0.28	(19)	5.75					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 7.14  
Cancer of the Corpus and Uterus, NOS (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.26	0.02		
High Five States				
D.C.	7.69	0.69	(01)	80.64 <sup>c</sup>
New Jersey	5.37	0.14	(02)	26.11 <sup>c</sup>
New York	5.22	0.09	(03)	22.59 <sup>c</sup>
Delaware	5.08	0.43	(04)	19.44 <sup>c</sup>
Maryland	4.96	0.17	(05)	16.59 <sup>c</sup>
Low Five States				
Wyoming	3.25	0.48	(47)	-23.70 <sup>c</sup>
New Mexico	3.18	0.24	(48)	-25.38 <sup>c</sup>
Arizona	3.09	0.13	(49)	-27.36 <sup>c</sup>
Arkansas	2.84	0.18	(50)	-33.27 <sup>c</sup>
Alaska	2.72	0.51	(51)	-36.01 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.41	0.15	(44)	-19.92 <sup>c</sup>	Montana	3.96	0.36	(30)	-7.04
Alaska	2.72	0.51	(51)	-36.01 <sup>c</sup>	Nebraska	4.70	0.30	(12)	10.46
Arizona	3.09	0.13	(49)	-27.36 <sup>c</sup>	Nevada	3.25	0.22	(46)	-23.61 <sup>c</sup>
Arkansas	2.84	0.18	(50)	-33.27 <sup>c</sup>	New Hampshire	4.62	0.34	(13)	8.47
California	4.19	0.07	(25)	-1.52	New Jersey	5.37	0.14	(02)	26.11 <sup>c</sup>
Colorado	3.44	0.17	(43)	-19.08 <sup>c</sup>	New Mexico	3.18	0.24	(48)	-25.38 <sup>c</sup>
Connecticut	4.37	0.20	(16)	2.63	New York	5.22	0.09	(03)	22.59 <sup>c</sup>
Delaware	5.08	0.43	(04)	19.44 <sup>c</sup>	North Carolina	4.04	0.12	(29)	-5.05
D.C.	7.69	0.69	(01)	80.64 <sup>c</sup>	North Dakota	4.49	0.47	(15)	5.51
Florida	3.82	0.08	(35)	-10.31	Ohio	4.82	0.12	(09)	13.32
Georgia	3.76	0.13	(37)	-11.66	Oklahoma	3.83	0.19	(33)	-9.99
Hawaii	3.91	0.31	(31)	-8.01	Oregon	3.89	0.19	(32)	-8.54
Idaho	4.15	0.32	(26)	-2.40	Pennsylvania	4.96	0.11	(06)	16.46 <sup>c</sup>
Illinois	4.80	0.12	(10)	12.89	Rhode Island	3.77	0.33	(36)	-11.53
Indiana	4.36	0.15	(17)	2.38	South Carolina	4.20	0.18	(22)	-1.32
Iowa	4.88	0.22	(08)	14.65	South Dakota	4.22	0.42	(21)	-0.87
Kansas	3.74	0.21	(38)	-12.21	Tennessee	3.70	0.14	(39)	-13.16
Kentucky	3.65	0.17	(41)	-14.27	Texas	3.62	0.08	(42)	-14.91
Louisiana	3.69	0.17	(40)	-13.23	Utah	3.37	0.25	(45)	-20.71 <sup>c</sup>
Maine	4.15	0.30	(27)	-2.60	Vermont	4.53	0.48	(14)	6.52
Maryland	4.96	0.17	(05)	16.59 <sup>c</sup>	Virginia	4.30	0.14	(19)	1.13
Massachusetts	4.20	0.14	(23)	-1.34	Washington	4.19	0.15	(24)	-1.47
Michigan	4.94	0.13	(07)	16.04 <sup>c</sup>	West Virginia	4.30	0.26	(18)	1.15
Minnesota	4.10	0.17	(28)	-3.71	Wisconsin	4.71	0.17	(11)	10.62
Mississippi	3.82	0.21	(34)	-10.28	Wyoming	3.25	0.48	(47)	-23.70 <sup>c</sup>
Missouri	4.22	0.15	(20)	-0.81					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 8.18  
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.27	0.02		
High Five States				
New Hampshire	5.93	0.29	(01)	38.91 <sup>c</sup>
Maine	5.83	0.27	(02)	36.59 <sup>c</sup>
D.C.	5.56	0.45	(03)	30.31 <sup>c</sup>
Rhode Island	5.38	0.30	(04)	26.15 <sup>c</sup>
Ohio	5.12	0.09	(05)	20.10 <sup>c</sup>
Low Five States				
Texas	3.54	0.06	(47)	-17.07 <sup>c</sup>
California	3.50	0.05	(48)	-17.90 <sup>c</sup>
New Mexico	3.37	0.18	(49)	-21.01 <sup>c</sup>
Hawaii	3.23	0.21	(50)	-24.19 <sup>c</sup>
Utah	2.80	0.17	(51)	-34.30 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.04	0.12	(42)	-5.21	Montana	4.65	0.28	(17)	9.09
Alaska	4.13	0.44	(37)	-3.19	Nebraska	4.47	0.21	(25)	4.73
Arizona	3.99	0.11	(43)	-6.56	Nevada	4.67	0.20	(16)	9.54
Arkansas	3.84	0.15	(46)	-9.88	New Hampshire	5.93	0.29	(01)	38.91 <sup>c</sup>
California	3.50	0.05	(48)	-17.90 <sup>c</sup>	New Jersey	4.07	0.09	(40)	-4.50
Colorado	4.07	0.14	(41)	-4.52	New Mexico	3.37	0.18	(49)	-21.01 <sup>c</sup>
Connecticut	4.27	0.15	(31)	0.17	New York	4.18	0.06	(35)	-2.00
Delaware	4.19	0.29	(34)	-1.89	North Carolina	3.91	0.09	(44)	-8.32
D.C.	5.56	0.45	(03)	30.31 <sup>c</sup>	North Dakota	4.15	0.33	(36)	-2.78
Florida	4.09	0.06	(39)	-4.19	Ohio	5.12	0.09	(05)	20.10 <sup>c</sup>
Georgia	3.89	0.10	(45)	-8.92	Oklahoma	4.58	0.15	(19)	7.40
Hawaii	3.23	0.21	(50)	-24.19 <sup>c</sup>	Oregon	4.60	0.15	(18)	7.83
Idaho	4.55	0.25	(22)	6.64	Pennsylvania	4.95	0.08	(08)	15.95 <sup>c</sup>
Illinois	4.57	0.08	(20)	7.19	Rhode Island	5.38	0.30	(04)	26.15 <sup>c</sup>
Indiana	5.01	0.12	(07)	17.47 <sup>c</sup>	South Carolina	4.33	0.13	(26)	1.50
Iowa	4.77	0.17	(14)	11.93	South Dakota	4.85	0.33	(11)	13.78
Kansas	4.20	0.17	(33)	-1.58	Tennessee	4.13	0.11	(38)	-3.29
Kentucky	4.50	0.14	(23)	5.57	Texas	3.54	0.06	(47)	-17.07 <sup>c</sup>
Louisiana	4.28	0.14	(30)	0.31	Utah	2.80	0.17	(51)	-34.30 <sup>c</sup>
Maine	5.83	0.27	(02)	36.59 <sup>c</sup>	Vermont	4.74	0.36	(15)	11.17
Maryland	4.29	0.12	(29)	0.49	Virginia	4.27	0.10	(32)	0.09
Massachusetts	5.10	0.12	(06)	19.59 <sup>c</sup>	Washington	4.87	0.12	(10)	14.19
Michigan	4.78	0.09	(13)	12.12	West Virginia	4.82	0.20	(12)	12.93
Minnesota	4.50	0.13	(24)	5.48	Wisconsin	4.93	0.13	(09)	15.50 <sup>c</sup>
Mississippi	4.32	0.17	(27)	1.22	Wyoming	4.56	0.41	(21)	6.88
Missouri	4.31	0.11	(28)	1.04					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 8.19  
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.61	0.03		
High Five States				
Maine	10.94	0.55	(01)	43.80 <sup>c</sup>
New Hampshire	10.59	0.59	(02)	39.23 <sup>c</sup>
Rhode Island	9.81	0.62	(03)	28.89 <sup>c</sup>
D.C.	9.36	0.90	(04)	23.01 <sup>c</sup>
Ohio	9.33	0.18	(05)	22.64 <sup>c</sup>
Low Five States				
Alaska	6.36	0.79	(47)	-16.41 <sup>c</sup>
California	6.07	0.09	(48)	-20.18 <sup>c</sup>
Hawaii	5.84	0.41	(49)	-23.20 <sup>c</sup>
New Mexico	5.81	0.35	(50)	-23.66 <sup>c</sup>
Utah	4.95	0.33	(51)	-34.98 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.49	0.26	(36)	-1.55	Montana	8.23	0.56	(18)	8.14
Alaska	6.36	0.79	(47)	-16.41 <sup>c</sup>	Nebraska	8.04	0.43	(22)	5.68
Arizona	7.03	0.22	(43)	-7.65	Nevada	8.11	0.38	(21)	6.64
Arkansas	6.66	0.30	(45)	-12.47	New Hampshire	10.59	0.59	(02)	39.23 <sup>c</sup>
California	6.07	0.09	(48)	-20.18 <sup>c</sup>	New Jersey	7.08	0.19	(42)	-6.92
Colorado	7.21	0.28	(40)	-5.22	New Mexico	5.81	0.35	(50)	-23.66 <sup>c</sup>
Connecticut	7.67	0.30	(31)	0.75	New York	7.40	0.13	(38)	-2.74
Delaware	7.83	0.61	(25)	2.89	North Carolina	7.14	0.19	(41)	-6.12
D.C.	9.36	0.90	(04)	23.01 <sup>c</sup>	North Dakota	7.60	0.67	(33)	-0.09
Florida	7.23	0.12	(39)	-4.98	Ohio	9.33	0.18	(05)	22.64 <sup>c</sup>
Georgia	6.82	0.20	(44)	-10.36	Oklahoma	8.28	0.31	(16)	8.79
Hawaii	5.84	0.41	(49)	-23.20 <sup>c</sup>	Oregon	8.17	0.30	(20)	7.38
Idaho	7.93	0.48	(23)	4.17	Pennsylvania	8.98	0.16	(08)	17.99 <sup>c</sup>
Illinois	8.27	0.17	(17)	8.73	Rhode Island	9.81	0.62	(03)	28.89 <sup>c</sup>
Indiana	9.17	0.25	(06)	20.53 <sup>c</sup>	South Carolina	7.64	0.27	(32)	0.45
Iowa	8.79	0.34	(11)	15.55 <sup>c</sup>	South Dakota	8.73	0.66	(12)	14.75
Kansas	7.82	0.34	(26)	2.78	Tennessee	7.43	0.23	(37)	-2.34
Kentucky	8.22	0.29	(19)	8.04	Texas	6.40	0.12	(46)	-15.83 <sup>c</sup>
Louisiana	7.54	0.28	(35)	-0.95	Utah	4.95	0.33	(51)	-34.98 <sup>c</sup>
Maine	10.94	0.55	(01)	43.80 <sup>c</sup>	Vermont	8.34	0.73	(14)	9.59
Maryland	7.73	0.25	(27)	1.65	Virginia	7.71	0.21	(30)	1.34
Massachusetts	9.04	0.24	(07)	18.86 <sup>c</sup>	Washington	8.28	0.24	(15)	8.83
Michigan	8.53	0.19	(13)	12.08	West Virginia	8.93	0.42	(09)	17.42 <sup>c</sup>
Minnesota	7.73	0.25	(28)	1.57	Wisconsin	8.82	0.25	(10)	15.93 <sup>c</sup>
Mississippi	7.86	0.35	(24)	3.32	Wyoming	7.71	0.79	(29)	1.37
Missouri	7.60	0.23	(34)	-0.17					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 8.20  
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.60	0.01		
High Five States				
D.C.	2.67	0.41	(01)	66.58 <sup>c</sup>
New Hampshire	2.10	0.23	(02)	31.18 <sup>c</sup>
Rhode Island	2.07	0.24	(03)	28.82 <sup>c</sup>
Washington	2.05	0.11	(04)	28.00 <sup>c</sup>
Massachusetts	2.04	0.10	(05)	27.23 <sup>c</sup>
Low Five States				
Kansas	1.27	0.12	(47)	-20.82 <sup>c</sup>
North Dakota	1.25	0.25	(48)	-22.18 <sup>c</sup>
Texas	1.24	0.05	(49)	-22.45 <sup>c</sup>
Hawaii	1.05	0.16	(50)	-34.48 <sup>c</sup>
Utah	0.97	0.13	(51)	-39.41 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.31	0.09	(45)	-18.07 <sup>c</sup>	Montana	1.52	0.22	(33)	-5.18
Alaska	1.89	0.43	(07)	17.67 <sup>c</sup>	Nebraska	1.54	0.17	(30)	-4.04
Arizona	1.42	0.09	(42)	-11.61	Nevada	1.62	0.16	(25)	0.81
Arkansas	1.52	0.13	(32)	-5.02	New Hampshire	2.10	0.23	(02)	31.18 <sup>c</sup>
California	1.46	0.04	(40)	-9.23	New Jersey	1.73	0.08	(15)	7.79
Colorado	1.51	0.11	(35)	-5.57	New Mexico	1.29	0.15	(46)	-19.76 <sup>c</sup>
Connecticut	1.67	0.12	(22)	4.43	New York	1.75	0.05	(14)	8.98
Delaware	1.33	0.22	(44)	-17.36 <sup>c</sup>	North Carolina	1.41	0.07	(43)	-12.04
D.C.	2.67	0.41	(01)	66.58 <sup>c</sup>	North Dakota	1.25	0.25	(48)	-22.18 <sup>c</sup>
Florida	1.50	0.05	(37)	-6.76	Ohio	1.86	0.07	(09)	15.70 <sup>c</sup>
Georgia	1.61	0.08	(26)	0.17	Oklahoma	1.54	0.12	(31)	-4.16
Hawaii	1.05	0.16	(50)	-34.48 <sup>c</sup>	Oregon	1.72	0.12	(18)	7.29
Idaho	1.58	0.20	(28)	-1.23	Pennsylvania	1.79	0.06	(11)	11.64
Illinois	1.75	0.07	(13)	9.15	Rhode Island	2.07	0.24	(03)	28.82 <sup>c</sup>
Indiana	1.70	0.10	(20)	6.07	South Carolina	1.70	0.11	(19)	6.32
Iowa	1.51	0.12	(36)	-5.79	South Dakota	1.59	0.25	(27)	-0.58
Kansas	1.27	0.12	(47)	-20.82 <sup>c</sup>	Tennessee	1.48	0.09	(39)	-7.57
Kentucky	1.49	0.11	(38)	-6.79	Texas	1.24	0.05	(49)	-22.45 <sup>c</sup>
Louisiana	1.64	0.11	(23)	2.58	Utah	0.97	0.13	(51)	-39.41 <sup>c</sup>
Maine	1.68	0.19	(21)	4.84	Vermont	1.92	0.31	(06)	19.95 <sup>c</sup>
Maryland	1.63	0.10	(24)	1.60	Virginia	1.52	0.08	(34)	-5.28
Massachusetts	2.04	0.10	(05)	27.23 <sup>c</sup>	Washington	2.05	0.11	(04)	28.00 <sup>c</sup>
Michigan	1.77	0.08	(12)	10.08	West Virginia	1.45	0.15	(41)	-9.70
Minnesota	1.87	0.11	(08)	16.88 <sup>c</sup>	Wisconsin	1.73	0.10	(17)	7.70
Mississippi	1.55	0.13	(29)	-3.57	Wyoming	1.81	0.35	(10)	13.00
Missouri	1.73	0.10	(16)	7.75					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 9.18  
Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.40	0.01		
High Five States				
Vermont	0.73	0.15	(01)	81.41 <sup>c</sup>
Louisiana	0.53	0.05	(02)	31.36 <sup>c</sup>
Maine	0.52	0.09	(03)	31.14 <sup>c</sup>
West Virginia	0.46	0.07	(04)	15.20 <sup>c</sup>
Wisconsin	0.46	0.04	(05)	14.93
Low Five States				
Maryland	0.35	0.04	(40)	-12.92
Indiana	0.34	0.03	(41)	-14.67
Kansas	0.34	0.05	(42)	-15.56 <sup>c</sup>
Arizona	0.33	0.03	(43)	-17.27 <sup>c</sup>
Utah	0.31	0.06	(44)	-22.83 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.41	0.04	(23)	1.56	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.43	0.07	(12)	8.10
Arizona	0.33	0.03	(43)	-17.27 <sup>c</sup>	Nevada	0.44	0.06	(09)	9.01
Arkansas	0.43	0.05	(14)	7.34	New Hampshire	0.36	0.07	(36)	-8.93
California	0.40	0.02	(24)	0.19	New Jersey	0.42	0.03	(18)	5.88
Colorado	0.36	0.04	(38)	-10.20	New Mexico	0.40	0.06	(26)	-0.23
Connecticut	0.43	0.05	(13)	8.09	New York	0.42	0.02	(15)	6.22
Delaware	0.36	0.09	(39)	-11.11	North Carolina	0.42	0.03	(21)	4.13
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.40	0.02	(27)	-0.26	Ohio	0.44	0.03	(07)	9.91
Georgia	0.37	0.03	(33)	-7.41	Oklahoma	0.43	0.05	(11)	8.15
Hawaii	-	-	(-)	-	Oregon	0.42	0.05	(17)	5.89
Idaho	0.40	0.07	(28)	-0.76	Pennsylvania	0.44	0.03	(10)	8.83
Illinois	0.37	0.02	(32)	-6.71	Rhode Island	0.45	0.09	(06)	13.40
Indiana	0.34	0.03	(41)	-14.67	South Carolina	0.38	0.04	(31)	-4.43
Iowa	0.41	0.05	(22)	3.72	South Dakota	-	-	(-)	-
Kansas	0.34	0.05	(42)	-15.56 <sup>c</sup>	Tennessee	0.42	0.04	(16)	6.15
Kentucky	0.37	0.04	(35)	-8.16	Texas	0.40	0.02	(29)	-1.04
Louisiana	0.53	0.05	(02)	31.36 <sup>c</sup>	Utah	0.31	0.06	(44)	-22.83 <sup>c</sup>
Maine	0.52	0.09	(03)	31.14 <sup>c</sup>	Vermont	0.73	0.15	(01)	81.41 <sup>c</sup>
Maryland	0.35	0.04	(40)	-12.92	Virginia	0.39	0.03	(30)	-2.47
Massachusetts	0.40	0.03	(25)	-0.19	Washington	0.36	0.03	(37)	-9.54
Michigan	0.42	0.03	(20)	4.35	West Virginia	0.46	0.07	(04)	15.20 <sup>c</sup>
Minnesota	0.42	0.04	(19)	5.39	Wisconsin	0.46	0.04	(05)	14.93
Mississippi	0.37	0.05	(34)	-7.87	Wyoming	-	-	(-)	-
Missouri	0.44	0.04	(08)	9.02					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 9.19  
Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.50	0.01		
High Five States				
Rhode Island	0.85	0.18	(01)	70.64 <sup>c</sup>
Nebraska	0.66	0.12	(02)	32.77 <sup>c</sup>
Maine	0.66	0.14	(03)	32.01 <sup>c</sup>
Nevada	0.64	0.11	(04)	29.32 <sup>c</sup>
Louisiana	0.63	0.08	(05)	27.42 <sup>c</sup>
Low Five States				
Iowa	0.40	0.07	(38)	-18.58 <sup>c</sup>
Maryland	0.39	0.06	(39)	-20.95 <sup>c</sup>
Mississippi	0.39	0.08	(40)	-21.09 <sup>c</sup>
Utah	0.35	0.08	(41)	-29.43 <sup>c</sup>
Kansas	0.34	0.07	(42)	-32.34 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.51	0.07	(21)	1.68	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.66	0.12	(02)	32.77 <sup>c</sup>
Arizona	0.42	0.05	(37)	-14.63	Nevada	0.64	0.11	(04)	29.32 <sup>c</sup>
Arkansas	0.49	0.09	(25)	-2.05	New Hampshire	0.54	0.14	(13)	7.84
California	0.49	0.03	(24)	-1.83	New Jersey	0.55	0.05	(09)	9.93
Colorado	0.42	0.07	(36)	-14.59	New Mexico	0.52	0.11	(17)	4.41
Connecticut	0.52	0.08	(18)	4.12	New York	0.52	0.03	(19)	3.87
Delaware	-	-	(-)	-	North Carolina	0.54	0.05	(12)	8.18
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.50	0.03	(22)	0.77	Ohio	0.55	0.05	(10)	9.72
Georgia	0.48	0.05	(26)	-2.53	Oklahoma	0.54	0.08	(11)	8.18
Hawaii	-	-	(-)	-	Oregon	0.51	0.07	(20)	2.86
Idaho	0.44	0.11	(34)	-11.87	Pennsylvania	0.55	0.04	(08)	10.53
Illinois	0.45	0.04	(33)	-9.47	Rhode Island	0.85	0.18	(01)	70.64 <sup>c</sup>
Indiana	0.47	0.06	(29)	-4.57	South Carolina	0.43	0.07	(35)	-12.84
Iowa	0.40	0.07	(38)	-18.58 <sup>c</sup>	South Dakota	-	-	(-)	-
Kansas	0.34	0.07	(42)	-32.34 <sup>c</sup>	Tennessee	0.47	0.06	(31)	-6.07
Kentucky	0.50	0.07	(23)	0.27	Texas	0.52	0.03	(16)	4.70
Louisiana	0.63	0.08	(05)	27.42 <sup>c</sup>	Utah	0.35	0.08	(41)	-29.43 <sup>c</sup>
Maine	0.66	0.14	(03)	32.01 <sup>c</sup>	Vermont	-	-	(-)	-
Maryland	0.39	0.06	(39)	-20.95 <sup>c</sup>	Virginia	0.48	0.05	(28)	-2.66
Massachusetts	0.53	0.06	(14)	7.54	Washington	0.46	0.06	(32)	-7.60
Michigan	0.53	0.05	(15)	7.31	West Virginia	0.48	0.10	(27)	-2.54
Minnesota	0.56	0.07	(07)	12.92	Wisconsin	0.56	0.07	(06)	13.13
Mississippi	0.39	0.08	(40)	-21.09 <sup>c</sup>	Wyoming	-	-	(-)	-
Missouri	0.47	0.06	(30)	-4.68					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 9.20  
Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.32	0.01		
High Five States				
Louisiana	0.44	0.06	(01)	36.98 <sup>c</sup>
West Virginia	0.43	0.09	(02)	33.73 <sup>c</sup>
Iowa	0.42	0.07	(03)	32.31 <sup>c</sup>
Maine	0.41	0.10	(04)	28.36 <sup>c</sup>
Missouri	0.40	0.05	(05)	25.18 <sup>c</sup>
Low Five States				
New Mexico	0.28	0.07	(33)	-13.77
Nevada	0.26	0.06	(34)	-17.46 <sup>c</sup>
Kentucky	0.25	0.05	(35)	-20.22 <sup>c</sup>
Indiana	0.25	0.04	(36)	-22.17 <sup>c</sup>
Arizona	0.25	0.04	(37)	-22.69 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.32	0.05	(20)	-0.14	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	-	-	(-)	-
Arizona	0.25	0.04	(37)	-22.69 <sup>c</sup>	Nevada	0.26	0.06	(34)	-17.46 <sup>c</sup>
Arkansas	0.39	0.07	(06)	22.67 <sup>c</sup>	New Hampshire	-	-	(-)	-
California	0.33	0.02	(18)	4.11	New Jersey	0.32	0.04	(19)	0.49
Colorado	0.29	0.05	(30)	-10.50	New Mexico	0.28	0.07	(33)	-13.77
Connecticut	0.35	0.06	(10)	10.62	New York	0.35	0.03	(13)	9.92
Delaware	-	-	(-)	-	North Carolina	0.32	0.04	(22)	-0.87
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.31	0.02	(25)	-2.82	Ohio	0.35	0.03	(11)	10.57
Georgia	0.28	0.04	(31)	-11.32	Oklahoma	0.36	0.06	(09)	13.76
Hawaii	-	-	(-)	-	Oregon	0.35	0.06	(14)	9.32
Idaho	-	-	(-)	-	Pennsylvania	0.34	0.03	(16)	7.38
Illinois	0.31	0.03	(24)	-2.81	Rhode Island	-	-	(-)	-
Indiana	0.25	0.04	(36)	-22.17 <sup>c</sup>	South Carolina	0.34	0.05	(17)	5.60
Iowa	0.42	0.07	(03)	32.31 <sup>c</sup>	South Dakota	-	-	(-)	-
Kansas	0.34	0.07	(15)	8.06	Tennessee	0.38	0.05	(07)	20.48 <sup>c</sup>
Kentucky	0.25	0.05	(35)	-20.22 <sup>c</sup>	Texas	0.29	0.02	(29)	-8.40
Louisiana	0.44	0.06	(01)	36.98 <sup>c</sup>	Utah	-	-	(-)	-
Maine	0.41	0.10	(04)	28.36 <sup>c</sup>	Vermont	-	-	(-)	-
Maryland	0.30	0.04	(27)	-5.79	Virginia	0.31	0.04	(26)	-4.08
Massachusetts	0.30	0.04	(28)	-7.44	Washington	0.28	0.04	(32)	-12.27
Michigan	0.32	0.03	(21)	-0.27	West Virginia	0.43	0.09	(02)	33.73 <sup>c</sup>
Minnesota	0.31	0.05	(23)	-2.26	Wisconsin	0.37	0.05	(08)	16.18 <sup>c</sup>
Mississippi	0.35	0.07	(12)	10.15	Wyoming	-	-	(-)	-
Missouri	0.40	0.05	(05)	25.18 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 11.18  
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.97	0.02		
High Five States				
Oklahoma	5.15	0.16	(01)	29.65 <sup>c</sup>
Louisiana	5.15	0.15	(02)	29.64 <sup>c</sup>
North Dakota	5.14	0.37	(03)	29.41 <sup>c</sup>
West Virginia	4.72	0.20	(04)	18.88 <sup>c</sup>
Iowa	4.70	0.16	(05)	18.35 <sup>c</sup>
Low Five States				
Utah	3.40	0.19	(47)	-14.30
New York	3.39	0.06	(48)	-14.58
Connecticut	3.26	0.13	(49)	-17.88 <sup>c</sup>
D.C.	2.97	0.33	(50)	-25.13 <sup>c</sup>
Hawaii	2.70	0.19	(51)	-32.13 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.92	0.12	(30)	-1.40	Montana	3.56	0.25	(41)	-10.43
Alaska	4.61	0.48	(07)	16.13 <sup>c</sup>	Nebraska	4.28	0.21	(19)	7.83
Arizona	3.82	0.11	(35)	-3.92	Nevada	3.98	0.18	(27)	0.10
Arkansas	4.44	0.17	(14)	11.73	New Hampshire	3.56	0.22	(42)	-10.47
California	3.52	0.05	(44)	-11.39	New Jersey	3.48	0.09	(46)	-12.33
Colorado	3.64	0.13	(37)	-8.34	New Mexico	4.51	0.21	(12)	13.61
Connecticut	3.26	0.13	(49)	-17.88 <sup>c</sup>	New York	3.39	0.06	(48)	-14.58
Delaware	3.83	0.28	(34)	-3.72	North Carolina	4.08	0.09	(25)	2.69
D.C.	2.97	0.33	(50)	-25.13 <sup>c</sup>	North Dakota	5.14	0.37	(03)	29.41 <sup>c</sup>
Florida	3.52	0.05	(43)	-11.37	Ohio	4.41	0.08	(16)	10.92
Georgia	3.62	0.09	(39)	-9.01	Oklahoma	5.15	0.16	(01)	29.65 <sup>c</sup>
Hawaii	2.70	0.19	(51)	-32.13 <sup>c</sup>	Oregon	3.85	0.14	(33)	-3.22
Idaho	4.03	0.23	(26)	1.50	Pennsylvania	3.96	0.07	(28)	-0.30
Illinois	4.16	0.08	(22)	4.82	Rhode Island	3.70	0.25	(36)	-6.75
Indiana	4.54	0.12	(09)	14.21	South Carolina	4.15	0.13	(23)	4.35
Iowa	4.70	0.16	(05)	18.35 <sup>c</sup>	South Dakota	4.31	0.31	(18)	8.52
Kansas	4.58	0.17	(08)	15.35 <sup>c</sup>	Tennessee	4.26	0.11	(20)	7.31
Kentucky	4.66	0.14	(06)	17.20 <sup>c</sup>	Texas	4.54	0.07	(10)	14.18
Louisiana	5.15	0.15	(02)	29.64 <sup>c</sup>	Utah	3.40	0.19	(47)	-14.30
Maine	4.13	0.22	(24)	4.02	Vermont	3.63	0.32	(38)	-8.52
Maryland	3.85	0.12	(32)	-3.20	Virginia	3.95	0.10	(29)	-0.69
Massachusetts	3.57	0.10	(40)	-10.06	Washington	3.88	0.11	(31)	-2.38
Michigan	4.18	0.09	(21)	5.11	West Virginia	4.72	0.20	(04)	18.88 <sup>c</sup>
Minnesota	4.45	0.13	(13)	12.07	Wisconsin	4.40	0.12	(17)	10.82
Mississippi	4.52	0.17	(11)	13.79	Wyoming	3.52	0.35	(45)	-11.47
Missouri	4.43	0.12	(15)	11.61					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 11.19  
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.77	0.03		
High Five States				
Oklahoma	7.58	0.30	(01)	31.38 <sup>c</sup>
Louisiana	7.19	0.28	(02)	24.65 <sup>c</sup>
North Dakota	7.14	0.65	(03)	23.66 <sup>c</sup>
Iowa	6.97	0.30	(04)	20.75 <sup>c</sup>
Mississippi	6.63	0.33	(05)	14.92
Low Five States				
Montana	4.94	0.44	(47)	-14.50
D.C.	4.84	0.63	(48)	-16.16 <sup>c</sup>
Connecticut	4.66	0.23	(49)	-19.24 <sup>c</sup>
Utah	4.64	0.33	(50)	-19.59 <sup>c</sup>
Hawaii	4.23	0.35	(51)	-26.74 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.40	0.23	(37)	-6.50	Montana	4.94	0.44	(47)	-14.50
Alaska	6.04	0.81	(21)	4.71	Nebraska	5.99	0.37	(22)	3.84
Arizona	5.45	0.19	(35)	-5.56	Nevada	5.67	0.33	(29)	-1.73
Arkansas	6.30	0.30	(15)	9.11	New Hampshire	5.41	0.42	(36)	-6.19
California	5.15	0.08	(44)	-10.77	New Jersey	5.34	0.16	(40)	-7.51
Colorado	5.39	0.24	(38)	-6.57	New Mexico	6.11	0.37	(19)	5.81
Connecticut	4.66	0.23	(49)	-19.24 <sup>c</sup>	New York	5.10	0.11	(46)	-11.68
Delaware	5.47	0.51	(33)	-5.23	North Carolina	6.28	0.18	(17)	8.73
D.C.	4.84	0.63	(48)	-16.16 <sup>c</sup>	North Dakota	7.14	0.65	(03)	23.66 <sup>c</sup>
Florida	5.16	0.10	(43)	-10.60	Ohio	6.27	0.15	(18)	8.67
Georgia	5.47	0.19	(34)	-5.29	Oklahoma	7.58	0.30	(01)	31.38 <sup>c</sup>
Hawaii	4.23	0.35	(51)	-26.74 <sup>c</sup>	Oregon	5.59	0.25	(30)	-3.18
Idaho	5.51	0.40	(32)	-4.49	Pennsylvania	5.68	0.13	(28)	-1.59
Illinois	5.90	0.15	(24)	2.19	Rhode Island	5.39	0.46	(39)	-6.61
Indiana	6.40	0.21	(12)	10.87	South Carolina	6.28	0.25	(16)	8.74
Iowa	6.97	0.30	(04)	20.75 <sup>c</sup>	South Dakota	5.79	0.53	(26)	0.32
Kansas	6.57	0.32	(09)	13.85	Tennessee	6.39	0.22	(13)	10.78
Kentucky	6.59	0.27	(08)	14.18	Texas	6.48	0.12	(10)	12.27
Louisiana	7.19	0.28	(02)	24.65 <sup>c</sup>	Utah	4.64	0.33	(50)	-19.59 <sup>c</sup>
Maine	6.08	0.42	(20)	5.41	Vermont	5.15	0.57	(45)	-10.86
Maryland	5.68	0.22	(27)	-1.53	Virginia	5.95	0.19	(23)	3.13
Massachusetts	5.25	0.19	(42)	-8.97	Washington	5.54	0.20	(31)	-4.05
Michigan	5.82	0.16	(25)	0.84	West Virginia	6.61	0.37	(06)	14.60
Minnesota	6.42	0.23	(11)	11.27	Wisconsin	6.37	0.22	(14)	10.45
Mississippi	6.63	0.33	(05)	14.92	Wyoming	5.25	0.64	(41)	-8.96
Missouri	6.60	0.22	(07)	14.38					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 11.20  
Cancer of the Kidney And Renal Pelvis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.58	0.02		
High Five States				
Louisiana	3.56	0.17	(01)	37.91 <sup>c</sup>
North Dakota	3.39	0.40	(02)	31.61 <sup>c</sup>
West Virginia	3.27	0.23	(03)	26.91 <sup>c</sup>
Oklahoma	3.23	0.17	(04)	25.20 <sup>c</sup>
New Mexico	3.22	0.24	(05)	24.78 <sup>c</sup>
Low Five States				
New Jersey	2.13	0.09	(47)	-17.36 <sup>c</sup>
New Hampshire	2.03	0.23	(48)	-21.13 <sup>c</sup>
Wyoming	2.03	0.37	(49)	-21.28 <sup>c</sup>
D.C.	1.45	0.30	(50)	-43.72 <sup>c</sup>
Hawaii	1.41	0.18	(51)	-45.22 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.77	0.14	(23)	7.43	Montana	2.40	0.28	(38)	-6.97
Alaska	3.22	0.56	(06)	24.74 <sup>c</sup>	Nebraska	2.86	0.23	(18)	11.07
Arizona	2.44	0.12	(35)	-5.22	Nevada	2.49	0.20	(30)	-3.42
Arkansas	2.97	0.18	(13)	15.24 <sup>c</sup>	New Hampshire	2.03	0.23	(48)	-21.13 <sup>c</sup>
California	2.23	0.05	(43)	-13.48	New Jersey	2.13	0.09	(47)	-17.36 <sup>c</sup>
Colorado	2.27	0.14	(42)	-12.08	New Mexico	3.22	0.24	(05)	24.78 <sup>c</sup>
Connecticut	2.19	0.14	(45)	-15.18 <sup>c</sup>	New York	2.16	0.06	(46)	-16.20 <sup>c</sup>
Delaware	2.62	0.31	(27)	1.76	North Carolina	2.48	0.10	(31)	-3.68
D.C.	1.45	0.30	(50)	-43.72 <sup>c</sup>	North Dakota	3.39	0.40	(02)	31.61 <sup>c</sup>
Florida	2.20	0.06	(44)	-14.49	Ohio	3.00	0.09	(10)	16.35 <sup>c</sup>
Georgia	2.27	0.10	(41)	-11.84	Oklahoma	3.23	0.17	(04)	25.20 <sup>c</sup>
Hawaii	1.41	0.18	(51)	-45.22 <sup>c</sup>	Oregon	2.45	0.15	(34)	-4.92
Idaho	2.80	0.27	(21)	8.55	Pennsylvania	2.64	0.08	(26)	2.57
Illinois	2.83	0.09	(19)	9.90	Rhode Island	2.48	0.27	(32)	-3.70
Indiana	3.07	0.13	(09)	19.00 <sup>c</sup>	South Carolina	2.54	0.14	(28)	-1.44
Iowa	2.96	0.17	(14)	14.91	South Dakota	3.09	0.35	(08)	19.68 <sup>c</sup>
Kansas	2.98	0.19	(12)	15.41 <sup>c</sup>	Tennessee	2.65	0.12	(25)	2.93
Kentucky	3.17	0.16	(07)	22.91 <sup>c</sup>	Texas	2.99	0.07	(11)	15.99 <sup>c</sup>
Louisiana	3.56	0.17	(01)	37.91 <sup>c</sup>	Utah	2.35	0.21	(40)	-8.95
Maine	2.75	0.24	(24)	6.62	Vermont	2.44	0.36	(36)	-5.49
Maryland	2.47	0.12	(33)	-4.08	Virginia	2.43	0.11	(37)	-5.64
Massachusetts	2.37	0.11	(39)	-8.25	Washington	2.53	0.12	(29)	-2.00
Michigan	2.91	0.10	(15)	12.79	West Virginia	3.27	0.23	(03)	26.91 <sup>c</sup>
Minnesota	2.87	0.14	(17)	11.21	Wisconsin	2.82	0.13	(20)	9.32
Mississippi	2.90	0.18	(16)	12.49	Wyoming	2.03	0.37	(49)	-21.28 <sup>c</sup>
Missouri	2.80	0.12	(22)	8.43					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.18  
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.13	0.01		
High Five States				
D.C.	1.76	0.25	(01)	55.45 <sup>c</sup>
West Virginia	1.59	0.12	(02)	39.88 <sup>c</sup>
Kentucky	1.49	0.08	(03)	31.28 <sup>c</sup>
Louisiana	1.49	0.08	(04)	30.93 <sup>c</sup>
Tennessee	1.41	0.06	(05)	24.48 <sup>c</sup>
Low Five States				
Kansas	0.77	0.07	(47)	-31.80 <sup>c</sup>
Oregon	0.75	0.06	(48)	-33.59 <sup>c</sup>
Colorado	0.68	0.06	(49)	-39.94 <sup>c</sup>
Wyoming	0.66	0.15	(50)	-42.02 <sup>c</sup>
Utah	0.43	0.07	(51)	-62.16 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.38	0.07	(07)	21.92 <sup>c</sup>	Montana	0.98	0.13	(33)	-13.61
Alaska	1.01	0.23	(32)	-11.39	Nebraska	0.96	0.10	(36)	-15.73 <sup>c</sup>
Arizona	0.89	0.05	(42)	-21.84 <sup>c</sup>	Nevada	1.17	0.10	(24)	2.80
Arkansas	1.15	0.08	(25)	1.45	New Hampshire	1.20	0.13	(21)	5.97
California	0.87	0.02	(44)	-23.29 <sup>c</sup>	New Jersey	1.21	0.05	(20)	6.31
Colorado	0.68	0.06	(49)	-39.94 <sup>c</sup>	New Mexico	0.97	0.10	(34)	-14.09
Connecticut	0.92	0.07	(37)	-18.96 <sup>c</sup>	New York	1.19	0.03	(22)	4.92
Delaware	1.35	0.16	(10)	19.26 <sup>c</sup>	North Carolina	1.24	0.05	(17)	9.05
D.C.	1.76	0.25	(01)	55.45 <sup>c</sup>	North Dakota	0.88	0.15	(43)	-22.45 <sup>c</sup>
Florida	1.29	0.03	(12)	13.56	Ohio	1.39	0.05	(06)	22.34 <sup>c</sup>
Georgia	1.11	0.05	(27)	-2.61	Oklahoma	1.08	0.07	(29)	-4.79
Hawaii	0.90	0.11	(40)	-20.90 <sup>c</sup>	Oregon	0.75	0.06	(48)	-33.59 <sup>c</sup>
Idaho	0.86	0.11	(45)	-23.95 <sup>c</sup>	Pennsylvania	1.23	0.04	(18)	7.98
Illinois	1.22	0.04	(19)	7.18	Rhode Island	1.26	0.15	(15)	10.93
Indiana	1.25	0.06	(16)	10.07	South Carolina	1.37	0.07	(08)	20.85 <sup>c</sup>
Iowa	0.96	0.07	(35)	-15.38 <sup>c</sup>	South Dakota	0.81	0.14	(46)	-28.41 <sup>c</sup>
Kansas	0.77	0.07	(47)	-31.80 <sup>c</sup>	Tennessee	1.41	0.06	(05)	24.48 <sup>c</sup>
Kentucky	1.49	0.08	(03)	31.28 <sup>c</sup>	Texas	1.13	0.03	(26)	-0.39
Louisiana	1.49	0.08	(04)	30.93 <sup>c</sup>	Utah	0.43	0.07	(51)	-62.16 <sup>c</sup>
Maine	1.27	0.12	(13)	11.72	Vermont	0.89	0.16	(41)	-21.21 <sup>c</sup>
Maryland	1.35	0.07	(11)	18.57 <sup>c</sup>	Virginia	1.10	0.05	(28)	-2.93
Massachusetts	1.08	0.05	(30)	-5.12	Washington	0.92	0.05	(39)	-19.25 <sup>c</sup>
Michigan	1.26	0.05	(14)	11.20	West Virginia	1.59	0.12	(02)	39.88 <sup>c</sup>
Minnesota	0.92	0.06	(38)	-19.06 <sup>c</sup>	Wisconsin	1.01	0.06	(31)	-11.04
Mississippi	1.36	0.09	(09)	19.44 <sup>c</sup>	Wyoming	0.66	0.15	(50)	-42.02 <sup>c</sup>
Missouri	1.17	0.06	(23)	3.41					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.19  
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	2.03	0.02		
High Five States				
D.C.	3.40	0.54	(01)	67.19 <sup>c</sup>
West Virginia	2.75	0.23	(02)	35.44 <sup>c</sup>
Kentucky	2.64	0.17	(03)	30.05 <sup>c</sup>
Mississippi	2.62	0.20	(04)	29.09 <sup>c</sup>
Louisiana	2.58	0.17	(05)	26.90 <sup>c</sup>
Low Five States				
Idaho	1.44	0.20	(46)	-28.92 <sup>c</sup>
Kansas	1.37	0.14	(47)	-32.77 <sup>c</sup>
Oregon	1.26	0.12	(48)	-38.12 <sup>c</sup>
Colorado	1.07	0.11	(49)	-47.45 <sup>c</sup>
Utah	0.72	0.13	(50)	-64.79 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.56	0.15	(06)	25.94 <sup>c</sup>	Montana	1.77	0.26	(33)	-13.03
Alaska	1.79	0.46	(32)	-12.10	Nebraska	1.74	0.20	(34)	-14.41
Arizona	1.54	0.10	(42)	-24.35 <sup>c</sup>	Nevada	1.84	0.19	(31)	-9.24
Arkansas	2.05	0.17	(26)	0.81	New Hampshire	2.13	0.26	(21)	4.77
California	1.59	0.05	(41)	-21.72 <sup>c</sup>	New Jersey	2.10	0.10	(22)	3.42
Colorado	1.07	0.11	(49)	-47.45 <sup>c</sup>	New Mexico	1.69	0.19	(38)	-17.00 <sup>c</sup>
Connecticut	1.70	0.14	(37)	-16.56 <sup>c</sup>	New York	2.18	0.07	(19)	7.08
Delaware	2.25	0.32	(14)	10.82	North Carolina	2.27	0.11	(13)	11.67
D.C.	3.40	0.54	(01)	67.19 <sup>c</sup>	North Dakota	1.72	0.32	(36)	-15.52 <sup>c</sup>
Florida	2.30	0.06	(12)	13.00	Ohio	2.51	0.09	(07)	23.31 <sup>c</sup>
Georgia	2.08	0.11	(24)	2.35	Oklahoma	1.91	0.15	(29)	-5.85
Hawaii	1.72	0.22	(35)	-15.28 <sup>c</sup>	Oregon	1.26	0.12	(48)	-38.12 <sup>c</sup>
Idaho	1.44	0.20	(46)	-28.92 <sup>c</sup>	Pennsylvania	2.23	0.08	(16)	9.62
Illinois	2.21	0.09	(18)	8.53	Rhode Island	2.30	0.30	(11)	13.16
Indiana	2.04	0.12	(27)	0.34	South Carolina	2.48	0.16	(09)	21.96 <sup>c</sup>
Iowa	1.68	0.15	(39)	-17.46 <sup>c</sup>	South Dakota	1.46	0.27	(45)	-28.34 <sup>c</sup>
Kansas	1.37	0.14	(47)	-32.77 <sup>c</sup>	Tennessee	2.50	0.13	(08)	23.01 <sup>c</sup>
Kentucky	2.64	0.17	(03)	30.05 <sup>c</sup>	Texas	2.09	0.07	(23)	2.70
Louisiana	2.58	0.17	(05)	26.90 <sup>c</sup>	Utah	0.72	0.13	(50)	-64.79 <sup>c</sup>
Maine	2.22	0.25	(17)	9.38	Vermont	1.51	0.30	(44)	-25.59 <sup>c</sup>
Maryland	2.36	0.14	(10)	16.12 <sup>c</sup>	Virginia	2.07	0.11	(25)	1.68
Massachusetts	1.95	0.11	(28)	-4.02	Washington	1.54	0.10	(43)	-24.47 <sup>c</sup>
Michigan	2.24	0.10	(15)	9.96	West Virginia	2.75	0.23	(02)	35.44 <sup>c</sup>
Minnesota	1.65	0.12	(40)	-18.78 <sup>c</sup>	Wisconsin	1.88	0.12	(30)	-7.63
Mississippi	2.62	0.20	(04)	29.09 <sup>c</sup>	Wyoming	-	-	(-)	-
Missouri	2.17	0.12	(20)	6.68					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 12.20  
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.43	0.01		
High Five States				
Louisiana	0.65	0.07	(01)	50.41 <sup>c</sup>
Kentucky	0.62	0.07	(02)	43.35 <sup>c</sup>
West Virginia	0.62	0.10	(03)	41.76 <sup>c</sup>
Indiana	0.61	0.06	(04)	39.95 <sup>c</sup>
Delaware	0.60	0.15	(05)	38.73 <sup>c</sup>
Low Five States				
Connecticut	0.34	0.05	(37)	-22.39 <sup>c</sup>
Minnesota	0.33	0.05	(38)	-23.75 <sup>c</sup>
California	0.32	0.02	(39)	-27.26 <sup>c</sup>
Nebraska	0.31	0.08	(40)	-29.52 <sup>c</sup>
Kansas	0.29	0.06	(41)	-33.97 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.46	0.06	(18)	5.98	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.31	0.08	(40)	-29.52 <sup>c</sup>
Arizona	0.35	0.04	(35)	-19.35 <sup>c</sup>	Nevada	0.60	0.10	(06)	38.51 <sup>c</sup>
Arkansas	0.43	0.07	(22)	-0.40	New Hampshire	0.46	0.11	(19)	5.46
California	0.32	0.02	(39)	-27.26 <sup>c</sup>	New Jersey	0.54	0.04	(09)	24.86 <sup>c</sup>
Colorado	0.38	0.06	(31)	-12.17	New Mexico	0.37	0.08	(33)	-14.46
Connecticut	0.34	0.05	(37)	-22.39 <sup>c</sup>	New York	0.46	0.03	(16)	6.40
Delaware	0.60	0.15	(05)	38.73 <sup>c</sup>	North Carolina	0.47	0.04	(15)	8.21
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.45	0.03	(21)	4.73	Ohio	0.52	0.04	(13)	18.65 <sup>c</sup>
Georgia	0.38	0.04	(29)	-12.03	Oklahoma	0.41	0.06	(27)	-4.82
Hawaii	-	-	(-)	-	Oregon	0.36	0.06	(34)	-18.07 <sup>c</sup>
Idaho	-	-	(-)	-	Pennsylvania	0.46	0.03	(20)	5.32
Illinois	0.46	0.04	(17)	6.19	Rhode Island	0.53	0.13	(10)	22.00 <sup>c</sup>
Indiana	0.61	0.06	(04)	39.95 <sup>c</sup>	South Carolina	0.53	0.06	(11)	21.67 <sup>c</sup>
Iowa	0.43	0.07	(23)	-0.97	South Dakota	-	-	(-)	-
Kansas	0.29	0.06	(41)	-33.97 <sup>c</sup>	Tennessee	0.57	0.06	(07)	32.11 <sup>c</sup>
Kentucky	0.62	0.07	(02)	43.35 <sup>c</sup>	Texas	0.37	0.03	(32)	-13.61
Louisiana	0.65	0.07	(01)	50.41 <sup>c</sup>	Utah	-	-	(-)	-
Maine	0.52	0.11	(12)	20.36 <sup>c</sup>	Vermont	-	-	(-)	-
Maryland	0.55	0.06	(08)	27.73 <sup>c</sup>	Virginia	0.38	0.04	(28)	-11.42
Massachusetts	0.42	0.05	(25)	-2.36	Washington	0.43	0.05	(24)	-1.74
Michigan	0.50	0.04	(14)	14.30	West Virginia	0.62	0.10	(03)	41.76 <sup>c</sup>
Minnesota	0.33	0.05	(38)	-23.75 <sup>c</sup>	Wisconsin	0.35	0.05	(36)	-20.08 <sup>c</sup>
Mississippi	0.38	0.07	(30)	-12.16	Wyoming	-	-	(-)	-
Missouri	0.42	0.05	(26)	-3.54					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 13.26  
Leukemia

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	7.07	0.02		
High Five States				
Wyoming	8.24	0.55	(01)	16.52 <sup>c</sup>
Wisconsin	8.02	0.16	(02)	13.49
North Dakota	7.92	0.45	(03)	11.99
Minnesota	7.80	0.17	(04)	10.36
Oklahoma	7.76	0.20	(05)	9.75
Low Five States				
Arizona	6.50	0.14	(47)	-8.05
Nevada	6.48	0.24	(48)	-8.35
New Mexico	6.40	0.25	(49)	-9.53
D.C.	6.18	0.47	(50)	-12.65
Hawaii	5.35	0.27	(51)	-24.32 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.40	0.17	(22)	4.73	Montana	7.63	0.37	(13)	7.98
Alaska	7.68	0.64	(12)	8.64	Nebraska	6.90	0.27	(33)	-2.39
Arizona	6.50	0.14	(47)	-8.05	Nevada	6.48	0.24	(48)	-8.35
Arkansas	7.68	0.22	(11)	8.66	New Hampshire	6.95	0.32	(31)	-1.76
California	6.69	0.06	(43)	-5.31	New Jersey	6.66	0.12	(45)	-5.86
Colorado	6.73	0.18	(40)	-4.80	New Mexico	6.40	0.25	(49)	-9.53
Connecticut	6.79	0.19	(36)	-3.95	New York	6.66	0.08	(44)	-5.75
Delaware	7.00	0.38	(29)	-0.95	North Carolina	6.93	0.12	(32)	-2.03
D.C.	6.18	0.47	(50)	-12.65	North Dakota	7.92	0.45	(03)	11.99
Florida	6.71	0.08	(42)	-5.04	Ohio	7.43	0.11	(20)	5.08
Georgia	6.79	0.13	(37)	-4.00	Oklahoma	7.76	0.20	(05)	9.75
Hawaii	5.35	0.27	(51)	-24.32 <sup>c</sup>	Oregon	7.07	0.19	(27)	0.06
Idaho	7.28	0.32	(24)	2.90	Pennsylvania	7.47	0.10	(17)	5.72
Illinois	7.47	0.11	(18)	5.59	Rhode Island	7.20	0.35	(25)	1.84
Indiana	7.70	0.15	(10)	8.96	South Carolina	6.96	0.17	(30)	-1.54
Iowa	7.20	0.20	(26)	1.81	South Dakota	7.42	0.40	(21)	4.90
Kansas	7.73	0.23	(09)	9.37	Tennessee	7.55	0.15	(14)	6.76
Kentucky	7.44	0.18	(19)	5.17	Texas	6.80	0.08	(35)	-3.78
Louisiana	7.49	0.19	(16)	6.00	Utah	6.72	0.26	(41)	-4.93
Maine	6.87	0.29	(34)	-2.83	Vermont	7.04	0.45	(28)	-0.40
Maryland	6.64	0.16	(46)	-6.05	Virginia	6.78	0.13	(38)	-4.14
Massachusetts	6.73	0.14	(39)	-4.80	Washington	7.30	0.15	(23)	3.29
Michigan	7.76	0.12	(07)	9.72	West Virginia	7.51	0.26	(15)	6.16
Minnesota	7.80	0.17	(04)	10.36	Wisconsin	8.02	0.16	(02)	13.49
Mississippi	7.76	0.23	(08)	9.70	Wyoming	8.24	0.55	(01)	16.52 <sup>c</sup>
Missouri	7.76	0.16	(06)	9.72					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 13.27  
Leukemia

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	9.51	0.04		
High Five States				
North Dakota	11.29	0.82	(01)	18.77 <sup>c</sup>
Wisconsin	11.16	0.29	(02)	17.36 <sup>c</sup>
Wyoming	10.88	0.96	(03)	14.48
Alaska	10.81	1.19	(04)	13.73
Minnesota	10.80	0.31	(05)	13.56
Low Five States				
California	8.72	0.11	(47)	-8.26
New Hampshire	8.49	0.54	(48)	-10.67
Arizona	8.44	0.24	(49)	-11.27
New Mexico	8.16	0.43	(50)	-14.12
Hawaii	6.91	0.46	(51)	-27.28 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	10.27	0.33	(15)	7.97	Montana	10.03	0.64	(21)	5.49
Alaska	10.81	1.19	(04)	13.73	Nebraska	9.26	0.47	(34)	-2.66
Arizona	8.44	0.24	(49)	-11.27	Nevada	8.99	0.43	(42)	-5.45
Arkansas	10.47	0.40	(09)	10.08	New Hampshire	8.49	0.54	(48)	-10.67
California	8.72	0.11	(47)	-8.26	New Jersey	9.04	0.22	(39)	-4.94
Colorado	8.83	0.32	(45)	-7.13	New Mexico	8.16	0.43	(50)	-14.12
Connecticut	9.29	0.34	(33)	-2.25	New York	9.03	0.15	(40)	-5.07
Delaware	10.05	0.71	(20)	5.66	North Carolina	9.69	0.23	(30)	1.94
D.C.	8.83	0.90	(46)	-7.18	North Dakota	11.29	0.82	(01)	18.77 <sup>c</sup>
Florida	8.92	0.13	(44)	-6.13	Ohio	9.86	0.19	(25)	3.73
Georgia	9.40	0.25	(32)	-1.08	Oklahoma	10.67	0.37	(07)	12.27
Hawaii	6.91	0.46	(51)	-27.28 <sup>c</sup>	Oregon	9.25	0.32	(35)	-2.76
Idaho	9.76	0.55	(28)	2.63	Pennsylvania	9.94	0.18	(22)	4.57
Illinois	10.08	0.20	(18)	5.97	Rhode Island	10.07	0.64	(19)	5.94
Indiana	10.42	0.28	(12)	9.57	South Carolina	9.61	0.32	(31)	1.07
Iowa	9.74	0.36	(29)	2.42	South Dakota	9.92	0.71	(23)	4.37
Kansas	10.29	0.40	(14)	8.25	Tennessee	10.56	0.29	(08)	11.04
Kentucky	10.21	0.34	(16)	7.41	Texas	9.01	0.15	(41)	-5.28
Louisiana	10.13	0.34	(17)	6.53	Utah	9.19	0.47	(36)	-3.38
Maine	9.81	0.54	(27)	3.21	Vermont	10.32	0.84	(13)	8.51
Maryland	8.98	0.28	(43)	-5.56	Virginia	9.16	0.25	(37)	-3.63
Massachusetts	9.06	0.25	(38)	-4.74	Washington	9.88	0.27	(24)	3.88
Michigan	10.45	0.22	(10)	9.91	West Virginia	9.81	0.46	(26)	3.21
Minnesota	10.80	0.31	(05)	13.56	Wisconsin	11.16	0.29	(02)	17.36 <sup>c</sup>
Mississippi	10.43	0.43	(11)	9.74	Wyoming	10.88	0.96	(03)	14.48
Missouri	10.76	0.28	(06)	13.21					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
 SE Standard error of the rate.  
 PD Percent difference between state rate and total U.S. rate.  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 13.28  
Leukemia

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.31	0.02		
High Five States				
Wyoming	6.01	0.64	(01)	13.17
New Hampshire	5.97	0.40	(02)	12.42
Mississippi	5.95	0.27	(03)	11.96
Kansas	5.87	0.26	(04)	10.55
Indiana	5.86	0.18	(05)	10.36
Low Five States				
Delaware	4.69	0.41	(47)	-11.77
Vermont	4.57	0.49	(48)	-13.97
D.C.	4.47	0.53	(49)	-15.81 <sup>c</sup>
Nevada	4.46	0.27	(50)	-16.02 <sup>c</sup>
Hawaii	4.09	0.32	(51)	-23.04 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.49	0.20	(20)	3.37	Montana	5.81	0.45	(06)	9.41
Alaska	5.25	0.71	(28)	-1.19	Nebraska	5.12	0.31	(33)	-3.71
Arizona	4.97	0.17	(44)	-6.37	Nevada	4.46	0.27	(50)	-16.02 <sup>c</sup>
Arkansas	5.73	0.25	(09)	7.84	New Hampshire	5.97	0.40	(02)	12.42
California	5.17	0.07	(31)	-2.69	New Jersey	5.03	0.14	(38)	-5.25
Colorado	5.15	0.21	(32)	-3.02	New Mexico	5.01	0.30	(41)	-5.79
Connecticut	5.05	0.21	(37)	-5.04	New York	5.02	0.09	(40)	-5.60
Delaware	4.69	0.41	(47)	-11.77	North Carolina	5.06	0.14	(36)	-4.67
D.C.	4.47	0.53	(49)	-15.81 <sup>c</sup>	North Dakota	5.47	0.51	(21)	2.98
Florida	4.98	0.09	(43)	-6.21	Ohio	5.71	0.13	(12)	7.54
Georgia	5.03	0.15	(39)	-5.37	Oklahoma	5.64	0.23	(16)	6.09
Hawaii	4.09	0.32	(51)	-23.04 <sup>c</sup>	Oregon	5.45	0.22	(23)	2.55
Idaho	5.26	0.36	(27)	-1.08	Pennsylvania	5.72	0.12	(10)	7.71
Illinois	5.69	0.12	(13)	7.14	Rhode Island	5.32	0.40	(26)	0.20
Indiana	5.86	0.18	(05)	10.36	South Carolina	5.07	0.20	(35)	-4.60
Iowa	5.46	0.24	(22)	2.74	South Dakota	5.50	0.46	(18)	3.46
Kansas	5.87	0.26	(04)	10.55	Tennessee	5.49	0.17	(19)	3.42
Kentucky	5.42	0.21	(24)	1.94	Texas	5.19	0.09	(30)	-2.36
Louisiana	5.67	0.21	(14)	6.75	Utah	4.82	0.30	(45)	-9.20
Maine	4.73	0.33	(46)	-11.06	Vermont	4.57	0.49	(48)	-13.97
Maryland	5.00	0.18	(42)	-5.88	Virginia	5.07	0.15	(34)	-4.60
Massachusetts	5.23	0.16	(29)	-1.55	Washington	5.39	0.18	(25)	1.51
Michigan	5.77	0.14	(08)	8.63	West Virginia	5.79	0.31	(07)	8.97
Minnesota	5.66	0.19	(15)	6.50	Wisconsin	5.72	0.18	(11)	7.66
Mississippi	5.95	0.27	(03)	11.96	Wyoming	6.01	0.64	(01)	13.17
Missouri	5.61	0.18	(17)	5.54					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.18  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.60	0.02		
High Five States				
D.C.	8.11	0.53	(01)	44.94 <sup>c</sup>
Texas	7.27	0.08	(02)	29.94 <sup>c</sup>
Louisiana	7.18	0.18	(03)	28.36 <sup>c</sup>
Hawaii	7.02	0.30	(04)	25.52 <sup>c</sup>
California	6.86	0.06	(05)	22.66 <sup>c</sup>
Low Five States				
South Dakota	4.17	0.30	(47)	-25.44 <sup>c</sup>
Montana	4.10	0.27	(48)	-26.66 <sup>c</sup>
Idaho	3.85	0.23	(49)	-31.21 <sup>c</sup>
North Dakota	3.73	0.31	(50)	-33.37 <sup>c</sup>
Utah	3.72	0.19	(51)	-33.56 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.76	0.15	(13)	2.88	Montana	4.10	0.27	(48)	-26.66 <sup>c</sup>
Alaska	5.26	0.48	(26)	-6.05	Nebraska	4.19	0.21	(46)	-25.10 <sup>c</sup>
Arizona	5.21	0.13	(27)	-6.93	Nevada	5.94	0.22	(10)	6.09
Arkansas	5.91	0.19	(11)	5.70	New Hampshire	4.48	0.25	(43)	-19.95 <sup>c</sup>
California	6.86	0.06	(05)	22.66 <sup>c</sup>	New Jersey	5.26	0.10	(25)	-6.00
Colorado	4.73	0.15	(40)	-15.45 <sup>c</sup>	New Mexico	6.35	0.24	(07)	13.48
Connecticut	4.92	0.16	(36)	-12.03	New York	5.71	0.07	(14)	2.10
Delaware	5.68	0.34	(15)	1.45	North Carolina	5.04	0.10	(32)	-9.89
D.C.	8.11	0.53	(01)	44.94 <sup>c</sup>	North Dakota	3.73	0.31	(50)	-33.37 <sup>c</sup>
Florida	5.43	0.07	(18)	-2.96	Ohio	4.88	0.09	(37)	-12.83
Georgia	5.11	0.11	(31)	-8.68	Oklahoma	5.39	0.16	(20)	-3.67
Hawaii	7.02	0.30	(04)	25.52 <sup>c</sup>	Oregon	5.31	0.16	(23)	-5.08
Idaho	3.85	0.23	(49)	-31.21 <sup>c</sup>	Pennsylvania	5.16	0.08	(28)	-7.74
Illinois	5.35	0.09	(22)	-4.42	Rhode Island	5.77	0.31	(12)	3.08
Indiana	4.64	0.12	(41)	-17.06 <sup>c</sup>	South Carolina	5.40	0.15	(19)	-3.43
Iowa	4.30	0.16	(45)	-23.18 <sup>c</sup>	South Dakota	4.17	0.30	(47)	-25.44 <sup>c</sup>
Kansas	4.45	0.17	(44)	-20.52 <sup>c</sup>	Tennessee	5.49	0.13	(17)	-1.79
Kentucky	4.98	0.15	(34)	-10.93	Texas	7.27	0.08	(02)	29.94 <sup>c</sup>
Louisiana	7.18	0.18	(03)	28.36 <sup>c</sup>	Utah	3.72	0.19	(51)	-33.56 <sup>c</sup>
Maine	4.78	0.24	(39)	-14.60	Vermont	5.15	0.38	(29)	-7.99
Maryland	5.50	0.14	(16)	-1.72	Virginia	5.13	0.11	(30)	-8.25
Massachusetts	5.97	0.13	(08)	6.78	Washington	5.96	0.13	(09)	6.60
Michigan	5.27	0.10	(24)	-5.76	West Virginia	4.62	0.20	(42)	-17.48 <sup>c</sup>
Minnesota	4.78	0.13	(38)	-14.56	Wisconsin	4.95	0.13	(35)	-11.52
Mississippi	6.70	0.21	(06)	19.81 <sup>c</sup>	Wyoming	5.01	0.43	(33)	-10.51
Missouri	5.35	0.13	(21)	-4.40					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.19  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	8.27	0.03		
High Five States				
D.C.	13.44	1.05	(01)	62.47 <sup>c</sup>
Louisiana	10.96	0.34	(02)	32.47 <sup>c</sup>
Texas	10.72	0.15	(03)	29.57 <sup>c</sup>
Hawaii	10.34	0.54	(04)	25.00 <sup>c</sup>
Mississippi	10.16	0.41	(05)	22.82 <sup>c</sup>
Low Five States				
Idaho	5.45	0.40	(47)	-34.15 <sup>c</sup>
North Dakota	5.35	0.55	(48)	-35.37 <sup>c</sup>
Montana	5.24	0.44	(49)	-36.72 <sup>c</sup>
Utah	5.12	0.33	(50)	-38.10 <sup>c</sup>
South Dakota	5.07	0.49	(51)	-38.73 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	8.69	0.28	(12)	5.01	Montana	5.24	0.44	(49)	-36.72 <sup>c</sup>
Alaska	7.59	0.86	(29)	-8.32	Nebraska	6.16	0.38	(45)	-25.50 <sup>c</sup>
Arizona	7.27	0.22	(35)	-12.13	Nevada	8.37	0.38	(16)	1.16
Arkansas	8.57	0.35	(14)	3.59	New Hampshire	6.41	0.45	(43)	-22.58 <sup>c</sup>
California	9.94	0.11	(06)	20.17 <sup>c</sup>	New Jersey	7.73	0.19	(24)	-6.54
Colorado	6.86	0.26	(41)	-17.08 <sup>c</sup>	New Mexico	9.36	0.44	(08)	13.12
Connecticut	7.55	0.29	(30)	-8.80	New York	8.71	0.14	(11)	5.24
Delaware	9.10	0.64	(09)	9.98	North Carolina	7.65	0.19	(27)	-7.52
D.C.	13.44	1.05	(01)	62.47 <sup>c</sup>	North Dakota	5.35	0.55	(48)	-35.37 <sup>c</sup>
Florida	8.19	0.12	(18)	-0.98	Ohio	7.29	0.16	(33)	-11.85
Georgia	7.49	0.21	(31)	-9.50	Oklahoma	7.69	0.29	(25)	-7.08
Hawaii	10.34	0.54	(04)	25.00 <sup>c</sup>	Oregon	7.68	0.28	(26)	-7.25
Idaho	5.45	0.40	(47)	-34.15 <sup>c</sup>	Pennsylvania	7.76	0.15	(22)	-6.24
Illinois	7.84	0.17	(21)	-5.25	Rhode Island	8.93	0.58	(10)	7.88
Indiana	6.95	0.22	(40)	-16.03 <sup>c</sup>	South Carolina	8.17	0.28	(19)	-1.32
Iowa	6.15	0.28	(46)	-25.62 <sup>c</sup>	South Dakota	5.07	0.49	(51)	-38.73 <sup>c</sup>
Kansas	6.39	0.30	(44)	-22.75 <sup>c</sup>	Tennessee	8.46	0.24	(15)	2.26
Kentucky	7.29	0.27	(34)	-11.89	Texas	10.72	0.15	(03)	29.57 <sup>c</sup>
Louisiana	10.96	0.34	(02)	32.47 <sup>c</sup>	Utah	5.12	0.33	(50)	-38.10 <sup>c</sup>
Maine	7.02	0.44	(39)	-15.11 <sup>c</sup>	Vermont	7.33	0.68	(32)	-11.46
Maryland	8.33	0.26	(17)	0.69	Virginia	7.93	0.22	(20)	-4.14
Massachusetts	9.50	0.25	(07)	14.78	Washington	8.66	0.24	(13)	4.66
Michigan	7.60	0.18	(28)	-8.19	West Virginia	6.41	0.36	(42)	-22.51 <sup>c</sup>
Minnesota	7.03	0.24	(38)	-15.02 <sup>c</sup>	Wisconsin	7.04	0.23	(37)	-14.97
Mississippi	10.16	0.41	(05)	22.82 <sup>c</sup>	Wyoming	7.04	0.75	(36)	-14.88
Missouri	7.75	0.23	(23)	-6.33					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 14.20  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.36	0.02		
High Five States				
Texas	4.36	0.09	(01)	29.80 <sup>c</sup>
California	4.23	0.07	(02)	26.04 <sup>c</sup>
Louisiana	4.11	0.18	(03)	22.52 <sup>c</sup>
Mississippi	4.08	0.22	(04)	21.48 <sup>c</sup>
Hawaii	3.99	0.31	(05)	18.81 <sup>c</sup>
Low Five States				
Iowa	2.69	0.16	(47)	-19.96 <sup>c</sup>
Nebraska	2.56	0.22	(48)	-23.91 <sup>c</sup>
Utah	2.49	0.22	(49)	-25.97 <sup>c</sup>
Idaho	2.44	0.25	(50)	-27.22 <sup>c</sup>
North Dakota	2.26	0.32	(51)	-32.77 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.45	0.15	(11)	2.60	Montana	2.97	0.32	(35)	-11.50
Alaska	3.01	0.52	(34)	-10.47	Nebraska	2.56	0.22	(48)	-23.91 <sup>c</sup>
Arizona	3.38	0.14	(13)	0.64	Nevada	3.68	0.24	(09)	9.54
Arkansas	3.73	0.20	(07)	11.00	New Hampshire	2.79	0.27	(45)	-16.79 <sup>c</sup>
California	4.23	0.07	(02)	26.04 <sup>c</sup>	New Jersey	3.29	0.11	(20)	-2.07
Colorado	2.89	0.16	(39)	-13.85	New Mexico	3.72	0.26	(08)	10.83
Connecticut	2.80	0.16	(42)	-16.50 <sup>c</sup>	New York	3.33	0.07	(17)	-0.97
Delaware	2.80	0.32	(44)	-16.76 <sup>c</sup>	North Carolina	2.90	0.10	(38)	-13.68
D.C.	3.96	0.50	(06)	18.03 <sup>c</sup>	North Dakota	2.26	0.32	(51)	-32.77 <sup>c</sup>
Florida	3.05	0.07	(32)	-9.12	Ohio	2.91	0.09	(37)	-13.25
Georgia	3.19	0.12	(25)	-5.11	Oklahoma	3.42	0.18	(12)	1.76
Hawaii	3.99	0.31	(05)	18.81 <sup>c</sup>	Oregon	3.20	0.17	(24)	-4.85
Idaho	2.44	0.25	(50)	-27.22 <sup>c</sup>	Pennsylvania	3.05	0.08	(33)	-9.24
Illinois	3.35	0.10	(14)	-0.21	Rhode Island	3.33	0.31	(16)	-0.93
Indiana	2.80	0.12	(43)	-16.57 <sup>c</sup>	South Carolina	3.13	0.15	(28)	-6.94
Iowa	2.69	0.16	(47)	-19.96 <sup>c</sup>	South Dakota	3.28	0.37	(21)	-2.31
Kansas	2.71	0.18	(46)	-19.30 <sup>c</sup>	Tennessee	3.10	0.13	(30)	-7.65
Kentucky	3.12	0.16	(29)	-7.13	Texas	4.36	0.09	(01)	29.80 <sup>c</sup>
Louisiana	4.11	0.18	(03)	22.52 <sup>c</sup>	Utah	2.49	0.22	(49)	-25.97 <sup>c</sup>
Maine	2.95	0.25	(36)	-12.23	Vermont	3.32	0.43	(19)	-1.27
Maryland	3.21	0.14	(22)	-4.34	Virginia	2.89	0.12	(40)	-14.05
Massachusetts	3.20	0.13	(23)	-4.81	Washington	3.58	0.14	(10)	6.65
Michigan	3.34	0.10	(15)	-0.54	West Virginia	3.09	0.22	(31)	-7.99
Minnesota	2.85	0.14	(41)	-15.14 <sup>c</sup>	Wisconsin	3.14	0.14	(27)	-6.54
Mississippi	4.08	0.22	(04)	21.48 <sup>c</sup>	Wyoming	3.18	0.47	(26)	-5.35
Missouri	3.32	0.13	(18)	-1.22					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
 SE Standard error of the rate.  
 PD Percent difference between state rate and total U.S. rate.  
 - Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 15.24  
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	49.52	0.06		
High Five States				
Kentucky	73.17	0.57	(01)	47.76 <sup>c</sup>
West Virginia	65.22	0.75	(02)	31.71 <sup>c</sup>
Arkansas	64.60	0.63	(03)	30.47 <sup>c</sup>
Tennessee	64.50	0.44	(04)	30.26 <sup>c</sup>
Mississippi	64.29	0.65	(05)	29.84 <sup>c</sup>
Low Five States				
California	38.69	0.15	(47)	-21.87 <sup>c</sup>
Colorado	36.73	0.42	(48)	-25.82 <sup>c</sup>
Hawaii	35.96	0.69	(49)	-27.38 <sup>c</sup>
New Mexico	35.11	0.58	(50)	-29.09 <sup>c</sup>
Utah	21.60	0.47	(51)	-56.39 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	60.35	0.48	(08)	21.87 <sup>c</sup>	Montana	45.07	0.90	(38)	-8.99 <sup>c</sup>
Alaska	52.76	1.63	(18)	6.55 <sup>c</sup>	Nebraska	46.20	0.69	(35)	-6.70 <sup>c</sup>
Arizona	40.57	0.35	(46)	-18.06 <sup>c</sup>	Nevada	52.39	0.67	(20)	5.80 <sup>c</sup>
Arkansas	64.60	0.63	(03)	30.47 <sup>c</sup>	New Hampshire	49.78	0.84	(26)	0.54
California	38.69	0.15	(47)	-21.87 <sup>c</sup>	New Jersey	44.79	0.31	(39)	-9.55 <sup>c</sup>
Colorado	36.73	0.42	(48)	-25.82 <sup>c</sup>	New Mexico	35.11	0.58	(50)	-29.09 <sup>c</sup>
Connecticut	44.17	0.47	(40)	-10.79 <sup>c</sup>	New York	43.52	0.20	(42)	-12.12 <sup>c</sup>
Delaware	56.42	1.07	(12)	13.95 <sup>c</sup>	North Carolina	55.68	0.34	(14)	12.45 <sup>c</sup>
D.C.	46.14	1.30	(36)	-6.83 <sup>c</sup>	North Dakota	41.53	1.05	(44)	-16.12 <sup>c</sup>
Florida	48.84	0.20	(31)	-1.38	Ohio	57.08	0.30	(11)	15.28 <sup>c</sup>
Georgia	52.92	0.36	(17)	6.88 <sup>c</sup>	Oklahoma	60.76	0.55	(06)	22.70 <sup>c</sup>
Hawaii	35.96	0.69	(49)	-27.38 <sup>c</sup>	Oregon	49.53	0.49	(29)	0.03
Idaho	41.01	0.75	(45)	-17.17 <sup>c</sup>	Pennsylvania	50.53	0.26	(25)	2.04
Illinois	51.79	0.28	(21)	4.59	Rhode Island	51.39	0.93	(22)	3.79
Indiana	60.46	0.43	(07)	22.10 <sup>c</sup>	South Carolina	55.56	0.48	(15)	12.20 <sup>c</sup>
Iowa	49.33	0.53	(30)	-0.39	South Dakota	45.98	1.01	(37)	-7.14 <sup>c</sup>
Kansas	51.23	0.59	(24)	3.45	Tennessee	64.50	0.44	(04)	30.26 <sup>c</sup>
Kentucky	73.17	0.57	(01)	47.76 <sup>c</sup>	Texas	46.22	0.21	(34)	-6.65 <sup>c</sup>
Louisiana	59.40	0.52	(09)	19.96 <sup>c</sup>	Utah	21.60	0.47	(51)	-56.39 <sup>c</sup>
Maine	56.11	0.83	(13)	13.32 <sup>c</sup>	Vermont	52.50	1.22	(19)	6.03 <sup>c</sup>
Maryland	49.56	0.42	(28)	0.08	Virginia	51.39	0.37	(23)	3.79
Massachusetts	49.66	0.37	(27)	0.29	Washington	48.46	0.39	(32)	-2.13
Michigan	54.12	0.32	(16)	9.29 <sup>c</sup>	West Virginia	65.22	0.75	(02)	31.71 <sup>c</sup>
Minnesota	43.86	0.41	(41)	-11.43 <sup>c</sup>	Wisconsin	46.89	0.39	(33)	-5.30 <sup>c</sup>
Mississippi	64.29	0.65	(05)	29.84 <sup>c</sup>	Wyoming	42.38	1.25	(43)	-14.41 <sup>c</sup>
Missouri	59.01	0.43	(10)	19.17 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 15.25  
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	63.48	0.10		
High Five States				
Kentucky	97.05	1.01	(01)	52.88 <sup>c</sup>
Mississippi	95.38	1.25	(02)	50.24 <sup>c</sup>
Arkansas	90.13	1.14	(03)	41.98 <sup>c</sup>
Tennessee	89.45	0.81	(04)	40.90 <sup>c</sup>
Alabama	87.37	0.91	(05)	37.63 <sup>c</sup>
Low Five States				
Hawaii	48.76	1.21	(47)	-23.20 <sup>c</sup>
California	47.22	0.26	(48)	-25.62 <sup>c</sup>
Colorado	44.20	0.70	(49)	-30.37 <sup>c</sup>
New Mexico	43.36	0.98	(50)	-31.70 <sup>c</sup>
Utah	27.54	0.80	(51)	-56.62 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	87.37	0.91	(05)	37.63 <sup>c</sup>	Montana	52.07	1.43	(43)	-17.98 <sup>c</sup>
Alaska	61.78	2.70	(27)	-2.68	Nebraska	60.33	1.20	(32)	-4.97
Arizona	49.82	0.58	(45)	-21.52 <sup>c</sup>	Nevada	59.66	1.06	(33)	-6.02 <sup>c</sup>
Arkansas	90.13	1.14	(03)	41.98 <sup>c</sup>	New Hampshire	59.00	1.40	(34)	-7.06 <sup>c</sup>
California	47.22	0.26	(48)	-25.62 <sup>c</sup>	New Jersey	55.78	0.53	(38)	-12.13 <sup>c</sup>
Colorado	44.20	0.70	(49)	-30.37 <sup>c</sup>	New Mexico	43.36	0.98	(50)	-31.70 <sup>c</sup>
Connecticut	52.72	0.79	(42)	-16.96 <sup>c</sup>	New York	54.31	0.35	(39)	-14.46 <sup>c</sup>
Delaware	68.19	1.80	(17)	7.42 <sup>c</sup>	North Carolina	76.61	0.63	(11)	20.68 <sup>c</sup>
D.C.	61.89	2.37	(26)	-2.51	North Dakota	54.14	1.80	(40)	-14.72 <sup>c</sup>
Florida	61.38	0.34	(29)	-3.32	Ohio	74.81	0.52	(13)	17.84 <sup>c</sup>
Georgia	73.12	0.68	(14)	15.18 <sup>c</sup>	Oklahoma	79.55	0.97	(08)	25.31 <sup>c</sup>
Hawaii	48.76	1.21	(47)	-23.20 <sup>c</sup>	Oregon	58.44	0.81	(35)	-7.94 <sup>c</sup>
Idaho	49.16	1.22	(46)	-22.56 <sup>c</sup>	Pennsylvania	65.83	0.45	(21)	3.70
Illinois	65.83	0.50	(20)	3.70	Rhode Island	64.92	1.61	(22)	2.26
Indiana	79.39	0.75	(09)	25.06 <sup>c</sup>	South Carolina	77.08	0.88	(10)	21.41 <sup>c</sup>
Iowa	64.06	0.91	(23)	0.90	South Dakota	61.91	1.76	(25)	-2.48
Kansas	66.77	1.02	(19)	5.18 <sup>c</sup>	Tennessee	89.45	0.81	(04)	40.90 <sup>c</sup>
Kentucky	97.05	1.01	(01)	52.88 <sup>c</sup>	Texas	60.71	0.38	(30)	-4.36
Louisiana	80.90	0.94	(07)	27.43 <sup>c</sup>	Utah	27.54	0.80	(51)	-56.62 <sup>c</sup>
Maine	69.94	1.41	(15)	10.16 <sup>c</sup>	Vermont	62.33	2.02	(24)	-1.81
Maryland	61.74	0.73	(28)	-2.75	Virginia	67.06	0.65	(18)	5.63 <sup>c</sup>
Massachusetts	60.61	0.64	(31)	-4.52	Washington	56.95	0.64	(37)	-10.29 <sup>c</sup>
Michigan	68.21	0.55	(16)	7.44 <sup>c</sup>	West Virginia	84.35	1.30	(06)	32.87 <sup>c</sup>
Minnesota	53.28	0.68	(41)	-16.07 <sup>c</sup>	Wisconsin	58.39	0.66	(36)	-8.03 <sup>c</sup>
Mississippi	95.38	1.25	(02)	50.24 <sup>c</sup>	Wyoming	49.90	2.03	(44)	-21.40 <sup>c</sup>
Missouri	76.49	0.74	(12)	20.48 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 15.26  
Cancer of the Lung And Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	39.18	0.07		
High Five States				
Kentucky	55.80	0.66	(01)	42.43 <sup>c</sup>
West Virginia	50.89	0.90	(02)	29.90 <sup>c</sup>
Delaware	47.66	1.32	(03)	21.66 <sup>c</sup>
Indiana	46.72	0.50	(04)	19.26 <sup>c</sup>
Oklahoma	46.57	0.65	(05)	18.88 <sup>c</sup>
Low Five States				
North Dakota	32.24	1.26	(47)	-17.72 <sup>c</sup>
Colorado	31.31	0.52	(48)	-20.07 <sup>c</sup>
New Mexico	28.57	0.71	(49)	-27.08 <sup>c</sup>
Hawaii	25.90	0.79	(50)	-33.89 <sup>c</sup>
Utah	16.76	0.56	(51)	-57.23 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	40.76	0.53	(24)	4.03	Montana	39.57	1.15	(29)	1.01
Alaska	45.84	2.07	(10)	17.01 <sup>c</sup>	Nebraska	35.47	0.82	(40)	-9.47 <sup>c</sup>
Arizona	33.19	0.43	(45)	-15.28 <sup>c</sup>	Nevada	46.48	0.86	(07)	18.63 <sup>c</sup>
Arkansas	45.36	0.71	(11)	15.79 <sup>c</sup>	New Hampshire	43.27	1.06	(16)	10.44 <sup>c</sup>
California	32.32	0.19	(46)	-17.50 <sup>c</sup>	New Jersey	37.25	0.37	(36)	-4.92
Colorado	31.31	0.52	(48)	-20.07 <sup>c</sup>	New Mexico	28.57	0.71	(49)	-27.08 <sup>c</sup>
Connecticut	38.23	0.59	(35)	-2.42	New York	36.06	0.25	(39)	-7.95 <sup>c</sup>
Delaware	47.66	1.32	(03)	21.66 <sup>c</sup>	North Carolina	40.67	0.39	(25)	3.82
D.C.	35.00	1.48	(42)	-10.66 <sup>c</sup>	North Dakota	32.24	1.26	(47)	-17.72 <sup>c</sup>
Florida	38.77	0.24	(31)	-1.04	Ohio	44.20	0.35	(13)	12.83 <sup>c</sup>
Georgia	38.72	0.41	(32)	-1.16	Oklahoma	46.57	0.65	(05)	18.88 <sup>c</sup>
Hawaii	25.90	0.79	(50)	-33.89 <sup>c</sup>	Oregon	42.75	0.62	(17)	9.12 <sup>c</sup>
Idaho	34.45	0.94	(43)	-12.06 <sup>c</sup>	Pennsylvania	39.59	0.30	(28)	1.05
Illinois	41.92	0.34	(22)	7.00 <sup>c</sup>	Rhode Island	42.56	1.13	(18)	8.62 <sup>c</sup>
Indiana	46.72	0.50	(04)	19.26 <sup>c</sup>	South Carolina	39.88	0.54	(27)	1.78
Iowa	38.34	0.63	(34)	-2.13	South Dakota	34.14	1.19	(44)	-12.87 <sup>c</sup>
Kansas	39.42	0.70	(30)	0.63	Tennessee	46.57	0.50	(06)	18.88 <sup>c</sup>
Kentucky	55.80	0.66	(01)	42.43 <sup>c</sup>	Texas	35.38	0.25	(41)	-9.68 <sup>c</sup>
Louisiana	43.63	0.59	(15)	11.37 <sup>c</sup>	Utah	16.76	0.56	(51)	-57.23 <sup>c</sup>
Maine	46.03	1.01	(09)	17.48 <sup>c</sup>	Vermont	45.09	1.52	(12)	15.08 <sup>c</sup>
Maryland	40.79	0.51	(23)	4.13	Virginia	40.02	0.43	(26)	2.15
Massachusetts	42.14	0.46	(20)	7.55 <sup>c</sup>	Washington	42.14	0.49	(19)	7.56 <sup>c</sup>
Michigan	43.87	0.38	(14)	11.98 <sup>c</sup>	West Virginia	50.89	0.90	(02)	29.90 <sup>c</sup>
Minnesota	37.01	0.50	(37)	-5.53 <sup>c</sup>	Wisconsin	38.40	0.48	(33)	-2.00
Mississippi	42.00	0.70	(21)	7.19 <sup>c</sup>	Wyoming	36.38	1.58	(38)	-7.13 <sup>c</sup>
Missouri	46.17	0.51	(08)	17.86 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 16.18  
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.73	0.01		
High Five States				
Utah	3.52	0.19	(01)	28.73 <sup>c</sup>
Colorado	3.49	0.13	(02)	27.88 <sup>c</sup>
Oklahoma	3.34	0.13	(03)	22.24 <sup>c</sup>
Vermont	3.30	0.31	(04)	20.84 <sup>c</sup>
Kentucky	3.28	0.12	(05)	19.96 <sup>c</sup>
Low Five States				
Louisiana	2.22	0.10	(47)	-18.70 <sup>c</sup>
New York	2.21	0.05	(48)	-19.27 <sup>c</sup>
North Dakota	2.10	0.24	(49)	-22.99 <sup>c</sup>
Hawaii	1.53	0.14	(50)	-43.86 <sup>c</sup>
D.C.	1.28	0.21	(51)	-53.19 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.92	0.11	(22)	7.01	Montana	2.72	0.22	(29)	-0.29
Alaska	2.28	0.31	(46)	-16.74 <sup>c</sup>	Nebraska	2.95	0.18	(20)	7.93
Arizona	2.95	0.10	(19)	8.08	Nevada	2.75	0.15	(27)	0.53
Arkansas	2.68	0.13	(32)	-1.89	New Hampshire	2.77	0.20	(26)	1.40
California	2.60	0.04	(38)	-4.98	New Jersey	2.63	0.07	(34)	-3.69
Colorado	3.49	0.13	(02)	27.88 <sup>c</sup>	New Mexico	2.89	0.17	(24)	5.90
Connecticut	2.51	0.11	(40)	-8.33	New York	2.21	0.05	(48)	-19.27 <sup>c</sup>
Delaware	2.74	0.24	(28)	0.29	North Carolina	3.03	0.08	(14)	10.70
D.C.	1.28	0.21	(51)	-53.19 <sup>c</sup>	North Dakota	2.10	0.24	(49)	-22.99 <sup>c</sup>
Florida	2.93	0.05	(21)	7.29	Ohio	2.90	0.07	(23)	6.13
Georgia	2.47	0.08	(42)	-9.78	Oklahoma	3.34	0.13	(03)	22.24 <sup>c</sup>
Hawaii	1.53	0.14	(50)	-43.86 <sup>c</sup>	Oregon	3.17	0.12	(09)	16.13 <sup>c</sup>
Idaho	3.23	0.21	(07)	18.25 <sup>c</sup>	Pennsylvania	2.87	0.06	(25)	4.99
Illinois	2.49	0.06	(41)	-9.01	Rhode Island	2.61	0.21	(35)	-4.33
Indiana	3.05	0.10	(11)	11.53	South Carolina	2.61	0.11	(36)	-4.53
Iowa	2.72	0.13	(30)	-0.53	South Dakota	2.60	0.24	(37)	-4.85
Kansas	3.03	0.14	(13)	11.06	Tennessee	3.01	0.10	(15)	10.05
Kentucky	3.28	0.12	(05)	19.96 <sup>c</sup>	Texas	2.54	0.05	(39)	-6.92
Louisiana	2.22	0.10	(47)	-18.70 <sup>c</sup>	Utah	3.52	0.19	(01)	28.73 <sup>c</sup>
Maine	3.04	0.20	(12)	11.11	Vermont	3.30	0.31	(04)	20.84 <sup>c</sup>
Maryland	2.69	0.10	(31)	-1.55	Virginia	3.00	0.09	(17)	9.84
Massachusetts	2.97	0.09	(18)	8.57	Washington	3.00	0.10	(16)	9.92
Michigan	2.41	0.07	(43)	-11.99	West Virginia	3.19	0.17	(08)	16.69 <sup>c</sup>
Minnesota	2.39	0.09	(44)	-12.41	Wisconsin	2.65	0.09	(33)	-3.07
Mississippi	2.36	0.13	(45)	-13.73	Wyoming	3.26	0.35	(06)	19.12 <sup>c</sup>
Missouri	3.11	0.10	(10)	13.88					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 16.19  
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	4.09	0.02		
High Five States				
Utah	5.42	0.35	(01)	32.50 <sup>c</sup>
Wyoming	5.11	0.65	(02)	24.97 <sup>c</sup>
Colorado	5.11	0.23	(03)	24.82 <sup>c</sup>
Vermont	5.01	0.58	(04)	22.41 <sup>c</sup>
Oklahoma	4.89	0.24	(05)	19.47 <sup>c</sup>
Low Five States				
North Dakota	3.28	0.44	(47)	-19.76 <sup>c</sup>
New York	3.28	0.09	(48)	-19.92 <sup>c</sup>
Alaska	2.84	0.52	(49)	-30.57 <sup>c</sup>
Hawaii	2.26	0.26	(50)	-44.88 <sup>c</sup>
D.C.	1.69	0.38	(51)	-58.66 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.61	0.21	(12)	12.69	Montana	3.43	0.36	(45)	-16.06 <sup>c</sup>
Alaska	2.84	0.52	(49)	-30.57 <sup>c</sup>	Nebraska	4.00	0.31	(31)	-2.29
Arizona	4.46	0.18	(17)	8.93	Nevada	4.28	0.28	(23)	4.52
Arkansas	4.04	0.25	(30)	-1.31	New Hampshire	4.08	0.36	(29)	-0.29
California	3.88	0.07	(37)	-5.28	New Jersey	3.84	0.14	(39)	-6.07
Colorado	5.11	0.23	(03)	24.82 <sup>c</sup>	New Mexico	4.24	0.31	(26)	3.71
Connecticut	3.62	0.21	(42)	-11.48	New York	3.28	0.09	(48)	-19.92 <sup>c</sup>
Delaware	4.31	0.46	(20)	5.32	North Carolina	4.57	0.16	(13)	11.68
D.C.	1.69	0.38	(51)	-58.66 <sup>c</sup>	North Dakota	3.28	0.44	(47)	-19.76 <sup>c</sup>
Florida	4.47	0.09	(16)	9.26	Ohio	4.38	0.13	(18)	6.96
Georgia	3.97	0.16	(32)	-3.02	Oklahoma	4.89	0.24	(05)	19.47 <sup>c</sup>
Hawaii	2.26	0.26	(50)	-44.88 <sup>c</sup>	Oregon	4.25	0.22	(25)	3.82
Idaho	4.82	0.38	(07)	17.76 <sup>c</sup>	Pennsylvania	4.24	0.11	(27)	3.53
Illinois	3.68	0.12	(40)	-10.13	Rhode Island	3.95	0.39	(34)	-3.55
Indiana	4.63	0.18	(10)	13.26	South Carolina	4.22	0.21	(28)	3.08
Iowa	3.96	0.23	(33)	-3.21	South Dakota	3.64	0.43	(41)	-10.98
Kansas	4.48	0.26	(15)	9.47	Tennessee	4.74	0.19	(08)	15.94 <sup>c</sup>
Kentucky	4.85	0.23	(06)	18.50 <sup>c</sup>	Texas	3.86	0.09	(38)	-5.71
Louisiana	3.40	0.20	(46)	-17.01 <sup>c</sup>	Utah	5.42	0.35	(01)	32.50 <sup>c</sup>
Maine	4.48	0.36	(14)	9.49	Vermont	5.01	0.58	(04)	22.41 <sup>c</sup>
Maryland	4.27	0.19	(24)	4.34	Virginia	4.31	0.16	(21)	5.24
Massachusetts	4.32	0.17	(19)	5.51	Washington	4.28	0.17	(22)	4.71
Michigan	3.55	0.12	(44)	-13.33	West Virginia	4.62	0.31	(11)	12.89
Minnesota	3.55	0.17	(43)	-13.24	Wisconsin	3.93	0.17	(35)	-3.86
Mississippi	3.88	0.26	(36)	-5.26	Wyoming	5.11	0.65	(02)	24.97 <sup>c</sup>
Missouri	4.67	0.18	(09)	14.07					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 16.20  
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.71	0.01		
High Five States				
Oregon	2.33	0.15	(01)	36.34 <sup>c</sup>
Colorado	2.22	0.13	(02)	29.36 <sup>c</sup>
Nebraska	2.16	0.21	(03)	25.95 <sup>c</sup>
Oklahoma	2.15	0.14	(04)	25.45 <sup>c</sup>
West Virginia	2.11	0.19	(05)	23.42 <sup>c</sup>
Low Five States				
Louisiana	1.39	0.11	(47)	-18.84 <sup>c</sup>
Mississippi	1.32	0.13	(48)	-22.91 <sup>c</sup>
North Dakota	1.09	0.22	(49)	-36.39 <sup>c</sup>
D.C.	1.03	0.25	(50)	-39.90 <sup>c</sup>
Hawaii	0.90	0.15	(51)	-47.67 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.70	0.11	(32)	-0.48	Montana	2.11	0.28	(06)	23.22 <sup>c</sup>
Alaska	1.73	0.37	(28)	1.30	Nebraska	2.16	0.21	(03)	25.95 <sup>c</sup>
Arizona	1.71	0.10	(31)	-0.28	Nevada	1.43	0.15	(46)	-16.50 <sup>c</sup>
Arkansas	1.63	0.14	(36)	-4.81	New Hampshire	1.71	0.21	(29)	-0.06
California	1.61	0.04	(37)	-6.03	New Jersey	1.74	0.08	(26)	1.43
Colorado	2.22	0.13	(02)	29.36 <sup>c</sup>	New Mexico	1.82	0.18	(21)	6.25
Connecticut	1.74	0.12	(27)	1.40	New York	1.44	0.05	(44)	-15.84 <sup>c</sup>
Delaware	1.58	0.24	(39)	-7.86	North Carolina	1.97	0.09	(13)	14.90
D.C.	1.03	0.25	(50)	-39.90 <sup>c</sup>	North Dakota	1.09	0.22	(49)	-36.39 <sup>c</sup>
Florida	1.71	0.05	(30)	-0.24	Ohio	1.84	0.07	(20)	7.28
Georgia	1.44	0.08	(45)	-16.06 <sup>c</sup>	Oklahoma	2.15	0.14	(04)	25.45 <sup>c</sup>
Hawaii	0.90	0.15	(51)	-47.67 <sup>c</sup>	Oregon	2.33	0.15	(01)	36.34 <sup>c</sup>
Idaho	1.85	0.22	(19)	7.79	Pennsylvania	1.85	0.07	(18)	8.09
Illinois	1.63	0.07	(35)	-4.67	Rhode Island	1.75	0.23	(25)	2.37
Indiana	1.87	0.10	(17)	9.45	South Carolina	1.47	0.11	(43)	-14.37
Iowa	1.77	0.14	(24)	3.65	South Dakota	1.78	0.27	(23)	3.85
Kansas	1.93	0.16	(16)	12.67	Tennessee	1.78	0.10	(22)	4.03
Kentucky	2.10	0.13	(07)	22.67 <sup>c</sup>	Texas	1.55	0.05	(40)	-9.77
Louisiana	1.39	0.11	(47)	-18.84 <sup>c</sup>	Utah	1.96	0.19	(14)	14.50
Maine	1.98	0.22	(11)	15.88 <sup>c</sup>	Vermont	2.07	0.33	(08)	21.16 <sup>c</sup>
Maryland	1.59	0.10	(38)	-6.95	Virginia	2.03	0.10	(09)	18.29 <sup>c</sup>
Massachusetts	1.98	0.10	(12)	15.57 <sup>c</sup>	Washington	2.00	0.11	(10)	16.86 <sup>c</sup>
Michigan	1.53	0.07	(41)	-10.41	West Virginia	2.11	0.19	(05)	23.42 <sup>c</sup>
Minnesota	1.50	0.10	(42)	-12.52	Wisconsin	1.64	0.10	(34)	-4.24
Mississippi	1.32	0.13	(48)	-22.91 <sup>c</sup>	Wyoming	1.69	0.33	(33)	-1.34
Missouri	1.95	0.11	(15)	13.78					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.18  
Myeloma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.38	0.01		
High Five States				
D.C.	6.12	0.47	(01)	80.83 <sup>c</sup>
Alabama	3.95	0.13	(02)	16.85 <sup>c</sup>
South Carolina	3.93	0.13	(03)	16.04 <sup>c</sup>
North Carolina	3.92	0.09	(04)	15.96 <sup>c</sup>
Virginia	3.91	0.10	(05)	15.53 <sup>c</sup>
Low Five States				
Alaska	2.92	0.40	(47)	-13.73
New Mexico	2.91	0.17	(48)	-13.99
Rhode Island	2.86	0.22	(49)	-15.55 <sup>c</sup>
Nevada	2.69	0.15	(50)	-20.44 <sup>c</sup>
Hawaii	2.39	0.18	(51)	-29.38 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.95	0.13	(02)	16.85 <sup>c</sup>	Montana	3.21	0.24	(38)	-5.21
Alaska	2.92	0.40	(47)	-13.73	Nebraska	3.32	0.18	(30)	-1.85
Arizona	2.98	0.10	(45)	-11.97	Nevada	2.69	0.15	(50)	-20.44 <sup>c</sup>
Arkansas	3.52	0.15	(18)	3.98	New Hampshire	3.23	0.21	(35)	-4.60
California	3.15	0.04	(40)	-6.83	New Jersey	3.27	0.08	(33)	-3.30
Colorado	3.22	0.12	(37)	-4.93	New Mexico	2.91	0.17	(48)	-13.99
Connecticut	2.98	0.12	(44)	-11.79	New York	3.14	0.05	(41)	-7.22
Delaware	3.46	0.27	(23)	2.41	North Carolina	3.92	0.09	(04)	15.96 <sup>c</sup>
D.C.	6.12	0.47	(01)	80.83 <sup>c</sup>	North Dakota	3.51	0.30	(19)	3.85
Florida	2.95	0.05	(46)	-12.75	Ohio	3.60	0.07	(13)	6.35
Georgia	3.51	0.10	(20)	3.74	Oklahoma	3.56	0.14	(17)	5.07
Hawaii	2.39	0.18	(51)	-29.38 <sup>c</sup>	Oregon	3.57	0.13	(16)	5.43
Idaho	3.41	0.22	(26)	0.71	Pennsylvania	3.35	0.07	(29)	-0.87
Illinois	3.47	0.07	(22)	2.65	Rhode Island	2.86	0.22	(49)	-15.55 <sup>c</sup>
Indiana	3.49	0.10	(21)	3.01	South Carolina	3.93	0.13	(03)	16.04 <sup>c</sup>
Iowa	3.22	0.13	(36)	-4.77	South Dakota	3.37	0.27	(27)	-0.26
Kansas	3.75	0.16	(10)	10.79	Tennessee	3.88	0.11	(06)	14.74
Kentucky	3.42	0.12	(25)	1.01	Texas	3.29	0.06	(32)	-2.90
Louisiana	3.81	0.13	(08)	12.66	Utah	3.36	0.19	(28)	-0.80
Maine	3.07	0.19	(42)	-9.19	Vermont	3.06	0.29	(43)	-9.67
Maryland	3.82	0.12	(07)	12.75	Virginia	3.91	0.10	(05)	15.53 <sup>c</sup>
Massachusetts	3.25	0.10	(34)	-4.09	Washington	3.29	0.10	(31)	-2.73
Michigan	3.75	0.08	(09)	10.93	West Virginia	3.45	0.17	(24)	2.09
Minnesota	3.60	0.12	(14)	6.27	Wisconsin	3.67	0.11	(11)	8.47
Mississippi	3.66	0.16	(12)	8.18	Wyoming	3.17	0.34	(39)	-6.41
Missouri	3.58	0.10	(15)	5.74					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.19  
Myeloma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	4.30	0.03		
High Five States				
D.C.	7.92	0.86	(01)	84.40 <sup>c</sup>
Tennessee	5.21	0.20	(02)	21.25 <sup>c</sup>
South Carolina	5.00	0.23	(03)	16.41 <sup>c</sup>
Alabama	4.99	0.22	(04)	16.07 <sup>c</sup>
North Carolina	4.89	0.16	(05)	13.93
Low Five States				
Florida	3.71	0.08	(47)	-13.76
Vermont	3.53	0.49	(48)	-17.89 <sup>c</sup>
Rhode Island	3.38	0.37	(49)	-21.39 <sup>c</sup>
Nevada	3.31	0.26	(50)	-23.02 <sup>c</sup>
Hawaii	3.23	0.31	(51)	-24.78 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.99	0.22	(04)	16.07 <sup>c</sup>	Montana	3.90	0.39	(41)	-9.31
Alaska	3.78	0.68	(43)	-12.11	Nebraska	4.09	0.31	(35)	-4.77
Arizona	3.77	0.16	(44)	-12.23	Nevada	3.31	0.26	(50)	-23.02 <sup>c</sup>
Arkansas	4.68	0.26	(12)	8.90	New Hampshire	3.96	0.36	(40)	-7.72
California	4.03	0.08	(37)	-6.28	New Jersey	4.21	0.15	(28)	-2.07
Colorado	4.19	0.22	(29)	-2.46	New Mexico	3.73	0.29	(46)	-13.28
Connecticut	3.85	0.21	(42)	-10.32	New York	3.99	0.10	(39)	-7.03
Delaware	4.69	0.48	(11)	9.10	North Carolina	4.89	0.16	(05)	13.93
D.C.	7.92	0.86	(01)	84.40 <sup>c</sup>	North Dakota	4.28	0.50	(25)	-0.44
Florida	3.71	0.08	(47)	-13.76	Ohio	4.51	0.13	(20)	4.87
Georgia	4.37	0.17	(23)	1.81	Oklahoma	4.65	0.24	(15)	8.13
Hawaii	3.23	0.31	(51)	-24.78 <sup>c</sup>	Oregon	4.68	0.23	(13)	8.86
Idaho	4.17	0.36	(30)	-3.02	Pennsylvania	4.16	0.11	(33)	-3.23
Illinois	4.50	0.13	(21)	4.77	Rhode Island	3.38	0.37	(49)	-21.39 <sup>c</sup>
Indiana	4.65	0.19	(14)	8.15	South Carolina	5.00	0.23	(03)	16.41 <sup>c</sup>
Iowa	4.08	0.23	(36)	-4.92	South Dakota	4.16	0.46	(32)	-3.13
Kansas	4.74	0.27	(09)	10.30	Tennessee	5.21	0.20	(02)	21.25 <sup>c</sup>
Kentucky	4.57	0.23	(19)	6.44	Texas	4.15	0.10	(34)	-3.37
Louisiana	4.83	0.24	(07)	12.34	Utah	4.23	0.32	(27)	-1.64
Maine	3.77	0.33	(45)	-12.29	Vermont	3.53	0.49	(48)	-17.89 <sup>c</sup>
Maryland	4.61	0.20	(17)	7.41	Virginia	4.88	0.18	(06)	13.55
Massachusetts	4.27	0.17	(26)	-0.61	Washington	4.37	0.18	(24)	1.75
Michigan	4.78	0.15	(08)	11.27	West Virginia	4.47	0.31	(22)	4.13
Minnesota	4.64	0.20	(16)	7.93	Wisconsin	4.73	0.19	(10)	10.20
Mississippi	4.17	0.27	(31)	-3.03	Wyoming	4.02	0.58	(38)	-6.35
Missouri	4.58	0.18	(18)	6.62					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 18.20  
Myeloma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.73	0.02		
High Five States				
D.C.	4.91	0.55	(01)	80.00 <sup>c</sup>
Mississippi	3.36	0.20	(02)	23.40 <sup>c</sup>
Maryland	3.27	0.14	(03)	20.11 <sup>c</sup>
North Carolina	3.27	0.11	(04)	19.83 <sup>c</sup>
Virginia	3.25	0.12	(05)	19.24 <sup>c</sup>
Low Five States				
Arizona	2.36	0.11	(47)	-13.42
New Mexico	2.28	0.20	(48)	-16.19 <sup>c</sup>
Nevada	2.18	0.19	(49)	-19.94 <sup>c</sup>
Alaska	2.15	0.46	(50)	-21.11 <sup>c</sup>
Hawaii	1.75	0.20	(51)	-35.69 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.21	0.15	(06)	17.79 <sup>c</sup>	Montana	2.65	0.30	(33)	-2.97
Alaska	2.15	0.46	(50)	-21.11 <sup>c</sup>	Nebraska	2.78	0.22	(20)	2.12
Arizona	2.36	0.11	(47)	-13.42	Nevada	2.18	0.19	(49)	-19.94 <sup>c</sup>
Arkansas	2.65	0.17	(32)	-2.94	New Hampshire	2.70	0.26	(27)	-1.06
California	2.51	0.05	(41)	-7.77	New Jersey	2.60	0.10	(35)	-4.47
Colorado	2.47	0.14	(43)	-9.48	New Mexico	2.28	0.20	(48)	-16.19 <sup>c</sup>
Connecticut	2.36	0.14	(46)	-13.40	New York	2.55	0.07	(37)	-6.53
Delaware	2.63	0.31	(34)	-3.66	North Carolina	3.27	0.11	(04)	19.83 <sup>c</sup>
D.C.	4.91	0.55	(01)	80.00 <sup>c</sup>	North Dakota	2.98	0.38	(11)	9.42
Florida	2.37	0.06	(45)	-13.13	Ohio	2.98	0.09	(12)	9.25
Georgia	2.94	0.11	(13)	7.78	Oklahoma	2.73	0.16	(23)	0.18
Hawaii	1.75	0.20	(51)	-35.69 <sup>c</sup>	Oregon	2.73	0.15	(24)	-0.02
Idaho	2.81	0.27	(18)	3.03	Pennsylvania	2.81	0.08	(17)	3.10
Illinois	2.77	0.09	(21)	1.68	Rhode Island	2.54	0.27	(38)	-6.77
Indiana	2.70	0.12	(26)	-0.98	South Carolina	3.21	0.15	(07)	17.68 <sup>c</sup>
Iowa	2.54	0.16	(39)	-6.79	South Dakota	2.74	0.33	(22)	0.61
Kansas	3.06	0.19	(09)	12.27	Tennessee	2.93	0.12	(14)	7.42
Kentucky	2.65	0.14	(29)	-2.74	Texas	2.65	0.07	(31)	-2.93
Louisiana	3.12	0.16	(08)	14.39	Utah	2.66	0.23	(28)	-2.35
Maine	2.55	0.23	(36)	-6.38	Vermont	2.72	0.37	(25)	-0.14
Maryland	3.27	0.14	(03)	20.11 <sup>c</sup>	Virginia	3.25	0.12	(05)	19.24 <sup>c</sup>
Massachusetts	2.53	0.11	(40)	-7.34	Washington	2.47	0.12	(44)	-9.50
Michigan	3.00	0.10	(10)	10.01	West Virginia	2.65	0.20	(30)	-2.78
Minnesota	2.79	0.14	(19)	2.17	Wisconsin	2.91	0.13	(15)	6.61
Mississippi	3.36	0.20	(02)	23.40 <sup>c</sup>	Wyoming	2.47	0.42	(42)	-9.45
Missouri	2.84	0.12	(16)	4.01					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 19.22  
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	6.43	0.02		
High Five States				
West Virginia	7.51	0.26	(01)	16.83 <sup>c</sup>
Oklahoma	7.26	0.19	(02)	12.88
Kentucky	7.14	0.18	(03)	10.96
Michigan	7.11	0.12	(04)	10.64
Ohio	7.10	0.11	(05)	10.37
Low Five States				
Georgia	5.75	0.12	(47)	-10.51
Maryland	5.72	0.14	(48)	-10.98
D.C.	5.69	0.45	(49)	-11.48
Hawaii	5.54	0.27	(50)	-13.90
New Mexico	5.43	0.23	(51)	-15.51 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.40	0.16	(22)	-0.41	Montana	6.24	0.33	(28)	-2.97
Alaska	6.38	0.59	(23)	-0.83	Nebraska	6.87	0.26	(13)	6.90
Arizona	6.07	0.14	(36)	-5.66	Nevada	5.79	0.23	(46)	-10.02
Arkansas	6.85	0.21	(14)	6.50	New Hampshire	5.79	0.29	(45)	-9.95
California	6.15	0.06	(35)	-4.34	New Jersey	6.27	0.12	(27)	-2.44
Colorado	5.92	0.17	(43)	-7.97	New Mexico	5.43	0.23	(51)	-15.51 <sup>c</sup>
Connecticut	6.05	0.17	(39)	-5.94	New York	6.16	0.08	(34)	-4.14
Delaware	6.32	0.36	(24)	-1.66	North Carolina	6.05	0.11	(37)	-5.87
D.C.	5.69	0.45	(49)	-11.48	North Dakota	5.84	0.39	(44)	-9.24
Florida	6.18	0.07	(32)	-3.87	Ohio	7.10	0.11	(05)	10.37
Georgia	5.75	0.12	(47)	-10.51	Oklahoma	7.26	0.19	(02)	12.88
Hawaii	5.54	0.27	(50)	-13.90	Oregon	6.66	0.18	(18)	3.59
Idaho	6.29	0.29	(25)	-2.13	Pennsylvania	6.97	0.09	(10)	8.34
Illinois	6.60	0.10	(20)	2.59	Rhode Island	5.95	0.31	(42)	-7.47
Indiana	7.05	0.15	(06)	9.65	South Carolina	6.01	0.16	(41)	-6.61
Iowa	6.92	0.20	(11)	7.68	South Dakota	6.41	0.37	(21)	-0.39
Kansas	6.91	0.21	(12)	7.47	Tennessee	7.01	0.15	(07)	9.03
Kentucky	7.14	0.18	(03)	10.96	Texas	6.21	0.08	(30)	-3.44
Louisiana	6.67	0.18	(17)	3.75	Utah	6.05	0.25	(38)	-5.90
Maine	6.70	0.29	(16)	4.19	Vermont	6.18	0.42	(33)	-3.88
Maryland	5.72	0.14	(48)	-10.98	Virginia	6.21	0.13	(29)	-3.39
Massachusetts	6.05	0.13	(40)	-5.96	Washington	6.76	0.15	(15)	5.14
Michigan	7.11	0.12	(04)	10.64	West Virginia	7.51	0.26	(01)	16.83 <sup>c</sup>
Minnesota	7.00	0.16	(08)	8.86	Wisconsin	6.99	0.15	(09)	8.68
Mississippi	6.21	0.21	(31)	-3.47	Wyoming	6.28	0.49	(26)	-2.41
Missouri	6.65	0.14	(19)	3.39					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 19.23  
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	8.20	0.04		
High Five States				
Minnesota	9.40	0.29	(01)	14.65
Indiana	9.30	0.26	(02)	13.52
Tennessee	9.18	0.27	(03)	12.04
Ohio	9.18	0.19	(04)	11.95
Kansas	9.14	0.38	(05)	11.51
Low Five States				
Hawaii	7.22	0.47	(47)	-11.94
Idaho	7.21	0.48	(48)	-11.98
Nevada	7.14	0.37	(49)	-12.86
North Dakota	6.67	0.63	(50)	-18.63 <sup>c</sup>
New Mexico	6.55	0.38	(51)	-20.05 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.98	0.28	(25)	-2.58	Montana	7.90	0.57	(28)	-3.66
Alaska	7.58	0.98	(42)	-7.55	Nebraska	8.43	0.45	(19)	2.87
Arizona	7.59	0.23	(41)	-7.44	Nevada	7.14	0.37	(49)	-12.86
Arkansas	8.57	0.36	(15)	4.57	New Hampshire	7.26	0.50	(46)	-11.44
California	7.79	0.10	(36)	-4.95	New Jersey	7.82	0.20	(33)	-4.57
Colorado	7.84	0.30	(31)	-4.30	New Mexico	6.55	0.38	(51)	-20.05 <sup>c</sup>
Connecticut	7.54	0.30	(43)	-8.00	New York	7.89	0.14	(29)	-3.75
Delaware	8.27	0.64	(23)	0.85	North Carolina	7.52	0.20	(44)	-8.26
D.C.	7.96	0.84	(26)	-2.93	North Dakota	6.67	0.63	(50)	-18.63 <sup>c</sup>
Florida	7.79	0.12	(37)	-5.01	Ohio	9.18	0.19	(04)	11.95
Georgia	7.62	0.23	(40)	-7.09	Oklahoma	9.00	0.33	(07)	9.81
Hawaii	7.22	0.47	(47)	-11.94	Oregon	8.46	0.31	(18)	3.27
Idaho	7.21	0.48	(48)	-11.98	Pennsylvania	8.87	0.17	(10)	8.27
Illinois	8.49	0.18	(17)	3.63	Rhode Island	8.29	0.58	(21)	1.17
Indiana	9.30	0.26	(02)	13.52	South Carolina	7.82	0.29	(34)	-4.59
Iowa	8.85	0.34	(11)	8.04	South Dakota	7.83	0.63	(32)	-4.50
Kansas	9.14	0.38	(05)	11.51	Tennessee	9.18	0.27	(03)	12.04
Kentucky	9.02	0.32	(06)	10.11	Texas	7.93	0.14	(27)	-3.19
Louisiana	8.59	0.31	(14)	4.85	Utah	7.66	0.43	(38)	-6.50
Maine	8.80	0.51	(12)	7.42	Vermont	7.81	0.73	(35)	-4.71
Maryland	7.43	0.26	(45)	-9.29	Virginia	8.16	0.23	(24)	-0.48
Massachusetts	7.85	0.23	(30)	-4.24	Washington	8.57	0.25	(16)	4.56
Michigan	8.95	0.20	(09)	9.17	West Virginia	8.75	0.43	(13)	6.73
Minnesota	9.40	0.29	(01)	14.65	Wisconsin	8.97	0.26	(08)	9.39
Mississippi	8.28	0.38	(22)	1.07	Wyoming	7.65	0.83	(39)	-6.68
Missouri	8.29	0.25	(20)	1.21					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 19.24  
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.11	0.02		
High Five States				
West Virginia	6.55	0.32	(01)	28.25 <sup>c</sup>
Oklahoma	5.97	0.23	(02)	16.80 <sup>c</sup>
Michigan	5.77	0.14	(03)	12.99
Kentucky	5.74	0.21	(04)	12.40
Nebraska	5.67	0.32	(05)	10.98
Low Five States				
New Mexico	4.45	0.28	(47)	-12.83
Colorado	4.44	0.19	(48)	-13.10
Rhode Island	4.36	0.35	(49)	-14.66
Hawaii	4.26	0.32	(50)	-16.55 <sup>c</sup>
D.C.	4.00	0.49	(51)	-21.71 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.25	0.19	(22)	2.83	Montana	4.88	0.40	(34)	-4.48
Alaska	5.46	0.74	(12)	6.84	Nebraska	5.67	0.32	(05)	10.98
Arizona	4.83	0.16	(36)	-5.53	Nevada	4.58	0.27	(44)	-10.30
Arkansas	5.54	0.25	(09)	8.48	New Hampshire	4.73	0.35	(41)	-7.47
California	4.90	0.07	(33)	-4.18	New Jersey	5.19	0.14	(23)	1.65
Colorado	4.44	0.19	(48)	-13.10	New Mexico	4.45	0.28	(47)	-12.83
Connecticut	4.95	0.20	(30)	-3.22	New York	4.93	0.09	(31)	-3.56
Delaware	4.82	0.42	(37)	-5.58	North Carolina	4.96	0.14	(29)	-2.99
D.C.	4.00	0.49	(51)	-21.71 <sup>c</sup>	North Dakota	5.11	0.48	(27)	-0.08
Florida	4.87	0.09	(35)	-4.63	Ohio	5.58	0.12	(07)	9.25
Georgia	4.46	0.14	(46)	-12.75	Oklahoma	5.97	0.23	(02)	16.80 <sup>c</sup>
Hawaii	4.26	0.32	(50)	-16.55 <sup>c</sup>	Oregon	5.31	0.21	(18)	3.88
Idaho	5.59	0.38	(06)	9.39	Pennsylvania	5.55	0.11	(08)	8.55
Illinois	5.26	0.12	(21)	2.90	Rhode Island	4.36	0.35	(49)	-14.66
Indiana	5.44	0.17	(13)	6.56	South Carolina	4.74	0.19	(40)	-7.33
Iowa	5.51	0.23	(11)	7.80	South Dakota	5.32	0.45	(17)	4.06
Kansas	5.26	0.25	(20)	2.91	Tennessee	5.37	0.17	(16)	5.02
Kentucky	5.74	0.21	(04)	12.40	Texas	4.91	0.09	(32)	-3.85
Louisiana	5.18	0.20	(25)	1.32	Utah	4.67	0.30	(43)	-8.65
Maine	5.12	0.33	(26)	0.13	Vermont	4.99	0.50	(28)	-2.38
Maryland	4.46	0.17	(45)	-12.64	Virginia	4.79	0.15	(39)	-6.30
Massachusetts	4.80	0.15	(38)	-6.15	Washington	5.37	0.17	(15)	5.11
Michigan	5.77	0.14	(03)	12.99	West Virginia	6.55	0.32	(01)	28.25 <sup>c</sup>
Minnesota	5.19	0.18	(24)	1.62	Wisconsin	5.51	0.18	(10)	7.89
Mississippi	4.69	0.24	(42)	-8.21	Wyoming	5.31	0.60	(19)	3.87
Missouri	5.44	0.17	(14)	6.56					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 20.24  
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.46	0.01		
High Five States				
D.C.	3.81	0.37	(01)	54.89 <sup>c</sup>
Mississippi	3.25	0.15	(02)	32.16 <sup>c</sup>
Louisiana	3.10	0.12	(03)	26.27 <sup>c</sup>
Tennessee	3.02	0.10	(04)	22.99 <sup>c</sup>
South Carolina	3.02	0.11	(05)	22.86 <sup>c</sup>
Low Five States				
Montana	1.97	0.19	(47)	-19.72 <sup>c</sup>
Minnesota	1.96	0.08	(48)	-20.08 <sup>c</sup>
Colorado	1.96	0.09	(49)	-20.17 <sup>c</sup>
South Dakota	1.91	0.21	(50)	-22.26 <sup>c</sup>
Utah	1.41	0.12	(51)	-42.65 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.89	0.11	(08)	17.60 <sup>c</sup>	Montana	1.97	0.19	(47)	-19.72 <sup>c</sup>
Alaska	2.06	0.31	(44)	-16.37 <sup>c</sup>	Nebraska	2.28	0.15	(34)	-7.42
Arizona	2.02	0.08	(45)	-18.01 <sup>c</sup>	Nevada	2.23	0.13	(37)	-9.39
Arkansas	2.98	0.14	(06)	21.19 <sup>c</sup>	New Hampshire	2.46	0.19	(26)	0.03
California	2.51	0.04	(22)	2.04	New Jersey	2.12	0.07	(42)	-13.63
Colorado	1.96	0.09	(49)	-20.17 <sup>c</sup>	New Mexico	2.16	0.14	(40)	-12.17
Connecticut	2.09	0.10	(43)	-14.94	New York	2.17	0.05	(39)	-11.78
Delaware	2.47	0.22	(25)	0.61	North Carolina	2.48	0.07	(24)	0.93
D.C.	3.81	0.37	(01)	54.89 <sup>c</sup>	North Dakota	2.19	0.24	(38)	-10.72
Florida	2.74	0.05	(10)	11.56	Ohio	2.62	0.06	(14)	6.77
Georgia	2.57	0.08	(16)	4.54	Oklahoma	2.54	0.11	(18)	3.52
Hawaii	2.82	0.19	(09)	14.55	Oregon	2.30	0.10	(33)	-6.31
Idaho	2.65	0.19	(13)	7.90	Pennsylvania	2.24	0.05	(36)	-8.95
Illinois	2.38	0.06	(29)	-3.17	Rhode Island	2.14	0.19	(41)	-12.77
Indiana	2.25	0.08	(35)	-8.60	South Carolina	3.02	0.11	(05)	22.86 <sup>c</sup>
Iowa	2.33	0.12	(30)	-5.11	South Dakota	1.91	0.21	(50)	-22.26 <sup>c</sup>
Kansas	2.01	0.11	(46)	-18.37 <sup>c</sup>	Tennessee	3.02	0.10	(04)	22.99 <sup>c</sup>
Kentucky	2.73	0.11	(11)	11.07	Texas	2.52	0.05	(20)	2.35
Louisiana	3.10	0.12	(03)	26.27 <sup>c</sup>	Utah	1.41	0.12	(51)	-42.65 <sup>c</sup>
Maine	2.44	0.17	(27)	-0.72	Vermont	2.72	0.27	(12)	10.73
Maryland	2.51	0.09	(21)	2.24	Virginia	2.31	0.08	(32)	-6.08
Massachusetts	2.57	0.08	(17)	4.39	Washington	2.90	0.09	(07)	17.99 <sup>c</sup>
Michigan	2.53	0.07	(19)	2.90	West Virginia	2.60	0.15	(15)	5.87
Minnesota	1.96	0.08	(48)	-20.08 <sup>c</sup>	Wisconsin	2.40	0.09	(28)	-2.45
Mississippi	3.25	0.15	(02)	32.16 <sup>c</sup>	Wyoming	2.49	0.30	(23)	1.12
Missouri	2.32	0.08	(31)	-5.62					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 20.25  
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	3.75	0.02		
High Five States				
D.C.	6.37	0.74	(01)	69.65 <sup>c</sup>
Mississippi	5.51	0.29	(02)	46.83 <sup>c</sup>
Louisiana	5.02	0.23	(03)	33.78 <sup>c</sup>
South Carolina	4.90	0.22	(04)	30.64 <sup>c</sup>
Tennessee	4.84	0.18	(05)	29.10 <sup>c</sup>
Low Five States				
South Dakota	2.80	0.37	(47)	-25.39 <sup>c</sup>
Montana	2.77	0.33	(48)	-26.10 <sup>c</sup>
Minnesota	2.77	0.15	(49)	-26.16 <sup>c</sup>
Colorado	2.71	0.17	(50)	-27.73 <sup>c</sup>
Utah	1.78	0.20	(51)	-52.64 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.62	0.21	(06)	23.23 <sup>c</sup>	Montana	2.77	0.33	(48)	-26.10 <sup>c</sup>
Alaska	2.93	0.54	(46)	-21.92 <sup>c</sup>	Nebraska	2.96	0.26	(45)	-21.21 <sup>c</sup>
Arizona	3.06	0.14	(43)	-18.48 <sup>c</sup>	Nevada	3.25	0.24	(41)	-13.40
Arkansas	4.59	0.26	(07)	22.42 <sup>c</sup>	New Hampshire	3.79	0.34	(22)	0.92
California	3.74	0.07	(24)	-0.36	New Jersey	3.08	0.12	(42)	-17.92 <sup>c</sup>
Colorado	2.71	0.17	(50)	-27.73 <sup>c</sup>	New Mexico	3.26	0.26	(40)	-13.21
Connecticut	3.27	0.19	(39)	-12.80	New York	3.42	0.09	(36)	-8.91
Delaware	3.87	0.42	(19)	3.26	North Carolina	3.92	0.14	(17)	4.35
D.C.	6.37	0.74	(01)	69.65 <sup>c</sup>	North Dakota	3.58	0.46	(29)	-4.66
Florida	4.23	0.09	(10)	12.79	Ohio	3.96	0.12	(15)	5.59
Georgia	4.21	0.16	(11)	12.31	Oklahoma	3.77	0.21	(23)	0.36
Hawaii	4.31	0.35	(08)	14.99	Oregon	3.53	0.19	(30)	-5.81
Idaho	3.69	0.33	(25)	-1.68	Pennsylvania	3.48	0.10	(34)	-7.38
Illinois	3.49	0.11	(31)	-7.04	Rhode Island	3.29	0.35	(38)	-12.32
Indiana	3.48	0.15	(33)	-7.25	South Carolina	4.90	0.22	(04)	30.64 <sup>c</sup>
Iowa	3.61	0.22	(28)	-3.76	South Dakota	2.80	0.37	(47)	-25.39 <sup>c</sup>
Kansas	3.03	0.21	(44)	-19.29 <sup>c</sup>	Tennessee	4.84	0.18	(05)	29.10 <sup>c</sup>
Kentucky	4.31	0.21	(09)	14.75	Texas	3.92	0.09	(16)	4.60
Louisiana	5.02	0.23	(03)	33.78 <sup>c</sup>	Utah	1.78	0.20	(51)	-52.64 <sup>c</sup>
Maine	3.48	0.31	(32)	-7.23	Vermont	4.19	0.52	(12)	11.74
Maryland	3.80	0.18	(21)	1.17	Virginia	3.64	0.15	(26)	-2.98
Massachusetts	3.91	0.16	(18)	4.17	Washington	4.12	0.17	(13)	9.74
Michigan	3.86	0.13	(20)	2.81	West Virginia	4.12	0.28	(14)	9.68
Minnesota	2.77	0.15	(49)	-26.16 <sup>c</sup>	Wisconsin	3.38	0.16	(37)	-9.92
Mississippi	5.51	0.29	(02)	46.83 <sup>c</sup>	Wyoming	3.64	0.53	(27)	-3.10
Missouri	3.42	0.15	(35)	-8.77					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 20.26  
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.39	0.01		
High Five States				
D.C.	1.88	0.35	(01)	35.50 <sup>c</sup>
Washington	1.84	0.10	(02)	32.65 <sup>c</sup>
Idaho	1.68	0.21	(03)	21.58 <sup>c</sup>
Arkansas	1.65	0.14	(04)	19.40 <sup>c</sup>
Nebraska	1.65	0.18	(05)	18.96 <sup>c</sup>
Low Five States				
Kansas	1.15	0.12	(46)	-16.85 <sup>c</sup>
Arizona	1.11	0.08	(47)	-19.74 <sup>c</sup>
South Dakota	1.07	0.20	(48)	-22.60 <sup>c</sup>
Utah	1.06	0.14	(49)	-23.59 <sup>c</sup>
North Dakota	1.06	0.24	(50)	-23.83 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.51	0.10	(12)	8.70	Montana	1.27	0.20	(35)	-8.42
Alaska	-	-	(-)	-	Nebraska	1.65	0.18	(05)	18.96 <sup>c</sup>
Arizona	1.11	0.08	(47)	-19.74 <sup>c</sup>	Nevada	1.30	0.14	(34)	-6.50
Arkansas	1.65	0.14	(04)	19.40 <sup>c</sup>	New Hampshire	1.30	0.19	(30)	-6.19
California	1.49	0.04	(16)	7.24	New Jersey	1.36	0.07	(26)	-1.69
Colorado	1.34	0.10	(28)	-3.56	New Mexico	1.18	0.14	(44)	-14.93
Connecticut	1.16	0.10	(45)	-16.64 <sup>c</sup>	New York	1.19	0.04	(43)	-14.14
Delaware	1.30	0.21	(33)	-6.44	North Carolina	1.33	0.07	(29)	-4.23
D.C.	1.88	0.35	(01)	35.50 <sup>c</sup>	North Dakota	1.06	0.24	(50)	-23.83 <sup>c</sup>
Florida	1.47	0.05	(19)	5.81	Ohio	1.51	0.06	(11)	8.71
Georgia	1.30	0.07	(32)	-6.35	Oklahoma	1.51	0.12	(13)	8.69
Hawaii	1.50	0.19	(14)	8.19	Oregon	1.26	0.10	(37)	-8.95
Idaho	1.68	0.21	(03)	21.58 <sup>c</sup>	Pennsylvania	1.25	0.05	(38)	-9.69
Illinois	1.46	0.06	(20)	5.66	Rhode Island	1.26	0.20	(36)	-8.77
Indiana	1.21	0.08	(42)	-12.82	South Carolina	1.48	0.11	(18)	6.79
Iowa	1.24	0.11	(40)	-10.20	South Dakota	1.07	0.20	(48)	-22.60 <sup>c</sup>
Kansas	1.15	0.12	(46)	-16.85 <sup>c</sup>	Tennessee	1.52	0.09	(10)	9.45
Kentucky	1.44	0.11	(23)	4.11	Texas	1.35	0.05	(27)	-2.53
Louisiana	1.54	0.11	(08)	11.07	Utah	1.06	0.14	(49)	-23.59 <sup>c</sup>
Maine	1.56	0.18	(07)	12.38	Vermont	1.53	0.27	(09)	10.35
Maryland	1.49	0.10	(15)	7.24	Virginia	1.25	0.08	(39)	-9.71
Massachusetts	1.48	0.09	(17)	7.05	Washington	1.84	0.10	(02)	32.65 <sup>c</sup>
Michigan	1.42	0.07	(24)	2.66	West Virginia	1.24	0.14	(41)	-10.38
Minnesota	1.30	0.09	(31)	-6.21	Wisconsin	1.58	0.10	(06)	14.10
Mississippi	1.45	0.13	(22)	4.40	Wyoming	1.46	0.31	(21)	5.11
Missouri	1.37	0.09	(25)	-1.24					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 21.14  
Cancer of the Ovary (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	8.08	0.03		
High Five States				
Washington	9.56	0.23	(01)	18.23 <sup>c</sup>
Alabama	9.24	0.25	(02)	14.37
Oregon	9.18	0.29	(03)	13.56
Wisconsin	9.08	0.23	(04)	12.34
Iowa	8.92	0.31	(05)	10.39
Low Five States				
Rhode Island	7.30	0.46	(47)	-9.67
Mississippi	7.27	0.29	(48)	-10.01
Nevada	6.91	0.33	(49)	-14.49
Alaska	6.66	0.76	(50)	-17.59 <sup>c</sup>
Hawaii	6.10	0.39	(51)	-24.54 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.24	0.25	(02)	14.37	Montana	8.74	0.53	(07)	8.20
Alaska	6.66	0.76	(50)	-17.59 <sup>c</sup>	Nebraska	7.72	0.38	(38)	-4.53
Arizona	7.77	0.21	(35)	-3.87	Nevada	6.91	0.33	(49)	-14.49
Arkansas	8.25	0.31	(17)	2.14	New Hampshire	7.95	0.45	(32)	-1.58
California	8.03	0.09	(27)	-0.67	New Jersey	8.71	0.18	(08)	7.72
Colorado	8.65	0.27	(09)	7.03	New Mexico	7.64	0.37	(39)	-5.49
Connecticut	7.93	0.27	(33)	-1.93	New York	8.06	0.12	(26)	-0.24
Delaware	8.07	0.54	(23)	-0.14	North Carolina	7.61	0.17	(40)	-5.88
D.C.	7.49	0.69	(44)	-7.33	North Dakota	8.37	0.65	(15)	3.52
Florida	7.34	0.11	(46)	-9.13	Ohio	8.15	0.15	(19)	0.80
Georgia	8.37	0.19	(14)	3.63	Oklahoma	8.02	0.27	(28)	-0.80
Hawaii	6.10	0.39	(51)	-24.54 <sup>c</sup>	Oregon	9.18	0.29	(03)	13.56
Idaho	7.78	0.44	(34)	-3.78	Pennsylvania	8.79	0.14	(06)	8.72
Illinois	8.08	0.15	(22)	-0.05	Rhode Island	7.30	0.46	(47)	-9.67
Indiana	8.43	0.21	(12)	4.32	South Carolina	8.11	0.25	(21)	0.37
Iowa	8.92	0.31	(05)	10.39	South Dakota	7.96	0.57	(30)	-1.46
Kansas	8.44	0.32	(10)	4.40	Tennessee	7.74	0.20	(36)	-4.24
Kentucky	8.12	0.25	(20)	0.43	Texas	7.53	0.11	(42)	-6.83
Louisiana	7.73	0.25	(37)	-4.29	Utah	7.60	0.38	(41)	-5.94
Maine	7.96	0.42	(31)	-1.46	Vermont	7.52	0.63	(43)	-6.98
Maryland	8.06	0.22	(25)	-0.22	Virginia	8.43	0.20	(11)	4.33
Massachusetts	8.07	0.20	(24)	-0.16	Washington	9.56	0.23	(01)	18.23 <sup>c</sup>
Michigan	8.42	0.17	(13)	4.15	West Virginia	8.33	0.36	(16)	3.02
Minnesota	8.01	0.23	(29)	-0.94	Wisconsin	9.08	0.23	(04)	12.34
Mississippi	7.27	0.29	(48)	-10.01	Wyoming	8.17	0.74	(18)	1.13
Missouri	7.46	0.20	(45)	-7.71					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.18  
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	10.90	0.03		
High Five States				
D.C.	14.06	0.71	(01)	28.99 <sup>c</sup>
Louisiana	12.50	0.24	(02)	14.74
Mississippi	11.93	0.28	(03)	9.46
Michigan	11.86	0.15	(04)	8.80
Connecticut	11.67	0.24	(05)	7.10
Low Five States				
Colorado	9.84	0.21	(47)	-9.66
Arizona	9.73	0.17	(48)	-10.73
New Mexico	9.56	0.30	(49)	-12.28
West Virginia	9.42	0.29	(50)	-13.59
Utah	9.12	0.31	(51)	-16.28 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	11.29	0.21	(15)	3.58	Montana	9.95	0.42	(46)	-8.65
Alaska	11.39	0.76	(12)	4.50	Nebraska	10.78	0.33	(31)	-1.05
Arizona	9.73	0.17	(48)	-10.73	Nevada	11.08	0.31	(21)	1.64
Arkansas	11.27	0.27	(16)	3.40	New Hampshire	11.65	0.40	(06)	6.87
California	10.44	0.08	(39)	-4.21	New Jersey	11.57	0.16	(07)	6.13
Colorado	9.84	0.21	(47)	-9.66	New Mexico	9.56	0.30	(49)	-12.28
Connecticut	11.67	0.24	(05)	7.10	New York	11.16	0.10	(19)	2.44
Delaware	10.99	0.47	(25)	0.86	North Carolina	10.69	0.15	(34)	-1.90
D.C.	14.06	0.71	(01)	28.99 <sup>c</sup>	North Dakota	10.33	0.52	(43)	-5.25
Florida	10.33	0.09	(42)	-5.18	Ohio	11.47	0.13	(10)	5.25
Georgia	10.57	0.16	(35)	-3.05	Oklahoma	10.46	0.23	(38)	-4.01
Hawaii	10.96	0.38	(27)	0.55	Oregon	11.03	0.23	(22)	1.23
Idaho	10.56	0.38	(36)	-3.11	Pennsylvania	11.54	0.12	(08)	5.89
Illinois	11.36	0.13	(13)	4.26	Rhode Island	10.35	0.41	(41)	-5.05
Indiana	10.98	0.18	(26)	0.79	South Carolina	11.13	0.22	(20)	2.10
Iowa	10.36	0.24	(40)	-4.90	South Dakota	10.07	0.47	(45)	-7.58
Kansas	10.77	0.27	(32)	-1.22	Tennessee	11.01	0.18	(24)	1.04
Kentucky	10.95	0.22	(28)	0.52	Texas	10.16	0.10	(44)	-6.79
Louisiana	12.50	0.24	(02)	14.74	Utah	9.12	0.31	(51)	-16.28 <sup>c</sup>
Maine	10.74	0.36	(33)	-1.49	Vermont	10.79	0.55	(30)	-0.94
Maryland	11.49	0.20	(09)	5.42	Virginia	11.01	0.17	(23)	1.07
Massachusetts	11.41	0.18	(11)	4.67	Washington	11.24	0.19	(17)	3.11
Michigan	11.86	0.15	(04)	8.80	West Virginia	9.42	0.29	(50)	-13.59
Minnesota	10.53	0.20	(37)	-3.35	Wisconsin	11.23	0.19	(18)	3.06
Mississippi	11.93	0.28	(03)	9.46	Wyoming	10.92	0.63	(29)	0.17
Missouri	11.29	0.19	(14)	3.64					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 22.19  
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	12.53	0.04		
High Five States				
D.C.	16.93	1.24	(01)	35.10 <sup>c</sup>
Louisiana	14.18	0.39	(02)	13.16 <sup>c</sup>
Mississippi	14.14	0.48	(03)	12.90 <sup>c</sup>
Connecticut	13.92	0.41	(04)	11.12 <sup>c</sup>
Michigan	13.92	0.25	(05)	11.09 <sup>c</sup>
Low Five States				
West Virginia	11.32	0.48	(47)	-9.63
South Dakota	11.14	0.74	(48)	-11.07 <sup>c</sup>
New Mexico	10.92	0.49	(49)	-12.84 <sup>c</sup>
Colorado	10.83	0.34	(50)	-13.54 <sup>c</sup>
Utah	9.96	0.48	(51)	-20.52 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	13.23	0.36	(08)	5.58	Montana	11.78	0.68	(43)	-6.01
Alaska	12.94	1.18	(18)	3.25	Nebraska	11.98	0.53	(40)	-4.40
Arizona	11.36	0.28	(46)	-9.29	Nevada	12.58	0.48	(27)	0.44
Arkansas	13.16	0.44	(10)	5.00	New Hampshire	13.22	0.66	(09)	5.50
California	11.69	0.13	(44)	-6.69	New Jersey	13.39	0.26	(07)	6.86
Colorado	10.83	0.34	(50)	-13.54 <sup>c</sup>	New Mexico	10.92	0.49	(49)	-12.84 <sup>c</sup>
Connecticut	13.92	0.41	(04)	11.12 <sup>c</sup>	New York	12.82	0.17	(20)	2.34
Delaware	13.03	0.79	(13)	4.03	North Carolina	12.02	0.25	(39)	-4.05
D.C.	16.93	1.24	(01)	35.10 <sup>c</sup>	North Dakota	12.55	0.86	(29)	0.21
Florida	12.11	0.15	(37)	-3.35	Ohio	13.15	0.22	(11)	4.96
Georgia	12.54	0.28	(30)	0.07	Oklahoma	12.22	0.38	(35)	-2.49
Hawaii	12.33	0.60	(33)	-1.61	Oregon	12.35	0.37	(32)	-1.45
Idaho	12.02	0.60	(38)	-4.03	Pennsylvania	13.39	0.20	(06)	6.87
Illinois	12.97	0.22	(15)	3.55	Rhode Island	12.22	0.69	(34)	-2.44
Indiana	12.97	0.30	(16)	3.53	South Carolina	12.71	0.36	(24)	1.46
Iowa	12.19	0.40	(36)	-2.68	South Dakota	11.14	0.74	(48)	-11.07 <sup>c</sup>
Kansas	12.58	0.44	(28)	0.40	Tennessee	13.08	0.31	(12)	4.38
Kentucky	12.74	0.37	(22)	1.67	Texas	11.82	0.17	(42)	-5.66
Louisiana	14.18	0.39	(02)	13.16 <sup>c</sup>	Utah	9.96	0.48	(51)	-20.52 <sup>c</sup>
Maine	11.65	0.57	(45)	-7.03	Vermont	12.35	0.88	(31)	-1.40
Maryland	12.95	0.33	(17)	3.33	Virginia	12.66	0.28	(26)	1.06
Massachusetts	12.68	0.29	(25)	1.24	Washington	12.71	0.30	(23)	1.48
Michigan	13.92	0.25	(05)	11.09 <sup>c</sup>	West Virginia	11.32	0.48	(47)	-9.63
Minnesota	11.94	0.32	(41)	-4.73	Wisconsin	12.76	0.31	(21)	1.86
Mississippi	14.14	0.48	(03)	12.90 <sup>c</sup>	Wyoming	12.94	1.03	(19)	3.25
Missouri	12.97	0.31	(14)	3.56					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 22.20  
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	9.56	0.03		
High Five States				
D.C.	12.02	0.86	(01)	25.75 <sup>c</sup>
Louisiana	11.10	0.30	(02)	16.12 <sup>c</sup>
New Hampshire	10.42	0.51	(03)	8.99
Massachusetts	10.36	0.22	(04)	8.39
Maryland	10.31	0.25	(05)	7.87
Low Five States				
Utah	8.36	0.40	(47)	-12.54
Montana	8.36	0.52	(48)	-12.61
Arizona	8.34	0.22	(49)	-12.74
North Dakota	8.34	0.62	(50)	-12.75
West Virginia	7.84	0.35	(51)	-18.04 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.71	0.26	(22)	1.53	Montana	8.36	0.52	(48)	-12.61
Alaska	9.76	0.96	(20)	2.07	Nebraska	9.67	0.42	(25)	1.18
Arizona	8.34	0.22	(49)	-12.74	Nevada	9.72	0.39	(21)	1.68
Arkansas	9.61	0.33	(26)	0.52	New Hampshire	10.42	0.51	(03)	8.99
California	9.40	0.10	(32)	-1.73	New Jersey	10.13	0.19	(07)	5.95
Colorado	8.98	0.27	(40)	-6.12	New Mexico	8.43	0.39	(46)	-11.80
Connecticut	9.94	0.29	(14)	3.92	New York	9.87	0.13	(17)	3.23
Delaware	9.46	0.59	(29)	-1.04	North Carolina	9.56	0.19	(27)	0.00
D.C.	12.02	0.86	(01)	25.75 <sup>c</sup>	North Dakota	8.34	0.62	(50)	-12.75
Florida	8.84	0.11	(43)	-7.48	Ohio	10.10	0.17	(09)	5.67
Georgia	9.09	0.20	(38)	-4.89	Oklahoma	8.95	0.29	(41)	-6.42
Hawaii	9.69	0.48	(23)	1.37	Oregon	9.90	0.29	(16)	3.53
Idaho	9.22	0.48	(36)	-3.55	Pennsylvania	10.02	0.15	(11)	4.82
Illinois	10.06	0.17	(10)	5.27	Rhode Island	8.89	0.50	(42)	-7.03
Indiana	9.42	0.22	(31)	-1.47	South Carolina	9.82	0.27	(18)	2.70
Iowa	8.84	0.30	(44)	-7.51	South Dakota	9.32	0.61	(34)	-2.47
Kansas	9.13	0.33	(37)	-4.48	Tennessee	9.45	0.22	(30)	-1.16
Kentucky	9.55	0.27	(28)	-0.07	Texas	8.83	0.12	(45)	-7.63
Louisiana	11.10	0.30	(02)	16.12 <sup>c</sup>	Utah	8.36	0.40	(47)	-12.54
Maine	9.78	0.46	(19)	2.28	Vermont	9.28	0.68	(35)	-2.91
Maryland	10.31	0.25	(05)	7.87	Virginia	9.68	0.21	(24)	1.28
Massachusetts	10.36	0.22	(04)	8.39	Washington	10.01	0.24	(12)	4.72
Michigan	10.18	0.18	(06)	6.44	West Virginia	7.84	0.35	(51)	-18.04 <sup>c</sup>
Minnesota	9.33	0.25	(33)	-2.39	Wisconsin	9.99	0.24	(13)	4.51
Mississippi	10.11	0.34	(08)	5.75	Wyoming	9.02	0.78	(39)	-5.63
Missouri	9.90	0.23	(15)	3.55					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 23.14  
Cancer of the Prostate (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	23.05	0.06		
High Five States				
D.C.	38.77	1.97	(01)	68.23 <sup>c</sup>
Mississippi	31.18	0.79	(02)	35.27 <sup>c</sup>
Alabama	28.90	0.58	(03)	25.39 <sup>c</sup>
South Carolina	27.48	0.59	(04)	19.24 <sup>c</sup>
Georgia	27.12	0.46	(05)	17.68 <sup>c</sup>
Low Five States				
Kansas	20.69	0.58	(47)	-10.24 <sup>c</sup>
Rhode Island	20.57	0.91	(48)	-10.75 <sup>c</sup>
Arizona	20.47	0.39	(49)	-11.17 <sup>c</sup>
Florida	20.15	0.20	(50)	-12.59 <sup>c</sup>
Hawaii	15.71	0.69	(51)	-31.85 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	28.90	0.58	(03)	25.39 <sup>c</sup>	Montana	26.44	1.06	(07)	14.72 <sup>c</sup>
Alaska	22.47	1.83	(35)	-2.49	Nebraska	22.94	0.76	(29)	-0.46
Arizona	20.47	0.39	(49)	-11.17 <sup>c</sup>	Nevada	23.91	0.75	(20)	3.77
Arkansas	25.27	0.65	(11)	9.64	New Hampshire	22.42	0.92	(36)	-2.72
California	22.33	0.18	(38)	-3.10	New Jersey	21.57	0.34	(44)	-6.42
Colorado	22.88	0.54	(31)	-0.73	New Mexico	24.41	0.78	(17)	5.91
Connecticut	22.26	0.52	(39)	-3.41	New York	21.83	0.23	(41)	-5.28
Delaware	23.67	1.14	(24)	2.72	North Carolina	25.84	0.40	(10)	12.14 <sup>c</sup>
D.C.	38.77	1.97	(01)	68.23 <sup>c</sup>	North Dakota	23.57	1.19	(25)	2.27
Florida	20.15	0.20	(50)	-12.59 <sup>c</sup>	Ohio	23.80	0.31	(23)	3.27
Georgia	27.12	0.46	(05)	17.68 <sup>c</sup>	Oklahoma	23.48	0.56	(26)	1.87
Hawaii	15.71	0.69	(51)	-31.85 <sup>c</sup>	Oregon	24.44	0.54	(16)	6.04
Idaho	25.91	0.94	(08)	12.44 <sup>c</sup>	Pennsylvania	22.74	0.27	(33)	-1.35
Illinois	24.25	0.32	(18)	5.24	Rhode Island	20.57	0.91	(48)	-10.75 <sup>c</sup>
Indiana	23.21	0.43	(28)	0.71	South Carolina	27.48	0.59	(04)	19.24 <sup>c</sup>
Iowa	22.71	0.55	(34)	-1.44	South Dakota	23.43	1.10	(27)	1.65
Kansas	20.69	0.58	(47)	-10.24 <sup>c</sup>	Tennessee	25.18	0.47	(12)	9.26
Kentucky	23.87	0.55	(21)	3.58	Texas	21.27	0.24	(45)	-7.70
Louisiana	26.63	0.59	(06)	15.53 <sup>c</sup>	Utah	25.90	0.83	(09)	12.38 <sup>c</sup>
Maine	22.94	0.85	(30)	-0.47	Vermont	22.13	1.28	(40)	-3.97
Maryland	25.00	0.49	(13)	8.48	Virginia	24.68	0.43	(14)	7.09
Massachusetts	22.39	0.40	(37)	-2.85	Washington	24.19	0.43	(19)	4.96
Michigan	21.81	0.32	(43)	-5.36	West Virginia	20.95	0.70	(46)	-9.12
Minnesota	23.86	0.47	(22)	3.52	Wisconsin	24.47	0.45	(15)	6.18
Mississippi	31.18	0.79	(02)	35.27 <sup>c</sup>	Wyoming	22.79	1.46	(32)	-1.11
Missouri	21.83	0.42	(42)	-5.30					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.18  
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.54	0.02		
High Five States				
Hawaii	6.03	0.28	(01)	70.08 <sup>c</sup>
Louisiana	4.80	0.15	(02)	35.46 <sup>c</sup>
Alaska	4.45	0.46	(03)	25.64 <sup>c</sup>
D.C.	4.43	0.40	(04)	24.96 <sup>c</sup>
California	4.43	0.05	(05)	24.90 <sup>c</sup>
Low Five States				
Montana	2.52	0.21	(47)	-28.85 <sup>c</sup>
Iowa	2.47	0.12	(48)	-30.38 <sup>c</sup>
Wyoming	2.44	0.30	(49)	-31.15 <sup>c</sup>
Utah	2.31	0.15	(50)	-34.83 <sup>c</sup>
Vermont	2.26	0.25	(51)	-36.23 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.69	0.12	(17)	4.19	Montana	2.52	0.21	(47)	-28.85 <sup>c</sup>
Alaska	4.45	0.46	(03)	25.64 <sup>c</sup>	Nebraska	2.54	0.16	(46)	-28.28 <sup>c</sup>
Arizona	2.99	0.10	(34)	-15.54 <sup>c</sup>	Nevada	3.41	0.17	(20)	-3.92
Arkansas	3.12	0.14	(31)	-12.06	New Hampshire	2.80	0.20	(40)	-21.08 <sup>c</sup>
California	4.43	0.05	(05)	24.90 <sup>c</sup>	New Jersey	4.16	0.09	(08)	17.44 <sup>c</sup>
Colorado	2.85	0.12	(37)	-19.60 <sup>c</sup>	New Mexico	4.27	0.20	(06)	20.50 <sup>c</sup>
Connecticut	3.94	0.14	(12)	11.28	New York	4.13	0.06	(10)	16.40 <sup>c</sup>
Delaware	4.18	0.30	(07)	17.85 <sup>c</sup>	North Carolina	3.24	0.08	(26)	-8.61
D.C.	4.43	0.40	(04)	24.96 <sup>c</sup>	North Dakota	3.22	0.29	(27)	-9.18
Florida	3.10	0.05	(32)	-12.45	Ohio	3.20	0.07	(29)	-9.64
Georgia	3.72	0.10	(16)	5.01	Oklahoma	3.16	0.13	(30)	-10.79
Hawaii	6.03	0.28	(01)	70.08 <sup>c</sup>	Oregon	2.61	0.11	(45)	-26.39 <sup>c</sup>
Idaho	2.73	0.19	(42)	-22.92 <sup>c</sup>	Pennsylvania	3.34	0.07	(24)	-5.80
Illinois	3.81	0.08	(14)	7.49	Rhode Island	3.37	0.23	(22)	-4.81
Indiana	2.72	0.09	(43)	-23.29 <sup>c</sup>	South Carolina	4.08	0.13	(11)	14.99
Iowa	2.47	0.12	(48)	-30.38 <sup>c</sup>	South Dakota	3.40	0.27	(21)	-4.05
Kansas	2.61	0.13	(44)	-26.25 <sup>c</sup>	Tennessee	3.46	0.10	(19)	-2.47
Kentucky	3.04	0.12	(33)	-14.27	Texas	3.84	0.06	(13)	8.37
Louisiana	4.80	0.15	(02)	35.46 <sup>c</sup>	Utah	2.31	0.15	(50)	-34.83 <sup>c</sup>
Maine	2.82	0.19	(39)	-20.55 <sup>c</sup>	Vermont	2.26	0.25	(51)	-36.23 <sup>c</sup>
Maryland	3.77	0.12	(15)	6.37	Virginia	3.26	0.09	(25)	-8.05
Massachusetts	3.55	0.10	(18)	0.10	Washington	3.21	0.10	(28)	-9.43
Michigan	3.34	0.08	(23)	-5.67	West Virginia	2.78	0.16	(41)	-21.70 <sup>c</sup>
Minnesota	2.85	0.10	(38)	-19.73 <sup>c</sup>	Wisconsin	2.98	0.10	(35)	-15.99 <sup>c</sup>
Mississippi	4.14	0.17	(09)	16.75 <sup>c</sup>	Wyoming	2.44	0.30	(49)	-31.15 <sup>c</sup>
Missouri	2.87	0.09	(36)	-18.97 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.19  
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	4.86	0.03		
High Five States				
Hawaii	8.14	0.50	(01)	67.38 <sup>c</sup>
Louisiana	6.80	0.27	(02)	39.74 <sup>c</sup>
D.C.	6.26	0.74	(03)	28.79 <sup>c</sup>
Delaware	6.13	0.54	(04)	25.98 <sup>c</sup>
New Mexico	6.02	0.37	(05)	23.84 <sup>c</sup>
Low Five States				
Montana	3.54	0.38	(47)	-27.26 <sup>c</sup>
Iowa	3.51	0.21	(48)	-27.92 <sup>c</sup>
Vermont	3.24	0.47	(49)	-33.41 <sup>c</sup>
Utah	2.99	0.26	(50)	-38.45 <sup>c</sup>
Wyoming	2.75	0.47	(51)	-43.36 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.99	0.22	(19)	2.68	Montana	3.54	0.38	(47)	-27.26 <sup>c</sup>
Alaska	5.45	0.76	(12)	12.14	Nebraska	3.57	0.29	(46)	-26.63 <sup>c</sup>
Arizona	3.93	0.17	(38)	-19.20 <sup>c</sup>	Nevada	4.52	0.30	(24)	-6.97
Arkansas	4.18	0.25	(33)	-13.98	New Hampshire	3.82	0.35	(41)	-21.45 <sup>c</sup>
California	5.85	0.09	(07)	20.20 <sup>c</sup>	New Jersey	5.76	0.17	(09)	18.40 <sup>c</sup>
Colorado	3.85	0.21	(40)	-20.93 <sup>c</sup>	New Mexico	6.02	0.37	(05)	23.84 <sup>c</sup>
Connecticut	5.77	0.26	(08)	18.65 <sup>c</sup>	New York	5.71	0.12	(10)	17.42 <sup>c</sup>
Delaware	6.13	0.54	(04)	25.98 <sup>c</sup>	North Carolina	4.52	0.15	(25)	-7.07
D.C.	6.26	0.74	(03)	28.79 <sup>c</sup>	North Dakota	4.64	0.52	(23)	-4.55
Florida	4.12	0.09	(35)	-15.20 <sup>c</sup>	Ohio	4.48	0.13	(26)	-7.94
Georgia	4.97	0.18	(20)	2.28	Oklahoma	4.45	0.23	(27)	-8.51
Hawaii	8.14	0.50	(01)	67.38 <sup>c</sup>	Oregon	3.58	0.20	(45)	-26.48 <sup>c</sup>
Idaho	3.72	0.34	(43)	-23.51 <sup>c</sup>	Pennsylvania	4.79	0.12	(21)	-1.42
Illinois	5.25	0.14	(14)	7.91	Rhode Island	5.24	0.45	(15)	7.71
Indiana	3.88	0.17	(39)	-20.16 <sup>c</sup>	South Carolina	5.87	0.25	(06)	20.67 <sup>c</sup>
Iowa	3.51	0.21	(48)	-27.92 <sup>c</sup>	South Dakota	4.43	0.47	(28)	-8.93
Kansas	3.67	0.24	(44)	-24.60 <sup>c</sup>	Tennessee	5.03	0.20	(18)	3.43
Kentucky	4.25	0.22	(32)	-12.52	Texas	5.08	0.11	(17)	4.46
Louisiana	6.80	0.27	(02)	39.74 <sup>c</sup>	Utah	2.99	0.26	(50)	-38.45 <sup>c</sup>
Maine	4.17	0.35	(34)	-14.22	Vermont	3.24	0.47	(49)	-33.41 <sup>c</sup>
Maryland	5.25	0.22	(13)	7.93	Virginia	4.42	0.17	(29)	-9.15
Massachusetts	5.09	0.19	(16)	4.77	Washington	4.35	0.18	(30)	-10.55
Michigan	4.79	0.15	(22)	-1.60	West Virginia	3.78	0.28	(42)	-22.31 <sup>c</sup>
Minnesota	4.10	0.19	(36)	-15.71 <sup>c</sup>	Wisconsin	4.33	0.18	(31)	-10.86
Mississippi	5.61	0.31	(11)	15.36 <sup>c</sup>	Wyoming	2.75	0.47	(51)	-43.36 <sup>c</sup>
Missouri	3.95	0.17	(37)	-18.84 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 24.20  
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.55	0.02		
High Five States				
Hawaii	4.42	0.33	(01)	73.73 <sup>c</sup>
Alaska	3.47	0.56	(02)	36.42 <sup>c</sup>
California	3.34	0.06	(03)	31.29 <sup>c</sup>
Louisiana	3.29	0.16	(04)	29.15 <sup>c</sup>
Mississippi	3.12	0.19	(05)	22.64 <sup>c</sup>
Low Five States				
Utah	1.74	0.18	(47)	-31.67 <sup>c</sup>
Nebraska	1.72	0.18	(48)	-32.40 <sup>c</sup>
Montana	1.68	0.23	(49)	-34.15 <sup>c</sup>
Iowa	1.66	0.13	(50)	-34.86 <sup>c</sup>
Vermont	1.55	0.28	(51)	-39.07 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.73	0.14	(14)	7.17	Montana	1.68	0.23	(49)	-34.15 <sup>c</sup>
Alaska	3.47	0.56	(02)	36.42 <sup>c</sup>	Nebraska	1.72	0.18	(48)	-32.40 <sup>c</sup>
Arizona	2.24	0.11	(29)	-11.93	Nevada	2.50	0.20	(19)	-1.83
Arkansas	2.30	0.16	(24)	-9.61	New Hampshire	1.92	0.22	(41)	-24.74 <sup>c</sup>
California	3.34	0.06	(03)	31.29 <sup>c</sup>	New Jersey	3.02	0.10	(07)	18.61 <sup>c</sup>
Colorado	2.04	0.13	(37)	-19.79 <sup>c</sup>	New Mexico	2.84	0.23	(10)	11.58
Connecticut	2.56	0.15	(17)	0.48	New York	2.99	0.07	(08)	17.55 <sup>c</sup>
Delaware	2.54	0.31	(18)	-0.12	North Carolina	2.27	0.09	(26)	-10.78
D.C.	3.06	0.44	(06)	20.11 <sup>c</sup>	North Dakota	2.15	0.32	(33)	-15.45 <sup>c</sup>
Florida	2.28	0.06	(25)	-10.47	Ohio	2.25	0.08	(27)	-11.46
Georgia	2.84	0.11	(11)	11.38	Oklahoma	2.16	0.14	(32)	-15.24 <sup>c</sup>
Hawaii	4.42	0.33	(01)	73.73 <sup>c</sup>	Oregon	1.88	0.13	(43)	-26.05 <sup>c</sup>
Idaho	1.89	0.22	(42)	-25.77 <sup>c</sup>	Pennsylvania	2.24	0.07	(30)	-12.06
Illinois	2.76	0.09	(13)	8.39	Rhode Island	2.19	0.25	(31)	-13.96
Indiana	1.84	0.10	(45)	-27.56 <sup>c</sup>	South Carolina	2.67	0.14	(15)	4.84
Iowa	1.66	0.13	(50)	-34.86 <sup>c</sup>	South Dakota	2.61	0.32	(16)	2.35
Kansas	1.86	0.15	(44)	-27.03 <sup>c</sup>	Tennessee	2.33	0.11	(22)	-8.49
Kentucky	2.13	0.13	(34)	-16.44 <sup>c</sup>	Texas	2.88	0.07	(09)	12.98
Louisiana	3.29	0.16	(04)	29.15 <sup>c</sup>	Utah	1.74	0.18	(47)	-31.67 <sup>c</sup>
Maine	1.76	0.20	(46)	-31.06 <sup>c</sup>	Vermont	1.55	0.28	(51)	-39.07 <sup>c</sup>
Maryland	2.77	0.13	(12)	8.73	Virginia	2.37	0.10	(21)	-6.71
Massachusetts	2.41	0.11	(20)	-5.45	Washington	2.31	0.11	(23)	-9.23
Michigan	2.25	0.09	(28)	-11.50	West Virginia	2.06	0.18	(36)	-19.17 <sup>c</sup>
Minnesota	1.92	0.11	(40)	-24.55 <sup>c</sup>	Wisconsin	1.97	0.11	(39)	-22.52 <sup>c</sup>
Mississippi	3.12	0.19	(05)	22.64 <sup>c</sup>	Wyoming	2.09	0.38	(35)	-17.72 <sup>c</sup>
Missouri	2.04	0.11	(38)	-20.03 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 25.14  
Cancer of the Testis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.25	0.01		
High Five States				
Missouri	0.34	0.05	(01)	39.05 <sup>c</sup>
New Mexico	0.34	0.09	(02)	36.70 <sup>c</sup>
Arizona	0.33	0.05	(03)	32.77 <sup>c</sup>
Pennsylvania	0.32	0.03	(04)	30.19 <sup>c</sup>
California	0.32	0.02	(05)	29.74 <sup>c</sup>
Low Five States				
Georgia	0.17	0.03	(25)	-32.44 <sup>c</sup>
Virginia	0.16	0.03	(26)	-35.81 <sup>c</sup>
South Carolina	0.15	0.04	(27)	-37.68 <sup>c</sup>
Massachusetts	0.14	0.03	(28)	-42.97 <sup>c</sup>
Minnesota	0.14	0.03	(29)	-44.46 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.31	0.05	(06)	26.63 <sup>c</sup>	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	-	-	(-)	-
Arizona	0.33	0.05	(03)	32.77 <sup>c</sup>	Nevada	-	-	(-)	-
Arkansas	0.30	0.07	(09)	20.90 <sup>c</sup>	New Hampshire	-	-	(-)	-
California	0.32	0.02	(05)	29.74 <sup>c</sup>	New Jersey	0.20	0.03	(23)	-18.46 <sup>c</sup>
Colorado	0.22	0.04	(20)	-8.68	New Mexico	0.34	0.09	(02)	36.70 <sup>c</sup>
Connecticut	-	-	(-)	-	New York	0.21	0.02	(22)	-13.17
Delaware	-	-	(-)	-	North Carolina	0.31	0.04	(07)	24.59 <sup>c</sup>
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.24	0.02	(15)	-0.43	Ohio	0.23	0.03	(19)	-7.12
Georgia	0.17	0.03	(25)	-32.44 <sup>c</sup>	Oklahoma	0.30	0.06	(08)	22.79 <sup>c</sup>
Hawaii	-	-	(-)	-	Oregon	-	-	(-)	-
Idaho	-	-	(-)	-	Pennsylvania	0.32	0.03	(04)	30.19 <sup>c</sup>
Illinois	0.25	0.03	(13)	3.46	Rhode Island	-	-	(-)	-
Indiana	0.23	0.04	(17)	-5.11	South Carolina	0.15	0.04	(27)	-37.68 <sup>c</sup>
Iowa	-	-	(-)	-	South Dakota	-	-	(-)	-
Kansas	-	-	(-)	-	Tennessee	0.25	0.04	(14)	2.56
Kentucky	0.23	0.05	(18)	-5.19	Texas	0.26	0.02	(11)	6.23
Louisiana	0.17	0.04	(24)	-29.52 <sup>c</sup>	Utah	-	-	(-)	-
Maine	-	-	(-)	-	Vermont	-	-	(-)	-
Maryland	0.26	0.04	(12)	4.62	Virginia	0.16	0.03	(26)	-35.81 <sup>c</sup>
Massachusetts	0.14	0.03	(28)	-42.97 <sup>c</sup>	Washington	0.24	0.04	(16)	-3.90
Michigan	0.27	0.03	(10)	8.00	West Virginia	-	-	(-)	-
Minnesota	0.14	0.03	(29)	-44.46 <sup>c</sup>	Wisconsin	0.22	0.04	(21)	-11.32
Mississippi	-	-	(-)	-	Wyoming	-	-	(-)	-
Missouri	0.34	0.05	(01)	39.05 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 26.18  
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.51	0.01		
High Five States				
Vermont	0.77	0.15	(01)	51.22 <sup>c</sup>
Idaho	0.70	0.10	(02)	38.00 <sup>c</sup>
Hawaii	0.66	0.09	(03)	29.80 <sup>c</sup>
Delaware	0.63	0.11	(04)	24.92 <sup>c</sup>
California	0.61	0.02	(05)	20.19 <sup>c</sup>
Low Five States				
Nebraska	0.44	0.07	(45)	-13.74
Maryland	0.42	0.04	(46)	-16.52 <sup>c</sup>
Georgia	0.40	0.03	(47)	-21.13 <sup>c</sup>
South Carolina	0.40	0.04	(48)	-21.15 <sup>c</sup>
Montana	0.34	0.08	(49)	-33.38 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.44	0.04	(41)	-12.80	Montana	0.34	0.08	(49)	-33.38 <sup>c</sup>
Alaska	-	-	(-)	-	Nebraska	0.44	0.07	(45)	-13.74
Arizona	0.52	0.04	(20)	1.69	Nevada	0.44	0.06	(42)	-12.86
Arkansas	0.48	0.05	(32)	-5.84	New Hampshire	0.52	0.08	(22)	1.55
California	0.61	0.02	(05)	20.19 <sup>c</sup>	New Jersey	0.53	0.03	(16)	3.52
Colorado	0.60	0.05	(07)	18.76 <sup>c</sup>	New Mexico	0.54	0.07	(13)	6.40
Connecticut	0.45	0.05	(38)	-10.55	New York	0.53	0.02	(15)	4.30
Delaware	0.63	0.11	(04)	24.92 <sup>c</sup>	North Carolina	0.46	0.03	(37)	-8.99
D.C.	-	-	(-)	-	North Dakota	0.44	0.11	(44)	-13.65
Florida	0.44	0.02	(43)	-13.30	Ohio	0.52	0.03	(19)	2.48
Georgia	0.40	0.03	(47)	-21.13 <sup>c</sup>	Oklahoma	0.49	0.05	(27)	-2.94
Hawaii	0.66	0.09	(03)	29.80 <sup>c</sup>	Oregon	0.48	0.05	(31)	-5.34
Idaho	0.70	0.10	(02)	38.00 <sup>c</sup>	Pennsylvania	0.51	0.03	(23)	1.11
Illinois	0.49	0.03	(28)	-2.98	Rhode Island	0.55	0.09	(11)	9.00
Indiana	0.48	0.04	(35)	-6.28	South Carolina	0.40	0.04	(48)	-21.15 <sup>c</sup>
Iowa	0.50	0.05	(26)	-1.89	South Dakota	0.58	0.11	(09)	13.64
Kansas	0.58	0.06	(08)	13.88	Tennessee	0.51	0.04	(24)	0.54
Kentucky	0.51	0.05	(25)	-0.53	Texas	0.52	0.02	(18)	3.02
Louisiana	0.48	0.05	(30)	-5.21	Utah	0.54	0.08	(14)	5.65
Maine	0.48	0.07	(33)	-5.97	Vermont	0.77	0.15	(01)	51.22 <sup>c</sup>
Maryland	0.42	0.04	(46)	-16.52 <sup>c</sup>	Virginia	0.45	0.03	(39)	-10.69
Massachusetts	0.52	0.04	(21)	1.61	Washington	0.56	0.04	(10)	9.34
Michigan	0.48	0.03	(34)	-6.11	West Virginia	0.49	0.07	(29)	-3.03
Minnesota	0.52	0.04	(17)	3.13	Wisconsin	0.55	0.04	(12)	7.67
Mississippi	0.45	0.05	(40)	-11.75	Wyoming	0.61	0.15	(06)	19.40 <sup>c</sup>
Missouri	0.47	0.04	(36)	-6.72					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 26.19  
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.50	0.01		
High Five States				
Kansas	0.73	0.11	(01)	46.62 <sup>c</sup>
Colorado	0.67	0.09	(02)	34.78 <sup>c</sup>
Utah	0.67	0.13	(03)	34.28 <sup>c</sup>
Idaho	0.66	0.14	(04)	32.71 <sup>c</sup>
New Hampshire	0.66	0.15	(05)	32.47 <sup>c</sup>
Low Five States				
Nevada	0.40	0.08	(36)	-20.48 <sup>c</sup>
Arkansas	0.39	0.08	(37)	-20.68 <sup>c</sup>
West Virginia	0.39	0.09	(38)	-21.17 <sup>c</sup>
Illinois	0.38	0.04	(39)	-23.37 <sup>c</sup>
Mississippi	0.35	0.07	(40)	-29.59 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.48	0.07	(23)	-4.15	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.49	0.11	(19)	-0.56
Arizona	0.50	0.06	(17)	0.11	Nevada	0.40	0.08	(36)	-20.48 <sup>c</sup>
Arkansas	0.39	0.08	(37)	-20.68 <sup>c</sup>	New Hampshire	0.66	0.15	(05)	32.47 <sup>c</sup>
California	0.55	0.03	(09)	10.61	New Jersey	0.54	0.05	(11)	9.39
Colorado	0.67	0.09	(02)	34.78 <sup>c</sup>	New Mexico	0.53	0.11	(13)	5.64
Connecticut	0.45	0.07	(27)	-8.73	New York	0.53	0.04	(12)	7.13
Delaware	-	-	(-)	-	North Carolina	0.45	0.05	(30)	-10.48
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.44	0.03	(32)	-11.40	Ohio	0.58	0.05	(07)	17.08 <sup>c</sup>
Georgia	0.43	0.05	(34)	-12.97	Oklahoma	0.50	0.08	(18)	-0.15
Hawaii	-	-	(-)	-	Oregon	0.45	0.07	(29)	-10.39
Idaho	0.66	0.14	(04)	32.71 <sup>c</sup>	Pennsylvania	0.48	0.04	(22)	-4.04
Illinois	0.38	0.04	(39)	-23.37 <sup>c</sup>	Rhode Island	-	-	(-)	-
Indiana	0.48	0.06	(21)	-3.92	South Carolina	0.43	0.07	(35)	-13.10
Iowa	0.59	0.09	(06)	19.39 <sup>c</sup>	South Dakota	-	-	(-)	-
Kansas	0.73	0.11	(01)	46.62 <sup>c</sup>	Tennessee	0.49	0.06	(20)	-0.73
Kentucky	0.45	0.07	(31)	-10.51	Texas	0.52	0.03	(15)	4.49
Louisiana	0.45	0.07	(28)	-10.36	Utah	0.67	0.13	(03)	34.28 <sup>c</sup>
Maine	-	-	(-)	-	Vermont	-	-	(-)	-
Maryland	0.46	0.06	(26)	-6.81	Virginia	0.54	0.06	(10)	9.48
Massachusetts	0.51	0.06	(16)	2.64	Washington	0.57	0.06	(08)	14.77
Michigan	0.44	0.04	(33)	-12.36	West Virginia	0.39	0.09	(38)	-21.17 <sup>c</sup>
Minnesota	0.52	0.06	(14)	5.30	Wisconsin	0.47	0.06	(24)	-4.99
Mississippi	0.35	0.07	(40)	-29.59 <sup>c</sup>	Wyoming	-	-	(-)	-
Missouri	0.47	0.06	(25)	-6.39					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.




Table 26.20  
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.51	0.01		
High Five States				
Vermont	1.03	0.24	(01)	102.34 <sup>c</sup>
Hawaii	0.89	0.15	(02)	74.17 <sup>c</sup>
Idaho	0.73	0.14	(03)	43.98 <sup>c</sup>
California	0.66	0.03	(04)	28.57 <sup>c</sup>
Delaware	0.62	0.15	(05)	22.08 <sup>c</sup>
Low Five States				
Maryland	0.39	0.05	(41)	-23.08 <sup>c</sup>
Virginia	0.39	0.04	(42)	-23.93 <sup>c</sup>
Georgia	0.37	0.04	(43)	-27.77 <sup>c</sup>
South Carolina	0.37	0.05	(44)	-28.20 <sup>c</sup>
Nebraska	0.36	0.08	(45)	-29.78 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.40	0.05	(40)	-20.88 <sup>c</sup>	Montana	-	-	(-)	-
Alaska	-	-	(-)	-	Nebraska	0.36	0.08	(45)	-29.78 <sup>c</sup>
Arizona	0.53	0.05	(16)	2.96	Nevada	0.47	0.09	(30)	-7.12
Arkansas	0.53	0.08	(11)	3.98	New Hampshire	0.42	0.10	(39)	-18.44 <sup>c</sup>
California	0.66	0.03	(04)	28.57 <sup>c</sup>	New Jersey	0.51	0.04	(20)	0.76
Colorado	0.53	0.07	(12)	3.79	New Mexico	0.53	0.10	(10)	4.14
Connecticut	0.46	0.06	(34)	-10.55	New York	0.53	0.03	(14)	3.54
Delaware	0.62	0.15	(05)	22.08 <sup>c</sup>	North Carolina	0.47	0.04	(31)	-7.20
D.C.	-	-	(-)	-	North Dakota	-	-	(-)	-
Florida	0.44	0.03	(36)	-14.52	Ohio	0.47	0.04	(32)	-8.44
Georgia	0.37	0.04	(43)	-27.77 <sup>c</sup>	Oklahoma	0.50	0.07	(27)	-2.75
Hawaii	0.89	0.15	(02)	74.17 <sup>c</sup>	Oregon	0.51	0.07	(22)	0.46
Idaho	0.73	0.14	(03)	43.98 <sup>c</sup>	Pennsylvania	0.54	0.04	(09)	5.49
Illinois	0.57	0.04	(07)	11.83	Rhode Island	0.50	0.11	(26)	-1.34
Indiana	0.45	0.05	(35)	-11.53	South Carolina	0.37	0.05	(44)	-28.20 <sup>c</sup>
Iowa	0.42	0.07	(38)	-17.23 <sup>c</sup>	South Dakota	-	-	(-)	-
Kansas	0.47	0.07	(29)	-7.07	Tennessee	0.51	0.05	(21)	0.72
Kentucky	0.54	0.07	(08)	6.07	Texas	0.52	0.03	(19)	1.21
Louisiana	0.50	0.06	(25)	-1.09	Utah	0.42	0.09	(37)	-16.98 <sup>c</sup>
Maine	0.53	0.10	(17)	2.90	Vermont	1.03	0.24	(01)	102.34 <sup>c</sup>
Maryland	0.39	0.05	(41)	-23.08 <sup>c</sup>	Virginia	0.39	0.04	(42)	-23.93 <sup>c</sup>
Massachusetts	0.52	0.05	(18)	1.92	Washington	0.53	0.05	(15)	3.53
Michigan	0.49	0.04	(28)	-3.81	West Virginia	0.53	0.09	(13)	3.68
Minnesota	0.51	0.06	(24)	-0.46	Wisconsin	0.61	0.06	(06)	19.19 <sup>c</sup>
Mississippi	0.51	0.08	(23)	-0.07	Wyoming	-	-	(-)	-
Missouri	0.46	0.05	(33)	-9.63					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 27.18  
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.41	0.02		
High Five States				
Maine	6.12	0.27	(01)	38.79 <sup>c</sup>
Vermont	5.62	0.40	(02)	27.39 <sup>c</sup>
Nevada	5.42	0.22	(03)	22.93 <sup>c</sup>
Massachusetts	5.19	0.12	(04)	17.74 <sup>c</sup>
Rhode Island	5.10	0.29	(05)	15.60 <sup>c</sup>
Low Five States				
Utah	3.62	0.20	(47)	-17.87 <sup>c</sup>
Mississippi	3.58	0.16	(48)	-18.81 <sup>c</sup>
Alaska	3.49	0.44	(49)	-20.84 <sup>c</sup>
New Mexico	3.39	0.18	(50)	-23.15 <sup>c</sup>
Hawaii	2.78	0.19	(51)	-37.09 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.82	0.12	(46)	-13.38	Montana	4.57	0.28	(21)	3.62
Alaska	3.49	0.44	(49)	-20.84 <sup>c</sup>	Nebraska	4.04	0.20	(39)	-8.54
Arizona	3.99	0.11	(41)	-9.57	Nevada	5.42	0.22	(03)	22.93 <sup>c</sup>
Arkansas	4.38	0.17	(27)	-0.79	New Hampshire	5.01	0.27	(07)	13.60
California	4.02	0.05	(40)	-8.96	New Jersey	4.93	0.10	(08)	11.73
Colorado	3.85	0.14	(43)	-12.84	New Mexico	3.39	0.18	(50)	-23.15 <sup>c</sup>
Connecticut	4.77	0.15	(13)	8.19	New York	4.48	0.07	(25)	1.65
Delaware	4.91	0.32	(09)	11.24	North Carolina	4.20	0.10	(34)	-4.69
D.C.	4.20	0.39	(35)	-4.74	North Dakota	4.65	0.34	(18)	5.38
Florida	4.51	0.06	(23)	2.24	Ohio	5.09	0.09	(06)	15.37 <sup>c</sup>
Georgia	4.18	0.11	(36)	-5.21	Oklahoma	4.29	0.15	(30)	-2.84
Hawaii	2.78	0.19	(51)	-37.09 <sup>c</sup>	Oregon	4.63	0.15	(19)	5.02
Idaho	4.32	0.24	(28)	-2.10	Pennsylvania	4.81	0.08	(11)	8.99
Illinois	4.49	0.08	(24)	1.89	Rhode Island	5.10	0.29	(05)	15.60 <sup>c</sup>
Indiana	4.63	0.12	(20)	4.96	South Carolina	4.14	0.13	(37)	-6.23
Iowa	4.51	0.16	(22)	2.30	South Dakota	3.90	0.29	(42)	-11.54
Kansas	3.82	0.16	(45)	-13.35	Tennessee	4.28	0.12	(31)	-3.03
Kentucky	4.74	0.15	(14)	7.45	Texas	3.85	0.06	(44)	-12.84
Louisiana	4.25	0.14	(32)	-3.60	Utah	3.62	0.20	(47)	-17.87 <sup>c</sup>
Maine	6.12	0.27	(01)	38.79 <sup>c</sup>	Vermont	5.62	0.40	(02)	27.39 <sup>c</sup>
Maryland	4.79	0.13	(12)	8.67	Virginia	4.23	0.11	(33)	-4.19
Massachusetts	5.19	0.12	(04)	17.74 <sup>c</sup>	Washington	4.72	0.12	(15)	7.01
Michigan	4.82	0.09	(10)	9.27	West Virginia	4.66	0.20	(17)	5.61
Minnesota	4.10	0.12	(38)	-7.13	Wisconsin	4.66	0.12	(16)	5.65
Mississippi	3.58	0.16	(48)	-18.81 <sup>c</sup>	Wyoming	4.43	0.40	(26)	0.38
Missouri	4.31	0.11	(29)	-2.35					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 27.19  
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.69	0.04		
High Five States				
Maine	10.54	0.56	(01)	37.02 <sup>c</sup>
Vermont	9.73	0.83	(02)	26.48 <sup>c</sup>
Massachusetts	9.22	0.25	(03)	19.90 <sup>c</sup>
Rhode Island	9.07	0.61	(04)	17.98 <sup>c</sup>
Ohio	9.04	0.19	(05)	17.56 <sup>c</sup>
Low Five States				
Mississippi	6.50	0.35	(47)	-15.45 <sup>c</sup>
Utah	6.40	0.40	(48)	-16.80 <sup>c</sup>
New Mexico	5.99	0.38	(49)	-22.17 <sup>c</sup>
Alaska	5.57	0.85	(50)	-27.61 <sup>c</sup>
Hawaii	4.78	0.38	(51)	-37.88 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.66	0.27	(44)	-13.40	Montana	8.28	0.59	(16)	7.61
Alaska	5.57	0.85	(50)	-27.61 <sup>c</sup>	Nebraska	6.95	0.41	(39)	-9.61
Arizona	6.95	0.23	(38)	-9.59	Nevada	8.77	0.44	(06)	14.09
Arkansas	7.34	0.34	(34)	-4.51	New Hampshire	8.73	0.56	(07)	13.58
California	6.83	0.10	(41)	-11.14	New Jersey	8.62	0.22	(08)	12.11
Colorado	6.73	0.28	(43)	-12.52	New Mexico	5.99	0.38	(49)	-22.17 <sup>c</sup>
Connecticut	8.31	0.32	(14)	8.11	New York	7.96	0.14	(21)	3.44
Delaware	8.30	0.65	(15)	7.94	North Carolina	7.36	0.21	(33)	-4.28
D.C.	6.52	0.79	(46)	-15.21 <sup>c</sup>	North Dakota	8.22	0.69	(17)	6.94
Florida	7.65	0.12	(26)	-0.57	Ohio	9.04	0.19	(05)	17.56 <sup>c</sup>
Georgia	7.51	0.24	(28)	-2.34	Oklahoma	7.55	0.31	(27)	-1.81
Hawaii	4.78	0.38	(51)	-37.88 <sup>c</sup>	Oregon	7.93	0.30	(22)	3.15
Idaho	7.43	0.49	(31)	-3.39	Pennsylvania	8.52	0.16	(10)	10.79
Illinois	7.87	0.18	(23)	2.37	Rhode Island	9.07	0.61	(04)	17.98 <sup>c</sup>
Indiana	8.07	0.25	(19)	4.90	South Carolina	6.94	0.28	(40)	-9.71
Iowa	8.34	0.33	(13)	8.40	South Dakota	7.03	0.60	(37)	-8.57
Kansas	6.63	0.33	(45)	-13.75	Tennessee	7.72	0.26	(25)	0.39
Kentucky	8.51	0.32	(11)	10.65	Texas	6.75	0.13	(42)	-12.19
Louisiana	7.23	0.30	(36)	-6.04	Utah	6.40	0.40	(48)	-16.80 <sup>c</sup>
Maine	10.54	0.56	(01)	37.02 <sup>c</sup>	Vermont	9.73	0.83	(02)	26.48 <sup>c</sup>
Maryland	8.34	0.28	(12)	8.43	Virginia	7.39	0.23	(32)	-3.88
Massachusetts	9.22	0.25	(03)	19.90 <sup>c</sup>	Washington	8.05	0.25	(20)	4.67
Michigan	8.58	0.20	(09)	11.52	West Virginia	7.76	0.41	(24)	0.90
Minnesota	7.24	0.26	(35)	-5.90	Wisconsin	8.21	0.26	(18)	6.80
Mississippi	6.50	0.35	(47)	-15.45 <sup>c</sup>	Wyoming	7.44	0.80	(30)	-3.27
Missouri	7.47	0.24	(29)	-2.87					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.


Table 27.20  
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.22	0.02		
High Five States				
Maine	3.03	0.25	(01)	36.89 <sup>c</sup>
Nevada	2.84	0.22	(02)	28.07 <sup>c</sup>
D.C.	2.77	0.41	(03)	25.16 <sup>c</sup>
Vermont	2.69	0.36	(04)	21.47 <sup>c</sup>
Massachusetts	2.63	0.11	(05)	18.70 <sup>c</sup>
Low Five States				
Arizona	1.75	0.10	(47)	-20.82 <sup>c</sup>
South Dakota	1.66	0.24	(48)	-25.00 <sup>c</sup>
Utah	1.52	0.17	(49)	-31.22 <sup>c</sup>
New Mexico	1.46	0.16	(50)	-33.94 <sup>c</sup>
Hawaii	1.30	0.18	(51)	-41.20 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.99	0.12	(39)	-10.22	Montana	1.87	0.24	(43)	-15.62 <sup>c</sup>
Alaska	1.77	0.43	(46)	-19.95 <sup>c</sup>	Nebraska	2.10	0.19	(33)	-5.47
Arizona	1.75	0.10	(47)	-20.82 <sup>c</sup>	Nevada	2.84	0.22	(02)	28.07 <sup>c</sup>
Arkansas	2.34	0.16	(17)	5.77	New Hampshire	2.53	0.25	(09)	13.95
California	2.07	0.05	(35)	-6.82	New Jersey	2.60	0.10	(06)	17.34 <sup>c</sup>
Colorado	1.81	0.12	(44)	-18.54 <sup>c</sup>	New Mexico	1.46	0.16	(50)	-33.94 <sup>c</sup>
Connecticut	2.46	0.14	(12)	11.22	New York	2.27	0.06	(24)	2.62
Delaware	2.41	0.29	(13)	8.94	North Carolina	2.25	0.09	(25)	1.67
D.C.	2.77	0.41	(03)	25.16 <sup>c</sup>	North Dakota	2.07	0.30	(34)	-6.81
Florida	2.19	0.06	(27)	-1.09	Ohio	2.50	0.08	(10)	12.73
Georgia	2.14	0.10	(29)	-3.39	Oklahoma	2.04	0.13	(36)	-7.75
Hawaii	1.30	0.18	(51)	-41.20 <sup>c</sup>	Oregon	2.22	0.13	(26)	0.38
Idaho	1.95	0.22	(40)	-12.22	Pennsylvania	2.41	0.07	(14)	8.92
Illinois	2.33	0.08	(20)	5.29	Rhode Island	2.59	0.26	(08)	16.81 <sup>c</sup>
Indiana	2.38	0.11	(16)	7.33	South Carolina	2.31	0.13	(22)	4.03
Iowa	2.00	0.14	(37)	-9.92	South Dakota	1.66	0.24	(48)	-25.00 <sup>c</sup>
Kansas	1.99	0.15	(38)	-10.05	Tennessee	2.10	0.11	(31)	-5.21
Kentucky	2.29	0.13	(23)	3.30	Texas	1.90	0.06	(42)	-14.31
Louisiana	2.34	0.14	(18)	5.57	Utah	1.52	0.17	(49)	-31.22 <sup>c</sup>
Maine	3.03	0.25	(01)	36.89 <sup>c</sup>	Vermont	2.69	0.36	(04)	21.47 <sup>c</sup>
Maryland	2.59	0.13	(07)	17.07 <sup>c</sup>	Virginia	2.13	0.10	(30)	-3.91
Massachusetts	2.63	0.11	(05)	18.70 <sup>c</sup>	Washington	2.40	0.12	(15)	8.26
Michigan	2.33	0.09	(19)	5.31	West Virginia	2.49	0.19	(11)	12.23
Minnesota	1.92	0.11	(41)	-13.49	Wisconsin	2.33	0.11	(21)	4.96
Mississippi	1.78	0.14	(45)	-19.52 <sup>c</sup>	Wyoming	2.10	0.37	(32)	-5.37
Missouri	2.17	0.11	(28)	-1.91					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.