

Table 1.10
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
Blacks, 2004-2013

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
All Sites	-1.3*	-2.4*	-0.3	-2.1*	-2.7*	-1.6*
Oral Cavity & Pharynx:	-2.0*	-2.0*	-2.1*	-2.9*	-3.4*	-1.8
Lip	-	-	-	-	-	-
Tongue	-1.9*	-1.3	-3.7*	-3.1*	-4.2*	-0.7
Salivary gland	-1.2	-2.1	-0.4	2.7	2.7	3.1
Floor of mouth	-2.9	-3.1	-2.0	-	-	-
Gum & other oral cavity	-2.3*	-2.2	-2.6*	-3.9*	-4.3*	-3.8*
Nasopharynx	-1.9	-2.6	-0.5	-4.1*	-5.3*	-2.2
Tonsil	-0.5	-0.5	0.1	-2.5*	-2.3*	-2.7
Oropharynx	-1.4	-2.3	-	-0.1	-0.5	0.7
Hypopharynx	-4.2*	-4.1*	-5.5	-2.5	-3.2	-
Other oral cavity & pharynx	-3.5	-2.6	-	-4.4*	-4.3*	-4.7*
Digestive System:	-1.6*	-1.7*	-1.7*	-1.6*	-1.5*	-1.8*
Esophagus	-4.0*	-4.5*	-3.4*	-4.4*	-4.6*	-4.4*
Stomach	-2.5*	-3.4*	-1.7	-3.1*	-3.4*	-3.0*
Small intestine	3.7*	3.3*	4.2*	0.5	0.5	0.5
Colon & Rectum:	-2.9*	-3.0*	-3.0*	-3.0*	-2.7*	-3.5*
Colon	-3.2*	-3.5*	-3.2*	-	-	-
Rectum	-2.0*	-1.7*	-2.5*	-	-	-
Anus, anal canal & anorectum	1.9*	2.1	1.5	3.8*	7.0*	0.3
Liver & intrahepatic bile duct	2.6*	2.6*	2.3*	2.5*	2.7*	2.0*
Gallbladder	0.6	1.4	0.5	1.1	2.0	0.8
Other biliary	0.4	-0.2	0.7	0.2	-1.2	1.2
Pancreas	-0.6	-0.3	-0.8	-0.4*	-0.5	-0.3
Retroperitoneum	1.9	-	2.8	-5.9	-	-
Peritoneum, omentum & mesentery	1.9	-	2.4	-0.1	-	0.5
Other digestive system	3.7*	3.4	3.9	2.7	2.9	2.3
Respiratory System:	-2.4*	-3.1*	-1.6*	-2.7*	-3.4*	-1.9*
Nose, nasal cavity & middle ear	-0.9	-1.6	0.2	-3.0	-5.2	-0.7
Larynx	-3.1*	-3.4*	-2.0	-3.7*	-4.0*	-2.7*
Lung & bronchus	-2.4*	-3.1*	-1.5*	-2.7*	-3.4*	-1.9*
Pleura	-	-	-	-	-	-
Trachea & other respiratory organs	-3.9	-	-	0.6	-	-
Bones & joints	-0.6	1.5	-2.1	-1.4	-1.2	-2.2
Soft tissue (including heart)	-0.6	-1.3	0.2	0.9	0.7	0.9
Skin (excl. basal & squamous):	-1.4	-1.6	-1.5	-2.1*	-2.1*	-2.0
Melanoma of the skin	-1.1	-2.8	-0.2	-1.7*	-2.1	-1.8
Other non-epithelial skin	-1.6	-0.4	-2.8	-2.4*	-2.1*	-2.6
Breast	0.5	3.6	0.6*	-1.5*	-0.1	-1.4*
Breast (<i>in situ</i>)	1.6*	-	1.7*	-	-	-

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

* The APC is significantly different from zero (p<.05).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.10 - continued
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
Blacks, 2004-2013

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
Female Genital System:	0.5	-	0.7	-0.5*	-	-0.3
Cervix uteri	-2.6*	-	-2.5*	-2.2*	-	-1.9*
Corpus uteri	2.7*	-	2.9*	0.2	-	0.5
Uterus, NOS	2.2	-	2.6	2.6*	-	2.8*
Ovary ^c	-1.4*	-	-1.1*	-2.0*	-	-1.8*
Vagina	-2.2	-	-2.5	-2.4*	-	-2.2*
Vulva	-0.7	-	-0.7	0.0	-	0.4
Other female genital system	5.2*	-	5.3*	4.0	-	3.8
Male Genital System:	-3.3*	-3.7*	-	-3.7*	-4.3*	-
Prostate	-3.4*	-3.8*	-	-3.7*	-4.3*	-
Testis	3.6*	3.6*	-	-4.4	-4.0	-
Penis	-0.2	-0.9	-	0.0	-0.3	-
Other male genital system	-	-	-	-	-	-
Urinary System:	0.4	0.2	0.4	-1.2*	-1.0*	-1.7*
Urinary bladder	-1.0*	-0.8	-1.9*	-0.9*	-0.6	-1.7*
Kidney & renal pelvis	1.6*	1.1	2.0*	-1.4*	-1.4*	-1.5*
Ureter	-4.9*	-	-	-0.2	-	-
Other urinary system	-1.1	-	-	-2.9	-	-5.0*
Eye & Orbit	2.0	-	-	-	-	-
Brain & Nervous System: ^d	0.2	1.4*	-1.0	0.4	0.7	0.1
Brain	0.7	1.9*	-0.5	-	-	-
Cranial nerves & other nervous system	-4.5	-3.9	-4.9*	-	-	-
Endocrine System:	4.0*	1.9	4.6*	0.2	2.0	-0.8
Thyroid	4.5*	2.5*	5.0*	0.5	1.4	0.0
Other endocrine & thymus	0.3	-0.2	0.7	0.0	2.5	-2.2
Lymphoma:	-0.7*	-1.0*	-0.5	-2.0*	-1.8*	-2.1*
Hodgkin lymphoma	-0.8*	-0.2	-1.6	-3.6*	-3.4	-3.7*
Non-Hodgkin lymphoma	-0.7	-1.1	-0.3	-1.9*	-1.6*	-2.0*
Myeloma	1.0*	0.7	1.2	-0.8*	-1.3*	-0.5
Leukemia:	0.4	0.2	0.4	-1.5*	-1.7*	-1.4*
Lymphocytic:	0.0	-0.4	0.0	-2.4*	-3.4*	-1.2
Acute lymphocytic	0.7	0.5	0.7	-0.8	-1.6	0.8
Chronic lymphocytic	-0.1	-0.6	-0.2	-2.7*	-3.3*	-1.9*
Other lymphocytic	-1.1	-	-	-5.2*	-9.7*	-
Myeloid & Monocytic:	1.6*	1.8*	1.3	-0.7	-0.1	-1.3
Acute myeloid	3.1*	3.8*	2.5	-0.1	0.4	-0.7
Chronic myeloid	-0.3	-1.4	0.4	-4.3*	-4.2*	-4.0
Acute monocytic	-	-	-	-	-	-
Other myeloid & monocytic	-	-	-	-0.9	1.9	-5.0
Other leukemia:	-5.4*	-5.9	-4.7	-1.8*	-2.1*	-1.6
Other acute leukemia	-3.7	-	-	-4.5*	-5.4*	-3.6*
Aleukemic, subleukemic & NOS	-6.4	-	-4.3	-0.7	-0.6	-0.8
Kaposi Sarcoma ^e	-1.4	-1.2	-	-	-	-
Mesothelioma ^e	-4.3*	-5.2*	-	-	-	-
Ill-defined & unspecified	-3.4*	-3.7*	-3.0*	-3.0*	-3.2*	-2.9*

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^d Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

^e Trend not shown for mortality. Category did not exist in mortality coding until 1999.

* The APC is significantly different from zero (p<.05).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.