

Table VII-4
CORPUS AND UTERUS, NOS CANCER (Invasive)

SEER INCIDENCE^a AND U.S. DEATH^b RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	Incidence ^a			US Death ^b		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<u>AGE-ADJUSTED RATES, 2001-2005</u>						
AGE AT DIAGNOSIS/DEATH:						
All ages	23.4	24.3	20.3	4.1	3.9	7.1
Under 65	14.7	15.3	10.9	1.4	1.3	2.3
65 and over	82.9	86.6	85.3	22.7	21.3	40.0
All ages(IARC world std) ^c	17.1	17.8	14.6	2.6	2.4	4.5
<u>AGE-SPECIFIC RATES, 2001-2005</u>						
AGE AT DIAGNOSIS/DEATH:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	0.4	0.3	-	-	-	-
25-29	1.3	1.2	1.4	0.1	0.0	-
30-34	3.6	3.5	3.3	0.2	0.2	0.3
35-39	7.1	6.8	5.9	0.4	0.4	0.5
40-44	12.8	13.1	9.0	0.9	0.8	1.2
45-49	23.2	23.7	13.2	1.8	1.7	2.2
50-54	43.1	45.8	25.1	3.7	3.5	5.1
55-59	68.0	71.8	46.2	6.8	6.5	10.2
60-64	82.9	86.4	80.9	11.4	10.2	22.7
65-69	90.4	94.6	97.0	16.1	14.3	33.7
70-74	85.4	89.5	90.8	20.5	18.8	40.4
75-79	84.6	88.5	88.6	23.7	22.3	42.7
80-84	80.2	84.1	69.9	28.5	27.6	43.9
85+	61.1	62.8	60.2	33.5	33.2	44.0

^a SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^c Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.