


Table 2.18  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	183.77	0.11		
High Five States				
Kentucky	216.54 <sup>b</sup>	1.00	(01)	17.83 <sup>c</sup>
Louisiana	211.47 <sup>b</sup>	0.99	(02)	15.07 <sup>c</sup>
West Virginia	209.83 <sup>b</sup>	1.38	(03)	14.18 <sup>c</sup>
Mississippi	207.73 <sup>b</sup>	1.20	(04)	13.04 <sup>c</sup>
Tennessee	206.04 <sup>b</sup>	0.82	(05)	12.12 <sup>c</sup>
Low Five States				
New Mexico	162.18 <sup>b</sup>	1.30	(47)	-11.75 <sup>c</sup>
Colorado	159.83 <sup>b</sup>	0.90	(48)	-13.03 <sup>c</sup>
Arizona	159.58 <sup>b</sup>	0.72	(49)	-13.17 <sup>c</sup>
Hawaii	150.77 <sup>b</sup>	1.46	(50)	-17.96 <sup>c</sup>
Utah	135.75 <sup>b</sup>	1.22	(51)	-26.13 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	201.52 <sup>b</sup>	0.91	(07)	9.66 <sup>c</sup>	Montana	179.27	1.85	(36)	-2.45
Alaska	180.85	3.15	(33)	-1.59	Nebraska	176.49 <sup>b</sup>	1.37	(39)	-3.96
Arizona	159.58 <sup>b</sup>	0.72	(49)	-13.17 <sup>c</sup>	Nevada	189.10 <sup>b</sup>	1.32	(20)	2.90
Arkansas	202.63 <sup>b</sup>	1.15	(06)	10.27 <sup>c</sup>	New Hampshire	187.81	1.68	(23)	2.20
California	166.86 <sup>b</sup>	0.32	(46)	-9.20 <sup>c</sup>	New Jersey	186.07 <sup>b</sup>	0.63	(27)	1.25
Colorado	159.83 <sup>b</sup>	0.90	(48)	-13.03 <sup>c</sup>	New Mexico	162.18 <sup>b</sup>	1.30	(47)	-11.75 <sup>c</sup>
Connecticut	179.34 <sup>b</sup>	0.96	(35)	-2.41	New York	171.94 <sup>b</sup>	0.41	(44)	-6.44 <sup>c</sup>
Delaware	198.42 <sup>b</sup>	2.10	(12)	7.97 <sup>c</sup>	North Carolina	192.35 <sup>b</sup>	0.67	(17)	4.67
D.C.	198.72 <sup>b</sup>	2.63	(11)	8.14 <sup>c</sup>	North Dakota	173.31 <sup>b</sup>	2.18	(43)	-5.69 <sup>c</sup>
Florida	174.08 <sup>b</sup>	0.39	(41)	-5.28 <sup>c</sup>	Ohio	200.16 <sup>b</sup>	0.57	(08)	8.92 <sup>c</sup>
Georgia	187.11 <sup>b</sup>	0.71	(25)	1.82	Oklahoma	196.94 <sup>b</sup>	1.02	(14)	7.16 <sup>c</sup>
Hawaii	150.77 <sup>b</sup>	1.46	(50)	-17.96 <sup>c</sup>	Oregon	185.78	0.98	(28)	1.09
Idaho	170.43 <sup>b</sup>	1.59	(45)	-7.26 <sup>c</sup>	Pennsylvania	193.81 <sup>b</sup>	0.51	(16)	5.46 <sup>c</sup>
Illinois	191.33 <sup>b</sup>	0.55	(18)	4.12	Rhode Island	187.47	1.77	(24)	2.02
Indiana	199.40 <sup>b</sup>	0.79	(10)	8.51 <sup>c</sup>	South Carolina	194.15 <sup>b</sup>	0.94	(15)	5.65 <sup>c</sup>
Iowa	182.07	1.03	(30)	-0.93	South Dakota	176.73	2.00	(38)	-3.83
Kansas	182.71	1.13	(29)	-0.58	Tennessee	206.04 <sup>b</sup>	0.82	(05)	12.12 <sup>c</sup>
Kentucky	216.54 <sup>b</sup>	1.00	(01)	17.83 <sup>c</sup>	Texas	177.54 <sup>b</sup>	0.43	(37)	-3.39
Louisiana	211.47 <sup>b</sup>	0.99	(02)	15.07 <sup>c</sup>	Utah	135.75 <sup>b</sup>	1.22	(51)	-26.13 <sup>c</sup>
Maine	199.69 <sup>b</sup>	1.60	(09)	8.66 <sup>c</sup>	Vermont	179.74	2.30	(34)	-2.19
Maryland	188.81 <sup>b</sup>	0.84	(21)	2.74	Virginia	188.63 <sup>b</sup>	0.72	(22)	2.64
Massachusetts	186.62 <sup>b</sup>	0.73	(26)	1.55	Washington	181.21	0.77	(32)	-1.39
Michigan	190.99 <sup>b</sup>	0.61	(19)	3.93	West Virginia	209.83 <sup>b</sup>	1.38	(03)	14.18 <sup>c</sup>
Minnesota	173.68 <sup>b</sup>	0.82	(42)	-5.49 <sup>c</sup>	Wisconsin	182.05	0.79	(31)	-0.94
Mississippi	207.73 <sup>b</sup>	1.20	(04)	13.04 <sup>c</sup>	Wyoming	176.28	2.63	(40)	-4.08
Missouri	197.47 <sup>b</sup>	0.80	(13)	7.45 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 2.19  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	225.44	0.19		
High Five States				
Mississippi	276.31 <sup>b</sup>	2.18	(01)	22.57 <sup>c</sup>
Kentucky	275.03 <sup>b</sup>	1.78	(02)	22.00 <sup>c</sup>
Louisiana	270.69 <sup>b</sup>	1.77	(03)	20.07 <sup>c</sup>
Tennessee	263.99 <sup>b</sup>	1.46	(04)	17.10 <sup>c</sup>
Alabama	263.80 <sup>b</sup>	1.64	(05)	17.02 <sup>c</sup>
Low Five States				
New Mexico	194.76 <sup>b</sup>	2.16	(47)	-13.61 <sup>c</sup>
Arizona	190.61 <sup>b</sup>	1.18	(48)	-15.45 <sup>c</sup>
Colorado	190.61 <sup>b</sup>	1.54	(49)	-15.45 <sup>c</sup>
Hawaii	187.27 <sup>b</sup>	2.44	(50)	-16.93 <sup>c</sup>
Utah	161.69 <sup>b</sup>	2.01	(51)	-28.28 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	263.80 <sup>b</sup>	1.64	(05)	17.02 <sup>c</sup>	Montana	211.23 <sup>b</sup>	3.03	(40)	-6.30 <sup>c</sup>
Alaska	213.05	5.27	(38)	-5.50 <sup>c</sup>	Nebraska	218.75	2.35	(33)	-2.96
Arizona	190.61 <sup>b</sup>	1.18	(48)	-15.45 <sup>c</sup>	Nevada	217.95 <sup>b</sup>	2.12	(35)	-3.32
Arkansas	255.50 <sup>b</sup>	1.98	(08)	13.34 <sup>c</sup>	New Hampshire	227.19	2.88	(25)	0.78
California	198.69 <sup>b</sup>	0.54	(46)	-11.86 <sup>c</sup>	New Jersey	222.48	1.09	(29)	-1.31
Colorado	190.61 <sup>b</sup>	1.54	(49)	-15.45 <sup>c</sup>	New Mexico	194.76 <sup>b</sup>	2.16	(47)	-13.61 <sup>c</sup>
Connecticut	218.19 <sup>b</sup>	1.66	(34)	-3.22	New York	205.96 <sup>b</sup>	0.70	(43)	-8.64 <sup>c</sup>
Delaware	241.42 <sup>b</sup>	3.58	(16)	7.09 <sup>c</sup>	North Carolina	244.81 <sup>b</sup>	1.19	(15)	8.59 <sup>c</sup>
D.C.	258.07 <sup>b</sup>	4.82	(07)	14.47 <sup>c</sup>	North Dakota	210.79 <sup>b</sup>	3.62	(42)	-6.50 <sup>c</sup>
Florida	210.96 <sup>b</sup>	0.64	(41)	-6.42 <sup>c</sup>	Ohio	249.06 <sup>b</sup>	0.99	(10)	10.48 <sup>c</sup>
Georgia	241.11 <sup>b</sup>	1.30	(17)	6.95 <sup>c</sup>	Oklahoma	246.00 <sup>b</sup>	1.77	(12)	9.12 <sup>c</sup>
Hawaii	187.27 <sup>b</sup>	2.44	(50)	-16.93 <sup>c</sup>	Oregon	219.85	1.62	(32)	-2.48
Idaho	202.25 <sup>b</sup>	2.61	(45)	-10.28 <sup>c</sup>	Pennsylvania	239.62 <sup>b</sup>	0.88	(18)	6.29 <sup>c</sup>
Illinois	235.38 <sup>b</sup>	0.96	(20)	4.41	Rhode Island	234.89	3.10	(21)	4.19
Indiana	249.41 <sup>b</sup>	1.39	(09)	10.64 <sup>c</sup>	South Carolina	248.96 <sup>b</sup>	1.67	(11)	10.44 <sup>c</sup>
Iowa	225.36	1.76	(26)	-0.03	South Dakota	220.31	3.37	(31)	-2.27
Kansas	225.22	1.92	(27)	-0.10	Tennessee	263.99 <sup>b</sup>	1.46	(04)	17.10 <sup>c</sup>
Kentucky	275.03 <sup>b</sup>	1.78	(02)	22.00 <sup>c</sup>	Texas	221.29 <sup>b</sup>	0.75	(30)	-1.84
Louisiana	270.69 <sup>b</sup>	1.77	(03)	20.07 <sup>c</sup>	Utah	161.69 <sup>b</sup>	2.01	(51)	-28.28 <sup>c</sup>
Maine	245.00 <sup>b</sup>	2.76	(14)	8.68 <sup>c</sup>	Vermont	215.29	3.92	(36)	-4.50
Maryland	230.79 <sup>b</sup>	1.47	(23)	2.38	Virginia	235.65 <sup>b</sup>	1.28	(19)	4.53
Massachusetts	230.49 <sup>b</sup>	1.27	(24)	2.24	Washington	213.97 <sup>b</sup>	1.29	(37)	-5.09 <sup>c</sup>
Michigan	232.81 <sup>b</sup>	1.04	(22)	3.27	West Virginia	259.40 <sup>b</sup>	2.38	(06)	15.07 <sup>c</sup>
Minnesota	211.87 <sup>b</sup>	1.40	(39)	-6.02 <sup>c</sup>	Wisconsin	223.48	1.34	(28)	-0.87
Mississippi	276.31 <sup>b</sup>	2.18	(01)	22.57 <sup>c</sup>	Wyoming	204.62 <sup>b</sup>	4.29	(44)	-9.24 <sup>c</sup>
Missouri	245.73 <sup>b</sup>	1.38	(13)	9.00 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 2.20  
All Cancer Sites (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates by State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	155.43	0.14		
High Five States				
Kentucky	177.55 <sup>b</sup>	1.20	(01)	14.23 <sup>c</sup>
West Virginia	175.76 <sup>b</sup>	1.70	(02)	13.07 <sup>c</sup>
Louisiana	171.55 <sup>b</sup>	1.18	(03)	10.37 <sup>c</sup>
Maine	169.50 <sup>b</sup>	1.98	(04)	9.05 <sup>c</sup>
Delaware	168.38 <sup>b</sup>	2.58	(05)	8.33 <sup>c</sup>
Low Five States				
Colorado	139.41 <sup>b</sup>	1.12	(47)	-10.31 <sup>c</sup>
New Mexico	137.97 <sup>b</sup>	1.62	(48)	-11.24 <sup>c</sup>
Arizona	135.42 <sup>b</sup>	0.90	(49)	-12.88 <sup>c</sup>
Hawaii	122.58 <sup>b</sup>	1.79	(50)	-21.13 <sup>c</sup>
Utah	116.33 <sup>b</sup>	1.52	(51)	-25.16 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	159.92 <sup>b</sup>	1.07	(22)	2.89	Montana	156.50	2.35	(30)	0.68
Alaska	155.99	3.94	(31)	0.36	Nebraska	148.04 <sup>b</sup>	1.69	(40)	-4.76
Arizona	135.42 <sup>b</sup>	0.90	(49)	-12.88 <sup>c</sup>	Nevada	165.21 <sup>b</sup>	1.69	(09)	6.29 <sup>c</sup>
Arkansas	164.91 <sup>b</sup>	1.39	(10)	6.10 <sup>c</sup>	New Hampshire	162.23	2.08	(18)	4.37
California	145.18 <sup>b</sup>	0.40	(46)	-6.60 <sup>c</sup>	New Jersey	163.24 <sup>b</sup>	0.79	(14)	5.02 <sup>c</sup>
Colorado	139.41 <sup>b</sup>	1.12	(47)	-10.31 <sup>c</sup>	New Mexico	137.97 <sup>b</sup>	1.62	(48)	-11.24 <sup>c</sup>
Connecticut	154.74	1.19	(33)	-0.45	New York	150.47 <sup>b</sup>	0.51	(38)	-3.19
Delaware	168.38 <sup>b</sup>	2.58	(05)	8.33 <sup>c</sup>	North Carolina	157.69	0.80	(27)	1.45
D.C.	162.25	3.11	(17)	4.38	North Dakota	147.88	2.74	(41)	-4.86
Florida	145.35 <sup>b</sup>	0.49	(45)	-6.49 <sup>c</sup>	Ohio	168.35 <sup>b</sup>	0.69	(06)	8.31 <sup>c</sup>
Georgia	152.98	0.83	(37)	-1.58	Oklahoma	162.64 <sup>b</sup>	1.25	(16)	4.63
Hawaii	122.58 <sup>b</sup>	1.79	(50)	-21.13 <sup>c</sup>	Oregon	161.70 <sup>b</sup>	1.23	(20)	4.03
Idaho	146.62 <sup>b</sup>	2.00	(43)	-5.67 <sup>c</sup>	Pennsylvania	164.01 <sup>b</sup>	0.63	(12)	5.51 <sup>c</sup>
Illinois	163.45 <sup>b</sup>	0.68	(13)	5.16 <sup>c</sup>	Rhode Island	158.56	2.16	(24)	2.01
Indiana	166.69 <sup>b</sup>	0.96	(08)	7.24 <sup>c</sup>	South Carolina	156.67	1.12	(29)	0.80
Iowa	153.32	1.28	(36)	-1.36	South Dakota	145.75 <sup>b</sup>	2.47	(44)	-6.23 <sup>c</sup>
Kansas	154.15	1.39	(35)	-0.83	Tennessee	167.13 <sup>b</sup>	0.97	(07)	7.52 <sup>c</sup>
Kentucky	177.55 <sup>b</sup>	1.20	(01)	14.23 <sup>c</sup>	Texas	147.31 <sup>b</sup>	0.52	(42)	-5.23 <sup>c</sup>
Louisiana	171.55 <sup>b</sup>	1.18	(03)	10.37 <sup>c</sup>	Utah	116.33 <sup>b</sup>	1.52	(51)	-25.16 <sup>c</sup>
Maine	169.50 <sup>b</sup>	1.98	(04)	9.05 <sup>c</sup>	Vermont	156.75	2.88	(28)	0.84
Maryland	161.68 <sup>b</sup>	1.02	(21)	4.02	Virginia	157.96	0.87	(26)	1.62
Massachusetts	159.41 <sup>b</sup>	0.89	(23)	2.56	Washington	158.48	0.97	(25)	1.96
Michigan	163.09 <sup>b</sup>	0.75	(15)	4.92	West Virginia	175.76 <sup>b</sup>	1.70	(02)	13.07 <sup>c</sup>
Minnesota	149.08 <sup>b</sup>	1.02	(39)	-4.09	Wisconsin	154.50	0.97	(34)	-0.60
Mississippi	162.02 <sup>b</sup>	1.40	(19)	4.24	Wyoming	155.23	3.35	(32)	-0.13
Missouri	164.89 <sup>b</sup>	0.97	(11)	6.08 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 3.18  
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.29	0.02		
High Five States				
Idaho	5.50 <sup>b</sup>	0.28	(01)	28.17 <sup>c</sup>
Iowa	5.35 <sup>b</sup>	0.18	(02)	24.79 <sup>c</sup>
Washington	5.30 <sup>b</sup>	0.13	(03)	23.55 <sup>c</sup>
Oregon	5.26 <sup>b</sup>	0.17	(04)	22.65 <sup>c</sup>
Tennessee	5.26 <sup>b</sup>	0.13	(05)	22.55 <sup>c</sup>
Low Five States				
New Jersey	3.73 <sup>b</sup>	0.09	(47)	-13.06
New Mexico	3.71	0.20	(48)	-13.58
New York	3.68 <sup>b</sup>	0.06	(49)	-14.11
D.C.	2.98 <sup>b</sup>	0.33	(50)	-30.51 <sup>c</sup>
Hawaii	2.58 <sup>b</sup>	0.19	(51)	-39.79 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.33	0.13	(31)	0.91	Montana	4.98	0.31	(08)	16.05 <sup>c</sup>
Alaska	4.27	0.46	(34)	-0.50	Nebraska	4.92	0.23	(10)	14.72
Arizona	4.06	0.12	(41)	-5.26	Nevada	4.25	0.19	(35)	-0.87
Arkansas	4.94 <sup>b</sup>	0.18	(09)	15.18 <sup>c</sup>	New Hampshire	4.79	0.27	(14)	11.63
California	4.28	0.05	(32)	-0.19	New Jersey	3.73 <sup>b</sup>	0.09	(47)	-13.06
Colorado	4.46	0.14	(27)	3.99	New Mexico	3.71	0.20	(48)	-13.58
Connecticut	4.14	0.15	(39)	-3.47	New York	3.68 <sup>b</sup>	0.06	(49)	-14.11
Delaware	4.20	0.31	(36)	-2.16	North Carolina	4.18	0.10	(37)	-2.51
D.C.	2.98 <sup>b</sup>	0.33	(50)	-30.51 <sup>c</sup>	North Dakota	4.88	0.39	(11)	13.87
Florida	3.96 <sup>b</sup>	0.06	(42)	-7.55	Ohio	4.42	0.09	(28)	3.11
Georgia	3.74 <sup>b</sup>	0.10	(46)	-12.76	Oklahoma	4.73	0.16	(15)	10.20
Hawaii	2.58 <sup>b</sup>	0.19	(51)	-39.79 <sup>c</sup>	Oregon	5.26 <sup>b</sup>	0.17	(04)	22.65 <sup>c</sup>
Idaho	5.50 <sup>b</sup>	0.28	(01)	28.17 <sup>c</sup>	Pennsylvania	4.14	0.08	(40)	-3.50
Illinois	3.89 <sup>b</sup>	0.08	(44)	-9.19	Rhode Island	4.60	0.28	(22)	7.27
Indiana	4.62	0.12	(19)	7.64	South Carolina	4.47	0.14	(26)	4.24
Iowa	5.35 <sup>b</sup>	0.18	(02)	24.79 <sup>c</sup>	South Dakota	5.10	0.35	(07)	19.04 <sup>c</sup>
Kansas	5.16 <sup>b</sup>	0.19	(06)	20.40 <sup>c</sup>	Tennessee	5.26 <sup>b</sup>	0.13	(05)	22.55 <sup>c</sup>
Kentucky	4.40	0.14	(30)	2.62	Texas	4.28	0.07	(33)	-0.29
Louisiana	4.67	0.15	(18)	8.93	Utah	4.54	0.22	(23)	5.93
Maine	4.88	0.25	(12)	13.69	Vermont	4.51	0.37	(25)	5.23
Maryland	3.92	0.12	(43)	-8.57	Virginia	3.85 <sup>b</sup>	0.10	(45)	-10.16
Massachusetts	4.16	0.11	(38)	-3.10	Washington	5.30 <sup>b</sup>	0.13	(03)	23.55 <sup>c</sup>
Michigan	4.71 <sup>b</sup>	0.10	(16)	9.75	West Virginia	4.61	0.21	(20)	7.49
Minnesota	4.40	0.13	(29)	2.63	Wisconsin	4.61	0.13	(21)	7.46
Mississippi	4.84	0.18	(13)	12.89	Wyoming	4.70	0.43	(17)	9.58
Missouri	4.54	0.12	(24)	5.82					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 3.19  
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.23	0.03		
High Five States				
South Dakota	6.85	0.60	(01)	30.96 <sup>c</sup>
Iowa	6.51 <sup>b</sup>	0.30	(02)	24.57 <sup>c</sup>
Washington	6.47 <sup>b</sup>	0.21	(03)	23.79 <sup>c</sup>
Oregon	6.46 <sup>b</sup>	0.27	(04)	23.55 <sup>c</sup>
Kansas	6.36 <sup>b</sup>	0.32	(05)	21.65 <sup>c</sup>
Low Five States				
Illinois	4.60 <sup>b</sup>	0.13	(47)	-12.00
New Mexico	4.53	0.32	(48)	-13.24
Georgia	4.45 <sup>b</sup>	0.16	(49)	-14.92
D.C.	3.72	0.55	(50)	-28.84 <sup>c</sup>
Hawaii	3.06 <sup>b</sup>	0.31	(51)	-41.50 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.28	0.22	(30)	1.03	Montana	5.95	0.49	(10)	13.90
Alaska	4.80	0.69	(42)	-8.15	Nebraska	5.81	0.37	(16)	11.14
Arizona	4.77	0.18	(43)	-8.71	Nevada	5.05	0.31	(37)	-3.47
Arkansas	5.89	0.30	(13)	12.66	New Hampshire	6.09	0.45	(07)	16.42 <sup>c</sup>
California	5.23	0.08	(33)	0.15	New Jersey	4.64 <sup>b</sup>	0.15	(46)	-11.28
Colorado	5.38	0.24	(26)	2.89	New Mexico	4.53	0.32	(48)	-13.24
Connecticut	4.81	0.24	(41)	-7.98	New York	4.69 <sup>b</sup>	0.10	(45)	-10.19
Delaware	5.28	0.52	(31)	0.98	North Carolina	5.11	0.16	(36)	-2.25
D.C.	3.72	0.55	(50)	-28.84 <sup>c</sup>	North Dakota	5.86	0.62	(14)	12.19
Florida	4.76 <sup>b</sup>	0.10	(44)	-8.86	Ohio	5.35	0.14	(28)	2.39
Georgia	4.45 <sup>b</sup>	0.16	(49)	-14.92	Oklahoma	5.78	0.26	(17)	10.63
Hawaii	3.06 <sup>b</sup>	0.31	(51)	-41.50 <sup>c</sup>	Oregon	6.46 <sup>b</sup>	0.27	(04)	23.55 <sup>c</sup>
Idaho	5.91	0.42	(12)	13.07	Pennsylvania	5.02	0.13	(38)	-3.90
Illinois	4.60 <sup>b</sup>	0.13	(47)	-12.00	Rhode Island	5.69	0.47	(18)	8.91
Indiana	5.46	0.20	(24)	4.47	South Carolina	5.51	0.24	(22)	5.41
Iowa	6.51 <sup>b</sup>	0.30	(02)	24.57 <sup>c</sup>	South Dakota	6.85	0.60	(01)	30.96 <sup>c</sup>
Kansas	6.36 <sup>b</sup>	0.32	(05)	21.65 <sup>c</sup>	Tennessee	6.17 <sup>b</sup>	0.21	(06)	17.97 <sup>c</sup>
Kentucky	5.36	0.24	(27)	2.52	Texas	5.15	0.11	(35)	-1.47
Louisiana	5.97	0.25	(08)	14.18	Utah	5.51	0.35	(23)	5.32
Maine	5.92	0.42	(11)	13.24	Vermont	5.42	0.59	(25)	3.66
Maryland	4.85	0.20	(40)	-7.16	Virginia	4.98	0.18	(39)	-4.63
Massachusetts	5.25	0.19	(32)	0.36	Washington	6.47 <sup>b</sup>	0.21	(03)	23.79 <sup>c</sup>
Michigan	5.83 <sup>b</sup>	0.16	(15)	11.52	West Virginia	5.53	0.34	(20)	5.89
Minnesota	5.35	0.21	(29)	2.32	Wisconsin	5.55	0.21	(19)	6.19
Mississippi	5.97	0.31	(09)	14.16	Wyoming	5.18	0.65	(34)	-0.98
Missouri	5.52	0.20	(21)	5.70					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 3.20  
Cancer of the Brain and Other Nervous System (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.50	0.02		
High Five States				
Idaho	5.03 <sup>b</sup>	0.38	(01)	43.50 <sup>c</sup>
Tennessee	4.50 <sup>b</sup>	0.16	(02)	28.40 <sup>c</sup>
Iowa	4.37 <sup>b</sup>	0.23	(03)	24.70 <sup>c</sup>
Washington	4.33 <sup>b</sup>	0.16	(04)	23.58 <sup>c</sup>
Wyoming	4.25	0.56	(05)	21.21 <sup>c</sup>
Low Five States				
Virginia	2.97 <sup>b</sup>	0.12	(47)	-15.12 <sup>c</sup>
New Jersey	2.97 <sup>b</sup>	0.11	(48)	-15.33 <sup>c</sup>
New York	2.87 <sup>b</sup>	0.07	(49)	-18.04 <sup>c</sup>
D.C.	2.33	0.39	(50)	-33.52 <sup>c</sup>
Hawaii	2.17 <sup>b</sup>	0.25	(51)	-37.94 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.52	0.16	(33)	0.44	Montana	4.07	0.39	(10)	16.28 <sup>c</sup>
Alaska	3.67	0.61	(23)	4.69	Nebraska	4.08	0.29	(09)	16.41 <sup>c</sup>
Arizona	3.44	0.15	(37)	-1.69	Nevada	3.51	0.24	(34)	0.14
Arkansas	4.15	0.23	(07)	18.56 <sup>c</sup>	New Hampshire	3.64	0.32	(26)	3.84
California	3.48	0.06	(36)	-0.68	New Jersey	2.97 <sup>b</sup>	0.11	(48)	-15.33 <sup>c</sup>
Colorado	3.65	0.18	(24)	4.10	New Mexico	2.99	0.24	(46)	-14.72
Connecticut	3.50	0.19	(35)	0.05	New York	2.87 <sup>b</sup>	0.07	(49)	-18.04 <sup>c</sup>
Delaware	3.30	0.37	(41)	-5.80	North Carolina	3.41	0.12	(39)	-2.56
D.C.	2.33	0.39	(50)	-33.52 <sup>c</sup>	North Dakota	3.96	0.47	(12)	13.06
Florida	3.28	0.08	(42)	-6.47	Ohio	3.65	0.11	(25)	4.09
Georgia	3.16	0.12	(45)	-9.81	Oklahoma	3.76	0.19	(17)	7.47
Hawaii	2.17 <sup>b</sup>	0.25	(51)	-37.94 <sup>c</sup>	Oregon	4.22	0.20	(06)	20.53 <sup>c</sup>
Idaho	5.03 <sup>b</sup>	0.38	(01)	43.50 <sup>c</sup>	Pennsylvania	3.42	0.10	(38)	-2.49
Illinois	3.34	0.10	(40)	-4.62	Rhode Island	3.76	0.35	(19)	7.40
Indiana	3.92	0.15	(13)	12.00	South Carolina	3.70	0.18	(21)	5.63
Iowa	4.37 <sup>b</sup>	0.23	(03)	24.70 <sup>c</sup>	South Dakota	3.61	0.41	(29)	3.06
Kansas	4.13	0.23	(08)	17.87 <sup>c</sup>	Tennessee	4.50 <sup>b</sup>	0.16	(02)	28.40 <sup>c</sup>
Kentucky	3.57	0.17	(31)	2.02	Texas	3.55	0.08	(32)	1.43
Louisiana	3.62	0.17	(28)	3.44	Utah	3.63	0.27	(27)	3.63
Maine	4.00	0.31	(11)	14.27	Vermont	3.68	0.46	(22)	5.05
Maryland	3.17	0.14	(44)	-9.37	Virginia	2.97 <sup>b</sup>	0.12	(47)	-15.12 <sup>c</sup>
Massachusetts	3.27	0.13	(43)	-6.78	Washington	4.33 <sup>b</sup>	0.16	(04)	23.58 <sup>c</sup>
Michigan	3.76	0.12	(18)	7.46	West Virginia	3.82	0.26	(16)	9.00
Minnesota	3.59	0.16	(30)	2.47	Wisconsin	3.82	0.16	(15)	9.01
Mississippi	3.91	0.22	(14)	11.73	Wyoming	4.25	0.56	(05)	21.21 <sup>c</sup>
Missouri	3.73	0.15	(20)	6.46					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.




Table 4.24  
Cancer of the Female Breast (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	23.96	0.05		
High Five States				
D.C.	28.25	1.30	(01)	17.89 <sup>c</sup>
Louisiana	27.74 <sup>b</sup>	0.48	(02)	15.78 <sup>c</sup>
New Jersey	26.96 <sup>b</sup>	0.32	(03)	12.48 <sup>c</sup>
Ohio	26.58 <sup>b</sup>	0.28	(04)	10.91 <sup>c</sup>
Missouri	25.84 <sup>b</sup>	0.39	(05)	7.85
Low Five States				
Idaho	21.64	0.77	(47)	-9.70
Montana	21.44	0.87	(48)	-10.55 <sup>c</sup>
Colorado	21.33 <sup>b</sup>	0.43	(49)	-10.98 <sup>c</sup>
Arizona	21.29 <sup>b</sup>	0.36	(50)	-11.16 <sup>c</sup>
Hawaii	17.86 <sup>b</sup>	0.70	(51)	-25.48 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	24.81	0.43	(12)	3.54	Montana	21.44	0.87	(48)	-10.55 <sup>c</sup>
Alaska	22.67	1.41	(37)	-5.42	Nebraska	22.39	0.66	(41)	-6.56
Arizona	21.29 <sup>b</sup>	0.36	(50)	-11.16 <sup>c</sup>	Nevada	23.90	0.64	(24)	-0.25
Arkansas	24.43	0.54	(18)	1.95	New Hampshire	23.14	0.78	(30)	-3.42
California	22.83 <sup>b</sup>	0.16	(34)	-4.72	New Jersey	26.96 <sup>b</sup>	0.32	(03)	12.48 <sup>c</sup>
Colorado	21.33 <sup>b</sup>	0.43	(49)	-10.98 <sup>c</sup>	New Mexico	22.09	0.65	(43)	-7.83
Connecticut	23.68	0.47	(26)	-1.19	New York	23.88	0.20	(25)	-0.33
Delaware	24.60	1.00	(15)	2.64	North Carolina	24.78	0.32	(13)	3.40
D.C.	28.25	1.30	(01)	17.89 <sup>c</sup>	North Dakota	22.44	1.08	(40)	-6.36
Florida	22.07 <sup>b</sup>	0.20	(44)	-7.89	Ohio	26.58 <sup>b</sup>	0.28	(04)	10.91 <sup>c</sup>
Georgia	23.92	0.33	(23)	-0.17	Oklahoma	24.67	0.49	(14)	2.94
Hawaii	17.86 <sup>b</sup>	0.70	(51)	-25.48 <sup>c</sup>	Oregon	23.21	0.47	(29)	-3.16
Idaho	21.64	0.77	(47)	-9.70	Pennsylvania	25.61 <sup>b</sup>	0.25	(08)	6.85
Illinois	25.22 <sup>b</sup>	0.27	(11)	5.25	Rhode Island	22.82	0.83	(35)	-4.76
Indiana	24.50	0.37	(17)	2.24	South Carolina	24.36	0.44	(19)	1.64
Iowa	21.96 <sup>b</sup>	0.49	(45)	-8.34	South Dakota	22.26	0.98	(42)	-7.10
Kansas	24.03	0.55	(22)	0.27	Tennessee	25.45 <sup>b</sup>	0.38	(10)	6.19
Kentucky	24.21	0.45	(21)	1.03	Texas	23.02 <sup>b</sup>	0.20	(31)	-3.94
Louisiana	27.74 <sup>b</sup>	0.48	(02)	15.78 <sup>c</sup>	Utah	22.81	0.67	(36)	-4.83
Maine	22.60	0.73	(39)	-5.70	Vermont	23.50	1.11	(27)	-1.92
Maryland	25.83 <sup>b</sup>	0.41	(06)	7.80	Virginia	25.58 <sup>b</sup>	0.35	(09)	6.73
Massachusetts	22.95	0.34	(33)	-4.25	Washington	23.00	0.37	(32)	-4.03
Michigan	24.52	0.29	(16)	2.32	West Virginia	24.25	0.64	(20)	1.21
Minnesota	21.83 <sup>b</sup>	0.39	(46)	-8.91	Wisconsin	22.64 <sup>b</sup>	0.37	(38)	-5.51
Mississippi	25.78	0.56	(07)	7.58	Wyoming	23.44	1.31	(28)	-2.18
Missouri	25.84 <sup>b</sup>	0.39	(05)	7.85					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 5.14  
Cancer of the Cervix Uteri (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.43	0.02		
High Five States				
Mississippi	4.02 <sup>b</sup>	0.23	(01)	65.27 <sup>c</sup>
D.C.	3.63	0.49	(02)	49.22 <sup>c</sup>
Arkansas	3.27 <sup>b</sup>	0.21	(03)	34.62 <sup>c</sup>
West Virginia	3.19	0.25	(04)	31.32 <sup>c</sup>
Louisiana	3.16 <sup>b</sup>	0.17	(05)	30.08 <sup>c</sup>
Low Five States				
Vermont	1.63	0.30	(47)	-32.84 <sup>c</sup>
Minnesota	1.60 <sup>b</sup>	0.11	(48)	-34.33 <sup>c</sup>
South Dakota	1.55	0.27	(49)	-36.38 <sup>c</sup>
Utah	1.52 <sup>b</sup>	0.17	(50)	-37.40 <sup>c</sup>
Massachusetts	1.40 <sup>b</sup>	0.09	(51)	-42.34 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.03 <sup>b</sup>	0.15	(07)	24.47 <sup>c</sup>	Montana	1.93	0.27	(36)	-20.43 <sup>c</sup>
Alaska	2.11	0.42	(31)	-13.18	Nebraska	1.70 <sup>b</sup>	0.19	(45)	-30.22 <sup>c</sup>
Arizona	2.07	0.12	(33)	-14.93	Nevada	2.60	0.21	(18)	6.99
Arkansas	3.27 <sup>b</sup>	0.21	(03)	34.62 <sup>c</sup>	New Hampshire	2.28	0.25	(24)	-6.35
California	2.35	0.05	(23)	-3.15	New Jersey	2.59	0.10	(19)	6.55
Colorado	1.81 <sup>b</sup>	0.12	(41)	-25.56 <sup>c</sup>	New Mexico	2.62	0.23	(16)	7.81
Connecticut	1.68 <sup>b</sup>	0.13	(46)	-30.81 <sup>c</sup>	New York	2.53	0.07	(20)	4.18
Delaware	2.26	0.31	(26)	-6.96	North Carolina	2.40	0.10	(22)	-1.28
D.C.	3.63	0.49	(02)	49.22 <sup>c</sup>	North Dakota	1.75	0.32	(43)	-28.11 <sup>c</sup>
Florida	2.66	0.07	(15)	9.61	Ohio	2.45	0.09	(21)	0.73
Georgia	2.71	0.11	(11)	11.59	Oklahoma	2.72	0.17	(10)	12.00
Hawaii	2.07	0.25	(32)	-14.73	Oregon	1.76 <sup>b</sup>	0.13	(42)	-27.47 <sup>c</sup>
Idaho	1.89	0.23	(38)	-22.42 <sup>c</sup>	Pennsylvania	2.24	0.08	(28)	-8.05
Illinois	2.69	0.09	(13)	10.73	Rhode Island	1.70	0.24	(44)	-30.21 <sup>c</sup>
Indiana	2.61	0.13	(17)	7.26	South Carolina	2.71	0.15	(12)	11.38
Iowa	2.27	0.17	(25)	-6.77	South Dakota	1.55	0.27	(49)	-36.38 <sup>c</sup>
Kansas	2.25	0.18	(27)	-7.57	Tennessee	2.83	0.13	(09)	16.43 <sup>c</sup>
Kentucky	2.86	0.16	(08)	17.73 <sup>c</sup>	Texas	3.10 <sup>b</sup>	0.08	(06)	27.41 <sup>c</sup>
Louisiana	3.16 <sup>b</sup>	0.17	(05)	30.08 <sup>c</sup>	Utah	1.52 <sup>b</sup>	0.17	(50)	-37.40 <sup>c</sup>
Maine	1.93	0.22	(37)	-20.46 <sup>c</sup>	Vermont	1.63	0.30	(47)	-32.84 <sup>c</sup>
Maryland	2.23	0.12	(29)	-8.43	Virginia	2.18	0.10	(30)	-10.22
Massachusetts	1.40 <sup>b</sup>	0.09	(51)	-42.34 <sup>c</sup>	Washington	1.97 <sup>b</sup>	0.11	(35)	-19.08 <sup>c</sup>
Michigan	1.99 <sup>b</sup>	0.09	(34)	-18.11 <sup>c</sup>	West Virginia	3.19	0.25	(04)	31.32 <sup>c</sup>
Minnesota	1.60 <sup>b</sup>	0.11	(48)	-34.33 <sup>c</sup>	Wisconsin	1.86 <sup>b</sup>	0.11	(39)	-23.58 <sup>c</sup>
Mississippi	4.02 <sup>b</sup>	0.23	(01)	65.27 <sup>c</sup>	Wyoming	1.85	0.38	(40)	-23.81 <sup>c</sup>
Missouri	2.67	0.13	(14)	9.66					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.




Table 6.25  
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	17.60	0.03		
High Five States				
West Virginia	21.00 <sup>b</sup>	0.44	(01)	19.31 <sup>c</sup>
Louisiana	20.86 <sup>b</sup>	0.31	(02)	18.48 <sup>c</sup>
Kentucky	20.78 <sup>b</sup>	0.31	(03)	18.04 <sup>c</sup>
Mississippi	20.29 <sup>b</sup>	0.37	(04)	15.26 <sup>c</sup>
D.C.	20.28	0.84	(05)	15.19 <sup>c</sup>
Low Five States				
Washington	15.69 <sup>b</sup>	0.23	(47)	-10.88 <sup>c</sup>
Idaho	15.30 <sup>b</sup>	0.48	(48)	-13.08 <sup>c</sup>
Hawaii	15.26 <sup>b</sup>	0.46	(49)	-13.32 <sup>c</sup>
Arizona	15.23 <sup>b</sup>	0.22	(50)	-13.49 <sup>c</sup>
Utah	12.82 <sup>b</sup>	0.37	(51)	-27.20 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	18.60 <sup>b</sup>	0.28	(17)	5.66	Montana	16.05	0.55	(41)	-8.82
Alaska	17.08	1.00	(36)	-2.96	Nebraska	18.85	0.44	(15)	7.10
Arizona	15.23 <sup>b</sup>	0.22	(50)	-13.49 <sup>c</sup>	Nevada	19.13 <sup>b</sup>	0.43	(11)	8.65
Arkansas	19.12 <sup>b</sup>	0.35	(12)	8.63	New Hampshire	17.50	0.51	(30)	-0.61
California	15.81 <sup>b</sup>	0.10	(46)	-10.21 <sup>c</sup>	New Jersey	19.45 <sup>b</sup>	0.20	(08)	10.49 <sup>c</sup>
Colorado	16.02 <sup>b</sup>	0.29	(42)	-9.00	New Mexico	15.95 <sup>b</sup>	0.41	(43)	-9.40
Connecticut	16.08 <sup>b</sup>	0.29	(40)	-8.68	New York	17.44	0.13	(31)	-0.91
Delaware	18.91	0.65	(14)	7.45	North Carolina	17.34	0.20	(34)	-1.47
D.C.	20.28	0.84	(05)	15.19 <sup>c</sup>	North Dakota	17.60	0.68	(29)	-0.04
Florida	15.91 <sup>b</sup>	0.12	(44)	-9.65	Ohio	19.62 <sup>b</sup>	0.18	(06)	11.46 <sup>c</sup>
Georgia	17.37	0.22	(33)	-1.35	Oklahoma	18.54	0.31	(18)	5.32
Hawaii	15.26 <sup>b</sup>	0.46	(49)	-13.32 <sup>c</sup>	Oregon	16.78	0.29	(38)	-4.68
Idaho	15.30 <sup>b</sup>	0.48	(48)	-13.08 <sup>c</sup>	Pennsylvania	19.20 <sup>b</sup>	0.16	(09)	9.07
Illinois	19.51 <sup>b</sup>	0.18	(07)	10.84 <sup>c</sup>	Rhode Island	17.10	0.53	(35)	-2.89
Indiana	19.19 <sup>b</sup>	0.24	(10)	9.01	South Carolina	17.92	0.29	(22)	1.78
Iowa	18.52	0.32	(19)	5.18	South Dakota	17.90	0.63	(23)	1.66
Kansas	17.88	0.35	(24)	1.59	Tennessee	18.97 <sup>b</sup>	0.25	(13)	7.74
Kentucky	20.78 <sup>b</sup>	0.31	(03)	18.04 <sup>c</sup>	Texas	16.98 <sup>b</sup>	0.13	(37)	-3.54
Louisiana	20.86 <sup>b</sup>	0.31	(02)	18.48 <sup>c</sup>	Utah	12.82 <sup>b</sup>	0.37	(51)	-27.20 <sup>c</sup>
Maine	18.21	0.48	(21)	3.46	Vermont	17.67	0.72	(27)	0.37
Maryland	18.64 <sup>b</sup>	0.26	(16)	5.88	Virginia	17.61	0.22	(28)	0.05
Massachusetts	17.40	0.22	(32)	-1.18	Washington	15.69 <sup>b</sup>	0.23	(47)	-10.88 <sup>c</sup>
Michigan	17.85	0.19	(26)	1.43	West Virginia	21.00 <sup>b</sup>	0.44	(01)	19.31 <sup>c</sup>
Minnesota	15.89 <sup>b</sup>	0.25	(45)	-9.75	Wisconsin	16.50 <sup>b</sup>	0.23	(39)	-6.26
Mississippi	20.29 <sup>b</sup>	0.37	(04)	15.26 <sup>c</sup>	Wyoming	17.87	0.84	(25)	1.51
Missouri	18.51 <sup>b</sup>	0.24	(20)	5.14					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 6.26  
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	21.19	0.06		
High Five States				
Louisiana	26.35 <sup>b</sup>	0.55	(01)	24.33 <sup>c</sup>
West Virginia	25.32 <sup>b</sup>	0.75	(02)	19.47 <sup>c</sup>
Kentucky	25.22 <sup>b</sup>	0.55	(03)	18.99 <sup>c</sup>
Mississippi	24.85 <sup>b</sup>	0.65	(04)	17.28 <sup>c</sup>
D.C.	24.37	1.48	(05)	15.00 <sup>c</sup>
Low Five States				
Arizona	18.51 <sup>b</sup>	0.37	(47)	-12.66 <sup>c</sup>
Washington	18.34 <sup>b</sup>	0.38	(48)	-13.44 <sup>c</sup>
Montana	18.14	0.89	(49)	-14.39 <sup>c</sup>
Idaho	17.22 <sup>b</sup>	0.76	(50)	-18.73 <sup>c</sup>
Utah	15.07 <sup>b</sup>	0.61	(51)	-28.87 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	23.56 <sup>b</sup>	0.49	(10)	11.16 <sup>c</sup>	Montana	18.14	0.89	(49)	-14.39 <sup>c</sup>
Alaska	20.96	1.69	(33)	-1.08	Nebraska	23.10	0.76	(15)	8.99
Arizona	18.51 <sup>b</sup>	0.37	(47)	-12.66 <sup>c</sup>	Nevada	21.98	0.68	(20)	3.72
Arkansas	23.36 <sup>b</sup>	0.60	(11)	10.24 <sup>c</sup>	New Hampshire	20.97	0.88	(32)	-1.05
California	18.79 <sup>b</sup>	0.17	(45)	-11.35 <sup>c</sup>	New Jersey	23.31 <sup>b</sup>	0.36	(12)	9.98
Colorado	18.68 <sup>b</sup>	0.48	(46)	-11.84 <sup>c</sup>	New Mexico	19.24	0.68	(41)	-9.19
Connecticut	18.81 <sup>b</sup>	0.49	(44)	-11.25 <sup>c</sup>	New York	20.84	0.22	(34)	-1.65
Delaware	22.32	1.09	(18)	5.31	North Carolina	21.24	0.35	(27)	0.21
D.C.	24.37	1.48	(05)	15.00 <sup>c</sup>	North Dakota	21.31	1.14	(25)	0.58
Florida	18.89 <sup>b</sup>	0.19	(42)	-10.87 <sup>c</sup>	Ohio	23.62 <sup>b</sup>	0.31	(08)	11.45 <sup>c</sup>
Georgia	21.27	0.39	(26)	0.36	Oklahoma	23.14 <sup>b</sup>	0.54	(13)	9.21
Hawaii	19.85	0.79	(39)	-6.31	Oregon	19.27 <sup>b</sup>	0.48	(40)	-9.05
Idaho	17.22 <sup>b</sup>	0.76	(50)	-18.73 <sup>c</sup>	Pennsylvania	23.57 <sup>b</sup>	0.28	(09)	11.23 <sup>c</sup>
Illinois	23.94 <sup>b</sup>	0.31	(07)	12.98 <sup>c</sup>	Rhode Island	21.03	0.93	(30)	-0.76
Indiana	24.05 <sup>b</sup>	0.43	(06)	13.48 <sup>c</sup>	South Carolina	21.43	0.49	(23)	1.13
Iowa	22.10	0.55	(19)	4.28	South Dakota	21.43	1.05	(24)	1.11
Kansas	21.56	0.59	(22)	1.76	Tennessee	23.13 <sup>b</sup>	0.43	(14)	9.15
Kentucky	25.22 <sup>b</sup>	0.55	(03)	18.99 <sup>c</sup>	Texas	21.03	0.23	(31)	-0.77
Louisiana	26.35 <sup>b</sup>	0.55	(01)	24.33 <sup>c</sup>	Utah	15.07 <sup>b</sup>	0.61	(51)	-28.87 <sup>c</sup>
Maine	20.81	0.81	(35)	-1.82	Vermont	20.54	1.21	(36)	-3.09
Maryland	22.78	0.46	(16)	7.51	Virginia	21.66	0.39	(21)	2.22
Massachusetts	21.06	0.39	(29)	-0.64	Washington	18.34 <sup>b</sup>	0.38	(48)	-13.44 <sup>c</sup>
Michigan	21.08	0.32	(28)	-0.51	West Virginia	25.32 <sup>b</sup>	0.75	(02)	19.47 <sup>c</sup>
Minnesota	18.83 <sup>b</sup>	0.42	(43)	-11.13 <sup>c</sup>	Wisconsin	19.86	0.40	(38)	-6.27
Mississippi	24.85 <sup>b</sup>	0.65	(04)	17.28 <sup>c</sup>	Wyoming	20.13	1.36	(37)	-5.02
Missouri	22.48	0.42	(17)	6.07					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 6.27  
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	14.89	0.04		
High Five States				
West Virginia	17.86 <sup>b</sup>	0.53	(01)	19.89 <sup>c</sup>
Kentucky	17.64 <sup>b</sup>	0.37	(02)	18.46 <sup>c</sup>
D.C.	17.57	1.01	(03)	17.94 <sup>c</sup>
Louisiana	16.89 <sup>b</sup>	0.37	(04)	13.42 <sup>c</sup>
Mississippi	16.89 <sup>b</sup>	0.45	(05)	13.38 <sup>c</sup>
Low Five States				
Florida	13.40 <sup>b</sup>	0.15	(47)	-10.02 <sup>c</sup>
New Mexico	13.33	0.50	(48)	-10.50 <sup>c</sup>
Arizona	12.51 <sup>b</sup>	0.27	(49)	-16.03 <sup>c</sup>
Hawaii	11.39 <sup>b</sup>	0.54	(50)	-23.56 <sup>c</sup>
Utah	10.87 <sup>b</sup>	0.46	(51)	-27.01 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	15.11	0.33	(24)	1.47	Montana	14.33	0.70	(37)	-3.78
Alaska	13.80	1.20	(42)	-7.36	Nebraska	15.66	0.53	(18)	5.17
Arizona	12.51 <sup>b</sup>	0.27	(49)	-16.03 <sup>c</sup>	Nevada	16.54	0.54	(08)	11.05 <sup>c</sup>
Arkansas	15.74	0.42	(16)	5.66	New Hampshire	14.84	0.62	(29)	-0.34
California	13.48 <sup>b</sup>	0.12	(45)	-9.47	New Jersey	16.73 <sup>b</sup>	0.25	(07)	12.31 <sup>c</sup>
Colorado	13.88	0.35	(40)	-6.84	New Mexico	13.33	0.50	(48)	-10.50 <sup>c</sup>
Connecticut	14.16	0.35	(38)	-4.90	New York	15.04	0.16	(26)	0.97
Delaware	16.02	0.79	(13)	7.57	North Carolina	14.48	0.24	(36)	-2.82
D.C.	17.57	1.01	(03)	17.94 <sup>c</sup>	North Dakota	14.83	0.83	(31)	-0.43
Florida	13.40 <sup>b</sup>	0.15	(47)	-10.02 <sup>c</sup>	Ohio	16.78 <sup>b</sup>	0.21	(06)	12.68 <sup>c</sup>
Georgia	14.73	0.26	(33)	-1.10	Oklahoma	15.02	0.37	(27)	0.86
Hawaii	11.39 <sup>b</sup>	0.54	(50)	-23.56 <sup>c</sup>	Oregon	14.76	0.37	(32)	-0.89
Idaho	13.69	0.61	(44)	-8.07	Pennsylvania	16.14 <sup>b</sup>	0.19	(11)	8.38
Illinois	16.46 <sup>b</sup>	0.21	(09)	10.50 <sup>c</sup>	Rhode Island	14.56	0.64	(35)	-2.23
Indiana	15.73	0.29	(17)	5.62	South Carolina	15.20	0.35	(23)	2.05
Iowa	15.82	0.40	(15)	6.24	South Dakota	15.06	0.77	(25)	1.13
Kansas	15.01	0.42	(28)	0.75	Tennessee	15.87	0.30	(14)	6.53
Kentucky	17.64 <sup>b</sup>	0.37	(02)	18.46 <sup>c</sup>	Texas	13.85 <sup>b</sup>	0.16	(41)	-7.01
Louisiana	16.89 <sup>b</sup>	0.37	(04)	13.42 <sup>c</sup>	Utah	10.87 <sup>b</sup>	0.46	(51)	-27.01 <sup>c</sup>
Maine	16.26	0.60	(10)	9.16	Vermont	15.36	0.88	(22)	3.14
Maryland	15.61	0.31	(19)	4.80	Virginia	14.70	0.26	(34)	-1.33
Massachusetts	14.84	0.26	(30)	-0.37	Washington	13.44 <sup>b</sup>	0.28	(46)	-9.75
Michigan	15.43	0.23	(21)	3.60	West Virginia	17.86 <sup>b</sup>	0.53	(01)	19.89 <sup>c</sup>
Minnesota	13.71 <sup>b</sup>	0.30	(43)	-7.93	Wisconsin	13.96	0.28	(39)	-6.27
Mississippi	16.89 <sup>b</sup>	0.45	(05)	13.38 <sup>c</sup>	Wyoming	16.04	1.07	(12)	7.67
Missouri	15.56	0.29	(20)	4.45					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 7.14  
Cancer of the Corpus and Uterus, NOS (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.15	0.02		
High Five States				
D.C.	5.56	0.57	(01)	34.11 <sup>c</sup>
Nebraska	5.14	0.31	(02)	23.93 <sup>c</sup>
New Jersey	5.14 <sup>b</sup>	0.14	(03)	23.81 <sup>c</sup>
New York	5.03 <sup>b</sup>	0.09	(04)	21.30 <sup>c</sup>
Pennsylvania	5.00 <sup>b</sup>	0.11	(05)	20.66 <sup>c</sup>
Low Five States				
Arkansas	3.24 <sup>b</sup>	0.19	(47)	-21.88 <sup>c</sup>
Colorado	3.19 <sup>b</sup>	0.17	(48)	-22.99 <sup>c</sup>
Nevada	3.01 <sup>b</sup>	0.23	(49)	-27.53 <sup>c</sup>
Arizona	2.91 <sup>b</sup>	0.13	(50)	-29.75 <sup>c</sup>
Alaska	2.28 <sup>b</sup>	0.47	(51)	-45.03 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.37 <sup>b</sup>	0.15	(44)	-18.65 <sup>c</sup>	Montana	4.67	0.41	(10)	12.63
Alaska	2.28 <sup>b</sup>	0.47	(51)	-45.03 <sup>c</sup>	Nebraska	5.14	0.31	(02)	23.93 <sup>c</sup>
Arizona	2.91 <sup>b</sup>	0.13	(50)	-29.75 <sup>c</sup>	Nevada	3.01 <sup>b</sup>	0.23	(49)	-27.53 <sup>c</sup>
Arkansas	3.24 <sup>b</sup>	0.19	(47)	-21.88 <sup>c</sup>	New Hampshire	3.83	0.32	(32)	-7.60
California	3.96	0.07	(29)	-4.60	New Jersey	5.14 <sup>b</sup>	0.14	(03)	23.81 <sup>c</sup>
Colorado	3.19 <sup>b</sup>	0.17	(48)	-22.99 <sup>c</sup>	New Mexico	3.49	0.26	(43)	-15.89 <sup>c</sup>
Connecticut	4.59	0.21	(12)	10.77	New York	5.03 <sup>b</sup>	0.09	(04)	21.30 <sup>c</sup>
Delaware	4.84	0.43	(07)	16.66 <sup>c</sup>	North Carolina	4.14	0.13	(24)	-0.27
D.C.	5.56	0.57	(01)	34.11 <sup>c</sup>	North Dakota	4.85	0.49	(06)	16.99 <sup>c</sup>
Florida	3.53 <sup>b</sup>	0.08	(42)	-14.79	Ohio	4.51	0.11	(14)	8.85
Georgia	3.68 <sup>b</sup>	0.13	(37)	-11.32	Oklahoma	3.98	0.19	(28)	-4.16
Hawaii	3.70	0.32	(36)	-10.88	Oregon	3.83	0.19	(31)	-7.55
Idaho	3.79	0.32	(33)	-8.56	Pennsylvania	5.00 <sup>b</sup>	0.11	(05)	20.66 <sup>c</sup>
Illinois	4.47	0.11	(17)	7.88	Rhode Island	3.62	0.33	(38)	-12.81
Indiana	4.55	0.16	(13)	9.68	South Carolina	4.12	0.18	(25)	-0.70
Iowa	4.73	0.22	(09)	14.02	South Dakota	4.00	0.41	(27)	-3.49
Kansas	3.56	0.21	(40)	-14.16	Tennessee	3.70	0.14	(35)	-10.71
Kentucky	3.61	0.17	(39)	-12.91	Texas	3.56 <sup>b</sup>	0.08	(41)	-14.24
Louisiana	3.70	0.17	(34)	-10.70	Utah	3.25	0.26	(46)	-21.66 <sup>c</sup>
Maine	4.49	0.32	(15)	8.18	Vermont	4.49	0.48	(16)	8.14
Maryland	4.42	0.17	(18)	6.46	Virginia	4.17	0.14	(23)	0.60
Massachusetts	4.20	0.15	(22)	1.27	Washington	4.05	0.16	(26)	-2.28
Michigan	4.74 <sup>b</sup>	0.13	(08)	14.29	West Virginia	4.39	0.27	(19)	5.78
Minnesota	4.37	0.18	(20)	5.35	Wisconsin	4.64	0.17	(11)	11.89
Mississippi	3.94	0.22	(30)	-5.08	Wyoming	3.30	0.49	(45)	-20.41 <sup>c</sup>
Missouri	4.28	0.16	(21)	3.27					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 8.18<h2>  
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.37	0.02		
High Five States				
Maine	6.25 <sup>b</sup>	0.28	(01)	43.04 <sup>c</sup>
New Hampshire	6.03 <sup>b</sup>	0.30	(02)	38.02 <sup>c</sup>
D.C.	5.56	0.44	(03)	27.17 <sup>c</sup>
Rhode Island	5.55 <sup>b</sup>	0.31	(04)	27.04 <sup>c</sup>
Wyoming	5.49	0.46	(05)	25.61 <sup>c</sup>
Low Five States				
Kansas	3.80 <sup>b</sup>	0.16	(47)	-13.12
Texas	3.72 <sup>b</sup>	0.06	(48)	-14.89
California	3.60 <sup>b</sup>	0.05	(49)	-17.53 <sup>c</sup>
Utah	3.22 <sup>b</sup>	0.19	(50)	-26.33 <sup>c</sup>
Hawaii	2.90 <sup>b</sup>	0.20	(51)	-33.52 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.04	0.13	(41)	-7.59	Montana	4.76	0.30	(13)	9.03
Alaska	4.68	0.48	(19)	7.13	Nebraska	4.37	0.22	(32)	-0.02
Arizona	3.99	0.11	(44)	-8.79	Nevada	4.27	0.19	(35)	-2.34
Arkansas	3.87	0.16	(46)	-11.39	New Hampshire	6.03 <sup>b</sup>	0.30	(02)	38.02 <sup>c</sup>
California	3.60 <sup>b</sup>	0.05	(49)	-17.53 <sup>c</sup>	New Jersey	4.23	0.10	(36)	-3.11
Colorado	4.02	0.14	(42)	-8.09	New Mexico	3.89	0.20	(45)	-11.03
Connecticut	4.55	0.15	(23)	4.11	New York	4.37	0.07	(33)	-0.07
Delaware	4.85	0.33	(12)	11.12	North Carolina	4.09	0.10	(40)	-6.44
D.C.	5.56	0.44	(03)	27.17 <sup>c</sup>	North Dakota	4.01	0.33	(43)	-8.31
Florida	4.10 <sup>b</sup>	0.06	(39)	-6.23	Ohio	5.19 <sup>b</sup>	0.09	(07)	18.68 <sup>c</sup>
Georgia	4.18	0.10	(37)	-4.36	Oklahoma	4.28	0.15	(34)	-2.15
Hawaii	2.90 <sup>b</sup>	0.20	(51)	-33.52 <sup>c</sup>	Oregon	5.06 <sup>b</sup>	0.16	(10)	15.90 <sup>c</sup>
Idaho	4.40	0.25	(28)	0.75	Pennsylvania	5.12 <sup>b</sup>	0.08	(09)	17.17 <sup>c</sup>
Illinois	4.60	0.09	(22)	5.32	Rhode Island	5.55 <sup>b</sup>	0.31	(04)	27.04 <sup>c</sup>
Indiana	4.74	0.12	(15)	8.45	South Carolina	4.64	0.14	(21)	6.12
Iowa	4.71	0.17	(17)	7.78	South Dakota	4.76	0.33	(14)	9.00
Kansas	3.80 <sup>b</sup>	0.16	(47)	-13.12	Tennessee	4.13	0.11	(38)	-5.56
Kentucky	4.54	0.14	(24)	3.95	Texas	3.72 <sup>b</sup>	0.06	(48)	-14.89
Louisiana	4.40	0.14	(29)	0.66	Utah	3.22 <sup>b</sup>	0.19	(50)	-26.33 <sup>c</sup>
Maine	6.25 <sup>b</sup>	0.28	(01)	43.04 <sup>c</sup>	Vermont	4.73	0.37	(16)	8.31
Maryland	4.54	0.13	(25)	3.92	Virginia	4.38	0.11	(31)	0.20
Massachusetts	5.45 <sup>b</sup>	0.13	(06)	24.71 <sup>c</sup>	Washington	4.69	0.12	(18)	7.27
Michigan	4.97 <sup>b</sup>	0.10	(11)	13.71	West Virginia	4.67	0.21	(20)	6.91
Minnesota	4.39	0.13	(30)	0.53	Wisconsin	5.14 <sup>b</sup>	0.13	(08)	17.72 <sup>c</sup>
Mississippi	4.42	0.17	(27)	1.17	Wyoming	5.49	0.46	(05)	25.61 <sup>c</sup>
Missouri	4.48	0.12	(26)	2.48					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 8.19  
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.79	0.03		
High Five States				
Maine	11.66 <sup>b</sup>	0.59	(01)	49.69 <sup>c</sup>
New Hampshire	10.67 <sup>b</sup>	0.61	(02)	36.96 <sup>c</sup>
D.C.	10.57	0.95	(03)	35.66 <sup>c</sup>
Massachusetts	9.98 <sup>b</sup>	0.26	(04)	28.08 <sup>c</sup>
Rhode Island	9.84	0.63	(05)	26.36 <sup>c</sup>
Low Five States				
New Mexico	6.79	0.39	(47)	-12.88
Texas	6.71 <sup>b</sup>	0.13	(48)	-13.82
California	6.21 <sup>b</sup>	0.09	(49)	-20.21 <sup>c</sup>
Utah	5.81 <sup>b</sup>	0.37	(50)	-25.37 <sup>c</sup>
Hawaii	5.05 <sup>b</sup>	0.40	(51)	-35.20 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.62	0.27	(34)	-2.15	Montana	7.89	0.57	(31)	1.26
Alaska	7.91	0.95	(29)	1.55	Nebraska	8.11	0.45	(24)	4.08
Arizona	7.09	0.22	(44)	-8.91	Nevada	7.21	0.37	(41)	-7.47
Arkansas	6.80	0.32	(46)	-12.67	New Hampshire	10.67 <sup>b</sup>	0.61	(02)	36.96 <sup>c</sup>
California	6.21 <sup>b</sup>	0.09	(49)	-20.21 <sup>c</sup>	New Jersey	7.42	0.20	(37)	-4.72
Colorado	7.21	0.29	(40)	-7.38	New Mexico	6.79	0.39	(47)	-12.88
Connecticut	8.25	0.32	(18)	5.87	New York	7.59	0.13	(35)	-2.58
Delaware	8.96	0.68	(10)	15.00 <sup>c</sup>	North Carolina	7.24	0.20	(39)	-7.02
D.C.	10.57	0.95	(03)	35.66 <sup>c</sup>	North Dakota	6.98	0.65	(45)	-10.32
Florida	7.13 <sup>b</sup>	0.12	(43)	-8.51	Ohio	9.52 <sup>b</sup>	0.19	(07)	22.29 <sup>c</sup>
Georgia	7.44	0.22	(36)	-4.54	Oklahoma	7.89	0.31	(30)	1.28
Hawaii	5.05 <sup>b</sup>	0.40	(51)	-35.20 <sup>c</sup>	Oregon	8.95 <sup>b</sup>	0.32	(11)	14.96
Idaho	7.80	0.50	(32)	0.18	Pennsylvania	9.33 <sup>b</sup>	0.17	(08)	19.79 <sup>c</sup>
Illinois	8.33	0.18	(16)	6.95	Rhode Island	9.84	0.63	(05)	26.36 <sup>c</sup>
Indiana	8.64	0.25	(13)	10.90	South Carolina	8.30	0.29	(17)	6.50
Iowa	8.63	0.34	(14)	10.85	South Dakota	8.49	0.66	(15)	9.05
Kansas	7.20	0.34	(42)	-7.50	Tennessee	7.41	0.24	(38)	-4.87
Kentucky	8.21	0.30	(21)	5.39	Texas	6.71 <sup>b</sup>	0.13	(48)	-13.82
Louisiana	7.99	0.29	(27)	2.63	Utah	5.81 <sup>b</sup>	0.37	(50)	-25.37 <sup>c</sup>
Maine	11.66 <sup>b</sup>	0.59	(01)	49.69 <sup>c</sup>	Vermont	8.24	0.76	(19)	5.73
Maryland	8.06	0.27	(26)	3.45	Virginia	7.79	0.22	(33)	0.07
Massachusetts	9.98 <sup>b</sup>	0.26	(04)	28.08 <sup>c</sup>	Washington	8.18	0.25	(23)	4.99
Michigan	8.84 <sup>b</sup>	0.20	(12)	13.51	West Virginia	8.22	0.41	(20)	5.49
Minnesota	7.92	0.27	(28)	1.66	Wisconsin	9.17 <sup>b</sup>	0.27	(09)	17.72 <sup>c</sup>
Mississippi	8.20	0.36	(22)	5.34	Wyoming	9.53	0.91	(06)	22.30 <sup>c</sup>
Missouri	8.08	0.25	(25)	3.70					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.




Table 8.20  
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.67	0.01		
High Five States				
New Hampshire	2.37	0.25	(01)	41.32 <sup>c</sup>
Rhode Island	2.30	0.26	(02)	37.55 <sup>c</sup>
Montana	2.13	0.27	(03)	27.04 <sup>c</sup>
Massachusetts	2.05 <sup>b</sup>	0.10	(04)	22.46 <sup>c</sup>
Vermont	2.01	0.33	(05)	19.86 <sup>c</sup>
Low Five States				
Arizona	1.34 <sup>b</sup>	0.09	(47)	-20.26 <sup>c</sup>
Alabama	1.27 <sup>b</sup>	0.09	(48)	-24.34 <sup>c</sup>
Kansas	1.10 <sup>b</sup>	0.12	(49)	-34.50 <sup>c</sup>
Hawaii	1.08 <sup>b</sup>	0.17	(50)	-35.29 <sup>c</sup>
Utah	0.96 <sup>b</sup>	0.14	(51)	-42.55 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.27 <sup>b</sup>	0.09	(48)	-24.34 <sup>c</sup>	Montana	2.13	0.27	(03)	27.04 <sup>c</sup>
Alaska	1.80	0.42	(17)	7.67	Nebraska	1.38	0.16	(44)	-17.79 <sup>c</sup>
Arizona	1.34 <sup>b</sup>	0.09	(47)	-20.26 <sup>c</sup>	Nevada	1.58	0.16	(34)	-5.94
Arkansas	1.48	0.13	(39)	-11.62	New Hampshire	2.37	0.25	(01)	41.32 <sup>c</sup>
California	1.55	0.04	(37)	-7.67	New Jersey	1.78	0.08	(20)	6.14
Colorado	1.40	0.11	(42)	-16.51 <sup>c</sup>	New Mexico	1.46	0.17	(40)	-13.10
Connecticut	1.74	0.13	(23)	3.70	New York	1.95 <sup>b</sup>	0.06	(07)	16.38 <sup>c</sup>
Delaware	1.68	0.26	(26)	0.58	North Carolina	1.62	0.08	(31)	-3.48
D.C.	1.96	0.34	(06)	17.20 <sup>c</sup>	North Dakota	1.66	0.29	(28)	-1.16
Florida	1.57	0.05	(35)	-6.12	Ohio	1.88	0.07	(15)	12.44
Georgia	1.74	0.09	(22)	3.72	Oklahoma	1.36	0.11	(45)	-18.67 <sup>c</sup>
Hawaii	1.08 <sup>b</sup>	0.17	(50)	-35.29 <sup>c</sup>	Oregon	1.92	0.13	(10)	14.52
Idaho	1.38	0.19	(43)	-17.58 <sup>c</sup>	Pennsylvania	1.90	0.07	(13)	13.46
Illinois	1.78	0.07	(18)	6.52	Rhode Island	2.30	0.26	(02)	37.55 <sup>c</sup>
Indiana	1.70	0.10	(24)	1.26	South Carolina	1.78	0.12	(19)	6.25
Iowa	1.64	0.13	(29)	-1.91	South Dakota	1.62	0.26	(30)	-3.07
Kansas	1.10 <sup>b</sup>	0.12	(49)	-34.50 <sup>c</sup>	Tennessee	1.53	0.09	(38)	-8.88
Kentucky	1.60	0.11	(32)	-4.46	Texas	1.35 <sup>b</sup>	0.05	(46)	-19.29 <sup>c</sup>
Louisiana	1.56	0.11	(36)	-6.80	Utah	0.96 <sup>b</sup>	0.14	(51)	-42.55 <sup>c</sup>
Maine	1.91	0.21	(12)	13.74	Vermont	2.01	0.33	(05)	19.86 <sup>c</sup>
Maryland	1.87	0.11	(16)	11.35	Virginia	1.68	0.09	(25)	0.59
Massachusetts	2.05 <sup>b</sup>	0.10	(04)	22.46 <sup>c</sup>	Washington	1.89	0.11	(14)	12.75
Michigan	1.93	0.08	(09)	15.31 <sup>c</sup>	West Virginia	1.77	0.17	(21)	5.86
Minnesota	1.58	0.11	(33)	-5.73	Wisconsin	1.91	0.11	(11)	14.02
Mississippi	1.45	0.13	(41)	-13.15	Wyoming	1.94	0.38	(08)	15.72 <sup>c</sup>
Missouri	1.66	0.10	(27)	-1.00					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 9.18  
Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.43	0.01		
High Five States				
Arkansas	0.56	0.06	(01)	28.59 <sup>c</sup>
Louisiana	0.53	0.05	(02)	23.57 <sup>c</sup>
Mississippi	0.53	0.06	(03)	21.96 <sup>c</sup>
Pennsylvania	0.53	0.03	(04)	21.68 <sup>c</sup>
New Jersey	0.52	0.03	(05)	20.53 <sup>c</sup>
Low Five States				
Arizona	0.35	0.03	(47)	-19.50 <sup>c</sup>
Utah	0.35	0.06	(48)	-20.06 <sup>c</sup>
Alaska	0.34	0.12	(49)	-20.15 <sup>c</sup>
Hawaii	0.27	0.06	(50)	-38.04 <sup>c</sup>
New Hampshire	0.25	0.06	(51)	-42.15 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.43	0.04	(25)	-1.03	Montana	0.36	0.09	(46)	-16.69 <sup>c</sup>
Alaska	0.34	0.12	(49)	-20.15 <sup>c</sup>	Nebraska	0.46	0.07	(15)	7.07
Arizona	0.35	0.03	(47)	-19.50 <sup>c</sup>	Nevada	0.45	0.06	(18)	4.36
Arkansas	0.56	0.06	(01)	28.59 <sup>c</sup>	New Hampshire	0.25	0.06	(51)	-42.15 <sup>c</sup>
California	0.43	0.02	(24)	-0.32	New Jersey	0.52	0.03	(05)	20.53 <sup>c</sup>
Colorado	0.40	0.04	(34)	-8.30	New Mexico	0.37	0.06	(40)	-13.38
Connecticut	0.41	0.05	(31)	-5.73	New York	0.44	0.02	(20)	1.12
Delaware	0.37	0.09	(42)	-14.70	North Carolina	0.42	0.03	(29)	-3.25
D.C.	0.43	0.12	(22)	0.71	North Dakota	0.42	0.12	(28)	-2.92
Florida	0.38	0.02	(39)	-12.68	Ohio	0.50	0.03	(06)	16.11 <sup>c</sup>
Georgia	0.37	0.03	(41)	-14.17	Oklahoma	0.47	0.05	(10)	9.48
Hawaii	0.27	0.06	(50)	-38.04 <sup>c</sup>	Oregon	0.44	0.05	(19)	1.58
Idaho	0.36	0.07	(45)	-16.50 <sup>c</sup>	Pennsylvania	0.53	0.03	(04)	21.68 <sup>c</sup>
Illinois	0.42	0.03	(30)	-3.73	Rhode Island	0.47	0.09	(12)	8.70
Indiana	0.36	0.03	(44)	-16.33 <sup>c</sup>	South Carolina	0.38	0.04	(38)	-12.53
Iowa	0.45	0.05	(17)	4.39	South Dakota	0.49	0.11	(08)	13.47
Kansas	0.40	0.05	(33)	-7.79	Tennessee	0.47	0.04	(14)	7.91
Kentucky	0.47	0.05	(11)	8.95	Texas	0.46	0.02	(16)	5.63
Louisiana	0.53	0.05	(02)	23.57 <sup>c</sup>	Utah	0.35	0.06	(48)	-20.06 <sup>c</sup>
Maine	0.39	0.07	(35)	-9.05	Vermont	0.41	0.11	(32)	-5.98
Maryland	0.42	0.04	(26)	-2.06	Virginia	0.39	0.03	(36)	-10.65
Massachusetts	0.36	0.03	(43)	-16.11 <sup>c</sup>	Washington	0.42	0.04	(27)	-2.57
Michigan	0.47	0.03	(13)	8.55	West Virginia	0.48	0.07	(09)	11.51
Minnesota	0.49	0.04	(07)	13.68	Wisconsin	0.44	0.04	(21)	1.06
Mississippi	0.53	0.06	(03)	21.96 <sup>c</sup>	Wyoming	0.38	0.12	(37)	-11.44
Missouri	0.43	0.04	(23)	0.14					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 9.19  
Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.53	0.01		
High Five States				
Arkansas	0.72	0.10	(01)	36.10 <sup>c</sup>
Nebraska	0.69	0.13	(02)	30.43 <sup>c</sup>
Pennsylvania	0.67	0.05	(03)	27.01 <sup>c</sup>
Louisiana	0.65	0.09	(04)	22.25 <sup>c</sup>
Kentucky	0.65	0.08	(05)	21.91 <sup>c</sup>
Low Five States				
Idaho	0.35	0.10	(47)	-34.96 <sup>c</sup>
New Hampshire	0.34	0.12	(48)	-35.57 <sup>c</sup>
Hawaii	0.33	0.10	(49)	-38.68 <sup>c</sup>
Alaska	0.29	0.12	(50)	-44.60 <sup>c</sup>
Vermont	0.26	0.13	(51)	-50.79 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.55	0.07	(15)	2.63	Montana	0.48	0.15	(32)	-9.20
Alaska	0.29	0.12	(50)	-44.60 <sup>c</sup>	Nebraska	0.69	0.13	(02)	30.43 <sup>c</sup>
Arizona	0.47	0.06	(37)	-10.84	Nevada	0.52	0.10	(24)	-2.40
Arkansas	0.72	0.10	(01)	36.10 <sup>c</sup>	New Hampshire	0.34	0.12	(48)	-35.57 <sup>c</sup>
California	0.54	0.03	(19)	0.71	New Jersey	0.62	0.06	(07)	16.29 <sup>c</sup>
Colorado	0.54	0.08	(16)	2.55	New Mexico	0.47	0.10	(36)	-10.66
Connecticut	0.60	0.09	(08)	12.36	New York	0.52	0.03	(25)	-2.53
Delaware	0.35	0.14	(46)	-33.84 <sup>c</sup>	North Carolina	0.58	0.06	(10)	9.68
D.C.	0.50	0.19	(28)	-6.13	North Dakota	0.52	0.19	(23)	-1.72
Florida	0.47	0.03	(38)	-11.07	Ohio	0.63	0.05	(06)	18.81 <sup>c</sup>
Georgia	0.45	0.05	(41)	-14.76	Oklahoma	0.53	0.08	(20)	0.07
Hawaii	0.33	0.10	(49)	-38.68 <sup>c</sup>	Oregon	0.49	0.08	(29)	-7.00
Idaho	0.35	0.10	(47)	-34.96 <sup>c</sup>	Pennsylvania	0.67	0.05	(03)	27.01 <sup>c</sup>
Illinois	0.51	0.04	(26)	-3.58	Rhode Island	0.54	0.15	(17)	2.16
Indiana	0.48	0.06	(34)	-9.71	South Carolina	0.45	0.07	(42)	-15.17 <sup>c</sup>
Iowa	0.39	0.07	(44)	-26.05 <sup>c</sup>	South Dakota	0.53	0.17	(21)	-0.82
Kansas	0.47	0.09	(39)	-11.39	Tennessee	0.48	0.06	(31)	-9.08
Kentucky	0.65	0.08	(05)	21.91 <sup>c</sup>	Texas	0.57	0.04	(12)	7.84
Louisiana	0.65	0.09	(04)	22.25 <sup>c</sup>	Utah	0.46	0.10	(40)	-14.34
Maine	0.54	0.13	(18)	1.03	Vermont	0.26	0.13	(51)	-50.79 <sup>c</sup>
Maryland	0.47	0.06	(35)	-10.63	Virginia	0.48	0.05	(30)	-8.88
Massachusetts	0.44	0.05	(43)	-17.32 <sup>c</sup>	Washington	0.52	0.06	(22)	-1.71
Michigan	0.59	0.05	(09)	11.61	West Virginia	0.50	0.11	(27)	-5.63
Minnesota	0.57	0.07	(13)	7.39	Wisconsin	0.55	0.07	(14)	2.87
Mississippi	0.58	0.10	(11)	8.59	Wyoming	0.37	0.19	(45)	-30.93 <sup>c</sup>
Missouri	0.48	0.06	(33)	-9.45					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 9.20  
Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.35	0.01		
High Five States				
Vermont	0.55	0.19	(01)	57.79 <sup>c</sup>
Mississippi	0.50	0.08	(02)	41.81 <sup>c</sup>
Iowa	0.49	0.08	(03)	40.59 <sup>c</sup>
South Dakota	0.46	0.15	(04)	30.69 <sup>c</sup>
Louisiana	0.46	0.06	(05)	30.63 <sup>c</sup>
Low Five States				
Montana	0.25	0.10	(47)	-27.45 <sup>c</sup>
Connecticut	0.24	0.05	(48)	-29.90 <sup>c</sup>
New Hampshire	0.23	0.08	(49)	-33.34 <sup>c</sup>
Arizona	0.23	0.04	(50)	-33.48 <sup>c</sup>
Hawaii	0.22	0.07	(51)	-38.02 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.32	0.05	(34)	-7.48	Montana	0.25	0.10	(47)	-27.45 <sup>c</sup>
Alaska	0.36	0.19	(24)	2.81	Nebraska	0.27	0.07	(44)	-23.25 <sup>c</sup>
Arizona	0.23	0.04	(50)	-33.48 <sup>c</sup>	Nevada	0.37	0.08	(20)	6.78
Arkansas	0.40	0.07	(15)	14.39	New Hampshire	0.23	0.08	(49)	-33.34 <sup>c</sup>
California	0.34	0.02	(27)	-1.26	New Jersey	0.44	0.04	(07)	27.35 <sup>c</sup>
Colorado	0.26	0.05	(45)	-26.58 <sup>c</sup>	New Mexico	0.28	0.07	(42)	-19.41 <sup>c</sup>
Connecticut	0.24	0.05	(48)	-29.90 <sup>c</sup>	New York	0.37	0.03	(21)	6.27
Delaware	0.40	0.14	(14)	15.76 <sup>c</sup>	North Carolina	0.29	0.03	(40)	-18.01 <sup>c</sup>
D.C.	0.34	0.14	(30)	-2.35	North Dakota	0.33	0.15	(31)	-4.51
Florida	0.29	0.02	(39)	-16.23 <sup>c</sup>	Ohio	0.40	0.03	(16)	14.32
Georgia	0.29	0.04	(37)	-15.54 <sup>c</sup>	Oklahoma	0.43	0.07	(10)	24.19 <sup>c</sup>
Hawaii	0.22	0.07	(51)	-38.02 <sup>c</sup>	Oregon	0.40	0.06	(17)	13.81
Idaho	0.37	0.10	(19)	7.01	Pennsylvania	0.41	0.03	(13)	16.02 <sup>c</sup>
Illinois	0.34	0.03	(29)	-1.93	Rhode Island	0.43	0.12	(11)	23.69 <sup>c</sup>
Indiana	0.28	0.04	(43)	-20.77 <sup>c</sup>	South Carolina	0.32	0.05	(35)	-9.66
Iowa	0.49	0.08	(03)	40.59 <sup>c</sup>	South Dakota	0.46	0.15	(04)	30.69 <sup>c</sup>
Kansas	0.35	0.07	(26)	-0.09	Tennessee	0.45	0.05	(06)	29.41 <sup>c</sup>
Kentucky	0.32	0.05	(33)	-7.28	Texas	0.36	0.03	(25)	1.98
Louisiana	0.46	0.06	(05)	30.63 <sup>c</sup>	Utah	0.26	0.07	(46)	-26.65 <sup>c</sup>
Maine	0.28	0.08	(41)	-18.72 <sup>c</sup>	Vermont	0.55	0.19	(01)	57.79 <sup>c</sup>
Maryland	0.37	0.05	(18)	7.14	Virginia	0.30	0.04	(36)	-14.23
Massachusetts	0.29	0.04	(38)	-15.73 <sup>c</sup>	Washington	0.33	0.04	(32)	-4.64
Michigan	0.36	0.04	(23)	4.27	West Virginia	0.44	0.09	(08)	27.06 <sup>c</sup>
Minnesota	0.44	0.06	(09)	25.46 <sup>c</sup>	Wisconsin	0.34	0.05	(28)	-1.89
Mississippi	0.50	0.08	(02)	41.81 <sup>c</sup>	Wyoming	0.41	0.17	(12)	16.49 <sup>c</sup>
Missouri	0.37	0.05	(22)	6.17					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.



Table 11.18  
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.08	0.02		
High Five States				
Louisiana	5.27 <sup>b</sup>	0.16	(01)	29.22 <sup>c</sup>
North Dakota	5.15	0.38	(02)	26.22 <sup>c</sup>
Oklahoma	5.09 <sup>b</sup>	0.16	(03)	24.96 <sup>c</sup>
South Dakota	4.89	0.33	(04)	19.96 <sup>c</sup>
Iowa	4.87 <sup>b</sup>	0.17	(05)	19.45 <sup>c</sup>
Low Five States				
Connecticut	3.49 <sup>b</sup>	0.13	(47)	-14.40
New York	3.44 <sup>b</sup>	0.06	(48)	-15.60 <sup>c</sup>
Utah	3.23 <sup>b</sup>	0.19	(49)	-20.87 <sup>c</sup>
D.C.	2.91 <sup>b</sup>	0.32	(50)	-28.62 <sup>c</sup>
Hawaii	2.59 <sup>b</sup>	0.19	(51)	-36.48 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.99	0.13	(34)	-2.19	Montana	4.00	0.28	(33)	-1.89
Alaska	4.20	0.48	(26)	3.13	Nebraska	4.52	0.22	(12)	10.98
Arizona	3.77	0.11	(42)	-7.44	Nevada	3.94	0.19	(36)	-3.35
Arkansas	4.32	0.17	(19)	6.03	New Hampshire	4.21	0.25	(25)	3.32
California	3.65 <sup>b</sup>	0.05	(44)	-10.45	New Jersey	3.63 <sup>b</sup>	0.09	(45)	-11.04
Colorado	3.86	0.14	(39)	-5.35	New Mexico	4.45	0.21	(15)	9.04
Connecticut	3.49 <sup>b</sup>	0.13	(47)	-14.40	New York	3.44 <sup>b</sup>	0.06	(48)	-15.60 <sup>c</sup>
Delaware	4.07	0.30	(30)	-0.17	North Carolina	4.27	0.10	(21)	4.72
D.C.	2.91 <sup>b</sup>	0.32	(50)	-28.62 <sup>c</sup>	North Dakota	5.15	0.38	(02)	26.22 <sup>c</sup>
Florida	3.51 <sup>b</sup>	0.06	(46)	-14.01	Ohio	4.56 <sup>b</sup>	0.09	(10)	11.76
Georgia	3.97	0.10	(35)	-2.50	Oklahoma	5.09 <sup>b</sup>	0.16	(03)	24.96 <sup>c</sup>
Hawaii	2.59 <sup>b</sup>	0.19	(51)	-36.48 <sup>c</sup>	Oregon	3.75	0.14	(43)	-8.03
Idaho	4.13	0.25	(29)	1.30	Pennsylvania	4.14	0.07	(28)	1.60
Illinois	4.29	0.08	(20)	5.17	Rhode Island	4.22	0.27	(24)	3.42
Indiana	4.75 <sup>b</sup>	0.12	(06)	16.63 <sup>c</sup>	South Carolina	4.25	0.14	(23)	4.21
Iowa	4.87 <sup>b</sup>	0.17	(05)	19.45 <sup>c</sup>	South Dakota	4.89	0.33	(04)	19.96 <sup>c</sup>
Kansas	4.54	0.18	(11)	11.30	Tennessee	4.37	0.12	(18)	7.12
Kentucky	4.71 <sup>b</sup>	0.15	(07)	15.52 <sup>c</sup>	Texas	4.67 <sup>b</sup>	0.07	(08)	14.52
Louisiana	5.27 <sup>b</sup>	0.16	(01)	29.22 <sup>c</sup>	Utah	3.23 <sup>b</sup>	0.19	(49)	-20.87 <sup>c</sup>
Maine	4.16	0.23	(27)	1.99	Vermont	4.02	0.34	(32)	-1.28
Maryland	3.80	0.12	(40)	-6.88	Virginia	3.92	0.10	(37)	-3.81
Massachusetts	3.88	0.10	(38)	-4.84	Washington	4.05	0.11	(31)	-0.77
Michigan	4.26	0.09	(22)	4.48	West Virginia	4.49	0.20	(14)	10.14
Minnesota	4.43	0.13	(17)	8.57	Wisconsin	4.44	0.12	(16)	8.95
Mississippi	4.52	0.18	(13)	10.91	Wyoming	3.78	0.38	(41)	-7.39
Missouri	4.60 <sup>b</sup>	0.12	(09)	12.87					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.



Table 11.19  
Cancer of the Kidney and Renal Pelvis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.89	0.03		
High Five States				
North Dakota	7.46	0.68	(01)	26.80 <sup>c</sup>
Louisiana	7.44 <sup>b</sup>	0.29	(02)	26.39 <sup>c</sup>
South Dakota	7.42	0.62	(03)	26.06 <sup>c</sup>
Oklahoma	7.25 <sup>b</sup>	0.30	(04)	23.10 <sup>c</sup>
Kentucky	7.03 <sup>b</sup>	0.28	(05)	19.37 <sup>c</sup>
Low Five States				
New York	5.06 <sup>b</sup>	0.11	(47)	-14.08
Connecticut	4.97 <sup>b</sup>	0.25	(48)	-15.54 <sup>c</sup>
Utah	4.34 <sup>b</sup>	0.32	(49)	-26.26 <sup>c</sup>
D.C.	3.98	0.58	(50)	-32.45 <sup>c</sup>
Hawaii	3.70 <sup>b</sup>	0.34	(51)	-37.12 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.66	0.24	(37)	-3.79	Montana	5.79	0.50	(32)	-1.62
Alaska	5.99	0.85	(27)	1.70	Nebraska	6.14	0.39	(22)	4.33
Arizona	5.27	0.19	(44)	-10.44	Nevada	5.56	0.33	(39)	-5.59
Arkansas	6.09	0.30	(24)	3.38	New Hampshire	5.85	0.46	(31)	-0.59
California	5.32 <sup>b</sup>	0.09	(42)	-9.54	New Jersey	5.31	0.17	(43)	-9.84
Colorado	5.73	0.26	(34)	-2.69	New Mexico	6.21	0.38	(20)	5.54
Connecticut	4.97 <sup>b</sup>	0.25	(48)	-15.54 <sup>c</sup>	New York	5.06 <sup>b</sup>	0.11	(47)	-14.08
Delaware	6.21	0.57	(19)	5.55	North Carolina	6.43	0.19	(13)	9.30
D.C.	3.98	0.58	(50)	-32.45 <sup>c</sup>	North Dakota	7.46	0.68	(01)	26.80 <sup>c</sup>
Florida	5.12 <sup>b</sup>	0.10	(46)	-12.98	Ohio	6.54 <sup>b</sup>	0.16	(11)	11.16
Georgia	5.78	0.20	(33)	-1.80	Oklahoma	7.25 <sup>b</sup>	0.30	(04)	23.10 <sup>c</sup>
Hawaii	3.70 <sup>b</sup>	0.34	(51)	-37.12 <sup>c</sup>	Oregon	5.45	0.25	(41)	-7.34
Idaho	5.65	0.43	(38)	-4.05	Pennsylvania	5.90	0.14	(29)	0.26
Illinois	6.33	0.16	(15)	7.57	Rhode Island	5.88	0.49	(30)	-0.10
Indiana	6.67	0.22	(08)	13.23	South Carolina	6.20	0.26	(21)	5.41
Iowa	6.82	0.30	(06)	15.92 <sup>c</sup>	South Dakota	7.42	0.62	(03)	26.06 <sup>c</sup>
Kansas	6.64	0.33	(10)	12.76	Tennessee	6.45	0.23	(12)	9.58
Kentucky	7.03 <sup>b</sup>	0.28	(05)	19.37 <sup>c</sup>	Texas	6.66 <sup>b</sup>	0.13	(09)	13.13
Louisiana	7.44 <sup>b</sup>	0.29	(02)	26.39 <sup>c</sup>	Utah	4.34 <sup>b</sup>	0.32	(49)	-26.26 <sup>c</sup>
Maine	5.70	0.41	(35)	-3.09	Vermont	5.23	0.60	(45)	-11.21
Maryland	5.54	0.22	(40)	-5.90	Virginia	6.04	0.20	(25)	2.59
Massachusetts	5.97	0.20	(28)	1.50	Washington	5.67	0.21	(36)	-3.72
Michigan	5.99	0.17	(26)	1.76	West Virginia	6.11	0.36	(23)	3.88
Minnesota	6.32	0.24	(16)	7.31	Wisconsin	6.29	0.22	(17)	6.85
Mississippi	6.40	0.33	(14)	8.76	Wyoming	6.26	0.73	(18)	6.28
Missouri	6.70 <sup>b</sup>	0.23	(07)	13.84					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.





Table 11.20  
Cancer of the Kidney And Renal Pelvis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.69	0.02		
High Five States				
Louisiana	3.60 <sup>b</sup>	0.17	(01)	34.08 <sup>c</sup>
Iowa	3.37 <sup>b</sup>	0.19	(02)	25.29 <sup>c</sup>
Oklahoma	3.34 <sup>b</sup>	0.18	(03)	24.38 <sup>c</sup>
North Dakota	3.34	0.42	(04)	24.29 <sup>c</sup>
Indiana	3.28 <sup>b</sup>	0.13	(05)	22.17 <sup>c</sup>
Low Five States				
New York	2.26 <sup>b</sup>	0.06	(47)	-15.80 <sup>c</sup>
Florida	2.19 <sup>b</sup>	0.06	(48)	-18.64 <sup>c</sup>
D.C.	1.98	0.34	(49)	-26.14 <sup>c</sup>
Hawaii	1.67 <sup>b</sup>	0.21	(50)	-37.82 <sup>c</sup>
Wyoming	1.60	0.34	(51)	-40.52 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.67	0.14	(31)	-0.57	Montana	2.56	0.30	(35)	-4.72
Alaska	2.56	0.52	(34)	-4.72	Nebraska	3.25	0.25	(06)	21.13 <sup>c</sup>
Arizona	2.50	0.12	(37)	-6.84	Nevada	2.50	0.21	(38)	-6.85
Arkansas	2.92	0.18	(21)	8.79	New Hampshire	2.98	0.28	(17)	10.82
California	2.35 <sup>b</sup>	0.05	(44)	-12.48	New Jersey	2.40	0.09	(43)	-10.65
Colorado	2.44	0.15	(39)	-9.16	New Mexico	3.00	0.24	(15)	11.69
Connecticut	2.41	0.15	(41)	-10.12	New York	2.26 <sup>b</sup>	0.06	(47)	-15.80 <sup>c</sup>
Delaware	2.56	0.32	(33)	-4.57	North Carolina	2.67	0.10	(32)	-0.74
D.C.	1.98	0.34	(49)	-26.14 <sup>c</sup>	North Dakota	3.34	0.42	(04)	24.29 <sup>c</sup>
Florida	2.19 <sup>b</sup>	0.06	(48)	-18.64 <sup>c</sup>	Ohio	3.10 <sup>b</sup>	0.09	(08)	15.57 <sup>c</sup>
Georgia	2.70	0.11	(30)	0.70	Oklahoma	3.34 <sup>b</sup>	0.18	(03)	24.38 <sup>c</sup>
Hawaii	1.67 <sup>b</sup>	0.21	(50)	-37.82 <sup>c</sup>	Oregon	2.40	0.15	(42)	-10.63
Idaho	2.82	0.28	(25)	5.05	Pennsylvania	2.80	0.08	(26)	4.18
Illinois	2.75	0.09	(28)	2.52	Rhode Island	3.08	0.30	(11)	14.68
Indiana	3.28 <sup>b</sup>	0.13	(05)	22.17 <sup>c</sup>	South Carolina	2.80	0.15	(27)	4.12
Iowa	3.37 <sup>b</sup>	0.19	(02)	25.29 <sup>c</sup>	South Dakota	2.85	0.35	(23)	6.15
Kansas	2.92	0.19	(20)	8.88	Tennessee	2.82	0.13	(24)	5.14
Kentucky	2.95	0.15	(19)	9.82	Texas	3.10 <sup>b</sup>	0.08	(10)	15.43 <sup>c</sup>
Louisiana	3.60 <sup>b</sup>	0.17	(01)	34.08 <sup>c</sup>	Utah	2.27	0.21	(46)	-15.45 <sup>c</sup>
Maine	3.02	0.27	(14)	12.33	Vermont	3.08	0.41	(12)	14.68
Maryland	2.52	0.13	(36)	-6.13	Virginia	2.33	0.11	(45)	-13.21
Massachusetts	2.41	0.11	(40)	-10.11	Washington	2.73	0.13	(29)	1.55
Michigan	2.96	0.10	(18)	10.24	West Virginia	3.19	0.22	(07)	18.73 <sup>c</sup>
Minnesota	2.91	0.14	(22)	8.28	Wisconsin	2.98	0.13	(16)	10.94
Mississippi	3.10	0.19	(09)	15.45 <sup>c</sup>	Wyoming	1.60	0.34	(51)	-40.52 <sup>c</sup>
Missouri	3.06	0.13	(13)	13.76					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 12.18  
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.22	0.01		
High Five States				
D.C.	2.07	0.27	(01)	68.90 <sup>c</sup>
West Virginia	1.84 <sup>b</sup>	0.13	(02)	50.48 <sup>c</sup>
Louisiana	1.79 <sup>b</sup>	0.09	(03)	46.29 <sup>c</sup>
Kentucky	1.55 <sup>b</sup>	0.08	(04)	27.03 <sup>c</sup>
Tennessee	1.54 <sup>b</sup>	0.07	(05)	25.77 <sup>c</sup>
Low Five States				
Hawaii	0.86	0.11	(47)	-29.30 <sup>c</sup>
Oregon	0.80 <sup>b</sup>	0.06	(48)	-34.39 <sup>c</sup>
Colorado	0.77 <sup>b</sup>	0.06	(49)	-37.35 <sup>c</sup>
Alaska	0.43 <sup>b</sup>	0.15	(50)	-64.92 <sup>c</sup>
Utah	0.41 <sup>b</sup>	0.07	(51)	-66.41 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.42	0.08	(10)	16.02 <sup>c</sup>	Montana	0.92	0.13	(43)	-24.65 <sup>c</sup>
Alaska	0.43 <sup>b</sup>	0.15	(50)	-64.92 <sup>c</sup>	Nebraska	1.02	0.11	(34)	-16.61 <sup>c</sup>
Arizona	0.93 <sup>b</sup>	0.06	(42)	-23.75 <sup>c</sup>	Nevada	1.16	0.10	(28)	-4.86
Arkansas	1.23	0.09	(22)	0.47	New Hampshire	1.13	0.13	(29)	-7.47
California	0.96 <sup>b</sup>	0.02	(40)	-21.30 <sup>c</sup>	New Jersey	1.30	0.05	(16)	6.11
Colorado	0.77 <sup>b</sup>	0.06	(49)	-37.35 <sup>c</sup>	New Mexico	1.02	0.10	(35)	-16.97 <sup>c</sup>
Connecticut	1.03	0.07	(33)	-15.84 <sup>c</sup>	New York	1.27	0.04	(18)	3.85
Delaware	1.53	0.18	(06)	24.97 <sup>c</sup>	North Carolina	1.37	0.06	(14)	11.74
D.C.	2.07	0.27	(01)	68.90 <sup>c</sup>	North Dakota	0.97	0.17	(39)	-20.61 <sup>c</sup>
Florida	1.44 <sup>b</sup>	0.04	(08)	17.34 <sup>c</sup>	Ohio	1.43 <sup>b</sup>	0.05	(09)	16.53 <sup>c</sup>
Georgia	1.27	0.06	(19)	3.54	Oklahoma	1.21	0.08	(25)	-1.07
Hawaii	0.86	0.11	(47)	-29.30 <sup>c</sup>	Oregon	0.80 <sup>b</sup>	0.06	(48)	-34.39 <sup>c</sup>
Idaho	0.98	0.12	(38)	-20.09 <sup>c</sup>	Pennsylvania	1.33	0.04	(15)	8.36
Illinois	1.29	0.05	(17)	5.36	Rhode Island	1.11	0.14	(30)	-9.17
Indiana	1.17	0.06	(27)	-4.23	South Carolina	1.40	0.08	(11)	14.45
Iowa	1.04	0.08	(32)	-15.26 <sup>c</sup>	South Dakota	0.88	0.14	(46)	-27.88 <sup>c</sup>
Kansas	0.99	0.08	(36)	-19.22 <sup>c</sup>	Tennessee	1.54 <sup>b</sup>	0.07	(05)	25.77 <sup>c</sup>
Kentucky	1.55 <sup>b</sup>	0.08	(04)	27.03 <sup>c</sup>	Texas	1.24	0.04	(21)	1.74
Louisiana	1.79 <sup>b</sup>	0.09	(03)	46.29 <sup>c</sup>	Utah	0.41 <sup>b</sup>	0.07	(51)	-66.41 <sup>c</sup>
Maine	1.25	0.13	(20)	2.55	Vermont	0.99	0.17	(37)	-19.34 <sup>c</sup>
Maryland	1.44	0.07	(07)	17.37 <sup>c</sup>	Virginia	1.18	0.06	(26)	-3.13
Massachusetts	1.22	0.06	(23)	-0.54	Washington	0.94 <sup>b</sup>	0.06	(41)	-23.48 <sup>c</sup>
Michigan	1.40	0.05	(12)	14.42	West Virginia	1.84 <sup>b</sup>	0.13	(02)	50.48 <sup>c</sup>
Minnesota	0.90 <sup>b</sup>	0.06	(45)	-26.46 <sup>c</sup>	Wisconsin	1.11	0.06	(31)	-9.53
Mississippi	1.38	0.10	(13)	13.14	Wyoming	0.91	0.19	(44)	-25.63 <sup>c</sup>
Missouri	1.21	0.06	(24)	-0.92					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.



Table 12.19  
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	2.21	0.02		
High Five States				
D.C.	4.13	0.60	(01)	86.57 <sup>c</sup>
Louisiana	3.22 <sup>b</sup>	0.19	(02)	45.29 <sup>c</sup>
West Virginia	3.13 <sup>b</sup>	0.25	(03)	41.12 <sup>c</sup>
Tennessee	2.90 <sup>b</sup>	0.15	(04)	30.92 <sup>c</sup>
Delaware	2.84	0.37	(05)	28.08 <sup>c</sup>
Low Five States				
Montana	1.35 <sup>b</sup>	0.24	(47)	-39.11 <sup>c</sup>
Colorado	1.34 <sup>b</sup>	0.13	(48)	-39.35 <sup>c</sup>
Oregon	1.29 <sup>b</sup>	0.12	(49)	-41.57 <sup>c</sup>
Alaska	0.73 <sup>b</sup>	0.27	(50)	-66.88 <sup>c</sup>
Utah	0.64 <sup>b</sup>	0.12	(51)	-71.27 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.62	0.16	(10)	18.23 <sup>c</sup>	Montana	1.35 <sup>b</sup>	0.24	(47)	-39.11 <sup>c</sup>
Alaska	0.73 <sup>b</sup>	0.27	(50)	-66.88 <sup>c</sup>	Nebraska	1.88	0.21	(33)	-15.13 <sup>c</sup>
Arizona	1.60 <sup>b</sup>	0.11	(43)	-27.78 <sup>c</sup>	Nevada	1.93	0.20	(31)	-12.78
Arkansas	2.30	0.19	(21)	3.75	New Hampshire	1.92	0.26	(32)	-13.43
California	1.78 <sup>b</sup>	0.05	(37)	-19.74 <sup>c</sup>	New Jersey	2.41	0.11	(15)	8.70
Colorado	1.34 <sup>b</sup>	0.13	(48)	-39.35 <sup>c</sup>	New Mexico	1.86	0.21	(34)	-16.07 <sup>c</sup>
Connecticut	1.96	0.16	(29)	-11.48	New York	2.35	0.07	(17)	6.29
Delaware	2.84	0.37	(05)	28.08 <sup>c</sup>	North Carolina	2.55	0.12	(12)	15.32 <sup>c</sup>
D.C.	4.13	0.60	(01)	86.57 <sup>c</sup>	North Dakota	1.47	0.30	(46)	-33.49 <sup>c</sup>
Florida	2.51 <sup>b</sup>	0.07	(13)	13.46	Ohio	2.58 <sup>b</sup>	0.10	(11)	16.35 <sup>c</sup>
Georgia	2.35	0.12	(18)	5.94	Oklahoma	2.05	0.16	(27)	-7.38
Hawaii	1.66	0.23	(39)	-24.90 <sup>c</sup>	Oregon	1.29 <sup>b</sup>	0.12	(49)	-41.57 <sup>c</sup>
Idaho	1.61	0.23	(42)	-27.19 <sup>c</sup>	Pennsylvania	2.39	0.09	(16)	8.00
Illinois	2.31	0.09	(20)	4.50	Rhode Island	2.03	0.29	(28)	-8.42
Indiana	2.15	0.13	(25)	-3.12	South Carolina	2.63	0.17	(09)	18.67 <sup>c</sup>
Iowa	1.84	0.16	(35)	-16.84 <sup>c</sup>	South Dakota	1.64	0.29	(40)	-25.92 <sup>c</sup>
Kansas	1.83	0.17	(36)	-17.32 <sup>c</sup>	Tennessee	2.90 <sup>b</sup>	0.15	(04)	30.92 <sup>c</sup>
Kentucky	2.83	0.18	(07)	27.71 <sup>c</sup>	Texas	2.34	0.08	(19)	5.83
Louisiana	3.22 <sup>b</sup>	0.19	(02)	45.29 <sup>c</sup>	Utah	0.64 <sup>b</sup>	0.12	(51)	-71.27 <sup>c</sup>
Maine	2.19	0.26	(24)	-1.12	Vermont	1.77	0.33	(38)	-20.11 <sup>c</sup>
Maryland	2.69	0.15	(08)	21.27 <sup>c</sup>	Virginia	2.26	0.12	(23)	2.11
Massachusetts	2.11	0.12	(26)	-4.72	Washington	1.57 <sup>b</sup>	0.11	(44)	-28.96 <sup>c</sup>
Michigan	2.46	0.10	(14)	11.16	West Virginia	3.13 <sup>b</sup>	0.25	(03)	41.12 <sup>c</sup>
Minnesota	1.63 <sup>b</sup>	0.12	(41)	-26.21 <sup>c</sup>	Wisconsin	1.96	0.12	(30)	-11.57
Mississippi	2.83	0.21	(06)	27.77 <sup>c</sup>	Wyoming	1.54	0.36	(45)	-30.62 <sup>c</sup>
Missouri	2.27	0.13	(22)	2.46					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.



Table 12.20  
Cancer of the Larynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.46	0.01		
High Five States				
West Virginia	0.79	0.12	(01)	71.12 <sup>c</sup>
Louisiana	0.70	0.08	(02)	53.17 <sup>c</sup>
D.C.	0.68	0.20	(03)	47.18 <sup>c</sup>
Kentucky	0.60	0.07	(04)	31.41 <sup>c</sup>
Montana	0.59	0.15	(05)	27.96 <sup>c</sup>
Low Five States				
Vermont	0.28	0.12	(47)	-38.53 <sup>c</sup>
South Dakota	0.25	0.11	(48)	-44.45 <sup>c</sup>
Utah	0.23	0.07	(49)	-50.01 <sup>c</sup>
Hawaii	0.20 <sup>b</sup>	0.07	(50)	-57.24 <sup>c</sup>
Alaska	0.12	0.12	(51)	-73.19 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.51	0.06	(17)	11.81	Montana	0.59	0.15	(05)	27.96 <sup>c</sup>
Alaska	0.12	0.12	(51)	-73.19 <sup>c</sup>	Nebraska	0.35	0.09	(39)	-23.15 <sup>c</sup>
Arizona	0.37	0.05	(37)	-19.56 <sup>c</sup>	Nevada	0.51	0.09	(16)	12.01
Arkansas	0.39	0.07	(35)	-14.13	New Hampshire	0.51	0.12	(18)	10.14
California	0.35 <sup>b</sup>	0.02	(41)	-24.33 <sup>c</sup>	New Jersey	0.48	0.04	(24)	5.46
Colorado	0.32	0.05	(45)	-30.65 <sup>c</sup>	New Mexico	0.32	0.08	(44)	-30.45 <sup>c</sup>
Connecticut	0.35	0.06	(40)	-23.64 <sup>c</sup>	New York	0.47	0.03	(25)	1.61
Delaware	0.43	0.13	(31)	-6.79	North Carolina	0.50	0.04	(19)	9.82
D.C.	0.68	0.20	(03)	47.18 <sup>c</sup>	North Dakota	0.56	0.17	(06)	22.61 <sup>c</sup>
Florida	0.54	0.03	(11)	17.78 <sup>c</sup>	Ohio	0.54	0.04	(10)	17.91 <sup>c</sup>
Georgia	0.48	0.05	(23)	5.47	Oklahoma	0.55	0.07	(09)	19.43 <sup>c</sup>
Hawaii	0.20 <sup>b</sup>	0.07	(50)	-57.24 <sup>c</sup>	Oregon	0.40	0.06	(33)	-11.87
Idaho	0.50	0.12	(20)	8.91	Pennsylvania	0.53	0.04	(13)	16.06 <sup>c</sup>
Illinois	0.51	0.04	(15)	12.07	Rhode Island	0.49	0.12	(21)	7.42
Indiana	0.46	0.05	(27)	-0.45	South Carolina	0.46	0.06	(26)	-0.27
Iowa	0.43	0.07	(30)	-6.25	South Dakota	0.25	0.11	(48)	-44.45 <sup>c</sup>
Kansas	0.33	0.06	(42)	-27.90 <sup>c</sup>	Tennessee	0.49	0.05	(22)	6.19
Kentucky	0.60	0.07	(04)	31.41 <sup>c</sup>	Texas	0.40	0.03	(34)	-13.12
Louisiana	0.70	0.08	(02)	53.17 <sup>c</sup>	Utah	0.23	0.07	(49)	-50.01 <sup>c</sup>
Maine	0.54	0.11	(12)	17.20 <sup>c</sup>	Vermont	0.28	0.12	(47)	-38.53 <sup>c</sup>
Maryland	0.52	0.06	(14)	13.66	Virginia	0.38	0.04	(36)	-17.81 <sup>c</sup>
Massachusetts	0.55	0.05	(08)	19.65 <sup>c</sup>	Washington	0.44	0.05	(28)	-4.80
Michigan	0.56	0.04	(07)	22.43 <sup>c</sup>	West Virginia	0.79	0.12	(01)	71.12 <sup>c</sup>
Minnesota	0.32	0.05	(43)	-29.94 <sup>c</sup>	Wisconsin	0.43	0.05	(29)	-5.95
Mississippi	0.29	0.06	(46)	-37.02 <sup>c</sup>	Wyoming	0.36	0.16	(38)	-21.89 <sup>c</sup>
Missouri	0.41	0.05	(32)	-11.55					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 13.26  
Leukemia

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	7.20	0.02		
High Five States				
Idaho	8.33	0.35	(01)	15.78 <sup>c</sup>
Wisconsin	8.22 <sup>b</sup>	0.17	(02)	14.25
Iowa	8.14 <sup>b</sup>	0.22	(03)	13.14
Missouri	7.95 <sup>b</sup>	0.16	(04)	10.47
Kansas	7.92	0.23	(05)	10.06
Low Five States				
D.C.	6.63	0.48	(47)	-7.82
Nevada	6.59	0.25	(48)	-8.43
New Mexico	6.36	0.26	(49)	-11.66
Arizona	6.35 <sup>b</sup>	0.14	(50)	-11.68
Hawaii	4.96 <sup>b</sup>	0.27	(51)	-31.13 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.37	0.18	(23)	2.40	Montana	7.74	0.39	(12)	7.52
Alaska	7.29	0.65	(26)	1.37	Nebraska	7.27	0.28	(27)	1.01
Arizona	6.35 <sup>b</sup>	0.14	(50)	-11.68	Nevada	6.59	0.25	(48)	-8.43
Arkansas	7.68	0.22	(14)	6.70	New Hampshire	6.79	0.32	(42)	-5.57
California	6.76 <sup>b</sup>	0.06	(44)	-6.09	New Jersey	7.06	0.12	(33)	-1.87
Colorado	6.74	0.19	(45)	-6.30	New Mexico	6.36	0.26	(49)	-11.66
Connecticut	6.97	0.19	(36)	-3.19	New York	6.76 <sup>b</sup>	0.08	(43)	-6.03
Delaware	6.99	0.40	(34)	-2.91	North Carolina	7.09	0.13	(31)	-1.52
D.C.	6.63	0.48	(47)	-7.82	North Dakota	7.15	0.44	(29)	-0.63
Florida	6.87 <sup>b</sup>	0.08	(40)	-4.46	Ohio	7.71 <sup>b</sup>	0.11	(13)	7.12
Georgia	7.07	0.14	(32)	-1.73	Oklahoma	7.74	0.20	(11)	7.57
Hawaii	4.96 <sup>b</sup>	0.27	(51)	-31.13 <sup>c</sup>	Oregon	7.25	0.19	(28)	0.79
Idaho	8.33	0.35	(01)	15.78 <sup>c</sup>	Pennsylvania	7.56 <sup>b</sup>	0.10	(18)	5.02
Illinois	7.55	0.11	(19)	4.95	Rhode Island	7.35	0.35	(24)	2.10
Indiana	7.77 <sup>b</sup>	0.16	(09)	8.04	South Carolina	6.95	0.18	(37)	-3.37
Iowa	8.14 <sup>b</sup>	0.22	(03)	13.14	South Dakota	7.81	0.42	(07)	8.61
Kansas	7.92	0.23	(05)	10.06	Tennessee	7.55	0.16	(20)	4.88
Kentucky	7.50	0.19	(21)	4.22	Texas	6.97	0.09	(35)	-3.09
Louisiana	7.12	0.18	(30)	-1.04	Utah	6.72	0.27	(46)	-6.58
Maine	7.61	0.32	(17)	5.69	Vermont	7.64	0.48	(16)	6.14
Maryland	6.87	0.16	(41)	-4.57	Virginia	6.92	0.14	(38)	-3.81
Massachusetts	6.91	0.14	(39)	-4.00	Washington	7.46	0.16	(22)	3.69
Michigan	7.79 <sup>b</sup>	0.12	(08)	8.27	West Virginia	7.64	0.27	(15)	6.21
Minnesota	7.82 <sup>b</sup>	0.18	(06)	8.63	Wisconsin	8.22 <sup>b</sup>	0.17	(02)	14.25
Mississippi	7.75	0.23	(10)	7.74	Wyoming	7.33	0.54	(25)	1.81
Missouri	7.95 <sup>b</sup>	0.16	(04)	10.47					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 13.27  
Leukemia

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	9.68	0.04		
High Five States				
Wisconsin	11.26 <sup>b</sup>	0.30	(01)	16.37 <sup>c</sup>
Iowa	11.20 <sup>b</sup>	0.39	(02)	15.69 <sup>c</sup>
Missouri	11.02 <sup>b</sup>	0.29	(03)	13.89
South Dakota	10.87	0.75	(04)	12.33
Oklahoma	10.85	0.38	(05)	12.07
Low Five States				
Nevada	8.83	0.43	(47)	-8.78
D.C.	8.53	0.87	(48)	-11.82
New Mexico	8.28	0.45	(49)	-14.42
Arizona	8.17 <sup>b</sup>	0.24	(50)	-15.61 <sup>c</sup>
Hawaii	6.78 <sup>b</sup>	0.47	(51)	-29.91 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.91	0.32	(24)	2.43	Montana	10.60	0.69	(09)	9.57
Alaska	9.32	1.14	(37)	-3.70	Nebraska	9.92	0.50	(23)	2.48
Arizona	8.17 <sup>b</sup>	0.24	(50)	-15.61 <sup>c</sup>	Nevada	8.83	0.43	(47)	-8.78
Arkansas	10.29	0.40	(15)	6.28	New Hampshire	9.02	0.58	(43)	-6.78
California	8.91 <sup>b</sup>	0.11	(45)	-7.95	New Jersey	9.57	0.23	(30)	-1.12
Colorado	9.00	0.34	(44)	-7.00	New Mexico	8.28	0.45	(49)	-14.42
Connecticut	9.51	0.35	(32)	-1.72	New York	9.18	0.15	(41)	-5.17
Delaware	9.29	0.71	(38)	-4.00	North Carolina	9.82	0.24	(27)	1.42
D.C.	8.53	0.87	(48)	-11.82	North Dakota	10.18	0.80	(19)	5.16
Florida	9.09 <sup>b</sup>	0.13	(42)	-6.12	Ohio	10.28	0.20	(16)	6.23
Georgia	9.44	0.26	(33)	-2.42	Oklahoma	10.85	0.38	(05)	12.07
Hawaii	6.78 <sup>b</sup>	0.47	(51)	-29.91 <sup>c</sup>	Oregon	9.57	0.34	(31)	-1.16
Idaho	10.38	0.59	(14)	7.22	Pennsylvania	10.15	0.18	(20)	4.84
Illinois	10.38	0.20	(13)	7.28	Rhode Island	9.25	0.62	(39)	-4.43
Indiana	10.74 <sup>b</sup>	0.29	(08)	10.95	South Carolina	9.59	0.33	(28)	-0.91
Iowa	11.20 <sup>b</sup>	0.39	(02)	15.69 <sup>c</sup>	South Dakota	10.87	0.75	(04)	12.33
Kansas	10.53	0.42	(12)	8.81	Tennessee	10.59	0.30	(10)	9.45
Kentucky	9.85	0.34	(26)	1.82	Texas	9.41	0.16	(34)	-2.78
Louisiana	9.58	0.33	(29)	-1.03	Utah	8.90	0.47	(46)	-8.04
Maine	10.79	0.58	(06)	11.50	Vermont	10.24	0.86	(18)	5.85
Maryland	9.32	0.30	(36)	-3.66	Virginia	9.18	0.26	(40)	-5.13
Massachusetts	9.37	0.26	(35)	-3.13	Washington	10.26	0.28	(17)	5.98
Michigan	10.58 <sup>b</sup>	0.22	(11)	9.35	West Virginia	10.00	0.47	(22)	3.37
Minnesota	10.75	0.32	(07)	11.10	Wisconsin	11.26 <sup>b</sup>	0.30	(01)	16.37 <sup>c</sup>
Mississippi	10.14	0.42	(21)	4.75	Wyoming	9.86	0.95	(25)	1.84
Missouri	11.02 <sup>b</sup>	0.29	(03)	13.89					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.




Table 13.28  
Leukemia

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.42	0.03		
High Five States				
Idaho	6.65	0.42	(01)	22.53 <sup>c</sup>
Mississippi	6.15	0.27	(02)	13.34
Rhode Island	6.12	0.43	(03)	12.85
Kansas	6.10	0.27	(04)	12.55
Iowa	6.09	0.25	(05)	12.23
Low Five States				
Utah	5.00	0.31	(47)	-7.91
New Mexico	4.87	0.31	(48)	-10.31
Arizona	4.85	0.17	(49)	-10.63
Nevada	4.74	0.29	(50)	-12.69
Hawaii	3.46 <sup>b</sup>	0.30	(51)	-36.27 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.68	0.20	(18)	4.74	Montana	5.56	0.44	(20)	2.55
Alaska	5.71	0.76	(16)	5.33	Nebraska	5.33	0.31	(32)	-1.66
Arizona	4.85	0.17	(49)	-10.63	Nevada	4.74	0.29	(50)	-12.69
Arkansas	5.84	0.26	(09)	7.69	New Hampshire	5.29	0.38	(34)	-2.39
California	5.17 <sup>b</sup>	0.08	(40)	-4.74	New Jersey	5.34	0.14	(31)	-1.60
Colorado	5.15	0.21	(41)	-5.04	New Mexico	4.87	0.31	(48)	-10.31
Connecticut	5.24	0.22	(36)	-3.38	New York	5.11	0.09	(45)	-5.82
Delaware	5.41	0.47	(26)	-0.30	North Carolina	5.21	0.14	(38)	-3.86
D.C.	5.15	0.55	(42)	-5.09	North Dakota	5.13	0.51	(44)	-5.50
Florida	5.10 <sup>b</sup>	0.09	(46)	-6.06	Ohio	5.94 <sup>b</sup>	0.13	(07)	9.55
Georgia	5.51	0.16	(22)	1.53	Oklahoma	5.50	0.23	(23)	1.31
Hawaii	3.46 <sup>b</sup>	0.30	(51)	-36.27 <sup>c</sup>	Oregon	5.47	0.23	(24)	0.88
Idaho	6.65	0.42	(01)	22.53 <sup>c</sup>	Pennsylvania	5.77	0.12	(15)	6.29
Illinois	5.66	0.12	(19)	4.44	Rhode Island	6.12	0.43	(03)	12.85
Indiana	5.81	0.18	(12)	7.18	South Carolina	5.15	0.20	(43)	-5.14
Iowa	6.09	0.25	(05)	12.23	South Dakota	5.46	0.47	(25)	0.59
Kansas	6.10	0.27	(04)	12.55	Tennessee	5.55	0.18	(21)	2.41
Kentucky	5.82	0.22	(11)	7.36	Texas	5.22	0.10	(37)	-3.74
Louisiana	5.40	0.21	(27)	-0.43	Utah	5.00	0.31	(47)	-7.91
Maine	5.34	0.35	(30)	-1.55	Vermont	5.71	0.56	(17)	5.30
Maryland	5.17	0.18	(39)	-4.74	Virginia	5.36	0.16	(29)	-1.23
Massachusetts	5.32	0.16	(33)	-1.93	Washington	5.40	0.18	(28)	-0.44
Michigan	5.78	0.14	(14)	6.49	West Virginia	5.93	0.31	(08)	9.39
Minnesota	5.83	0.20	(10)	7.54	Wisconsin	6.01	0.19	(06)	10.71
Mississippi	6.15	0.27	(02)	13.34	Wyoming	5.25	0.60	(35)	-3.14
Missouri	5.79	0.18	(13)	6.67					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.



Table 14.18  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.21	0.02		
High Five States				
Hawaii	7.31 <sup>b</sup>	0.32	(01)	40.24 <sup>c</sup>
Texas	6.88 <sup>b</sup>	0.08	(02)	32.01 <sup>c</sup>
Louisiana	6.81 <sup>b</sup>	0.18	(03)	30.58 <sup>c</sup>
California	6.46 <sup>b</sup>	0.06	(04)	23.99 <sup>c</sup>
Mississippi	6.29 <sup>b</sup>	0.21	(05)	20.65 <sup>c</sup>
Low Five States				
Iowa	3.86 <sup>b</sup>	0.15	(47)	-25.88 <sup>c</sup>
Nebraska	3.83 <sup>b</sup>	0.20	(48)	-26.49 <sup>c</sup>
Montana	3.79 <sup>b</sup>	0.27	(49)	-27.28 <sup>c</sup>
Vermont	3.60 <sup>b</sup>	0.33	(50)	-30.99 <sup>c</sup>
Utah	3.46 <sup>b</sup>	0.19	(51)	-33.54 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.23	0.15	(15)	0.30	Montana	3.79 <sup>b</sup>	0.27	(49)	-27.28 <sup>c</sup>
Alaska	6.08	0.57	(08)	16.67 <sup>c</sup>	Nebraska	3.83 <sup>b</sup>	0.20	(48)	-26.49 <sup>c</sup>
Arizona	4.93	0.13	(21)	-5.46	Nevada	5.87	0.23	(09)	12.63
Arkansas	5.70	0.19	(10)	9.44	New Hampshire	4.53	0.26	(35)	-13.05
California	6.46 <sup>b</sup>	0.06	(04)	23.99 <sup>c</sup>	New Jersey	5.06	0.10	(17)	-2.97
Colorado	4.50 <sup>b</sup>	0.15	(37)	-13.60	New Mexico	6.15 <sup>b</sup>	0.25	(07)	18.01 <sup>c</sup>
Connecticut	4.74	0.16	(28)	-9.10	New York	5.27	0.07	(13)	1.16
Delaware	4.73	0.32	(30)	-9.34	North Carolina	4.57 <sup>b</sup>	0.10	(33)	-12.35
D.C.	6.24	0.47	(06)	19.62 <sup>c</sup>	North Dakota	4.05	0.34	(43)	-22.23 <sup>c</sup>
Florida	5.03	0.07	(18)	-3.49	Ohio	4.47 <sup>b</sup>	0.08	(39)	-14.17
Georgia	4.50 <sup>b</sup>	0.11	(38)	-13.66	Oklahoma	4.79	0.16	(27)	-8.06
Hawaii	7.31 <sup>b</sup>	0.32	(01)	40.24 <sup>c</sup>	Oregon	4.58 <sup>b</sup>	0.15	(32)	-12.20
Idaho	3.89 <sup>b</sup>	0.24	(46)	-25.45 <sup>c</sup>	Pennsylvania	4.89 <sup>b</sup>	0.08	(24)	-6.25
Illinois	5.09	0.09	(16)	-2.44	Rhode Island	5.42	0.30	(12)	4.03
Indiana	4.64 <sup>b</sup>	0.12	(31)	-11.00	South Carolina	4.84	0.15	(26)	-7.13
Iowa	3.86 <sup>b</sup>	0.15	(47)	-25.88 <sup>c</sup>	South Dakota	3.99 <sup>b</sup>	0.31	(45)	-23.46 <sup>c</sup>
Kansas	4.03 <sup>b</sup>	0.17	(44)	-22.67 <sup>c</sup>	Tennessee	4.88	0.12	(25)	-6.36
Kentucky	4.92	0.15	(22)	-5.59	Texas	6.88 <sup>b</sup>	0.08	(02)	32.01 <sup>c</sup>
Louisiana	6.81 <sup>b</sup>	0.18	(03)	30.58 <sup>c</sup>	Utah	3.46 <sup>b</sup>	0.19	(51)	-33.54 <sup>c</sup>
Maine	4.33 <sup>b</sup>	0.24	(41)	-16.87 <sup>c</sup>	Vermont	3.60 <sup>b</sup>	0.33	(50)	-30.99 <sup>c</sup>
Maryland	4.91	0.13	(23)	-5.79	Virginia	4.56 <sup>b</sup>	0.11	(34)	-12.47
Massachusetts	5.24	0.12	(14)	0.47	Washington	5.49	0.13	(11)	5.28
Michigan	4.96	0.10	(20)	-4.93	West Virginia	4.41 <sup>b</sup>	0.20	(40)	-15.48 <sup>c</sup>
Minnesota	4.53 <sup>b</sup>	0.13	(36)	-13.13	Wisconsin	4.73 <sup>b</sup>	0.13	(29)	-9.33
Mississippi	6.29 <sup>b</sup>	0.21	(05)	20.65 <sup>c</sup>	Wyoming	4.24	0.40	(42)	-18.57 <sup>c</sup>
Missouri	4.98	0.13	(19)	-4.44					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.



Table 14.19  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.65	0.03		
High Five States				
Louisiana	10.35 <sup>b</sup>	0.34	(01)	35.28 <sup>c</sup>
D.C.	10.23	0.92	(02)	33.62 <sup>c</sup>
Hawaii	10.18 <sup>b</sup>	0.56	(03)	33.05 <sup>c</sup>
Texas	10.18 <sup>b</sup>	0.16	(04)	33.01 <sup>c</sup>
Mississippi	9.56 <sup>b</sup>	0.40	(05)	24.95 <sup>c</sup>
Low Five States				
North Dakota	5.32 <sup>b</sup>	0.58	(47)	-30.44 <sup>c</sup>
South Dakota	5.30 <sup>b</sup>	0.52	(48)	-30.70 <sup>c</sup>
Montana	5.05 <sup>b</sup>	0.45	(49)	-33.99 <sup>c</sup>
Vermont	4.89 <sup>b</sup>	0.57	(50)	-36.08 <sup>c</sup>
Utah	4.64 <sup>b</sup>	0.33	(51)	-39.36 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.61	0.27	(17)	-0.55	Montana	5.05 <sup>b</sup>	0.45	(49)	-33.99 <sup>c</sup>
Alaska	8.64	1.02	(08)	12.87	Nebraska	5.75 <sup>b</sup>	0.37	(42)	-24.91 <sup>c</sup>
Arizona	7.00	0.22	(28)	-8.51	Nevada	8.22	0.39	(11)	7.40
Arkansas	7.96	0.34	(12)	3.95	New Hampshire	6.62	0.48	(35)	-13.51
California	9.29 <sup>b</sup>	0.11	(06)	21.44 <sup>c</sup>	New Jersey	7.54	0.20	(18)	-1.43
Colorado	6.30 <sup>b</sup>	0.27	(39)	-17.72 <sup>c</sup>	New Mexico	8.74	0.44	(07)	14.17
Connecticut	7.28	0.30	(23)	-4.86	New York	7.94	0.13	(13)	3.78
Delaware	7.86	0.63	(14)	2.69	North Carolina	6.86 <sup>b</sup>	0.19	(31)	-10.42
D.C.	10.23	0.92	(02)	33.62 <sup>c</sup>	North Dakota	5.32 <sup>b</sup>	0.58	(47)	-30.44 <sup>c</sup>
Florida	7.65	0.12	(16)	-0.01	Ohio	6.58 <sup>b</sup>	0.16	(36)	-13.96
Georgia	6.68 <sup>b</sup>	0.21	(34)	-12.69	Oklahoma	7.12	0.29	(26)	-6.94
Hawaii	10.18 <sup>b</sup>	0.56	(03)	33.05 <sup>c</sup>	Oregon	6.34 <sup>b</sup>	0.27	(38)	-17.16 <sup>c</sup>
Idaho	5.60 <sup>b</sup>	0.42	(45)	-26.84 <sup>c</sup>	Pennsylvania	7.29	0.15	(21)	-4.81
Illinois	7.39	0.17	(20)	-3.48	Rhode Island	8.50	0.58	(09)	11.12
Indiana	6.81 <sup>b</sup>	0.22	(33)	-11.03	South Carolina	7.28	0.28	(22)	-4.85
Iowa	5.52 <sup>b</sup>	0.27	(46)	-27.92 <sup>c</sup>	South Dakota	5.30 <sup>b</sup>	0.52	(48)	-30.70 <sup>c</sup>
Kansas	5.73 <sup>b</sup>	0.30	(43)	-25.13 <sup>c</sup>	Tennessee	7.20	0.23	(25)	-5.89
Kentucky	6.90	0.28	(30)	-9.85	Texas	10.18 <sup>b</sup>	0.16	(04)	33.01 <sup>c</sup>
Louisiana	10.35 <sup>b</sup>	0.34	(01)	35.28 <sup>c</sup>	Utah	4.64 <sup>b</sup>	0.33	(51)	-39.36 <sup>c</sup>
Maine	6.19	0.43	(40)	-19.15 <sup>c</sup>	Vermont	4.89 <sup>b</sup>	0.57	(50)	-36.08 <sup>c</sup>
Maryland	7.41	0.25	(19)	-3.13	Virginia	6.90 <sup>b</sup>	0.21	(29)	-9.84
Massachusetts	8.36	0.24	(10)	9.20	Washington	7.74	0.24	(15)	1.07
Michigan	7.03 <sup>b</sup>	0.18	(27)	-8.14	West Virginia	6.08 <sup>b</sup>	0.36	(41)	-20.60 <sup>c</sup>
Minnesota	6.57 <sup>b</sup>	0.24	(37)	-14.21	Wisconsin	6.83 <sup>b</sup>	0.23	(32)	-10.72
Mississippi	9.56 <sup>b</sup>	0.40	(05)	24.95 <sup>c</sup>	Wyoming	5.60	0.70	(44)	-26.82 <sup>c</sup>
Missouri	7.26	0.23	(24)	-5.18					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.



Table 14.20  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.20	0.02		
High Five States				
Hawaii	4.70 <sup>b</sup>	0.35	(01)	46.58 <sup>c</sup>
Texas	4.18 <sup>b</sup>	0.09	(02)	30.34 <sup>c</sup>
California	4.06 <sup>b</sup>	0.07	(03)	26.81 <sup>c</sup>
Louisiana	3.99 <sup>b</sup>	0.18	(04)	24.51 <sup>c</sup>
New Mexico	3.89	0.27	(05)	21.43 <sup>c</sup>
Low Five States				
Vermont	2.50	0.37	(47)	-22.00 <sup>c</sup>
Utah	2.43 <sup>b</sup>	0.22	(48)	-24.13 <sup>c</sup>
Idaho	2.38	0.26	(49)	-25.57 <sup>c</sup>
Nebraska	2.25 <sup>b</sup>	0.21	(50)	-29.74 <sup>c</sup>
Delaware	2.08 <sup>b</sup>	0.29	(51)	-35.06 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.37	0.15	(11)	5.04	Montana	2.64	0.31	(44)	-17.53 <sup>c</sup>
Alaska	3.80	0.62	(08)	18.66 <sup>c</sup>	Nebraska	2.25 <sup>b</sup>	0.21	(50)	-29.74 <sup>c</sup>
Arizona	3.12	0.14	(18)	-2.59	Nevada	3.66	0.25	(09)	14.14
Arkansas	3.83	0.21	(07)	19.55 <sup>c</sup>	New Hampshire	2.77	0.27	(39)	-13.47
California	4.06 <sup>b</sup>	0.07	(03)	26.81 <sup>c</sup>	New Jersey	3.10	0.11	(19)	-3.13
Colorado	2.98	0.16	(27)	-6.93	New Mexico	3.89	0.27	(05)	21.43 <sup>c</sup>
Connecticut	2.74	0.16	(42)	-14.40	New York	3.16	0.07	(15)	-1.32
Delaware	2.08 <sup>b</sup>	0.29	(51)	-35.06 <sup>c</sup>	North Carolina	2.75 <sup>b</sup>	0.11	(41)	-14.23
D.C.	3.16	0.43	(16)	-1.38	North Dakota	2.99	0.39	(26)	-6.83
Florida	2.77 <sup>b</sup>	0.07	(40)	-13.71	Ohio	2.81 <sup>b</sup>	0.09	(36)	-12.35
Georgia	2.80 <sup>b</sup>	0.11	(37)	-12.77	Oklahoma	2.87	0.17	(31)	-10.44
Hawaii	4.70 <sup>b</sup>	0.35	(01)	46.58 <sup>c</sup>	Oregon	3.04	0.17	(22)	-5.10
Idaho	2.38	0.26	(49)	-25.57 <sup>c</sup>	Pennsylvania	3.00	0.08	(24)	-6.53
Illinois	3.26	0.10	(13)	1.79	Rhode Island	3.05	0.29	(21)	-4.81
Indiana	2.93	0.13	(29)	-8.67	South Carolina	2.82	0.15	(34)	-12.14
Iowa	2.53 <sup>b</sup>	0.16	(46)	-21.12 <sup>c</sup>	South Dakota	2.82	0.35	(33)	-11.88
Kansas	2.57 <sup>b</sup>	0.18	(45)	-19.92 <sup>c</sup>	Tennessee	2.99	0.13	(25)	-6.69
Kentucky	3.37	0.17	(12)	5.04	Texas	4.18 <sup>b</sup>	0.09	(02)	30.34 <sup>c</sup>
Louisiana	3.99 <sup>b</sup>	0.18	(04)	24.51 <sup>c</sup>	Utah	2.43 <sup>b</sup>	0.22	(48)	-24.13 <sup>c</sup>
Maine	2.81	0.25	(35)	-12.30	Vermont	2.50	0.37	(47)	-22.00 <sup>c</sup>
Maryland	2.94	0.14	(28)	-8.33	Virginia	2.72 <sup>b</sup>	0.11	(43)	-15.07 <sup>c</sup>
Massachusetts	2.77 <sup>b</sup>	0.12	(38)	-13.46	Washington	3.51	0.14	(10)	9.42
Michigan	3.24	0.10	(14)	0.99	West Virginia	3.02	0.22	(23)	-5.70
Minnesota	2.85	0.14	(32)	-10.98	Wisconsin	2.91	0.13	(30)	-9.19
Mississippi	3.84	0.22	(06)	19.80 <sup>c</sup>	Wyoming	3.05	0.46	(20)	-4.79
Missouri	3.13	0.13	(17)	-2.28					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 15.24  
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	52.51	0.06		
High Five States				
Kentucky	76.54 <sup>b</sup>	0.59	(01)	45.76 <sup>c</sup>
Tennessee	67.66 <sup>b</sup>	0.47	(02)	28.84 <sup>c</sup>
West Virginia	67.56 <sup>b</sup>	0.78	(03)	28.66 <sup>c</sup>
Arkansas	67.06 <sup>b</sup>	0.66	(04)	27.70 <sup>c</sup>
Mississippi	66.49 <sup>b</sup>	0.68	(05)	26.62 <sup>c</sup>
Low Five States				
California	41.67 <sup>b</sup>	0.16	(47)	-20.64 <sup>c</sup>
Colorado	39.05 <sup>b</sup>	0.45	(48)	-25.64 <sup>c</sup>
Hawaii	38.15 <sup>b</sup>	0.74	(49)	-27.36 <sup>c</sup>
New Mexico	37.09 <sup>b</sup>	0.62	(50)	-29.37 <sup>c</sup>
Utah	23.85 <sup>b</sup>	0.51	(51)	-54.58 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	62.75 <sup>b</sup>	0.51	(08)	19.49 <sup>c</sup>	Montana	50.84	0.99	(32)	-3.19
Alaska	53.81	1.72	(22)	2.46	Nebraska	48.32 <sup>b</sup>	0.72	(39)	-7.98 <sup>c</sup>
Arizona	43.55 <sup>b</sup>	0.38	(45)	-17.07 <sup>c</sup>	Nevada	57.44 <sup>b</sup>	0.73	(16)	9.38 <sup>c</sup>
Arkansas	67.06 <sup>b</sup>	0.66	(04)	27.70 <sup>c</sup>	New Hampshire	53.45	0.90	(25)	1.79
California	41.67 <sup>b</sup>	0.16	(47)	-20.64 <sup>c</sup>	New Jersey	48.53 <sup>b</sup>	0.33	(37)	-7.59 <sup>c</sup>
Colorado	39.05 <sup>b</sup>	0.45	(48)	-25.64 <sup>c</sup>	New Mexico	37.09 <sup>b</sup>	0.62	(50)	-29.37 <sup>c</sup>
Connecticut	48.03 <sup>b</sup>	0.50	(40)	-8.53 <sup>c</sup>	New York	45.25 <sup>b</sup>	0.21	(43)	-13.83 <sup>c</sup>
Delaware	61.41 <sup>b</sup>	1.16	(11)	16.95 <sup>c</sup>	North Carolina	58.85 <sup>b</sup>	0.37	(15)	12.07 <sup>c</sup>
D.C.	48.60	1.31	(36)	-7.45 <sup>c</sup>	North Dakota	45.06 <sup>b</sup>	1.13	(44)	-14.20 <sup>c</sup>
Florida	52.08	0.21	(30)	-0.83	Ohio	59.82 <sup>b</sup>	0.31	(13)	13.92 <sup>c</sup>
Georgia	56.90 <sup>b</sup>	0.39	(17)	8.35 <sup>c</sup>	Oklahoma	63.24 <sup>b</sup>	0.58	(07)	20.42 <sup>c</sup>
Hawaii	38.15 <sup>b</sup>	0.74	(49)	-27.36 <sup>c</sup>	Oregon	53.52	0.53	(24)	1.92
Idaho	43.31 <sup>b</sup>	0.81	(46)	-17.52 <sup>c</sup>	Pennsylvania	53.25	0.27	(26)	1.41
Illinois	54.09 <sup>b</sup>	0.30	(21)	3.01	Rhode Island	53.03	0.95	(27)	0.98
Indiana	62.62 <sup>b</sup>	0.45	(09)	19.24 <sup>c</sup>	South Carolina	58.96 <sup>b</sup>	0.52	(14)	12.28 <sup>c</sup>
Iowa	52.49	0.56	(28)	-0.04	South Dakota	48.89	1.06	(35)	-6.89 <sup>c</sup>
Kansas	54.29	0.62	(20)	3.38	Tennessee	67.66 <sup>b</sup>	0.47	(02)	28.84 <sup>c</sup>
Kentucky	76.54 <sup>b</sup>	0.59	(01)	45.76 <sup>c</sup>	Texas	50.59 <sup>b</sup>	0.23	(33)	-3.67
Louisiana	64.05 <sup>b</sup>	0.55	(06)	21.98 <sup>c</sup>	Utah	23.85 <sup>b</sup>	0.51	(51)	-54.58 <sup>c</sup>
Maine	60.45 <sup>b</sup>	0.88	(12)	15.11 <sup>c</sup>	Vermont	50.55	1.22	(34)	-3.73
Maryland	53.76	0.45	(23)	2.37	Virginia	55.65 <sup>b</sup>	0.39	(19)	5.97 <sup>c</sup>
Massachusetts	52.47	0.39	(29)	-0.08	Washington	51.62	0.42	(31)	-1.69
Michigan	55.98 <sup>b</sup>	0.33	(18)	6.60 <sup>c</sup>	West Virginia	67.56 <sup>b</sup>	0.78	(03)	28.66 <sup>c</sup>
Minnesota	45.78 <sup>b</sup>	0.43	(42)	-12.81 <sup>c</sup>	Wisconsin	48.50 <sup>b</sup>	0.41	(38)	-7.63 <sup>c</sup>
Mississippi	66.49 <sup>b</sup>	0.68	(05)	26.62 <sup>c</sup>	Wyoming	45.90 <sup>b</sup>	1.34	(41)	-12.60 <sup>c</sup>
Missouri	62.51 <sup>b</sup>	0.45	(10)	19.04 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 15.25  
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	68.81	0.10		
High Five States				
Kentucky	105.22 <sup>b</sup>	1.08	(01)	52.91 <sup>c</sup>
Mississippi	99.40 <sup>b</sup>	1.30	(02)	44.44 <sup>c</sup>
Tennessee	95.72 <sup>b</sup>	0.87	(03)	39.11 <sup>c</sup>
Arkansas	93.17 <sup>b</sup>	1.19	(04)	35.39 <sup>c</sup>
Alabama	92.19 <sup>b</sup>	0.96	(05)	33.98 <sup>c</sup>
Low Five States				
California	51.38 <sup>b</sup>	0.28	(47)	-25.34 <sup>c</sup>
Hawaii	51.19 <sup>b</sup>	1.28	(48)	-25.61 <sup>c</sup>
Colorado	47.82 <sup>b</sup>	0.77	(49)	-30.50 <sup>c</sup>
New Mexico	46.15 <sup>b</sup>	1.05	(50)	-32.94 <sup>c</sup>
Utah	31.41 <sup>b</sup>	0.88	(51)	-54.35 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	92.19 <sup>b</sup>	0.96	(05)	33.98 <sup>c</sup>	Montana	60.70 <sup>b</sup>	1.61	(39)	-11.79 <sup>c</sup>
Alaska	64.83	2.83	(33)	-5.79 <sup>c</sup>	Nebraska	65.63	1.28	(28)	-4.63
Arizona	54.11 <sup>b</sup>	0.63	(45)	-21.36 <sup>c</sup>	Nevada	64.96	1.14	(32)	-5.60 <sup>c</sup>
Arkansas	93.17 <sup>b</sup>	1.19	(04)	35.39 <sup>c</sup>	New Hampshire	65.49	1.52	(30)	-4.83
California	51.38 <sup>b</sup>	0.28	(47)	-25.34 <sup>c</sup>	New Jersey	61.48 <sup>b</sup>	0.57	(38)	-10.66 <sup>c</sup>
Colorado	47.82 <sup>b</sup>	0.77	(49)	-30.50 <sup>c</sup>	New Mexico	46.15 <sup>b</sup>	1.05	(50)	-32.94 <sup>c</sup>
Connecticut	59.65 <sup>b</sup>	0.87	(40)	-13.32 <sup>c</sup>	New York	57.67 <sup>b</sup>	0.37	(43)	-16.20 <sup>c</sup>
Delaware	76.39 <sup>b</sup>	1.99	(16)	11.02 <sup>c</sup>	North Carolina	82.49 <sup>b</sup>	0.68	(12)	19.87 <sup>c</sup>
D.C.	69.13	2.48	(24)	0.46	North Dakota	58.57 <sup>b</sup>	1.92	(41)	-14.89 <sup>c</sup>
Florida	66.22 <sup>b</sup>	0.36	(27)	-3.76	Ohio	80.32 <sup>b</sup>	0.56	(14)	16.73 <sup>c</sup>
Georgia	81.58 <sup>b</sup>	0.74	(13)	18.55 <sup>c</sup>	Oklahoma	84.98 <sup>b</sup>	1.03	(08)	23.50 <sup>c</sup>
Hawaii	51.19 <sup>b</sup>	1.28	(48)	-25.61 <sup>c</sup>	Oregon	64.16 <sup>b</sup>	0.87	(34)	-6.76 <sup>c</sup>
Idaho	53.38 <sup>b</sup>	1.34	(46)	-22.43 <sup>c</sup>	Pennsylvania	71.45 <sup>b</sup>	0.48	(20)	3.83
Illinois	71.07 <sup>b</sup>	0.53	(21)	3.28	Rhode Island	69.75	1.69	(23)	1.37
Indiana	83.85 <sup>b</sup>	0.80	(11)	21.86 <sup>c</sup>	South Carolina	83.92 <sup>b</sup>	0.95	(10)	21.95 <sup>c</sup>
Iowa	70.82	0.98	(22)	2.91	South Dakota	65.55	1.84	(29)	-4.75
Kansas	72.29	1.09	(19)	5.06 <sup>c</sup>	Tennessee	95.72 <sup>b</sup>	0.87	(03)	39.11 <sup>c</sup>
Kentucky	105.22 <sup>b</sup>	1.08	(01)	52.91 <sup>c</sup>	Texas	68.26	0.42	(26)	-0.80
Louisiana	89.71 <sup>b</sup>	1.00	(07)	30.37 <sup>c</sup>	Utah	31.41 <sup>b</sup>	0.88	(51)	-54.35 <sup>c</sup>
Maine	76.88 <sup>b</sup>	1.53	(15)	11.73 <sup>c</sup>	Vermont	61.84	2.07	(36)	-10.14 <sup>c</sup>
Maryland	69.09	0.80	(25)	0.41	Virginia	74.53 <sup>b</sup>	0.71	(17)	8.31 <sup>c</sup>
Massachusetts	65.46 <sup>b</sup>	0.68	(31)	-4.88	Washington	61.51 <sup>b</sup>	0.69	(37)	-10.61 <sup>c</sup>
Michigan	72.50 <sup>b</sup>	0.58	(18)	5.36 <sup>c</sup>	West Virginia	91.03 <sup>b</sup>	1.39	(06)	32.28 <sup>c</sup>
Minnesota	58.28 <sup>b</sup>	0.73	(42)	-15.31 <sup>c</sup>	Wisconsin	62.35 <sup>b</sup>	0.71	(35)	-9.39 <sup>c</sup>
Mississippi	99.40 <sup>b</sup>	1.30	(02)	44.44 <sup>c</sup>	Wyoming	55.56 <sup>b</sup>	2.22	(44)	-19.25 <sup>c</sup>
Missouri	84.35 <sup>b</sup>	0.80	(09)	22.57 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.




Table 15.26  
Cancer of the Lung And Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	40.56	0.07		
High Five States				
Kentucky	56.03 <sup>b</sup>	0.68	(01)	38.16 <sup>c</sup>
Nevada	50.87 <sup>b</sup>	0.93	(02)	25.43 <sup>c</sup>
West Virginia	50.47 <sup>b</sup>	0.91	(03)	24.45 <sup>c</sup>
Delaware	50.37 <sup>b</sup>	1.41	(04)	24.19 <sup>c</sup>
Maine	48.71 <sup>b</sup>	1.06	(05)	20.09 <sup>c</sup>
Low Five States				
California	34.51 <sup>b</sup>	0.20	(47)	-14.90 <sup>c</sup>
Colorado	32.85 <sup>b</sup>	0.55	(48)	-19.00 <sup>c</sup>
New Mexico	29.92 <sup>b</sup>	0.75	(49)	-26.24 <sup>c</sup>
Hawaii	27.70 <sup>b</sup>	0.85	(50)	-31.70 <sup>c</sup>
Utah	17.61 <sup>b</sup>	0.60	(51)	-56.57 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	41.69	0.55	(27)	2.79	Montana	43.28	1.24	(19)	6.71 <sup>c</sup>
Alaska	44.41	2.13	(15)	9.51 <sup>c</sup>	Nebraska	35.61 <sup>b</sup>	0.84	(42)	-12.19 <sup>c</sup>
Arizona	34.92 <sup>b</sup>	0.46	(45)	-13.89 <sup>c</sup>	Nevada	50.87 <sup>b</sup>	0.93	(02)	25.43 <sup>c</sup>
Arkansas	47.46 <sup>b</sup>	0.75	(08)	17.02 <sup>c</sup>	New Hampshire	44.69 <sup>b</sup>	1.10	(14)	10.18 <sup>c</sup>
California	34.51 <sup>b</sup>	0.20	(47)	-14.90 <sup>c</sup>	New Jersey	39.70	0.39	(34)	-2.13
Colorado	32.85 <sup>b</sup>	0.55	(48)	-19.00 <sup>c</sup>	New Mexico	29.92 <sup>b</sup>	0.75	(49)	-26.24 <sup>c</sup>
Connecticut	40.10	0.61	(32)	-1.13	New York	36.68 <sup>b</sup>	0.25	(40)	-9.57 <sup>c</sup>
Delaware	50.37 <sup>b</sup>	1.41	(04)	24.19 <sup>c</sup>	North Carolina	42.03 <sup>b</sup>	0.41	(26)	3.64
D.C.	34.91 <sup>b</sup>	1.45	(46)	-13.93 <sup>c</sup>	North Dakota	35.11 <sup>b</sup>	1.36	(44)	-13.44 <sup>c</sup>
Florida	40.60	0.26	(30)	0.11	Ohio	45.34 <sup>b</sup>	0.36	(13)	11.78 <sup>c</sup>
Georgia	39.74	0.43	(33)	-2.01	Oklahoma	47.09 <sup>b</sup>	0.67	(09)	16.10 <sup>c</sup>
Hawaii	27.70 <sup>b</sup>	0.85	(50)	-31.70 <sup>c</sup>	Oregon	45.49 <sup>b</sup>	0.66	(12)	12.16 <sup>c</sup>
Idaho	35.44 <sup>b</sup>	0.99	(43)	-12.61 <sup>c</sup>	Pennsylvania	40.43	0.31	(31)	-0.30
Illinois	42.21 <sup>b</sup>	0.35	(23)	4.08	Rhode Island	42.04	1.12	(25)	3.65
Indiana	47.61 <sup>b</sup>	0.52	(06)	17.39 <sup>c</sup>	South Carolina	40.64	0.57	(29)	0.21
Iowa	39.15	0.65	(35)	-3.47	South Dakota	36.48	1.25	(41)	-10.04 <sup>c</sup>
Kansas	41.25	0.73	(28)	1.70	Tennessee	47.54 <sup>b</sup>	0.52	(07)	17.22 <sup>c</sup>
Kentucky	56.03 <sup>b</sup>	0.68	(01)	38.16 <sup>c</sup>	Texas	37.53 <sup>b</sup>	0.27	(38)	-7.47 <sup>c</sup>
Louisiana	45.58 <sup>b</sup>	0.61	(11)	12.37 <sup>c</sup>	Utah	17.61 <sup>b</sup>	0.60	(51)	-56.57 <sup>c</sup>
Maine	48.71 <sup>b</sup>	1.06	(05)	20.09 <sup>c</sup>	Vermont	42.45	1.51	(22)	4.68
Maryland	42.94 <sup>b</sup>	0.53	(21)	5.87 <sup>c</sup>	Virginia	42.11	0.45	(24)	3.83
Massachusetts	43.77 <sup>b</sup>	0.47	(18)	7.92 <sup>c</sup>	Washington	44.30 <sup>b</sup>	0.52	(16)	9.23 <sup>c</sup>
Michigan	44.07 <sup>b</sup>	0.39	(17)	8.67 <sup>c</sup>	West Virginia	50.47 <sup>b</sup>	0.91	(03)	24.45 <sup>c</sup>
Minnesota	36.96 <sup>b</sup>	0.52	(39)	-8.88 <sup>c</sup>	Wisconsin	38.55 <sup>b</sup>	0.49	(36)	-4.96
Mississippi	43.04	0.72	(20)	6.12 <sup>c</sup>	Wyoming	38.43	1.68	(37)	-5.25 <sup>c</sup>
Missouri	46.61 <sup>b</sup>	0.52	(10)	14.93 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 16.18  
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.69	0.01		
High Five States				
Oklahoma	3.47 <sup>b</sup>	0.14	(01)	28.75 <sup>c</sup>
Utah	3.38 <sup>b</sup>	0.19	(02)	25.39 <sup>c</sup>
Wyoming	3.34	0.36	(03)	24.05 <sup>c</sup>
Kentucky	3.27 <sup>b</sup>	0.12	(04)	21.52 <sup>c</sup>
Idaho	3.24	0.22	(05)	20.48 <sup>c</sup>
Low Five States				
Louisiana	2.09 <sup>b</sup>	0.10	(47)	-22.38 <sup>c</sup>
Alaska	2.01	0.30	(48)	-25.34 <sup>c</sup>
North Dakota	1.74 <sup>b</sup>	0.22	(49)	-35.26 <sup>c</sup>
Hawaii	1.73 <sup>b</sup>	0.16	(50)	-35.70 <sup>c</sup>
D.C.	1.51 <sup>b</sup>	0.22	(51)	-43.97 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.85	0.11	(25)	5.99	Montana	2.84	0.24	(26)	5.66
Alaska	2.01	0.30	(48)	-25.34 <sup>c</sup>	Nebraska	2.70	0.17	(32)	0.12
Arizona	2.81	0.10	(28)	4.27	Nevada	2.92	0.16	(20)	8.63
Arkansas	2.59	0.13	(35)	-3.77	New Hampshire	3.10	0.21	(08)	15.19 <sup>c</sup>
California	2.56	0.04	(37)	-4.83	New Jersey	2.68	0.08	(33)	-0.38
Colorado	3.01	0.12	(13)	11.71	New Mexico	2.73	0.17	(30)	1.41
Connecticut	2.86	0.12	(24)	6.04	New York	2.18 <sup>b</sup>	0.05	(46)	-19.17 <sup>c</sup>
Delaware	3.08	0.26	(09)	14.45	North Carolina	2.93	0.08	(19)	8.76
D.C.	1.51 <sup>b</sup>	0.22	(51)	-43.97 <sup>c</sup>	North Dakota	1.74 <sup>b</sup>	0.22	(49)	-35.26 <sup>c</sup>
Florida	2.94 <sup>b</sup>	0.05	(16)	9.19	Ohio	2.63	0.07	(34)	-2.44
Georgia	2.44	0.08	(41)	-9.54	Oklahoma	3.47 <sup>b</sup>	0.14	(01)	28.75 <sup>c</sup>
Hawaii	1.73 <sup>b</sup>	0.16	(50)	-35.70 <sup>c</sup>	Oregon	3.06	0.13	(11)	13.48
Idaho	3.24	0.22	(05)	20.48 <sup>c</sup>	Pennsylvania	2.88	0.06	(22)	7.05
Illinois	2.55	0.06	(39)	-5.19	Rhode Island	2.97	0.22	(15)	10.41
Indiana	2.88	0.10	(23)	6.89	South Carolina	2.57	0.11	(36)	-4.52
Iowa	2.42	0.12	(42)	-9.98	South Dakota	2.80	0.26	(29)	3.90
Kansas	3.06	0.15	(10)	13.79	Tennessee	2.90	0.10	(21)	7.67
Kentucky	3.27 <sup>b</sup>	0.12	(04)	21.52 <sup>c</sup>	Texas	2.54	0.05	(40)	-5.57
Louisiana	2.09 <sup>b</sup>	0.10	(47)	-22.38 <sup>c</sup>	Utah	3.38 <sup>b</sup>	0.19	(02)	25.39 <sup>c</sup>
Maine	2.83	0.19	(27)	5.29	Vermont	2.93	0.29	(17)	8.80
Maryland	2.71	0.10	(31)	0.64	Virginia	2.93	0.09	(18)	8.77
Massachusetts	3.02 <sup>b</sup>	0.09	(12)	12.28	Washington	2.99	0.10	(14)	11.09
Michigan	2.39 <sup>b</sup>	0.07	(43)	-11.24	West Virginia	3.21	0.18	(06)	19.19 <sup>c</sup>
Minnesota	2.27 <sup>b</sup>	0.09	(45)	-15.74 <sup>c</sup>	Wisconsin	2.56	0.09	(38)	-4.99
Mississippi	2.30	0.13	(44)	-14.45	Wyoming	3.34	0.36	(03)	24.05 <sup>c</sup>
Missouri	3.12 <sup>b</sup>	0.10	(07)	15.74 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 16.19  
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	3.98	0.02		
High Five States				
Delaware	5.16	0.52	(01)	29.57 <sup>c</sup>
Utah	5.12	0.35	(02)	28.71 <sup>c</sup>
Oklahoma	5.03 <sup>b</sup>	0.25	(03)	26.43 <sup>c</sup>
Kentucky	4.87 <sup>b</sup>	0.23	(04)	22.43 <sup>c</sup>
Wyoming	4.76	0.65	(05)	19.52 <sup>c</sup>
Low Five States				
Minnesota	3.09 <sup>b</sup>	0.17	(47)	-22.36 <sup>c</sup>
North Dakota	2.50 <sup>b</sup>	0.39	(48)	-37.14 <sup>c</sup>
Alaska	2.49	0.47	(49)	-37.40 <sup>c</sup>
Hawaii	2.46 <sup>b</sup>	0.28	(50)	-38.17 <sup>c</sup>
D.C.	2.41	0.46	(51)	-39.50 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.27	0.21	(23)	7.26	Montana	3.37	0.38	(43)	-15.34 <sup>c</sup>
Alaska	2.49	0.47	(49)	-37.40 <sup>c</sup>	Nebraska	3.92	0.31	(31)	-1.55
Arizona	4.30	0.18	(21)	8.12	Nevada	4.46	0.29	(09)	12.00
Arkansas	3.86	0.24	(34)	-3.05	New Hampshire	4.64	0.40	(07)	16.63 <sup>c</sup>
California	3.91	0.07	(32)	-1.81	New Jersey	3.99	0.14	(30)	0.12
Colorado	4.40	0.22	(14)	10.45	New Mexico	3.74	0.30	(36)	-6.06
Connecticut	4.32	0.23	(20)	8.45	New York	3.19 <sup>b</sup>	0.09	(46)	-19.96 <sup>c</sup>
Delaware	5.16	0.52	(01)	29.57 <sup>c</sup>	North Carolina	4.20	0.15	(26)	5.45
D.C.	2.41	0.46	(51)	-39.50 <sup>c</sup>	North Dakota	2.50 <sup>b</sup>	0.39	(48)	-37.14 <sup>c</sup>
Florida	4.38 <sup>b</sup>	0.09	(15)	9.98	Ohio	3.88	0.12	(33)	-2.53
Georgia	3.68	0.16	(37)	-7.52	Oklahoma	5.03 <sup>b</sup>	0.25	(03)	26.43 <sup>c</sup>
Hawaii	2.46 <sup>b</sup>	0.28	(50)	-38.17 <sup>c</sup>	Oregon	4.40	0.22	(12)	10.53
Idaho	4.75	0.39	(06)	19.34 <sup>c</sup>	Pennsylvania	4.26	0.12	(24)	7.14
Illinois	3.66	0.12	(38)	-8.12	Rhode Island	4.34	0.42	(18)	9.11
Indiana	4.32	0.18	(19)	8.53	South Carolina	3.99	0.21	(29)	0.21
Iowa	3.31	0.21	(44)	-16.72 <sup>c</sup>	South Dakota	3.58	0.43	(42)	-10.16
Kansas	4.43	0.27	(11)	11.31	Tennessee	4.36	0.18	(16)	9.53
Kentucky	4.87 <sup>b</sup>	0.23	(04)	22.43 <sup>c</sup>	Texas	3.81	0.10	(35)	-4.30
Louisiana	3.26 <sup>b</sup>	0.19	(45)	-18.20 <sup>c</sup>	Utah	5.12	0.35	(02)	28.71 <sup>c</sup>
Maine	4.06	0.35	(27)	1.90	Vermont	4.34	0.55	(17)	9.12
Maryland	4.05	0.19	(28)	1.76	Virginia	4.23	0.17	(25)	6.15
Massachusetts	4.40	0.17	(13)	10.48	Washington	4.27	0.18	(22)	7.27
Michigan	3.63	0.13	(40)	-8.69	West Virginia	4.63	0.32	(08)	16.23 <sup>c</sup>
Minnesota	3.09 <sup>b</sup>	0.17	(47)	-22.36 <sup>c</sup>	Wisconsin	3.59	0.17	(41)	-9.73
Mississippi	3.64	0.25	(39)	-8.55	Wyoming	4.76	0.65	(05)	19.52 <sup>c</sup>
Missouri	4.45	0.18	(10)	11.92					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 16.20  
Melanoma of the Skin (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.73	0.01		
High Five States				
Montana	2.41	0.30	(01)	39.43 <sup>c</sup>
Oklahoma	2.23	0.15	(02)	29.15 <sup>c</sup>
Wyoming	2.18	0.40	(03)	26.03 <sup>c</sup>
South Dakota	2.17	0.31	(04)	25.48 <sup>c</sup>
West Virginia	2.13	0.20	(05)	22.94 <sup>c</sup>
Low Five States				
Mississippi	1.35	0.13	(47)	-21.84 <sup>c</sup>
Louisiana	1.22 <sup>b</sup>	0.10	(48)	-29.50 <sup>c</sup>
Hawaii	1.09 <sup>b</sup>	0.17	(49)	-36.67 <sup>c</sup>
North Dakota	1.09	0.24	(50)	-36.81 <sup>c</sup>
D.C.	0.99	0.24	(51)	-42.74 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.82	0.12	(25)	5.26	Montana	2.41	0.30	(01)	39.43 <sup>c</sup>
Alaska	1.46	0.35	(46)	-15.39 <sup>c</sup>	Nebraska	1.80	0.19	(27)	3.88
Arizona	1.55	0.10	(41)	-10.47	Nevada	1.55	0.16	(40)	-10.19
Arkansas	1.56	0.14	(39)	-9.61	New Hampshire	1.93	0.23	(16)	11.48
California	1.55 <sup>b</sup>	0.04	(42)	-10.49	New Jersey	1.74	0.08	(31)	0.91
Colorado	1.91	0.13	(18)	10.72	New Mexico	1.96	0.19	(15)	13.09
Connecticut	1.82	0.13	(24)	5.54	New York	1.49 <sup>b</sup>	0.05	(44)	-14.03
Delaware	1.56	0.25	(38)	-9.60	North Carolina	2.03	0.09	(10)	17.56 <sup>c</sup>
D.C.	0.99	0.24	(51)	-42.74 <sup>c</sup>	North Dakota	1.09	0.24	(50)	-36.81 <sup>c</sup>
Florida	1.78	0.06	(28)	2.72	Ohio	1.72	0.07	(32)	-0.30
Georgia	1.57	0.08	(37)	-9.08	Oklahoma	2.23	0.15	(02)	29.15 <sup>c</sup>
Hawaii	1.09 <sup>b</sup>	0.17	(49)	-36.67 <sup>c</sup>	Oregon	1.97	0.14	(14)	13.87
Idaho	1.91	0.23	(20)	10.42	Pennsylvania	1.88	0.07	(21)	8.67
Illinois	1.75	0.07	(30)	1.04	Rhode Island	2.00	0.25	(12)	15.78 <sup>c</sup>
Indiana	1.81	0.10	(26)	4.67	South Carolina	1.52	0.11	(43)	-12.28
Iowa	1.69	0.14	(35)	-2.23	South Dakota	2.17	0.31	(04)	25.48 <sup>c</sup>
Kansas	2.04	0.16	(09)	18.15 <sup>c</sup>	Tennessee	1.83	0.10	(23)	6.04
Kentucky	2.07	0.13	(08)	19.56 <sup>c</sup>	Texas	1.60	0.05	(36)	-7.33
Louisiana	1.22 <sup>b</sup>	0.10	(48)	-29.50 <sup>c</sup>	Utah	1.92	0.19	(17)	10.87
Maine	1.91	0.22	(19)	10.69	Vermont	1.87	0.32	(22)	8.34
Maryland	1.76	0.11	(29)	1.85	Virginia	2.02	0.10	(11)	16.75 <sup>c</sup>
Massachusetts	2.07	0.10	(07)	19.80 <sup>c</sup>	Washington	1.98	0.11	(13)	14.49
Michigan	1.48 <sup>b</sup>	0.07	(45)	-14.56	West Virginia	2.13	0.20	(05)	22.94 <sup>c</sup>
Minnesota	1.70	0.11	(34)	-1.83	Wisconsin	1.72	0.10	(33)	-0.71
Mississippi	1.35	0.13	(47)	-21.84 <sup>c</sup>	Wyoming	2.18	0.40	(03)	26.03 <sup>c</sup>
Missouri	2.10	0.11	(06)	21.26 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 18.18  
Myeloma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.55	0.02		
High Five States				
D.C.	5.08 <sup>b</sup>	0.42	(01)	42.97 <sup>c</sup>
Alabama	4.32 <sup>b</sup>	0.13	(02)	21.52 <sup>c</sup>
South Carolina	4.31 <sup>b</sup>	0.14	(03)	21.24 <sup>c</sup>
Tennessee	4.15 <sup>b</sup>	0.12	(04)	16.87 <sup>c</sup>
North Carolina	4.13 <sup>b</sup>	0.10	(05)	16.38 <sup>c</sup>
Low Five States				
Florida	3.09 <sup>b</sup>	0.05	(47)	-12.89
Arizona	3.09 <sup>b</sup>	0.10	(48)	-12.99
Alaska	2.90	0.40	(49)	-18.41 <sup>c</sup>
Nevada	2.87 <sup>b</sup>	0.16	(50)	-19.34 <sup>c</sup>
Hawaii	2.39 <sup>b</sup>	0.18	(51)	-32.68 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.32 <sup>b</sup>	0.13	(02)	21.52 <sup>c</sup>	Montana	3.68	0.26	(22)	3.71
Alaska	2.90	0.40	(49)	-18.41 <sup>c</sup>	Nebraska	3.35	0.19	(40)	-5.72
Arizona	3.09 <sup>b</sup>	0.10	(48)	-12.99	Nevada	2.87 <sup>b</sup>	0.16	(50)	-19.34 <sup>c</sup>
Arkansas	3.61	0.15	(26)	1.69	New Hampshire	3.62	0.23	(25)	1.96
California	3.27 <sup>b</sup>	0.05	(44)	-7.82	New Jersey	3.52	0.09	(29)	-0.88
Colorado	3.49	0.13	(32)	-1.83	New Mexico	3.31	0.19	(43)	-6.81
Connecticut	3.43	0.13	(36)	-3.41	New York	3.14 <sup>b</sup>	0.06	(46)	-11.47
Delaware	3.33	0.27	(42)	-6.24	North Carolina	4.13 <sup>b</sup>	0.10	(05)	16.38 <sup>c</sup>
D.C.	5.08 <sup>b</sup>	0.42	(01)	42.97 <sup>c</sup>	North Dakota	3.74	0.32	(18)	5.40
Florida	3.09 <sup>b</sup>	0.05	(47)	-12.89	Ohio	3.84 <sup>b</sup>	0.08	(11)	8.00
Georgia	3.74	0.10	(19)	5.32	Oklahoma	3.37	0.13	(39)	-5.13
Hawaii	2.39 <sup>b</sup>	0.18	(51)	-32.68 <sup>c</sup>	Oregon	3.84	0.14	(10)	8.21
Idaho	3.48	0.23	(33)	-2.04	Pennsylvania	3.52	0.07	(30)	-0.93
Illinois	3.71	0.08	(21)	4.47	Rhode Island	3.23	0.23	(45)	-9.04
Indiana	3.63	0.11	(24)	2.24	South Carolina	4.31 <sup>b</sup>	0.14	(03)	21.24 <sup>c</sup>
Iowa	3.42	0.14	(38)	-3.60	South Dakota	3.76	0.29	(17)	5.76
Kansas	3.95	0.17	(09)	11.29	Tennessee	4.15 <sup>b</sup>	0.12	(04)	16.87 <sup>c</sup>
Kentucky	3.73	0.13	(20)	4.97	Texas	3.53	0.06	(28)	-0.71
Louisiana	4.08 <sup>b</sup>	0.14	(06)	14.94	Utah	3.45	0.20	(35)	-2.89
Maine	3.45	0.21	(34)	-2.82	Vermont	3.43	0.32	(37)	-3.54
Maryland	4.04 <sup>b</sup>	0.12	(07)	13.67	Virginia	3.96 <sup>b</sup>	0.11	(08)	11.39
Massachusetts	3.35	0.10	(41)	-5.83	Washington	3.51	0.11	(31)	-1.11
Michigan	3.80	0.09	(13)	6.97	West Virginia	3.67	0.18	(23)	3.27
Minnesota	3.76	0.12	(16)	5.94	Wisconsin	3.57	0.11	(27)	0.50
Mississippi	3.77	0.16	(14)	6.15	Wyoming	3.83	0.39	(12)	7.73
Missouri	3.77	0.11	(15)	6.12					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 18.19  
Myeloma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	4.44	0.03		
High Five States				
D.C.	6.00	0.75	(01)	35.17 <sup>c</sup>
Tennessee	5.36 <sup>b</sup>	0.21	(02)	20.74 <sup>c</sup>
Louisiana	5.21	0.25	(03)	17.26 <sup>c</sup>
Alabama	5.15	0.23	(04)	16.06 <sup>c</sup>
South Carolina	5.09	0.24	(05)	14.50
Low Five States				
Arizona	3.80 <sup>b</sup>	0.17	(47)	-14.51
Rhode Island	3.79	0.40	(48)	-14.71
Nevada	3.53	0.27	(49)	-20.60 <sup>c</sup>
Hawaii	2.98 <sup>b</sup>	0.31	(50)	-32.85 <sup>c</sup>
Alaska	2.65	0.55	(51)	-40.26 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.15	0.23	(04)	16.06 <sup>c</sup>	Montana	4.91	0.46	(14)	10.58
Alaska	2.65	0.55	(51)	-40.26 <sup>c</sup>	Nebraska	3.97	0.32	(44)	-10.64
Arizona	3.80 <sup>b</sup>	0.17	(47)	-14.51	Nevada	3.53	0.27	(49)	-20.60 <sup>c</sup>
Arkansas	4.49	0.26	(27)	1.08	New Hampshire	4.23	0.39	(42)	-4.75
California	4.08 <sup>b</sup>	0.08	(43)	-8.22	New Jersey	4.32	0.15	(34)	-2.84
Colorado	4.41	0.24	(32)	-0.80	New Mexico	4.24	0.32	(41)	-4.53
Connecticut	4.25	0.23	(39)	-4.40	New York	3.96 <sup>b</sup>	0.10	(45)	-10.87
Delaware	4.28	0.48	(38)	-3.62	North Carolina	5.01	0.17	(09)	12.90
D.C.	6.00	0.75	(01)	35.17 <sup>c</sup>	North Dakota	4.62	0.53	(22)	3.97
Florida	3.94 <sup>b</sup>	0.09	(46)	-11.37	Ohio	4.67	0.14	(21)	5.24
Georgia	4.56	0.18	(24)	2.71	Oklahoma	4.41	0.24	(31)	-0.65
Hawaii	2.98 <sup>b</sup>	0.31	(50)	-32.85 <sup>c</sup>	Oregon	5.03	0.25	(08)	13.34
Idaho	4.49	0.39	(28)	1.08	Pennsylvania	4.30	0.12	(37)	-3.29
Illinois	4.44	0.13	(30)	-0.15	Rhode Island	3.79	0.40	(48)	-14.71
Indiana	4.86	0.20	(17)	9.44	South Carolina	5.09	0.24	(05)	14.50
Iowa	4.46	0.25	(29)	0.48	South Dakota	4.79	0.50	(19)	7.81
Kansas	4.96	0.29	(11)	11.61	Tennessee	5.36 <sup>b</sup>	0.21	(02)	20.74 <sup>c</sup>
Kentucky	4.90	0.24	(15)	10.38	Texas	4.37	0.11	(33)	-1.57
Louisiana	5.21	0.25	(03)	17.26 <sup>c</sup>	Utah	4.24	0.33	(40)	-4.48
Maine	4.51	0.38	(25)	1.53	Vermont	4.50	0.58	(26)	1.22
Maryland	4.93	0.22	(13)	10.95	Virginia	5.05	0.19	(07)	13.63
Massachusetts	4.30	0.17	(36)	-3.18	Washington	4.68	0.19	(20)	5.27
Michigan	4.99 <sup>b</sup>	0.15	(10)	12.40	West Virginia	4.80	0.33	(18)	8.13
Minnesota	4.87	0.21	(16)	9.61	Wisconsin	4.56	0.19	(23)	2.77
Mississippi	4.32	0.27	(35)	-2.85	Wyoming	5.05	0.67	(06)	13.78
Missouri	4.96	0.20	(12)	11.57					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.




Table 18.20  
Myeloma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.92	0.02		
High Five States				
D.C.	4.59	0.52	(01)	57.47 <sup>c</sup>
South Carolina	3.83 <sup>b</sup>	0.17	(02)	31.20 <sup>c</sup>
Alabama	3.72 <sup>b</sup>	0.16	(03)	27.36 <sup>c</sup>
North Carolina	3.53 <sup>b</sup>	0.12	(04)	21.06 <sup>c</sup>
Maryland	3.45 <sup>b</sup>	0.15	(05)	18.38 <sup>c</sup>
Low Five States				
New York	2.58 <sup>b</sup>	0.07	(47)	-11.65
Arizona	2.54	0.12	(48)	-12.78
Florida	2.43 <sup>b</sup>	0.06	(49)	-16.73 <sup>c</sup>
Nevada	2.30	0.20	(50)	-21.29 <sup>c</sup>
Hawaii	1.93 <sup>b</sup>	0.22	(51)	-33.86 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.72 <sup>b</sup>	0.16	(03)	27.36 <sup>c</sup>	Montana	2.74	0.31	(38)	-6.10
Alaska	3.03	0.56	(16)	3.94	Nebraska	2.95	0.23	(23)	1.08
Arizona	2.54	0.12	(48)	-12.78	Nevada	2.30	0.20	(50)	-21.29 <sup>c</sup>
Arkansas	2.96	0.19	(20)	1.60	New Hampshire	3.15	0.29	(14)	8.09
California	2.68 <sup>b</sup>	0.05	(42)	-8.03	New Jersey	2.97	0.11	(19)	1.93
Colorado	2.83	0.16	(32)	-2.91	New Mexico	2.59	0.22	(46)	-11.22
Connecticut	2.87	0.16	(29)	-1.73	New York	2.58 <sup>b</sup>	0.07	(47)	-11.65
Delaware	2.65	0.32	(44)	-9.26	North Carolina	3.53 <sup>b</sup>	0.12	(04)	21.06 <sup>c</sup>
D.C.	4.59	0.52	(01)	57.47 <sup>c</sup>	North Dakota	3.05	0.38	(15)	4.49
Florida	2.43 <sup>b</sup>	0.06	(49)	-16.73 <sup>c</sup>	Ohio	3.27 <sup>b</sup>	0.10	(09)	12.16
Georgia	3.18	0.12	(13)	9.18	Oklahoma	2.65	0.16	(45)	-9.29
Hawaii	1.93 <sup>b</sup>	0.22	(51)	-33.86 <sup>c</sup>	Oregon	2.95	0.17	(21)	1.21
Idaho	2.77	0.28	(36)	-4.91	Pennsylvania	2.99	0.08	(18)	2.41
Illinois	3.22	0.09	(11)	10.52	Rhode Island	2.89	0.29	(27)	-1.06
Indiana	2.81	0.12	(35)	-3.79	South Carolina	3.83 <sup>b</sup>	0.17	(02)	31.20 <sup>c</sup>
Iowa	2.69	0.17	(40)	-7.63	South Dakota	2.84	0.34	(31)	-2.59
Kansas	3.26	0.20	(10)	11.77	Tennessee	3.29	0.14	(08)	12.69
Kentucky	2.92	0.15	(25)	0.20	Texas	2.91	0.07	(26)	-0.29
Louisiana	3.30	0.16	(07)	13.21	Utah	2.85	0.24	(30)	-2.35
Maine	2.74	0.25	(39)	-6.21	Vermont	2.75	0.38	(37)	-5.73
Maryland	3.45 <sup>b</sup>	0.15	(05)	18.38 <sup>c</sup>	Virginia	3.22	0.12	(12)	10.33
Massachusetts	2.68	0.11	(41)	-8.02	Washington	2.65	0.13	(43)	-9.08
Michigan	2.95	0.10	(22)	1.11	West Virginia	2.81	0.21	(34)	-3.65
Minnesota	2.99	0.14	(17)	2.43	Wisconsin	2.87	0.13	(28)	-1.45
Mississippi	3.33	0.20	(06)	13.99	Wyoming	2.83	0.46	(33)	-3.05
Missouri	2.93	0.13	(24)	0.46					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 19.22  
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	6.89	0.02		
High Five States				
West Virginia	7.98 <sup>b</sup>	0.27	(01)	15.90 <sup>c</sup>
Michigan	7.70 <sup>b</sup>	0.12	(02)	11.88
Indiana	7.68 <sup>b</sup>	0.16	(03)	11.49
Oregon	7.63 <sup>b</sup>	0.20	(04)	10.81
Pennsylvania	7.61 <sup>b</sup>	0.10	(05)	10.48
Low Five States				
Nevada	6.16	0.24	(47)	-10.47
D.C.	6.11	0.46	(48)	-11.25
New Mexico	5.99 <sup>b</sup>	0.25	(49)	-12.99
Alaska	5.87	0.58	(50)	-14.77
Hawaii	5.74 <sup>b</sup>	0.28	(51)	-16.67 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.92	0.17	(25)	0.46	Montana	7.20	0.37	(18)	4.62
Alaska	5.87	0.58	(50)	-14.77	Nebraska	7.39	0.28	(13)	7.26
Arizona	6.21 <sup>b</sup>	0.14	(46)	-9.79	Nevada	6.16	0.24	(47)	-10.47
Arkansas	6.86	0.21	(27)	-0.31	New Hampshire	6.77	0.32	(29)	-1.63
California	6.55 <sup>b</sup>	0.06	(37)	-4.90	New Jersey	7.10	0.12	(22)	3.11
Colorado	6.43	0.18	(40)	-6.57	New Mexico	5.99 <sup>b</sup>	0.25	(49)	-12.99
Connecticut	6.83	0.19	(28)	-0.82	New York	6.36 <sup>b</sup>	0.08	(43)	-7.64
Delaware	6.75	0.39	(30)	-2.04	North Carolina	6.66	0.12	(32)	-3.28
D.C.	6.11	0.46	(48)	-11.25	North Dakota	6.66	0.42	(33)	-3.32
Florida	6.55 <sup>b</sup>	0.08	(36)	-4.87	Ohio	7.42 <sup>b</sup>	0.11	(12)	7.73
Georgia	6.32 <sup>b</sup>	0.13	(45)	-8.28	Oklahoma	7.38	0.20	(14)	7.11
Hawaii	5.74 <sup>b</sup>	0.28	(51)	-16.67 <sup>c</sup>	Oregon	7.63 <sup>b</sup>	0.20	(04)	10.81
Idaho	7.26	0.33	(16)	5.44	Pennsylvania	7.61 <sup>b</sup>	0.10	(05)	10.48
Illinois	7.12	0.11	(21)	3.39	Rhode Island	6.74	0.33	(31)	-2.06
Indiana	7.68 <sup>b</sup>	0.16	(03)	11.49	South Carolina	6.38	0.17	(42)	-7.41
Iowa	7.43	0.21	(10)	7.91	South Dakota	6.90	0.39	(26)	0.19
Kansas	7.51	0.23	(06)	9.03	Tennessee	7.43	0.16	(11)	7.87
Kentucky	7.49	0.19	(07)	8.73	Texas	6.59 <sup>b</sup>	0.08	(34)	-4.32
Louisiana	7.31	0.19	(15)	6.10	Utah	6.57	0.27	(35)	-4.52
Maine	7.18	0.31	(20)	4.34	Vermont	6.52	0.44	(38)	-5.32
Maryland	6.38	0.15	(41)	-7.40	Virginia	6.51	0.14	(39)	-5.47
Massachusetts	6.96	0.14	(23)	1.13	Washington	7.21	0.15	(17)	4.75
Michigan	7.70 <sup>b</sup>	0.12	(02)	11.88	West Virginia	7.98 <sup>b</sup>	0.27	(01)	15.90 <sup>c</sup>
Minnesota	7.19	0.17	(19)	4.44	Wisconsin	7.43	0.16	(09)	7.97
Mississippi	6.35	0.21	(44)	-7.78	Wyoming	7.48	0.54	(08)	8.65
Missouri	6.94	0.15	(24)	0.78					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 19.23  
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	8.72	0.04		
High Five States				
West Virginia	0.47	(01)	14.86	
Indiana <sup>b</sup>	0.28	(02)	14.74	
Kansas <sup>b</sup>	0.40	(03)	10.71	
Michigan	0.21	(04)	10.61	
Pennsylvania	0.18	(05)	10.25	
Low Five States				
Arizona <sup>b</sup>	0.24	(47)	-11.15	
Hawaii <sup>b</sup>	0.49	(48)	-14.04	
New Mexico	0.42	(49)	-14.92	
Alaska <sup>b</sup>	1.02	(50)	-15.57 <sup>c</sup>	
Nevada <sup>b</sup>	0.38	(51)	-19.46 <sup>c</sup>	

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	8.60	0.30	(30)	-1.37	Montana	8.65	0.62	(29)	-0.80
Alaska	7.36	1.02	(50)	-15.57 <sup>c</sup>	Nebraska	9.08	0.48	(18)	4.07
Arizona	7.75 <sup>b</sup>	0.24	(47)	-11.15	Nevada	7.02 <sup>b</sup>	0.38	(51)	-19.46 <sup>c</sup>
Arkansas	8.91	0.37	(22)	2.13	New Hampshire	8.66	0.57	(28)	-0.69
California	8.34 <sup>b</sup>	0.11	(36)	-4.40	New Jersey	8.94	0.22	(20)	2.56
Colorado	8.37	0.32	(33)	-3.96	New Mexico	7.42	0.42	(49)	-14.92
Connecticut	8.77	0.33	(27)	0.53	New York	7.97 <sup>b</sup>	0.14	(46)	-8.65
Delaware	9.17	0.70	(16)	5.12	North Carolina	8.29	0.22	(38)	-4.92
D.C.	8.84	0.88	(24)	1.39	North Dakota	8.36	0.72	(34)	-4.15
Florida	8.26 <sup>b</sup>	0.13	(40)	-5.31	Ohio	9.55 <sup>b</sup>	0.20	(07)	9.57
Georgia	8.21	0.24	(41)	-5.88	Oklahoma	9.34	0.35	(14)	7.10
Hawaii	7.50	0.49	(48)	-14.04	Oregon	9.53	0.34	(09)	9.33
Idaho	8.43	0.53	(32)	-3.31	Pennsylvania	9.61 <sup>b</sup>	0.18	(05)	10.25
Illinois	9.15	0.19	(17)	4.94	Rhode Island	8.83	0.60	(25)	1.28
Indiana	10.01 <sup>b</sup>	0.28	(02)	14.74	South Carolina	8.00	0.30	(45)	-8.26
Iowa	9.53	0.36	(10)	9.32	South Dakota	8.77	0.67	(26)	0.58
Kansas	9.65	0.40	(03)	10.71	Tennessee	9.55	0.28	(08)	9.46
Kentucky	9.61	0.34	(06)	10.16	Texas	8.30	0.15	(37)	-4.87
Louisiana	9.35	0.33	(13)	7.25	Utah	8.11	0.45	(43)	-6.95
Maine	9.35	0.54	(12)	7.28	Vermont	8.53	0.79	(31)	-2.23
Maryland	8.10	0.28	(44)	-7.07	Virginia	8.19	0.24	(42)	-6.07
Massachusetts	8.92	0.25	(21)	2.27	Washington	9.04	0.27	(19)	3.67
Michigan	9.64 <sup>b</sup>	0.21	(04)	10.61	West Virginia	10.02	0.47	(01)	14.86
Minnesota	9.52	0.30	(11)	9.18	Wisconsin	9.32	0.28	(15)	6.87
Mississippi	8.29	0.38	(39)	-4.98	Wyoming	8.34	0.88	(35)	-4.34
Missouri	8.88	0.26	(23)	1.81					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 19.24  
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.52	0.03		
High Five States				
Wyoming	6.79	0.69	(01)	23.10 <sup>c</sup>
West Virginia	6.44	0.32	(02)	16.73 <sup>c</sup>
Michigan	6.31 <sup>b</sup>	0.15	(03)	14.28
Idaho	6.21	0.41	(04)	12.52
Oregon	6.18	0.24	(05)	12.03
Low Five States				
Mississippi	4.92	0.24	(47)	-10.89
New Mexico	4.86	0.30	(48)	-11.92
Alaska	4.80	0.71	(49)	-13.12
Hawaii	4.31 <sup>b</sup>	0.33	(50)	-21.99 <sup>c</sup>
D.C.	4.09	0.48	(51)	-25.88 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.68	0.20	(21)	2.90	Montana	6.08	0.46	(08)	10.20
Alaska	4.80	0.71	(49)	-13.12	Nebraska	6.17	0.34	(06)	11.83
Arizona	4.92 <sup>b</sup>	0.17	(46)	-10.87	Nevada	5.26	0.31	(33)	-4.74
Arkansas	5.30	0.25	(31)	-4.03	New Hampshire	5.52	0.38	(25)	-0.03
California	5.19 <sup>b</sup>	0.08	(39)	-6.02	New Jersey	5.77	0.15	(19)	4.51
Colorado	4.98	0.21	(44)	-9.85	New Mexico	4.86	0.30	(48)	-11.92
Connecticut	5.47	0.22	(27)	-0.91	New York	5.19 <sup>b</sup>	0.09	(38)	-6.02
Delaware	4.96	0.44	(45)	-10.20	North Carolina	5.43	0.15	(28)	-1.55
D.C.	4.09	0.48	(51)	-25.88 <sup>c</sup>	North Dakota	5.21	0.49	(36)	-5.65
Florida	5.14 <sup>b</sup>	0.09	(40)	-6.84	Ohio	5.88	0.13	(15)	6.56
Georgia	4.99 <sup>b</sup>	0.15	(43)	-9.61	Oklahoma	5.88	0.23	(16)	6.52
Hawaii	4.31 <sup>b</sup>	0.33	(50)	-21.99 <sup>c</sup>	Oregon	6.18	0.24	(05)	12.03
Idaho	6.21	0.41	(04)	12.52	Pennsylvania	6.16 <sup>b</sup>	0.12	(07)	11.61
Illinois	5.70	0.12	(20)	3.32	Rhode Island	5.24	0.38	(34)	-5.08
Indiana	6.04	0.18	(09)	9.37	South Carolina	5.23	0.20	(35)	-5.17
Iowa	5.90	0.24	(13)	6.92	South Dakota	5.40	0.46	(29)	-2.16
Kansas	6.03	0.27	(10)	9.30	Tennessee	5.89	0.18	(14)	6.73
Kentucky	5.98	0.22	(12)	8.42	Texas	5.32	0.10	(30)	-3.53
Louisiana	5.83	0.22	(17)	5.56	Utah	5.21	0.32	(37)	-5.66
Maine	5.65	0.36	(23)	2.27	Vermont	5.13	0.52	(41)	-7.00
Maryland	5.08	0.18	(42)	-8.00	Virginia	5.26	0.16	(32)	-4.71
Massachusetts	5.67	0.16	(22)	2.65	Washington	5.78	0.18	(18)	4.78
Michigan	6.31 <sup>b</sup>	0.15	(03)	14.28	West Virginia	6.44	0.32	(02)	16.73 <sup>c</sup>
Minnesota	5.51	0.19	(26)	-0.20	Wisconsin	6.02	0.19	(11)	9.04
Mississippi	4.92	0.24	(47)	-10.89	Wyoming	6.79	0.69	(01)	23.10 <sup>c</sup>
Missouri	5.57	0.18	(24)	0.86					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 20.24  
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.55	0.01		
High Five States				
D.C.	3.83	0.37	(01)	50.13 <sup>c</sup>
Mississippi	3.43 <sup>b</sup>	0.15	(02)	34.58 <sup>c</sup>
Louisiana	3.27 <sup>b</sup>	0.12	(03)	28.23 <sup>c</sup>
South Carolina	3.14 <sup>b</sup>	0.12	(04)	23.09 <sup>c</sup>
Alabama	3.09 <sup>b</sup>	0.11	(05)	21.14 <sup>c</sup>
Low Five States				
Connecticut	2.05 <sup>b</sup>	0.10	(47)	-19.70 <sup>c</sup>
New Mexico	2.04 <sup>b</sup>	0.14	(48)	-20.07 <sup>c</sup>
Nebraska	1.99 <sup>b</sup>	0.15	(49)	-21.71 <sup>c</sup>
South Dakota	1.95	0.21	(50)	-23.62 <sup>c</sup>
Utah	1.60 <sup>b</sup>	0.13	(51)	-37.02 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.09 <sup>b</sup>	0.11	(05)	21.14 <sup>c</sup>	Montana	2.12	0.20	(45)	-16.64 <sup>c</sup>
Alaska	2.84	0.38	(10)	11.53	Nebraska	1.99 <sup>b</sup>	0.15	(49)	-21.71 <sup>c</sup>
Arizona	2.09 <sup>b</sup>	0.08	(46)	-18.10 <sup>c</sup>	Nevada	2.54	0.15	(24)	-0.21
Arkansas	3.04	0.14	(07)	19.36 <sup>c</sup>	New Hampshire	2.69	0.20	(16)	5.46
California	2.60	0.04	(22)	1.88	New Jersey	2.31	0.07	(35)	-9.46
Colorado	2.17 <sup>b</sup>	0.10	(42)	-14.87	New Mexico	2.04 <sup>b</sup>	0.14	(48)	-20.07 <sup>c</sup>
Connecticut	2.05 <sup>b</sup>	0.10	(47)	-19.70 <sup>c</sup>	New York	2.25 <sup>b</sup>	0.05	(38)	-11.57
Delaware	2.45	0.23	(31)	-3.95	North Carolina	2.70	0.08	(14)	5.77
D.C.	3.83	0.37	(01)	50.13 <sup>c</sup>	North Dakota	2.21	0.25	(39)	-13.19
Florida	2.81 <sup>b</sup>	0.05	(12)	10.18	Ohio	2.67	0.07	(18)	4.79
Georgia	2.82	0.09	(11)	10.60	Oklahoma	2.69	0.12	(17)	5.40
Hawaii	2.94	0.20	(09)	15.19 <sup>c</sup>	Oregon	2.46	0.11	(29)	-3.63
Idaho	2.54	0.19	(26)	-0.47	Pennsylvania	2.32 <sup>b</sup>	0.06	(34)	-9.10
Illinois	2.49	0.06	(28)	-2.36	Rhode Island	2.13	0.19	(44)	-16.49 <sup>c</sup>
Indiana	2.26	0.08	(37)	-11.19	South Carolina	3.14 <sup>b</sup>	0.12	(04)	23.09 <sup>c</sup>
Iowa	2.54	0.12	(25)	-0.32	South Dakota	1.95	0.21	(50)	-23.62 <sup>c</sup>
Kansas	2.13	0.12	(43)	-16.24 <sup>c</sup>	Tennessee	3.04 <sup>b</sup>	0.10	(06)	19.40 <sup>c</sup>
Kentucky	2.81	0.11	(13)	10.12	Texas	2.61	0.05	(20)	2.26
Louisiana	3.27 <sup>b</sup>	0.12	(03)	28.23 <sup>c</sup>	Utah	1.60 <sup>b</sup>	0.13	(51)	-37.02 <sup>c</sup>
Maine	2.64	0.18	(19)	3.45	Vermont	2.32	0.26	(33)	-8.79
Maryland	2.58	0.10	(23)	1.13	Virginia	2.29	0.08	(36)	-10.25
Massachusetts	2.50	0.08	(27)	-2.05	Washington	2.98 <sup>b</sup>	0.10	(08)	17.08 <sup>c</sup>
Michigan	2.60	0.07	(21)	2.09	West Virginia	2.69	0.16	(15)	5.52
Minnesota	2.20 <sup>b</sup>	0.09	(41)	-13.81	Wisconsin	2.45	0.09	(30)	-3.70
Mississippi	3.43 <sup>b</sup>	0.15	(02)	34.58 <sup>c</sup>	Wyoming	2.20	0.29	(40)	-13.71
Missouri	2.38	0.09	(32)	-6.41					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 20.25  
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	3.90	0.02		
High Five States				
D.C.	6.57 <sup>b</sup>	0.75	(01)	68.55 <sup>c</sup>
Mississippi	5.77 <sup>b</sup>	0.30	(02)	47.91 <sup>c</sup>
Louisiana	5.44 <sup>b</sup>	0.24	(03)	39.49 <sup>c</sup>
South Carolina	5.13 <sup>b</sup>	0.23	(04)	31.42 <sup>c</sup>
Arkansas	4.99 <sup>b</sup>	0.27	(05)	27.83 <sup>c</sup>
Low Five States				
Nebraska	2.90 <sup>b</sup>	0.27	(47)	-25.63 <sup>c</sup>
Wyoming	2.89	0.50	(48)	-25.81 <sup>c</sup>
Alaska	2.85	0.55	(49)	-27.03 <sup>c</sup>
South Dakota	2.83	0.38	(50)	-27.36 <sup>c</sup>
Utah	2.17 <sup>b</sup>	0.22	(51)	-44.23 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.93 <sup>b</sup>	0.22	(06)	26.52 <sup>c</sup>	Montana	3.20	0.37	(39)	-18.03 <sup>c</sup>
Alaska	2.85	0.55	(49)	-27.03 <sup>c</sup>	Nebraska	2.90 <sup>b</sup>	0.27	(47)	-25.63 <sup>c</sup>
Arizona	3.15 <sup>b</sup>	0.15	(42)	-19.30 <sup>c</sup>	Nevada	3.57	0.26	(32)	-8.57
Arkansas	4.99 <sup>b</sup>	0.27	(05)	27.83 <sup>c</sup>	New Hampshire	3.90	0.36	(22)	0.12
California	3.86	0.07	(23)	-1.03	New Jersey	3.53	0.13	(35)	-9.50
Colorado	2.91 <sup>b</sup>	0.18	(46)	-25.40 <sup>c</sup>	New Mexico	3.16	0.27	(41)	-19.10 <sup>c</sup>
Connecticut	3.13 <sup>b</sup>	0.19	(43)	-19.63 <sup>c</sup>	New York	3.49 <sup>b</sup>	0.09	(36)	-10.44
Delaware	3.79	0.44	(26)	-2.74	North Carolina	4.04	0.15	(17)	3.56
D.C.	6.57 <sup>b</sup>	0.75	(01)	68.55 <sup>c</sup>	North Dakota	3.24	0.45	(38)	-16.97 <sup>c</sup>
Florida	4.31 <sup>b</sup>	0.09	(12)	10.46	Ohio	4.08	0.12	(15)	4.67
Georgia	4.57 <sup>b</sup>	0.17	(08)	17.23 <sup>c</sup>	Oklahoma	4.08	0.22	(16)	4.58
Hawaii	4.43	0.37	(10)	13.72	Oregon	3.63	0.20	(30)	-6.82
Idaho	3.63	0.34	(31)	-7.00	Pennsylvania	3.64	0.11	(29)	-6.74
Illinois	3.80	0.12	(25)	-2.68	Rhode Island	3.08	0.35	(45)	-21.08 <sup>c</sup>
Indiana	3.47	0.16	(37)	-11.11	South Carolina	5.13 <sup>b</sup>	0.23	(04)	31.42 <sup>c</sup>
Iowa	3.67	0.22	(27)	-5.93	South Dakota	2.83	0.38	(50)	-27.36 <sup>c</sup>
Kansas	3.11 <sup>b</sup>	0.22	(44)	-20.24 <sup>c</sup>	Tennessee	4.85 <sup>b</sup>	0.19	(07)	24.31 <sup>c</sup>
Kentucky	4.45	0.22	(09)	14.23	Texas	4.11	0.10	(14)	5.30
Louisiana	5.44 <sup>b</sup>	0.24	(03)	39.49 <sup>c</sup>	Utah	2.17 <sup>b</sup>	0.22	(51)	-44.23 <sup>c</sup>
Maine	3.99	0.34	(19)	2.37	Vermont	3.97	0.52	(21)	1.92
Maryland	4.17	0.19	(13)	6.82	Virginia	3.56	0.15	(33)	-8.72
Massachusetts	3.85	0.16	(24)	-1.40	Washington	4.33	0.18	(11)	11.01
Michigan	3.98	0.13	(20)	2.05	West Virginia	4.02	0.29	(18)	3.20
Minnesota	3.19 <sup>b</sup>	0.17	(40)	-18.12 <sup>c</sup>	Wisconsin	3.64	0.17	(28)	-6.73
Mississippi	5.77 <sup>b</sup>	0.30	(02)	47.91 <sup>c</sup>	Wyoming	2.89	0.50	(48)	-25.81 <sup>c</sup>
Missouri	3.53	0.16	(34)	-9.41					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.





Table 20.26  
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.43	0.01		
High Five States				
Alaska	2.79	0.54	(01)	94.88 <sup>c</sup>
D.C.	1.87	0.34	(02)	30.13 <sup>c</sup>
Washington	1.84 <sup>b</sup>	0.11	(03)	28.69 <sup>c</sup>
New Hampshire	1.64	0.21	(04)	14.59
Alabama	1.63	0.11	(05)	13.81
Low Five States				
Connecticut	1.19	0.10	(47)	-17.27 <sup>c</sup>
Arizona	1.16	0.08	(48)	-18.81 <sup>c</sup>
Utah	1.07	0.15	(49)	-25.48 <sup>c</sup>
New Mexico	1.05	0.14	(50)	-26.47 <sup>c</sup>
Vermont	0.96	0.22	(51)	-32.76 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.63	0.11	(05)	13.81	Montana	1.20	0.21	(45)	-16.27 <sup>c</sup>
Alaska	2.79	0.54	(01)	94.88 <sup>c</sup>	Nebraska	1.27	0.16	(41)	-11.24
Arizona	1.16	0.08	(48)	-18.81 <sup>c</sup>	Nevada	1.61	0.17	(06)	12.20
Arkansas	1.42	0.13	(31)	-0.72	New Hampshire	1.64	0.21	(04)	14.59
California	1.56	0.04	(11)	8.83	New Jersey	1.33	0.07	(36)	-7.08
Colorado	1.58	0.12	(09)	10.24	New Mexico	1.05	0.14	(50)	-26.47 <sup>c</sup>
Connecticut	1.19	0.10	(47)	-17.27 <sup>c</sup>	New York	1.27 <sup>b</sup>	0.05	(42)	-11.35
Delaware	1.33	0.23	(38)	-7.49	North Carolina	1.57	0.08	(10)	9.43
D.C.	1.87	0.34	(02)	30.13 <sup>c</sup>	North Dakota	1.36	0.26	(35)	-5.33
Florida	1.51	0.05	(18)	5.26	Ohio	1.50	0.06	(20)	4.58
Georgia	1.49	0.08	(22)	3.99	Oklahoma	1.55	0.12	(12)	8.43
Hawaii	1.59	0.20	(08)	11.18	Oregon	1.46	0.12	(27)	1.76
Idaho	1.55	0.21	(14)	7.83	Pennsylvania	1.26	0.05	(44)	-11.99
Illinois	1.41	0.06	(32)	-1.35	Rhode Island	1.43	0.21	(29)	-0.53
Indiana	1.29	0.08	(40)	-9.79	South Carolina	1.52	0.11	(17)	5.79
Iowa	1.59	0.13	(07)	11.25	South Dakota	1.19	0.22	(46)	-17.11 <sup>c</sup>
Kansas	1.33	0.13	(37)	-7.28	Tennessee	1.55	0.09	(13)	8.15
Kentucky	1.46	0.11	(25)	2.13	Texas	1.37	0.05	(34)	-4.12
Louisiana	1.52	0.11	(16)	6.12	Utah	1.07	0.15	(49)	-25.48 <sup>c</sup>
Maine	1.50	0.18	(19)	4.78	Vermont	0.96	0.22	(51)	-32.76 <sup>c</sup>
Maryland	1.31	0.09	(39)	-8.70	Virginia	1.27	0.08	(43)	-11.44
Massachusetts	1.46	0.09	(28)	1.57	Washington	1.84 <sup>b</sup>	0.11	(03)	28.69 <sup>c</sup>
Michigan	1.48	0.07	(23)	3.19	West Virginia	1.50	0.16	(21)	4.49
Minnesota	1.39	0.10	(33)	-3.10	Wisconsin	1.46	0.09	(26)	1.87
Mississippi	1.47	0.13	(24)	2.23	Wyoming	1.53	0.33	(15)	6.77
Missouri	1.43	0.09	(30)	-0.55					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 21.14  
Cancer of the Ovary (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	8.59	0.03		
High Five States				
Washington	10.08 <sup>b</sup>	0.25	(01)	17.33 <sup>c</sup>
Oregon	9.76 <sup>b</sup>	0.30	(02)	13.60
Wisconsin	9.49 <sup>b</sup>	0.24	(03)	10.47
Wyoming	9.41	0.82	(04)	9.54
Alabama	9.33	0.26	(05)	8.59
Low Five States				
Florida	7.82 <sup>b</sup>	0.11	(47)	-9.01
Mississippi	7.62	0.30	(48)	-11.35
D.C.	7.54	0.68	(49)	-12.32
Hawaii	6.85 <sup>b</sup>	0.43	(50)	-20.27 <sup>c</sup>
Alaska	6.17	0.75	(51)	-28.16 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.33	0.26	(05)	8.59	Montana	9.09	0.56	(09)	5.73
Alaska	6.17	0.75	(51)	-28.16 <sup>c</sup>	Nebraska	8.41	0.41	(33)	-2.13
Arizona	8.59	0.23	(25)	-0.03	Nevada	8.17	0.37	(41)	-4.94
Arkansas	8.32	0.31	(37)	-3.23	New Hampshire	8.36	0.47	(35)	-2.69
California	8.63	0.10	(23)	0.44	New Jersey	9.01	0.19	(12)	4.79
Colorado	8.99	0.28	(13)	4.64	New Mexico	8.08	0.39	(42)	-5.93
Connecticut	8.57	0.28	(27)	-0.32	New York	8.59	0.12	(26)	-0.06
Delaware	8.93	0.60	(15)	3.90	North Carolina	8.29	0.18	(38)	-3.55
D.C.	7.54	0.68	(49)	-12.32	North Dakota	8.38	0.67	(34)	-2.51
Florida	7.82 <sup>b</sup>	0.11	(47)	-9.01	Ohio	8.61	0.16	(24)	0.16
Georgia	8.71	0.20	(22)	1.39	Oklahoma	8.44	0.28	(32)	-1.83
Hawaii	6.85 <sup>b</sup>	0.43	(50)	-20.27 <sup>c</sup>	Oregon	9.76 <sup>b</sup>	0.30	(02)	13.60
Idaho	8.53	0.49	(28)	-0.74	Pennsylvania	9.26 <sup>b</sup>	0.15	(06)	7.74
Illinois	8.72	0.16	(21)	1.47	Rhode Island	9.20	0.53	(07)	7.04
Indiana	8.87	0.22	(17)	3.19	South Carolina	8.25	0.26	(39)	-3.95
Iowa	9.07	0.31	(11)	5.55	South Dakota	8.84	0.62	(19)	2.82
Kansas	9.10	0.34	(08)	5.92	Tennessee	8.78	0.22	(20)	2.13
Kentucky	8.45	0.26	(30)	-1.62	Texas	7.85 <sup>b</sup>	0.12	(46)	-8.66
Louisiana	8.47	0.26	(29)	-1.47	Utah	8.34	0.41	(36)	-2.93
Maine	8.00	0.43	(44)	-6.92	Vermont	7.96	0.65	(45)	-7.41
Maryland	8.23	0.23	(40)	-4.23	Virginia	8.93	0.21	(16)	3.86
Massachusetts	8.85	0.21	(18)	3.03	Washington	10.08 <sup>b</sup>	0.25	(01)	17.33 <sup>c</sup>
Michigan	8.95	0.18	(14)	4.18	West Virginia	9.08	0.39	(10)	5.66
Minnesota	8.44	0.24	(31)	-1.80	Wisconsin	9.49 <sup>b</sup>	0.24	(03)	10.47
Mississippi	7.62	0.30	(48)	-11.35	Wyoming	9.41	0.82	(04)	9.54
Missouri	8.07	0.21	(43)	-6.12					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 22.18  
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	10.70	0.03		
High Five States				
D.C.	12.48	0.66	(01)	16.56 <sup>c</sup>
Louisiana	12.03 <sup>b</sup>	0.24	(02)	12.40
Connecticut	11.83 <sup>b</sup>	0.25	(03)	10.52
New Hampshire	11.69	0.42	(04)	9.23
Massachusetts	11.58 <sup>b</sup>	0.18	(05)	8.22
Low Five States				
Colorado	9.78 <sup>b</sup>	0.22	(47)	-8.61
Vermont	9.77	0.54	(48)	-8.70
West Virginia	9.24 <sup>b</sup>	0.29	(49)	-13.69
Arizona	9.23 <sup>b</sup>	0.17	(50)	-13.73
Utah	8.88 <sup>b</sup>	0.31	(51)	-17.07 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	10.75	0.21	(28)	0.47	Montana	10.40	0.44	(35)	-2.82
Alaska	10.44	0.76	(34)	-2.47	Nebraska	9.96	0.33	(46)	-6.96
Arizona	9.23 <sup>b</sup>	0.17	(50)	-13.73	Nevada	10.69	0.31	(29)	-0.10
Arkansas	10.82	0.27	(23)	1.13	New Hampshire	11.69	0.42	(04)	9.23
California	10.32 <sup>b</sup>	0.08	(37)	-3.56	New Jersey	11.21	0.16	(14)	4.71
Colorado	9.78 <sup>b</sup>	0.22	(47)	-8.61	New Mexico	10.05	0.32	(42)	-6.11
Connecticut	11.83 <sup>b</sup>	0.25	(03)	10.52	New York	10.83	0.10	(22)	1.23
Delaware	10.33	0.48	(36)	-3.48	North Carolina	11.05	0.16	(18)	3.25
D.C.	12.48	0.66	(01)	16.56 <sup>c</sup>	North Dakota	10.61	0.54	(32)	-0.84
Florida	10.00 <sup>b</sup>	0.09	(43)	-6.53	Ohio	11.06	0.13	(16)	3.35
Georgia	10.52	0.17	(33)	-1.68	Oklahoma	9.97	0.23	(45)	-6.88
Hawaii	10.64	0.39	(31)	-0.57	Oregon	11.10	0.24	(15)	3.74
Idaho	10.92	0.40	(20)	1.99	Pennsylvania	11.43 <sup>b</sup>	0.12	(07)	6.76
Illinois	11.25 <sup>b</sup>	0.13	(11)	5.07	Rhode Island	10.25	0.41	(40)	-4.23
Indiana	11.06	0.19	(17)	3.34	South Carolina	10.67	0.22	(30)	-0.32
Iowa	10.13	0.24	(41)	-5.34	South Dakota	10.28	0.48	(39)	-3.99
Kansas	10.82	0.27	(24)	1.10	Tennessee	10.78	0.19	(25)	0.73
Kentucky	10.77	0.22	(27)	0.65	Texas	9.97 <sup>b</sup>	0.10	(44)	-6.87
Louisiana	12.03 <sup>b</sup>	0.24	(02)	12.40	Utah	8.88 <sup>b</sup>	0.31	(51)	-17.07 <sup>c</sup>
Maine	11.24	0.38	(12)	4.99	Vermont	9.77	0.54	(48)	-8.70
Maryland	11.54 <sup>b</sup>	0.21	(06)	7.80	Virginia	11.23	0.18	(13)	4.92
Massachusetts	11.58 <sup>b</sup>	0.18	(05)	8.22	Washington	10.78	0.19	(26)	0.71
Michigan	11.33 <sup>b</sup>	0.15	(10)	5.89	West Virginia	9.24 <sup>b</sup>	0.29	(49)	-13.69
Minnesota	10.31	0.20	(38)	-3.68	Wisconsin	10.88	0.19	(21)	1.66
Mississippi	11.38	0.28	(09)	6.31	Wyoming	11.41	0.67	(08)	6.63
Missouri	11.02	0.19	(19)	3.00					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table 22.19  
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	12.35	0.04		
High Five States				
D.C.	15.46	1.18	(01)	25.21 <sup>c</sup>
Connecticut	14.15 <sup>b</sup>	0.42	(02)	14.63 <sup>c</sup>
Louisiana	13.60	0.39	(03)	10.16 <sup>c</sup>
Mississippi	13.50	0.48	(04)	9.37
Massachusetts	13.41	0.30	(05)	8.62
Low Five States				
New Mexico	11.16	0.51	(47)	-9.64
Arizona	11.02 <sup>b</sup>	0.28	(48)	-10.77 <sup>c</sup>
Colorado	10.93 <sup>b</sup>	0.36	(49)	-11.51 <sup>c</sup>
Vermont	10.83	0.86	(50)	-12.27 <sup>c</sup>
Utah	9.81 <sup>b</sup>	0.49	(51)	-20.52 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	12.74	0.36	(17)	3.23	Montana	12.13	0.71	(31)	-1.71
Alaska	11.74	1.22	(35)	-4.95	Nebraska	12.12	0.55	(32)	-1.86
Arizona	11.02 <sup>b</sup>	0.28	(48)	-10.77 <sup>c</sup>	Nevada	11.95	0.49	(33)	-3.24
Arkansas	12.58	0.44	(19)	1.86	New Hampshire	12.35	0.66	(26)	0.07
California	11.58 <sup>b</sup>	0.13	(40)	-6.19	New Jersey	13.06	0.26	(09)	5.77
Colorado	10.93 <sup>b</sup>	0.36	(49)	-11.51 <sup>c</sup>	New Mexico	11.16	0.51	(47)	-9.64
Connecticut	14.15 <sup>b</sup>	0.42	(02)	14.63 <sup>c</sup>	New York	12.39	0.17	(25)	0.40
Delaware	11.16	0.76	(46)	-9.60	North Carolina	12.84	0.27	(15)	4.00
D.C.	15.46	1.18	(01)	25.21 <sup>c</sup>	North Dakota	11.57	0.85	(41)	-6.27
Florida	11.69 <sup>b</sup>	0.15	(37)	-5.32	Ohio	12.90	0.22	(13)	4.49
Georgia	12.51	0.29	(22)	1.31	Oklahoma	11.77	0.38	(34)	-4.64
Hawaii	12.24	0.62	(29)	-0.86	Oregon	12.40	0.38	(24)	0.47
Idaho	11.48	0.61	(43)	-7.05	Pennsylvania	13.37 <sup>b</sup>	0.21	(07)	8.28
Illinois	13.00	0.22	(10)	5.32	Rhode Island	11.52	0.68	(42)	-6.72
Indiana	13.11	0.32	(08)	6.22	South Carolina	12.35	0.37	(27)	0.07
Iowa	11.67	0.40	(38)	-5.45	South Dakota	11.22	0.76	(45)	-9.10
Kansas	12.55	0.45	(21)	1.62	Tennessee	12.66	0.32	(18)	2.54
Kentucky	12.49	0.37	(23)	1.13	Texas	11.64 <sup>b</sup>	0.17	(39)	-5.74
Louisiana	13.60	0.39	(03)	10.16 <sup>c</sup>	Utah	9.81 <sup>b</sup>	0.49	(51)	-20.52 <sup>c</sup>
Maine	12.93	0.62	(12)	4.72	Vermont	10.83	0.86	(50)	-12.27 <sup>c</sup>
Maryland	12.79	0.34	(16)	3.62	Virginia	12.96	0.30	(11)	4.94
Massachusetts	13.41	0.30	(05)	8.62	Washington	12.19	0.30	(30)	-1.27
Michigan	13.40 <sup>b</sup>	0.25	(06)	8.51	West Virginia	11.35	0.49	(44)	-8.06
Minnesota	11.70	0.33	(36)	-5.24	Wisconsin	12.56	0.32	(20)	1.76
Mississippi	13.50	0.48	(04)	9.37	Wyoming	12.29	1.04	(28)	-0.49
Missouri	12.89	0.31	(14)	4.38					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 22.20  
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	9.36	0.03		
High Five States				
New Hampshire	11.22	0.54	(01)	19.91 <sup>c</sup>
Louisiana	10.74 <sup>b</sup>	0.29	(02)	14.79
Wyoming	10.72	0.88	(03)	14.58
Maryland	10.52 <sup>b</sup>	0.26	(04)	12.43
D.C.	10.44	0.78	(05)	11.64
Low Five States				
Oklahoma	8.52	0.28	(47)	-8.94
Nebraska	8.38	0.40	(48)	-10.48
Utah	8.08	0.40	(49)	-13.69
Arizona	7.68 <sup>b</sup>	0.21	(50)	-17.86 <sup>c</sup>
West Virginia	7.55 <sup>b</sup>	0.35	(51)	-19.33 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.20	0.25	(36)	-1.64	Montana	8.84	0.55	(43)	-5.56
Alaska	9.35	0.99	(31)	-0.10	Nebraska	8.38	0.40	(48)	-10.48
Arizona	7.68 <sup>b</sup>	0.21	(50)	-17.86 <sup>c</sup>	Nevada	9.53	0.41	(22)	-1.92
Arkansas	9.39	0.33	(28)	0.33	New Hampshire	11.22	0.54	(01)	19.91 <sup>c</sup>
California	9.25	0.10	(34)	-1.09	New Jersey	9.79	0.19	(14)	4.59
Colorado	8.77	0.28	(44)	-6.28	New Mexico	9.15	0.41	(38)	-2.25
Connecticut	10.01	0.30	(08)	6.99	New York	9.64	0.13	(17)	3.06
Delaware	9.52	0.61	(24)	1.71	North Carolina	9.58	0.20	(18)	2.44
D.C.	10.44	0.78	(05)	11.64	North Dakota	9.52	0.68	(23)	1.80
Florida	8.55 <sup>b</sup>	0.12	(46)	-8.61	Ohio	9.58	0.16	(19)	2.39
Georgia	9.06	0.20	(40)	-3.21	Oklahoma	8.52	0.28	(47)	-8.94
Hawaii	9.25	0.48	(35)	-1.09	Oregon	9.99	0.31	(09)	6.79
Idaho	10.27	0.52	(06)	9.82	Pennsylvania	9.85	0.15	(12)	5.32
Illinois	9.85	0.16	(11)	5.33	Rhode Island	9.31	0.52	(32)	-0.48
Indiana	9.42	0.23	(27)	0.71	South Carolina	9.30	0.27	(33)	-0.61
Iowa	8.89	0.30	(41)	-4.97	South Dakota	9.55	0.63	(21)	2.05
Kansas	9.38	0.34	(29)	0.22	Tennessee	9.19	0.23	(37)	-1.80
Kentucky	9.36	0.27	(30)	0.01	Texas	8.62 <sup>b</sup>	0.13	(45)	-7.86
Louisiana	10.74 <sup>b</sup>	0.29	(02)	14.79	Utah	8.08	0.40	(49)	-13.69
Maine	9.85	0.47	(10)	5.33	Vermont	8.88	0.69	(42)	-5.08
Maryland	10.52 <sup>b</sup>	0.26	(04)	12.43	Virginia	9.82	0.22	(13)	4.94
Massachusetts	10.18 <sup>b</sup>	0.22	(07)	8.80	Washington	9.57	0.24	(20)	2.30
Michigan	9.66	0.18	(16)	3.23	West Virginia	7.55 <sup>b</sup>	0.35	(51)	-19.33 <sup>c</sup>
Minnesota	9.14	0.25	(39)	-2.36	Wisconsin	9.45	0.24	(26)	0.99
Mississippi	9.76	0.34	(15)	4.34	Wyoming	10.72	0.88	(03)	14.58
Missouri	9.47	0.23	(25)	1.25					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 23.14  
Cancer of the Prostate (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	24.72	0.07		
High Five States				
D.C.	41.68 <sup>b</sup>	2.03	(01)	68.59 <sup>c</sup>
Mississippi	32.14 <sup>b</sup>	0.79	(02)	30.02 <sup>c</sup>
Alabama	30.11 <sup>b</sup>	0.59	(03)	21.81 <sup>c</sup>
South Carolina	28.89 <sup>b</sup>	0.61	(04)	16.85 <sup>c</sup>
Georgia	28.88 <sup>b</sup>	0.49	(05)	16.83 <sup>c</sup>
Low Five States				
Wyoming	21.82	1.47	(47)	-11.72 <sup>c</sup>
Arizona	20.99 <sup>b</sup>	0.40	(48)	-15.09 <sup>c</sup>
Alaska	20.60	1.84	(49)	-16.68 <sup>c</sup>
Florida	20.52 <sup>b</sup>	0.20	(50)	-16.99 <sup>c</sup>
Hawaii	17.09 <sup>b</sup>	0.74	(51)	-30.86 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	30.11 <sup>b</sup>	0.59	(03)	21.81 <sup>c</sup>	Montana	27.85	1.13	(07)	12.65 <sup>c</sup>
Alaska	20.60	1.84	(49)	-16.68 <sup>c</sup>	Nebraska	24.48	0.80	(35)	-0.99
Arizona	20.99 <sup>b</sup>	0.40	(48)	-15.09 <sup>c</sup>	Nevada	24.53	0.77	(34)	-0.78
Arkansas	27.06	0.67	(14)	9.46	New Hampshire	26.25	1.04	(17)	6.17
California	23.58 <sup>b</sup>	0.19	(42)	-4.63	New Jersey	23.87	0.37	(39)	-3.46
Colorado	24.46	0.59	(36)	-1.07	New Mexico	25.38	0.81	(29)	2.68
Connecticut	25.51	0.58	(26)	3.19	New York	23.46 <sup>b</sup>	0.25	(43)	-5.10
Delaware	26.07	1.24	(19)	5.47	North Carolina	27.74 <sup>b</sup>	0.43	(08)	12.19 <sup>c</sup>
D.C.	41.68 <sup>b</sup>	2.03	(01)	68.59 <sup>c</sup>	North Dakota	26.40	1.28	(15)	6.77
Florida	20.52 <sup>b</sup>	0.20	(50)	-16.99 <sup>c</sup>	Ohio	26.27 <sup>b</sup>	0.34	(16)	6.28
Georgia	28.88 <sup>b</sup>	0.49	(05)	16.83 <sup>c</sup>	Oklahoma	23.65	0.57	(40)	-4.36
Hawaii	17.09 <sup>b</sup>	0.74	(51)	-30.86 <sup>c</sup>	Oregon	25.99	0.57	(21)	5.13
Idaho	27.47	1.00	(11)	11.13 <sup>c</sup>	Pennsylvania	25.03	0.29	(32)	1.23
Illinois	26.22 <sup>b</sup>	0.34	(18)	6.07	Rhode Island	24.16	1.01	(37)	-2.29
Indiana	25.55	0.47	(25)	3.36	South Carolina	28.89 <sup>b</sup>	0.61	(04)	16.85 <sup>c</sup>
Iowa	25.88	0.60	(22)	4.70	South Dakota	26.04	1.16	(20)	5.33
Kansas	22.57	0.62	(45)	-8.69	Tennessee	27.61 <sup>b</sup>	0.51	(09)	11.70 <sup>c</sup>
Kentucky	25.82	0.59	(23)	4.45	Texas	23.14 <sup>b</sup>	0.26	(44)	-6.41
Louisiana	28.77 <sup>b</sup>	0.61	(06)	16.39 <sup>c</sup>	Utah	25.70	0.85	(24)	3.96
Maine	25.44	0.93	(27)	2.88	Vermont	25.25	1.41	(31)	2.13
Maryland	27.49 <sup>b</sup>	0.54	(10)	11.18 <sup>c</sup>	Virginia	27.30 <sup>b</sup>	0.47	(12)	10.41 <sup>c</sup>
Massachusetts	24.63	0.43	(33)	-0.39	Washington	25.40	0.46	(28)	2.75
Michigan	23.89	0.35	(38)	-3.35	West Virginia	22.29	0.74	(46)	-9.86
Minnesota	25.32	0.50	(30)	2.42	Wisconsin	27.08 <sup>b</sup>	0.48	(13)	9.56
Mississippi	32.14 <sup>b</sup>	0.79	(02)	30.02 <sup>c</sup>	Wyoming	21.82	1.47	(47)	-11.72 <sup>c</sup>
Missouri	23.62	0.45	(41)	-4.46					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.




Table 24.18  
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.83	0.02		
High Five States				
Hawaii	6.90 <sup>b</sup>	0.31	(01)	80.30 <sup>c</sup>
D.C.	5.04	0.42	(02)	31.55 <sup>c</sup>
Louisiana	4.92 <sup>b</sup>	0.15	(03)	28.38 <sup>c</sup>
New Mexico	4.80 <sup>b</sup>	0.22	(04)	25.28 <sup>c</sup>
New York	4.66 <sup>b</sup>	0.07	(05)	21.57 <sup>c</sup>
Low Five States				
Wyoming	2.84	0.33	(47)	-25.88 <sup>c</sup>
Iowa	2.73 <sup>b</sup>	0.13	(48)	-28.83 <sup>c</sup>
Nebraska	2.66 <sup>b</sup>	0.17	(49)	-30.58 <sup>c</sup>
Vermont	2.51 <sup>b</sup>	0.27	(50)	-34.53 <sup>c</sup>
Utah	2.50 <sup>b</sup>	0.16	(51)	-34.69 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.83	0.13	(17)	-0.04	Montana	2.88 <sup>b</sup>	0.23	(45)	-24.88 <sup>c</sup>
Alaska	4.20	0.46	(11)	9.79	Nebraska	2.66 <sup>b</sup>	0.17	(49)	-30.58 <sup>c</sup>
Arizona	3.11 <sup>b</sup>	0.10	(39)	-18.68 <sup>c</sup>	Nevada	3.69	0.19	(22)	-3.56
Arkansas	3.55	0.15	(26)	-7.34	New Hampshire	3.01 <sup>b</sup>	0.21	(41)	-21.27 <sup>c</sup>
California	4.65 <sup>b</sup>	0.05	(06)	21.48 <sup>c</sup>	New Jersey	4.59 <sup>b</sup>	0.10	(07)	19.78 <sup>c</sup>
Colorado	3.18 <sup>b</sup>	0.13	(37)	-17.02 <sup>c</sup>	New Mexico	4.80 <sup>b</sup>	0.22	(04)	25.28 <sup>c</sup>
Connecticut	4.30	0.15	(09)	12.26	New York	4.66 <sup>b</sup>	0.07	(05)	21.57 <sup>c</sup>
Delaware	3.89	0.29	(16)	1.59	North Carolina	3.57	0.09	(25)	-6.67
D.C.	5.04	0.42	(02)	31.55 <sup>c</sup>	North Dakota	3.75	0.32	(20)	-2.00
Florida	3.43 <sup>b</sup>	0.06	(27)	-10.42	Ohio	3.42 <sup>b</sup>	0.07	(28)	-10.66
Georgia	3.78	0.10	(19)	-1.41	Oklahoma	3.15 <sup>b</sup>	0.13	(38)	-17.82 <sup>c</sup>
Hawaii	6.90 <sup>b</sup>	0.31	(01)	80.30 <sup>c</sup>	Oregon	3.00 <sup>b</sup>	0.12	(43)	-21.72 <sup>c</sup>
Idaho	2.87 <sup>b</sup>	0.21	(46)	-24.98 <sup>c</sup>	Pennsylvania	3.71	0.07	(21)	-3.19
Illinois	4.20 <sup>b</sup>	0.08	(12)	9.67	Rhode Island	4.44	0.27	(08)	15.90 <sup>c</sup>
Indiana	3.00 <sup>b</sup>	0.10	(42)	-21.56 <sup>c</sup>	South Carolina	4.16	0.14	(13)	8.72
Iowa	2.73 <sup>b</sup>	0.13	(48)	-28.83 <sup>c</sup>	South Dakota	3.30	0.28	(32)	-13.81
Kansas	2.89 <sup>b</sup>	0.14	(44)	-24.43 <sup>c</sup>	Tennessee	3.40 <sup>b</sup>	0.11	(29)	-11.16
Kentucky	3.35 <sup>b</sup>	0.12	(31)	-12.57	Texas	4.12 <sup>b</sup>	0.07	(14)	7.52
Louisiana	4.92 <sup>b</sup>	0.15	(03)	28.38 <sup>c</sup>	Utah	2.50 <sup>b</sup>	0.16	(51)	-34.69 <sup>c</sup>
Maine	3.18	0.20	(36)	-16.92 <sup>c</sup>	Vermont	2.51 <sup>b</sup>	0.27	(50)	-34.53 <sup>c</sup>
Maryland	3.96	0.12	(15)	3.30	Virginia	3.58	0.10	(23)	-6.52
Massachusetts	3.82	0.10	(18)	-0.22	Washington	3.40 <sup>b</sup>	0.11	(30)	-11.16
Michigan	3.58	0.08	(24)	-6.60	West Virginia	3.26	0.17	(33)	-14.95
Minnesota	3.07 <sup>b</sup>	0.11	(40)	-19.70 <sup>c</sup>	Wisconsin	3.22 <sup>b</sup>	0.10	(34)	-15.83 <sup>c</sup>
Mississippi	4.22	0.17	(10)	10.31	Wyoming	2.84	0.33	(47)	-25.88 <sup>c</sup>
Missouri	3.21 <sup>b</sup>	0.10	(35)	-16.07 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 24.19  
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.30	0.03		
High Five States				
Hawaii	9.78 <sup>b</sup>	0.56	(01)	84.65 <sup>c</sup>
Louisiana	7.20 <sup>b</sup>	0.29	(02)	35.82 <sup>c</sup>
Rhode Island	7.15	0.54	(03)	35.01 <sup>c</sup>
D.C.	6.98	0.78	(04)	31.64 <sup>c</sup>
New Mexico	6.90 <sup>b</sup>	0.41	(05)	30.23 <sup>c</sup>
Low Five States				
Iowa	3.81 <sup>b</sup>	0.23	(47)	-28.12 <sup>c</sup>
Vermont	3.70	0.52	(48)	-30.17 <sup>c</sup>
Nebraska	3.56 <sup>b</sup>	0.30	(49)	-32.87 <sup>c</sup>
Wyoming	3.43	0.55	(50)	-35.29 <sup>c</sup>
Utah	3.13 <sup>b</sup>	0.28	(51)	-41.02 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.45	0.24	(15)	2.82	Montana	3.86	0.41	(46)	-27.16 <sup>c</sup>
Alaska	5.36	0.78	(17)	1.17	Nebraska	3.56 <sup>b</sup>	0.30	(49)	-32.87 <sup>c</sup>
Arizona	4.03 <sup>b</sup>	0.17	(44)	-23.98 <sup>c</sup>	Nevada	4.79	0.32	(26)	-9.63
Arkansas	4.56	0.26	(32)	-13.93	New Hampshire	4.45	0.40	(33)	-16.04 <sup>c</sup>
California	6.24 <sup>b</sup>	0.10	(08)	17.85 <sup>c</sup>	New Jersey	6.46 <sup>b</sup>	0.19	(06)	21.86 <sup>c</sup>
Colorado	4.37 <sup>b</sup>	0.23	(39)	-17.54 <sup>c</sup>	New Mexico	6.90 <sup>b</sup>	0.41	(05)	30.23 <sup>c</sup>
Connecticut	6.04	0.28	(09)	14.07	New York	6.37 <sup>b</sup>	0.12	(07)	20.18 <sup>c</sup>
Delaware	5.44	0.54	(16)	2.65	North Carolina	5.13	0.17	(21)	-3.15
D.C.	6.98	0.78	(04)	31.64 <sup>c</sup>	North Dakota	4.83	0.55	(25)	-8.91
Florida	4.62 <sup>b</sup>	0.10	(30)	-12.86	Ohio	4.89	0.14	(24)	-7.72
Georgia	5.23	0.19	(19)	-1.23	Oklahoma	4.24 <sup>b</sup>	0.23	(42)	-19.98 <sup>c</sup>
Hawaii	9.78 <sup>b</sup>	0.56	(01)	84.65 <sup>c</sup>	Oregon	3.96 <sup>b</sup>	0.22	(45)	-25.22 <sup>c</sup>
Idaho	4.15	0.37	(43)	-21.71 <sup>c</sup>	Pennsylvania	5.23	0.13	(20)	-1.37
Illinois	5.90 <sup>b</sup>	0.15	(12)	11.29	Rhode Island	7.15	0.54	(03)	35.01 <sup>c</sup>
Indiana	4.27 <sup>b</sup>	0.18	(41)	-19.33 <sup>c</sup>	South Carolina	6.01	0.26	(10)	13.41
Iowa	3.81 <sup>b</sup>	0.23	(47)	-28.12 <sup>c</sup>	South Dakota	4.40	0.48	(37)	-17.01 <sup>c</sup>
Kansas	4.33 <sup>b</sup>	0.26	(40)	-18.35 <sup>c</sup>	Tennessee	4.73	0.20	(28)	-10.68
Kentucky	4.75	0.24	(27)	-10.44	Texas	5.65	0.12	(14)	6.63
Louisiana	7.20 <sup>b</sup>	0.29	(02)	35.82 <sup>c</sup>	Utah	3.13 <sup>b</sup>	0.28	(51)	-41.02 <sup>c</sup>
Maine	4.41	0.37	(36)	-16.85 <sup>c</sup>	Vermont	3.70	0.52	(48)	-30.17 <sup>c</sup>
Maryland	5.24	0.22	(18)	-1.18	Virginia	5.12	0.19	(22)	-3.29
Massachusetts	5.68	0.20	(13)	7.20	Washington	4.44 <sup>b</sup>	0.19	(34)	-16.26 <sup>c</sup>
Michigan	5.03	0.15	(23)	-5.12	West Virginia	4.42	0.31	(35)	-16.55 <sup>c</sup>
Minnesota	4.38 <sup>b</sup>	0.20	(38)	-17.37 <sup>c</sup>	Wisconsin	4.73	0.20	(29)	-10.73
Mississippi	5.99	0.32	(11)	13.02	Wyoming	3.43	0.55	(50)	-35.29 <sup>c</sup>
Missouri	4.60 <sup>b</sup>	0.19	(31)	-13.17					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.



Table 24.20  
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.74	0.02		
High Five States				
Hawaii	4.73 <sup>b</sup>	0.35	(01)	72.69 <sup>c</sup>
D.C.	3.63	0.46	(02)	32.48 <sup>c</sup>
California	3.47 <sup>b</sup>	0.06	(03)	26.57 <sup>c</sup>
New York	3.46 <sup>b</sup>	0.08	(04)	26.24 <sup>c</sup>
New Jersey	3.29 <sup>b</sup>	0.11	(05)	20.00 <sup>c</sup>
Low Five States				
Iowa	1.92 <sup>b</sup>	0.14	(47)	-30.14 <sup>c</sup>
New Hampshire	1.87 <sup>b</sup>	0.22	(48)	-31.71 <sup>c</sup>
Idaho	1.81 <sup>b</sup>	0.22	(49)	-33.94 <sup>c</sup>
Kansas	1.79 <sup>b</sup>	0.15	(50)	-34.66 <sup>c</sup>
Vermont	1.61 <sup>b</sup>	0.29	(51)	-41.25 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.65	0.14	(21)	-3.32	Montana	2.10	0.27	(43)	-23.49 <sup>c</sup>
Alaska	3.12	0.54	(08)	13.79	Nebraska	1.99 <sup>b</sup>	0.19	(45)	-27.38 <sup>c</sup>
Arizona	2.39	0.12	(32)	-12.82	Nevada	2.74	0.22	(19)	-0.14
Arkansas	2.76	0.18	(15)	0.66	New Hampshire	1.87 <sup>b</sup>	0.22	(48)	-31.71 <sup>c</sup>
California	3.47 <sup>b</sup>	0.06	(03)	26.57 <sup>c</sup>	New Jersey	3.29 <sup>b</sup>	0.11	(05)	20.00 <sup>c</sup>
Colorado	2.30	0.14	(37)	-16.05 <sup>c</sup>	New Mexico	3.13	0.24	(07)	14.30
Connecticut	3.00	0.16	(10)	9.35	New York	3.46 <sup>b</sup>	0.08	(04)	26.24 <sup>c</sup>
Delaware	2.70	0.33	(20)	-1.41	North Carolina	2.44	0.10	(30)	-11.00
D.C.	3.63	0.46	(02)	32.48 <sup>c</sup>	North Dakota	2.90	0.38	(13)	5.81
Florida	2.48 <sup>b</sup>	0.06	(27)	-9.60	Ohio	2.38 <sup>b</sup>	0.08	(33)	-13.24
Georgia	2.76	0.11	(16)	0.57	Oklahoma	2.31	0.15	(35)	-15.82 <sup>c</sup>
Hawaii	4.73 <sup>b</sup>	0.35	(01)	72.69 <sup>c</sup>	Oregon	2.24	0.15	(38)	-18.27 <sup>c</sup>
Idaho	1.81 <sup>b</sup>	0.22	(49)	-33.94 <sup>c</sup>	Pennsylvania	2.60	0.08	(23)	-5.20
Illinois	2.99	0.09	(11)	9.15	Rhode Island	2.75	0.28	(18)	0.15
Indiana	2.08 <sup>b</sup>	0.11	(44)	-23.96 <sup>c</sup>	South Carolina	2.75	0.15	(17)	0.26
Iowa	1.92 <sup>b</sup>	0.14	(47)	-30.14 <sup>c</sup>	South Dakota	2.48	0.33	(26)	-9.57
Kansas	1.79 <sup>b</sup>	0.15	(50)	-34.66 <sup>c</sup>	Tennessee	2.42	0.12	(31)	-11.73
Kentucky	2.33	0.14	(34)	-15.07 <sup>c</sup>	Texas	2.95	0.07	(12)	7.44
Louisiana	3.22	0.16	(06)	17.49 <sup>c</sup>	Utah	1.98 <sup>b</sup>	0.20	(46)	-27.70 <sup>c</sup>
Maine	2.23	0.22	(39)	-18.59 <sup>c</sup>	Vermont	1.61 <sup>b</sup>	0.29	(51)	-41.25 <sup>c</sup>
Maryland	3.10	0.14	(09)	13.01	Virginia	2.49	0.11	(25)	-9.10
Massachusetts	2.53	0.11	(24)	-7.71	Washington	2.62	0.12	(22)	-4.40
Michigan	2.47	0.09	(28)	-10.02	West Virginia	2.45	0.20	(29)	-10.57
Minnesota	2.14 <sup>b</sup>	0.12	(41)	-21.98 <sup>c</sup>	Wisconsin	2.11 <sup>b</sup>	0.11	(42)	-23.00 <sup>c</sup>
Mississippi	2.89	0.19	(14)	5.40	Wyoming	2.31	0.40	(36)	-15.91 <sup>c</sup>
Missouri	2.18 <sup>b</sup>	0.11	(40)	-20.65 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 25.14  
Cancer of the Testis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.24	0.01		
High Five States				
Rhode Island	0.37	0.12	(01)	53.80 <sup>c</sup>
Idaho	0.35	0.10	(02)	45.29 <sup>c</sup>
North Dakota	0.33	0.14	(03)	36.15 <sup>c</sup>
Nevada	0.33	0.08	(04)	35.84 <sup>c</sup>
Wyoming	0.33	0.17	(05)	35.74 <sup>c</sup>
Low Five States				
Wisconsin	0.15	0.03	(47)	-39.02 <sup>c</sup>
Montana	0.15	0.07	(48)	-39.43 <sup>c</sup>
Hawaii	0.12	0.06	(49)	-52.48 <sup>c</sup>
Delaware	0.11	0.07	(50)	-56.32 <sup>c</sup>
Alaska	0.06	0.06	(51)	-75.66 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.28	0.05	(13)	15.75 <sup>c</sup>	Montana	0.15	0.07	(48)	-39.43 <sup>c</sup>
Alaska	0.06	0.06	(51)	-75.66 <sup>c</sup>	Nebraska	0.28	0.08	(11)	17.46 <sup>c</sup>
Arizona	0.26	0.04	(17)	5.99	Nevada	0.33	0.08	(04)	35.84 <sup>c</sup>
Arkansas	0.20	0.06	(36)	-16.85 <sup>c</sup>	New Hampshire	0.22	0.08	(29)	-7.71
California	0.31 <sup>b</sup>	0.02	(06)	27.70 <sup>c</sup>	New Jersey	0.20	0.03	(35)	-16.71 <sup>c</sup>
Colorado	0.20	0.04	(38)	-18.01 <sup>c</sup>	New Mexico	0.19	0.07	(40)	-22.18 <sup>c</sup>
Connecticut	0.21	0.05	(31)	-12.52	New York	0.21	0.02	(33)	-14.58
Delaware	0.11	0.07	(50)	-56.32 <sup>c</sup>	North Carolina	0.29	0.04	(09)	19.98 <sup>c</sup>
D.C.	0.28	0.14	(15)	15.21 <sup>c</sup>	North Dakota	0.33	0.14	(03)	36.15 <sup>c</sup>
Florida	0.22	0.02	(28)	-7.43	Ohio	0.24	0.03	(24)	-2.21
Georgia	0.16	0.03	(45)	-35.20 <sup>c</sup>	Oklahoma	0.24	0.05	(21)	0.66
Hawaii	0.12	0.06	(49)	-52.48 <sup>c</sup>	Oregon	0.25	0.05	(19)	4.00
Idaho	0.35	0.10	(02)	45.29 <sup>c</sup>	Pennsylvania	0.29	0.03	(10)	19.81 <sup>c</sup>
Illinois	0.22	0.03	(30)	-9.04	Rhode Island	0.37	0.12	(01)	53.80 <sup>c</sup>
Indiana	0.19	0.04	(41)	-22.76 <sup>c</sup>	South Carolina	0.15	0.04	(46)	-37.56 <sup>c</sup>
Iowa	0.19	0.05	(39)	-20.88 <sup>c</sup>	South Dakota	0.21	0.10	(32)	-13.77
Kansas	0.23	0.06	(27)	-6.97	Tennessee	0.26	0.04	(18)	5.78
Kentucky	0.23	0.05	(26)	-4.81	Texas	0.28	0.02	(14)	15.57 <sup>c</sup>
Louisiana	0.24	0.05	(25)	-2.59	Utah	0.20	0.06	(37)	-17.69 <sup>c</sup>
Maine	0.25	0.09	(20)	2.57	Vermont	0.17	0.10	(43)	-28.18 <sup>c</sup>
Maryland	0.26	0.04	(16)	7.58	Virginia	0.16	0.03	(44)	-32.00 <sup>c</sup>
Massachusetts	0.18	0.03	(42)	-25.03 <sup>c</sup>	Washington	0.24	0.04	(23)	-0.69
Michigan	0.24	0.03	(22)	-0.40	West Virginia	0.28	0.08	(12)	16.57 <sup>c</sup>
Minnesota	0.21	0.04	(34)	-14.83	Wisconsin	0.15	0.03	(47)	-39.02 <sup>c</sup>
Mississippi	0.30	0.07	(08)	24.71 <sup>c</sup>	Wyoming	0.33	0.17	(05)	35.74 <sup>c</sup>
Missouri	0.31	0.05	(07)	26.07 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 26.18  
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.48	0.01		
High Five States				
Kansas	0.60	0.07	(01)	26.77 <sup>c</sup>
Idaho	0.58	0.09	(02)	20.95 <sup>c</sup>
Arizona	0.55	0.04	(03)	15.59 <sup>c</sup>
Hawaii	0.55	0.09	(04)	15.06 <sup>c</sup>
New Mexico	0.55	0.08	(05)	14.80
Low Five States				
Nebraska	0.37	0.06	(47)	-22.97 <sup>c</sup>
Alaska	0.36	0.15	(48)	-23.71 <sup>c</sup>
Vermont	0.36	0.11	(49)	-23.99 <sup>c</sup>
South Carolina	0.36	0.04	(50)	-24.32 <sup>c</sup>
Alabama	0.35	0.04	(51)	-25.65 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.35	0.04	(51)	-25.65 <sup>c</sup>	Montana	0.40	0.09	(40)	-15.91 <sup>c</sup>
Alaska	0.36	0.15	(48)	-23.71 <sup>c</sup>	Nebraska	0.37	0.06	(47)	-22.97 <sup>c</sup>
Arizona	0.55	0.04	(03)	15.59 <sup>c</sup>	Nevada	0.43	0.06	(37)	-9.36
Arkansas	0.39	0.05	(44)	-19.07 <sup>c</sup>	New Hampshire	0.50	0.09	(16)	5.01
California	0.54	0.02	(09)	12.25	New Jersey	0.54	0.03	(07)	13.71
Colorado	0.52	0.05	(12)	7.99	New Mexico	0.55	0.08	(05)	14.80
Connecticut	0.50	0.05	(21)	3.99	New York	0.50	0.02	(19)	4.42
Delaware	0.49	0.11	(23)	3.07	North Carolina	0.41	0.03	(39)	-14.50
D.C.	0.39	0.12	(42)	-17.96 <sup>c</sup>	North Dakota	0.50	0.12	(22)	3.94
Florida	0.40 <sup>b</sup>	0.02	(41)	-16.79 <sup>c</sup>	Ohio	0.50	0.03	(20)	4.29
Georgia	0.39	0.03	(43)	-18.39 <sup>c</sup>	Oklahoma	0.47	0.05	(28)	-0.74
Hawaii	0.55	0.09	(04)	15.06 <sup>c</sup>	Oregon	0.43	0.05	(36)	-9.33
Idaho	0.58	0.09	(02)	20.95 <sup>c</sup>	Pennsylvania	0.48	0.03	(26)	0.55
Illinois	0.50	0.03	(18)	4.45	Rhode Island	0.49	0.09	(24)	2.87
Indiana	0.46	0.04	(32)	-4.56	South Carolina	0.36	0.04	(50)	-24.32 <sup>c</sup>
Iowa	0.50	0.05	(15)	5.41	South Dakota	0.44	0.10	(34)	-7.50
Kansas	0.60	0.07	(01)	26.77 <sup>c</sup>	Tennessee	0.45	0.04	(33)	-5.11
Kentucky	0.54	0.05	(06)	13.80	Texas	0.50	0.02	(14)	5.75
Louisiana	0.49	0.05	(25)	2.50	Utah	0.46	0.07	(30)	-2.60
Maine	0.44	0.07	(35)	-8.42	Vermont	0.36	0.11	(49)	-23.99 <sup>c</sup>
Maryland	0.54	0.04	(08)	12.98	Virginia	0.37	0.03	(46)	-21.97 <sup>c</sup>
Massachusetts	0.48	0.04	(27)	0.10	Washington	0.51	0.04	(13)	7.02
Michigan	0.47	0.03	(29)	-1.23	West Virginia	0.38	0.06	(45)	-21.25 <sup>c</sup>
Minnesota	0.50	0.04	(17)	4.46	Wisconsin	0.53	0.04	(11)	10.20
Mississippi	0.43	0.05	(38)	-10.07	Wyoming	0.53	0.14	(10)	10.59
Missouri	0.46	0.04	(31)	-3.65					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 26.19  
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.47	0.01		
High Five States				
D.C.	0.71	0.25	(01)	50.88 <sup>c</sup>
Kansas	0.63	0.10	(02)	33.63 <sup>c</sup>
Utah	0.62	0.12	(03)	33.33 <sup>c</sup>
Idaho	0.59	0.14	(04)	25.87 <sup>c</sup>
Colorado	0.59	0.08	(05)	25.17 <sup>c</sup>
Low Five States				
Arkansas	0.37	0.08	(47)	-20.40 <sup>c</sup>
Alabama	0.37	0.06	(48)	-21.57 <sup>c</sup>
South Carolina	0.37	0.06	(49)	-21.82 <sup>c</sup>
West Virginia	0.34	0.09	(50)	-27.68 <sup>c</sup>
Nebraska	0.29	0.08	(51)	-38.54 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.37	0.06	(48)	-21.57 <sup>c</sup>	Montana	0.40	0.13	(43)	-15.22 <sup>c</sup>
Alaska	0.45	0.29	(34)	-4.07	Nebraska	0.29	0.08	(51)	-38.54 <sup>c</sup>
Arizona	0.51	0.06	(14)	8.77	Nevada	0.42	0.09	(37)	-9.85
Arkansas	0.37	0.08	(47)	-20.40 <sup>c</sup>	New Hampshire	0.58	0.15	(06)	24.42 <sup>c</sup>
California	0.51	0.03	(13)	9.20	New Jersey	0.51	0.05	(15)	8.50
Colorado	0.59	0.08	(05)	25.17 <sup>c</sup>	New Mexico	0.52	0.11	(12)	10.10
Connecticut	0.45	0.07	(32)	-3.55	New York	0.48	0.03	(23)	1.79
Delaware	0.54	0.17	(09)	15.04 <sup>c</sup>	North Carolina	0.46	0.05	(28)	-2.24
D.C.	0.71	0.25	(01)	50.88 <sup>c</sup>	North Dakota	0.47	0.17	(26)	0.93
Florida	0.41	0.03	(38)	-13.10	Ohio	0.52	0.04	(11)	11.27
Georgia	0.40	0.05	(42)	-14.83	Oklahoma	0.46	0.08	(27)	-1.44
Hawaii	0.44	0.12	(36)	-5.52	Oregon	0.50	0.08	(19)	5.89
Idaho	0.59	0.14	(04)	25.87 <sup>c</sup>	Pennsylvania	0.48	0.04	(22)	3.39
Illinois	0.41	0.04	(39)	-13.34	Rhode Island	0.45	0.14	(35)	-4.60
Indiana	0.50	0.06	(18)	7.23	South Carolina	0.37	0.06	(49)	-21.82 <sup>c</sup>
Iowa	0.58	0.09	(07)	23.93 <sup>c</sup>	South Dakota	0.50	0.16	(17)	7.71
Kansas	0.63	0.10	(02)	33.63 <sup>c</sup>	Tennessee	0.40	0.06	(41)	-14.28
Kentucky	0.57	0.08	(08)	20.96 <sup>c</sup>	Texas	0.49	0.04	(20)	4.44
Louisiana	0.38	0.06	(46)	-19.04 <sup>c</sup>	Utah	0.62	0.12	(03)	33.33 <sup>c</sup>
Maine	0.45	0.11	(33)	-4.06	Vermont	0.40	0.16	(44)	-15.24 <sup>c</sup>
Maryland	0.53	0.07	(10)	13.37	Virginia	0.40	0.05	(40)	-14.18
Massachusetts	0.47	0.06	(25)	1.03	Washington	0.45	0.06	(31)	-3.51
Michigan	0.47	0.05	(24)	1.29	West Virginia	0.34	0.09	(50)	-27.68 <sup>c</sup>
Minnesota	0.39	0.06	(45)	-17.60 <sup>c</sup>	Wisconsin	0.45	0.06	(29)	-2.92
Mississippi	0.45	0.09	(30)	-3.38	Wyoming	0.51	0.21	(16)	7.83
Missouri	0.48	0.06	(21)	3.47					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.





Table 26.20  
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.48	0.01		
High Five States				
Hawaii	0.65	0.13	(01)	34.88 <sup>c</sup>
Kansas	0.60	0.09	(02)	25.77 <sup>c</sup>
Arizona	0.59	0.06	(03)	23.31 <sup>c</sup>
New Mexico	0.57	0.10	(04)	19.02 <sup>c</sup>
Wisconsin	0.57	0.06	(05)	18.65 <sup>c</sup>
Low Five States				
Virginia	0.35	0.04	(47)	-27.20 <sup>c</sup>
Alabama	0.33	0.05	(48)	-31.64 <sup>c</sup>
Utah	0.32	0.08	(49)	-33.76 <sup>c</sup>
Vermont	0.31	0.13	(50)	-34.41 <sup>c</sup>
D.C.	0.14 <sup>b</sup>	0.08	(51)	-69.77 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.33	0.05	(48)	-31.64 <sup>c</sup>	Montana	0.39	0.11	(38)	-18.37 <sup>c</sup>
Alaska	0.36	0.19	(44)	-24.72 <sup>c</sup>	Nebraska	0.40	0.09	(36)	-15.67 <sup>c</sup>
Arizona	0.59	0.06	(03)	23.31 <sup>c</sup>	Nevada	0.45	0.09	(30)	-6.73
Arkansas	0.38	0.06	(43)	-21.66 <sup>c</sup>	New Hampshire	0.47	0.11	(26)	-2.42
California	0.55	0.02	(10)	15.81 <sup>c</sup>	New Jersey	0.57	0.05	(07)	18.05 <sup>c</sup>
Colorado	0.45	0.06	(29)	-6.11	New Mexico	0.57	0.10	(04)	19.02 <sup>c</sup>
Connecticut	0.52	0.07	(15)	9.38	New York	0.51	0.03	(19)	6.32
Delaware	0.47	0.14	(27)	-2.55	North Carolina	0.36	0.04	(45)	-24.81 <sup>c</sup>
D.C.	0.14 <sup>b</sup>	0.08	(51)	-69.77 <sup>c</sup>	North Dakota	0.49	0.17	(20)	3.24
Florida	0.39 <sup>b</sup>	0.02	(41)	-19.46 <sup>c</sup>	Ohio	0.47	0.04	(25)	-2.21
Georgia	0.39	0.04	(40)	-19.36 <sup>c</sup>	Oklahoma	0.48	0.07	(21)	1.07
Hawaii	0.65	0.13	(01)	34.88 <sup>c</sup>	Oregon	0.39	0.06	(37)	-17.79 <sup>c</sup>
Idaho	0.55	0.12	(12)	15.07 <sup>c</sup>	Pennsylvania	0.47	0.03	(24)	-1.53
Illinois	0.57	0.04	(06)	18.60 <sup>c</sup>	Rhode Island	0.51	0.11	(18)	6.38
Indiana	0.41	0.05	(34)	-14.19	South Carolina	0.36	0.05	(46)	-24.86 <sup>c</sup>
Iowa	0.44	0.07	(31)	-8.12	South Dakota	0.38	0.12	(42)	-20.28 <sup>c</sup>
Kansas	0.60	0.09	(02)	25.77 <sup>c</sup>	Tennessee	0.48	0.05	(22)	-0.35
Kentucky	0.52	0.06	(16)	8.61	Texas	0.51	0.03	(17)	6.80
Louisiana	0.56	0.07	(09)	16.71 <sup>c</sup>	Utah	0.32	0.08	(49)	-33.76 <sup>c</sup>
Maine	0.41	0.09	(35)	-14.87	Vermont	0.31	0.13	(50)	-34.41 <sup>c</sup>
Maryland	0.53	0.06	(14)	10.40	Virginia	0.35	0.04	(47)	-27.20 <sup>c</sup>
Massachusetts	0.48	0.05	(23)	-0.38	Washington	0.55	0.06	(13)	14.56
Michigan	0.46	0.04	(28)	-4.75	West Virginia	0.39	0.08	(39)	-18.41 <sup>c</sup>
Minnesota	0.55	0.06	(11)	15.55 <sup>c</sup>	Wisconsin	0.57	0.06	(05)	18.65 <sup>c</sup>
Mississippi	0.41	0.07	(33)	-13.43	Wyoming	0.56	0.20	(08)	16.81 <sup>c</sup>
Missouri	0.44	0.05	(32)	-9.07					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 27.18  
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.34	0.02		
High Five States				
Maine	5.96 <sup>b</sup>	0.28	(01)	37.38 <sup>c</sup>
Delaware	5.56	0.35	(02)	28.34 <sup>c</sup>
Vermont	5.44	0.40	(03)	25.53 <sup>c</sup>
Rhode Island	5.31	0.29	(04)	22.37 <sup>c</sup>
Nevada	5.25 <sup>b</sup>	0.23	(05)	21.21 <sup>c</sup>
Low Five States				
Alabama	3.76 <sup>b</sup>	0.12	(47)	-13.29
New Mexico	3.51 <sup>b</sup>	0.19	(48)	-19.15 <sup>c</sup>
Mississippi	3.49 <sup>b</sup>	0.16	(49)	-19.39 <sup>c</sup>
Utah	3.38 <sup>b</sup>	0.19	(50)	-21.94 <sup>c</sup>
Hawaii	2.67 <sup>b</sup>	0.19	(51)	-38.40 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.76 <sup>b</sup>	0.12	(47)	-13.29	Montana	4.49	0.29	(20)	3.47
Alaska	4.34	0.52	(24)	0.00	Nebraska	3.86	0.20	(44)	-11.00
Arizona	3.99	0.11	(39)	-7.89	Nevada	5.25 <sup>b</sup>	0.23	(05)	21.21 <sup>c</sup>
Arkansas	3.89	0.16	(41)	-10.24	New Hampshire	4.94	0.27	(07)	13.94
California	4.12 <sup>b</sup>	0.05	(33)	-4.90	New Jersey	4.88 <sup>b</sup>	0.10	(10)	12.60
Colorado	3.99	0.14	(40)	-7.93	New Mexico	3.51 <sup>b</sup>	0.19	(48)	-19.15 <sup>c</sup>
Connecticut	4.90 <sup>b</sup>	0.16	(09)	12.94	New York	4.32	0.06	(26)	-0.28
Delaware	5.56	0.35	(02)	28.34 <sup>c</sup>	North Carolina	4.10	0.10	(35)	-5.48
D.C.	4.86	0.41	(11)	12.00	North Dakota	4.36	0.33	(23)	0.54
Florida	4.33	0.06	(25)	-0.10	Ohio	5.02 <sup>b</sup>	0.09	(06)	15.73 <sup>c</sup>
Georgia	3.82 <sup>b</sup>	0.10	(46)	-11.99	Oklahoma	4.11	0.15	(34)	-5.21
Hawaii	2.67 <sup>b</sup>	0.19	(51)	-38.40 <sup>c</sup>	Oregon	4.79	0.16	(13)	10.50
Idaho	4.24	0.25	(29)	-2.24	Pennsylvania	4.73 <sup>b</sup>	0.08	(14)	9.14
Illinois	4.46	0.08	(21)	2.86	Rhode Island	5.31	0.29	(04)	22.37 <sup>c</sup>
Indiana	4.45	0.12	(22)	2.61	South Carolina	3.88	0.13	(43)	-10.54
Iowa	4.21	0.15	(31)	-2.93	South Dakota	4.21	0.30	(30)	-2.81
Kansas	3.88	0.16	(42)	-10.41	Tennessee	4.13	0.12	(32)	-4.72
Kentucky	4.51	0.15	(18)	4.01	Texas	3.82 <sup>b</sup>	0.06	(45)	-11.89
Louisiana	4.06	0.14	(37)	-6.29	Utah	3.38 <sup>b</sup>	0.19	(50)	-21.94 <sup>c</sup>
Maine	5.96 <sup>b</sup>	0.28	(01)	37.38 <sup>c</sup>	Vermont	5.44	0.40	(03)	25.53 <sup>c</sup>
Maryland	4.49	0.13	(19)	3.51	Virginia	4.25	0.11	(28)	-2.02
Massachusetts	4.94 <sup>b</sup>	0.12	(08)	13.90	Washington	4.70	0.12	(15)	8.34
Michigan	4.67 <sup>b</sup>	0.10	(16)	7.66	West Virginia	4.66	0.20	(17)	7.52
Minnesota	4.03	0.12	(38)	-7.03	Wisconsin	4.81 <sup>b</sup>	0.13	(12)	10.92
Mississippi	3.49 <sup>b</sup>	0.16	(49)	-19.39 <sup>c</sup>	Wyoming	4.06	0.40	(36)	-6.29
Missouri	4.27	0.12	(27)	-1.51					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.



Table 27.19  
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.54	0.04		
High Five States				
Maine	9.98 <sup>b</sup>	0.57	(01)	32.40 <sup>c</sup>
Rhode Island	9.94 <sup>b</sup>	0.65	(02)	31.88 <sup>c</sup>
Vermont	9.81	0.85	(03)	30.14 <sup>c</sup>
Delaware	9.79	0.74	(04)	29.96 <sup>c</sup>
Ohio	9.11 <sup>b</sup>	0.20	(05)	20.93 <sup>c</sup>
Low Five States				
Wyoming	6.48	0.78	(47)	-13.99
Arkansas	6.36 <sup>b</sup>	0.32	(48)	-15.67 <sup>c</sup>
Utah	6.33	0.41	(49)	-16.05 <sup>c</sup>
New Mexico	5.55 <sup>b</sup>	0.37	(50)	-26.36 <sup>c</sup>
Hawaii	4.32 <sup>b</sup>	0.37	(51)	-42.66 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.70	0.27	(42)	-11.13	Montana	7.72	0.59	(22)	2.45
Alaska	8.07	1.11	(16)	7.12	Nebraska	6.68	0.42	(44)	-11.40
Arizona	6.70 <sup>b</sup>	0.22	(41)	-11.08	Nevada	8.55	0.44	(09)	13.42
Arkansas	6.36 <sup>b</sup>	0.32	(48)	-15.67 <sup>c</sup>	New Hampshire	8.72	0.58	(07)	15.75 <sup>c</sup>
California	7.07 <sup>b</sup>	0.10	(38)	-6.19	New Jersey	8.53 <sup>b</sup>	0.22	(10)	13.24
Colorado	7.07	0.31	(37)	-6.14	New Mexico	5.55 <sup>b</sup>	0.37	(50)	-26.36 <sup>c</sup>
Connecticut	8.57	0.33	(08)	13.74	New York	7.55	0.14	(27)	0.13
Delaware	9.79	0.74	(04)	29.96 <sup>c</sup>	North Carolina	7.08	0.21	(36)	-6.12
D.C.	7.37	0.84	(30)	-2.25	North Dakota	7.94	0.70	(18)	5.33
Florida	7.23	0.12	(32)	-4.00	Ohio	9.11 <sup>b</sup>	0.20	(05)	20.93 <sup>c</sup>
Georgia	6.87	0.23	(39)	-8.78	Oklahoma	7.09	0.31	(35)	-5.97
Hawaii	4.32 <sup>b</sup>	0.37	(51)	-42.66 <sup>c</sup>	Oregon	8.13	0.32	(15)	7.90
Idaho	7.41	0.51	(29)	-1.63	Pennsylvania	8.36 <sup>b</sup>	0.17	(11)	10.87
Illinois	7.72	0.18	(23)	2.43	Rhode Island	9.94 <sup>b</sup>	0.65	(02)	31.88 <sup>c</sup>
Indiana	7.72	0.25	(21)	2.48	South Carolina	6.68	0.29	(43)	-11.37
Iowa	7.64	0.33	(25)	1.41	South Dakota	8.20	0.65	(14)	8.87
Kansas	7.12	0.35	(33)	-5.48	Tennessee	7.28	0.25	(31)	-3.47
Kentucky	7.82	0.31	(19)	3.77	Texas	6.57 <sup>b</sup>	0.14	(45)	-12.82
Louisiana	6.85	0.29	(40)	-9.09	Utah	6.33	0.41	(49)	-16.05 <sup>c</sup>
Maine	9.98 <sup>b</sup>	0.57	(01)	32.40 <sup>c</sup>	Vermont	9.81	0.85	(03)	30.14 <sup>c</sup>
Maryland	7.69	0.28	(24)	2.02	Virginia	7.56	0.24	(26)	0.28
Massachusetts	8.96 <sup>b</sup>	0.26	(06)	18.93 <sup>c</sup>	Washington	8.06	0.26	(17)	6.88
Michigan	8.30 <sup>b</sup>	0.20	(13)	10.16	West Virginia	7.54	0.42	(28)	0.04
Minnesota	7.12	0.26	(34)	-5.56	Wisconsin	8.32	0.27	(12)	10.35
Mississippi	6.51	0.35	(46)	-13.61	Wyoming	6.48	0.78	(47)	-13.99
Missouri	7.77	0.25	(20)	3.14					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.


Table 27.20  
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2003-2007

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.21	0.02		
High Five States				
D.C.	3.28	0.42	(01)	48.59 <sup>c</sup>
Maine	3.15 <sup>b</sup>	0.26	(02)	42.69 <sup>c</sup>
West Virginia	2.70	0.20	(03)	22.17 <sup>c</sup>
Connecticut	2.61	0.15	(04)	17.90 <sup>c</sup>
New Jersey	2.57 <sup>b</sup>	0.10	(05)	16.48 <sup>c</sup>
Low Five States				
Mississippi	1.67 <sup>b</sup>	0.14	(47)	-24.49 <sup>c</sup>
South Dakota	1.47	0.24	(48)	-33.28 <sup>c</sup>
Hawaii	1.44 <sup>b</sup>	0.19	(49)	-35.05 <sup>c</sup>
Alaska	1.42	0.39	(50)	-35.66 <sup>c</sup>
Utah	1.13 <sup>b</sup>	0.15	(51)	-48.69 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.90	0.11	(43)	-13.98	Montana	2.14	0.26	(27)	-2.97
Alaska	1.42	0.39	(50)	-35.66 <sup>c</sup>	Nebraska	2.06	0.19	(34)	-6.68
Arizona	1.92	0.11	(40)	-12.99	Nevada	2.57	0.21	(06)	16.38 <sup>c</sup>
Arkansas	2.22	0.15	(26)	0.54	New Hampshire	2.41	0.25	(15)	8.96
California	2.09	0.05	(32)	-5.22	New Jersey	2.57 <sup>b</sup>	0.10	(05)	16.48 <sup>c</sup>
Colorado	1.91	0.13	(42)	-13.59	New Mexico	1.92	0.19	(41)	-13.01
Connecticut	2.61	0.15	(04)	17.90 <sup>c</sup>	New York	2.29	0.06	(21)	3.65
Delaware	2.49	0.31	(09)	12.87	North Carolina	2.25	0.09	(24)	1.94
D.C.	3.28	0.42	(01)	48.59 <sup>c</sup>	North Dakota	1.81	0.28	(45)	-18.18 <sup>c</sup>
Florida	2.12	0.06	(30)	-3.92	Ohio	2.44	0.08	(12)	10.41
Georgia	1.95	0.09	(39)	-11.89	Oklahoma	2.10	0.14	(31)	-4.85
Hawaii	1.44 <sup>b</sup>	0.19	(49)	-35.05 <sup>c</sup>	Oregon	2.40	0.15	(17)	8.45
Idaho	1.81	0.22	(46)	-18.24 <sup>c</sup>	Pennsylvania	2.44	0.07	(13)	10.33
Illinois	2.42	0.08	(14)	9.31	Rhode Island	2.52	0.26	(08)	14.11
Indiana	2.35	0.11	(20)	6.23	South Carolina	2.03	0.12	(35)	-8.14
Iowa	2.07	0.14	(33)	-6.27	South Dakota	1.47	0.24	(48)	-33.28 <sup>c</sup>
Kansas	1.81	0.14	(44)	-18.06 <sup>c</sup>	Tennessee	2.12	0.11	(29)	-3.89
Kentucky	2.39	0.14	(18)	8.24	Texas	1.99 <sup>b</sup>	0.06	(36)	-10.04
Louisiana	2.27	0.13	(23)	2.77	Utah	1.13 <sup>b</sup>	0.15	(51)	-48.69 <sup>c</sup>
Maine	3.15 <sup>b</sup>	0.26	(02)	42.69 <sup>c</sup>	Vermont	2.40	0.34	(16)	8.49
Maryland	2.48	0.13	(10)	12.13	Virginia	2.14	0.10	(28)	-3.22
Massachusetts	2.47	0.11	(11)	11.63	Washington	2.35	0.12	(19)	6.38
Michigan	2.27	0.09	(22)	2.92	West Virginia	2.70	0.20	(03)	22.17 <sup>c</sup>
Minnesota	1.98	0.12	(37)	-10.21	Wisconsin	2.56	0.12	(07)	15.89 <sup>c</sup>
Mississippi	1.67 <sup>b</sup>	0.14	(47)	-24.49 <sup>c</sup>	Wyoming	2.24	0.40	(25)	1.44
Missouri	1.98	0.10	(38)	-10.61					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.