

Table 33.1
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2005-2011
Both Sexes by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	1,791,391	67.6	0.0	236,943	62.3	0.1	141,518	66.2	0.1	9,229	59.7	0.6	221,531	67.8	0.1
Oral Cavity & Pharynx	42,526	68.7	0.3	4,844	50.1	0.9	3,584	69.4	0.9	238	65.6	3.5	4,068	64.5	0.9
Esophagus	17,067	21.0	0.4	2,251	15.4	1.0	930	19.3	1.7	98	21.2	4.7	1,517	19.5	1.3
Stomach	24,480	29.7	0.3	4,878	31.2	0.8	4,987	39.1	0.8	264	21.0	3.1	6,233	29.5	0.7
Colon & Rectum	163,859	65.7	0.1	25,087	59.8	0.4	16,772	68.9	0.4	1,103	61.1	1.7	21,016	65.5	0.4
Liver & Intrahepatic Bile Duct	27,309	19.9	0.3	4,940	15.8	0.7	6,562	27.1	0.7	379	18.2	2.5	7,371	20.7	0.6
Pancreas	44,919	7.5	0.2	6,865	7.6	0.4	3,939	9.6	0.6	284	5.7	1.7	5,676	8.7	0.5
Larynx	13,066	67.3	0.5	2,422	59.6	1.2	509	70.4	2.4	59	46.7	9.2	1,307	66.0	1.6
Lung & Bronchus	217,451	20.0	0.1	29,769	16.6	0.3	15,371	20.9	0.4	976	14.5	1.4	14,800	19.1	0.4
Melanoma of the Skin	89,524	89.6	0.1	466	69.8	2.5	598	73.3	2.2	120	87.5	3.3	3,002	83.8	0.8
Breast	262,853	88.8	0.1	35,276	79.9	0.3	25,301	91.5	0.2	1,334	85.3	1.2	33,551	87.8	0.2
Cervix uteri	16,579	71.5	0.4	3,016	61.2	1.0	1,956	71.6	1.2	157	64.1	4.4	5,004	74.9	0.7
Corpus & Uterus, NOS	55,612	83.4	0.2	6,628	64.0	0.7	5,206	84.0	0.6	350	83.3	2.4	7,377	81.9	0.5
Ovary ^d	27,916	45.9	0.4	2,816	37.7	1.1	2,586	56.6	1.2	168	43.7	4.7	4,084	53.6	1.0
Prostate	283,819	94.1	0.1	53,401	92.7	0.1	17,046	94.3	0.2	938	89.2	1.2	32,497	93.2	0.2
Testis	13,947	95.7	0.2	452	92.1	1.5	568	94.5	1.0	116	93.7	2.6	3,394	92.8	0.5
Urinary Bladder	77,368	78.9	0.2	4,720	67.8	0.8	3,394	78.7	0.8	179	69.2	3.8	5,205	78.1	0.7
Kidney & Renal Pelvis	57,651	74.7	0.2	7,799	75.9	0.6	3,513	73.5	0.9	568	69.5	2.2	9,269	75.8	0.5
Brain & Nervous System	28,372	33.8	0.3	2,306	42.1	1.2	1,764	42.9	1.4	137	47.9	4.7	4,408	46.9	0.9
Thyroid	51,551	97.1	0.1	4,191	96.3	0.3	6,418	96.4	0.3	267	96.1	1.2	9,453	97.2	0.2
Hodgkin Lymphoma	12,141	88.1	0.3	1,845	85.3	1.0	738	88.5	1.3	42	97.3	2.7	2,427	86.0	0.8
Non-Hodgkin Lymphoma	77,320	71.7	0.2	7,520	66.0	0.6	5,963	69.3	0.7	352	69.8	2.8	11,011	69.5	0.5
Myeloma	21,228	50.7	0.4	5,710	54.3	0.9	1,631	51.7	1.6	122	66.6	4.9	3,229	50.3	1.2
Leukemia	50,832	62.3	0.2	4,747	57.7	0.8	3,404	55.4	1.0	248	63.1	3.4	8,382	63.3	0.6
Mesothelioma	3,969	10.0	0.6	233	15.5	3.2	132	3.4	2.2	-	-	-	448	15.9	2.2

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2005-2011 with follow-up through 2012.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 33.2
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2005-2011
Males by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	929,636	67.5	0.1	126,321	64.5	0.2	65,916	61.9	0.2	4,433	56.9	0.8	109,251	66.0	0.2
Oral Cavity & Pharynx	30,395	68.4	0.3	3,329	45.6	1.0	2,342	66.5	1.2	169	66.3	4.0	2,771	60.7	1.1
Esophagus	13,694	21.1	0.4	1,547	13.6	1.1	708	18.5	1.9	67	20.5	5.6	1,218	19.2	1.4
Stomach	15,242	27.8	0.4	2,686	28.5	1.1	2,791	39.8	1.1	153	17.1	4.0	3,580	28.6	0.9
Colon & Rectum	84,792	66.1	0.2	11,984	58.1	0.5	8,667	69.7	0.6	583	61.5	2.4	11,349	65.1	0.5
Liver & Intrahepatic Bile Duct	20,026	19.8	0.4	3,661	14.6	0.8	4,579	27.4	0.8	267	19.9	3.0	5,373	20.2	0.7
Pancreas	22,704	7.3	0.2	3,151	7.4	0.6	1,909	9.7	0.9	146	-	-	2,800	7.8	0.7
Larynx	10,551	67.8	0.5	1,936	59.2	1.4	444	68.5	2.6	47	57.5	9.6	1,132	66.1	1.7
Lung & Bronchus	113,654	17.4	0.1	16,629	14.1	0.3	8,799	18.0	0.5	504	13.3	1.9	8,057	15.0	0.5
Melanoma of the Skin	50,347	87.3	0.2	198	62.8	4.0	305	67.7	3.4	52	81.7	6.0	1,270	78.5	1.4
Prostate	283,819	94.1	0.1	53,401	92.7	0.1	17,046	94.3	0.2	938	89.2	1.2	32,497	93.2	0.2
Testis	13,947	95.7	0.2	452	92.1	1.5	568	94.5	1.0	116	93.7	2.6	3,394	92.8	0.5
Urinary Bladder	58,729	80.1	0.2	3,117	73.1	0.9	2,556	79.6	1.0	144	71.8	4.3	3,866	80.1	0.8
Kidney & Renal Pelvis	35,848	74.2	0.3	4,631	74.2	0.8	2,209	73.3	1.1	336	68.9	2.9	5,509	74.1	0.7
Brain & Nervous System	15,920	33.0	0.4	1,174	38.7	1.6	958	41.0	1.9	74	45.2	6.2	2,443	46.1	1.2
Thyroid	12,171	94.6	0.2	741	94.1	1.0	1,271	94.1	0.7	56	92.7	3.5	1,692	94.4	0.7
Hodgkin Lymphoma	6,656	87.6	0.4	985	82.0	1.4	399	88.2	1.8	-	-	-	1,327	83.8	1.2
Non-Hodgkin Lymphoma	41,870	70.7	0.3	4,054	62.8	0.9	3,200	68.4	0.9	193	69.6	3.8	5,920	66.6	0.7
Myeloma	11,902	52.4	0.6	2,750	54.2	1.3	870	49.2	2.3	64	63.2	6.7	1,714	51.1	1.7
Leukemia	29,287	63.1	0.3	2,537	59.2	1.1	1,896	57.1	1.3	138	62.6	4.5	4,705	64.6	0.8
Mesothelioma	3,030	7.9	0.6	160	12.3	3.5	95	2.7	2.5	-	-	-	320	12.3	2.4

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2005-2011 with follow-up through 2012.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 33.3
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2005-2011
Females by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	861,755	67.8	0.1	110,622	59.8	0.2	75,602	70.0	0.2	4,796	62.3	0.8	112,280	69.6	0.2
Oral Cavity & Pharynx	12,131	69.4	0.5	1,515	59.7	1.5	1,242	74.5	1.4	69	64.4	6.8	1,297	72.4	1.4
Esophagus	3,373	20.6	0.8	704	19.7	1.9	222	21.8	3.7	31	23.5	8.3	299	20.7	2.9
Stomach	9,238	32.7	0.6	2,192	34.6	1.2	2,196	38.2	1.2	111	26.2	4.7	2,653	30.6	1.1
Colon & Rectum	79,067	65.3	0.2	13,103	61.2	0.5	8,105	68.0	0.6	520	60.7	2.5	9,667	65.9	0.6
Liver & Intrahepatic Bile Duct	7,283	20.0	0.6	1,279	19.4	1.5	1,983	26.3	1.2	112	13.1	4.7	1,998	22.1	1.2
Pancreas	22,215	7.7	0.2	3,714	7.7	0.6	2,030	9.6	0.8	138	5.0	2.3	2,876	9.5	0.7
Larynx	2,515	65.1	1.1	486	60.9	2.6	65	85.2	5.0	-	-	-	175	65.4	4.5
Lung & Bronchus	103,797	22.8	0.2	13,140	19.8	0.4	6,572	24.7	0.7	472	15.8	2.1	6,743	23.9	0.6
Melanoma of the Skin	39,177	92.4	0.2	268	75.0	3.2	293	79.0	2.9	68	92.2	3.4	1,732	87.7	0.9
Breast	260,879	88.9	0.1	34,961	80.0	0.3	25,163	91.5	0.2	1,329	85.2	1.2	33,376	87.8	0.2
Cervix uteri	16,579	71.5	0.4	3,016	61.2	1.0	1,956	71.6	1.2	157	64.1	4.4	5,004	74.9	0.7
Corpus & Uterus, NOS	55,612	83.4	0.2	6,628	64.0	0.7	5,206	84.0	0.6	350	83.3	2.4	7,377	81.9	0.5
Ovary ^d	27,916	45.9	0.4	2,816	37.7	1.1	2,586	56.6	1.2	168	43.7	4.7	4,084	53.6	1.0
Urinary Bladder	18,639	75.1	0.4	1,603	57.6	1.5	838	75.8	1.7	35	58.9	8.6	1,339	72.5	1.4
Kidney & Renal Pelvis	21,803	75.6	0.3	3,168	78.4	0.8	1,304	73.8	1.4	232	70.4	3.4	3,760	78.2	0.8
Brain & Nervous System	12,452	34.8	0.5	1,132	45.6	1.7	806	45.3	2.1	63	50.4	7.2	1,965	47.9	1.3
Thyroid	39,380	97.9	0.1	3,450	96.7	0.3	5,147	96.9	0.3	211	97.0	1.2	7,761	97.8	0.2
Hodgkin Lymphoma	5,485	88.6	0.5	860	89.0	1.2	339	89.0	1.8	-	-	-	1,100	88.5	1.1
Non-Hodgkin Lymphoma	35,450	72.7	0.3	3,466	69.6	0.9	2,763	70.3	1.0	159	70.2	4.0	5,091	72.8	0.7
Myeloma	9,326	48.5	0.7	2,960	54.5	1.2	761	54.5	2.4	58	70.6	7.1	1,515	49.5	1.7
Leukemia	21,545	61.2	0.4	2,210	55.9	1.3	1,508	53.4	1.5	110	63.9	5.2	3,677	61.7	0.9
Mesothelioma	939	16.6	1.5	73	26.6	5.8	37	5.2	4.9	-	-	-	128	24.6	4.8

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2005-2011 with follow-up through 2012.

^b Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 33.4

Cancer of the Female Breast

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2005-2011

Race/Ethnicity	All Ages				Ages <50				Ages 50+			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	324,960	100.0	88.1	0.1	79,205	100.0	89.8	0.1	245,755	100.0	87.6	0.1
White	260,879	80.3	88.9	0.1	59,292	74.9	90.9	0.1	201,587	82.0	88.2	0.1
White Non-Hispanic	228,800	70.4	89.0	0.1	47,719	60.2	91.6	0.2	181,081	73.7	88.3	0.1
White Hispanic	32,079	9.9	87.9	0.2	11,573	14.6	88.2	0.4	20,506	8.3	87.7	0.3
Black	34,961	10.8	80.0	0.3	10,558	13.3	81.3	0.5	24,403	9.9	79.4	0.3
American Indian/Alaska Native ^c	1,329	0.4	85.2	1.2	415	0.5	85.8	2.0	914	0.4	84.9	1.5
Asian	23,247	7.2	91.8	0.2	7,657	9.7	93.1	0.4	15,590	6.3	91.2	0.3
Asian Indian, Pakistani	2,145	0.7	90.6	0.8	835	1.1	91.3	1.3	1,310	0.5	90.2	1.0
Chinese	4,558	1.4	92.9	0.5	1,563	2.0	93.6	0.8	2,995	1.2	92.5	0.6
Filipino	6,380	2.0	90.1	0.5	1,819	2.3	91.5	0.8	4,561	1.9	89.6	0.6
Japanese	3,236	1.0	93.2	0.5	740	0.9	95.6	0.9	2,496	1.0	92.4	0.6
Korean	1,714	0.5	91.9	0.8	682	0.9	94.7	1.0	1,032	0.4	89.9	1.2
Vietnamese	1,501	0.5	91.3	0.9	632	0.8	93.5	1.2	869	0.4	89.6	1.3
Other Asian ^d	3,713	1.1	93.3	0.5	1,386	1.7	93.5	0.8	2,327	0.9	93.2	0.7
Pacific Islander	1,916	0.6	87.0	1.0	547	0.7	83.8	2.0	1,369	0.6	88.4	1.1
Hawaiian	1,216	0.4	89.7	1.1	327	0.4	86.9	2.3	889	0.4	90.8	1.2
Other Pacific Islander ^e	700	0.2	82.0	1.9	220	0.3	78.9	3.5	480	0.2	83.7	2.2
Hispanic ^f	33,376	10.3	87.8	0.2	12,049	15.2	88.1	0.4	21,327	8.7	87.7	0.3

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2005-2011 with follow-up through 2012.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorro, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.5

Cancer of the Colon and Rectum

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2005-2011

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	208,809	100.0	65.4	0.1	107,135	100.0	65.7	0.2	101,674	100.0	65.1	0.2
White	163,859	78.5	65.7	0.1	84,792	79.1	66.1	0.2	79,067	77.8	65.3	0.2
White Non-Hispanic	143,622	68.8	65.7	0.1	73,829	68.9	66.3	0.2	69,793	68.6	65.2	0.2
White Hispanic	20,237	9.7	65.4	0.4	10,963	10.2	65.0	0.6	9,274	9.1	66.0	0.6
Black	25,087	12.0	59.8	0.4	11,984	11.2	58.1	0.5	13,103	12.9	61.2	0.5
American Indian/Alaska Native ^c	1,103	0.5	61.1	1.7	583	0.5	61.5	2.4	520	0.5	60.7	2.5
Asian	15,760	7.5	69.4	0.4	8,090	7.6	70.3	0.6	7,670	7.5	68.4	0.6
Asian Indian, Pakistani	869	0.4	69.8	2.0	499	0.5	72.1	2.5	370	0.4	66.9	3.1
Chinese	3,592	1.7	69.8	0.9	1,797	1.7	71.7	1.3	1,795	1.8	67.8	1.3
Filipino	3,439	1.6	68.4	1.0	1,772	1.7	68.5	1.3	1,667	1.6	68.4	1.4
Japanese	2,847	1.4	69.7	1.0	1,456	1.4	70.1	1.4	1,391	1.4	69.2	1.4
Korean	1,531	0.7	67.4	1.4	738	0.7	68.4	2.0	793	0.8	66.5	2.0
Vietnamese	1,367	0.7	67.4	1.5	754	0.7	67.5	2.1	613	0.6	67.2	2.2
Other Asian ^d	2,115	1.0	72.5	1.2	1,074	1.0	74.0	1.6	1,041	1.0	71.0	1.7
Pacific Islander	1,012	0.5	60.6	1.9	577	0.5	59.8	2.6	435	0.4	61.5	2.8
Hawaiian	675	0.3	60.8	2.4	397	0.4	58.8	3.3	278	0.3	63.4	3.4
Other Pacific Islander ^e	337	0.2	60.3	3.2	180	0.2	62.0	4.2	157	0.2	58.0	4.9
Hispanic ^f	21,016	10.1	65.5	0.4	11,349	10.6	65.1	0.5	9,667	9.5	65.9	0.6

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2005-2011 with follow-up through 2012.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorroan, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.6

Cancer of the Liver and Intrahepatic Bile Duct

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2005-2011

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	39,459	100.0	20.7	0.3	28,726	100.0	20.5	0.3	10,733	100.0	21.2	0.5
White	27,309	69.2	19.9	0.3	20,026	69.7	19.8	0.4	7,283	67.9	20.0	0.6
White Non-Hispanic	20,164	51.1	19.7	0.4	14,819	51.6	19.7	0.4	5,345	49.8	19.5	0.7
White Hispanic	7,145	18.1	20.5	0.6	5,207	18.1	20.1	0.7	1,938	18.1	21.5	1.2
Black	4,940	12.5	15.8	0.7	3,661	12.7	14.6	0.8	1,279	11.9	19.4	1.5
American Indian/Alaska Native ^c	379	1.0	18.2	2.5	267	0.9	19.9	3.0	112	1.0	13.1	4.7
Asian	6,204	15.7	27.6	0.7	4,295	15.0	28.1	0.8	1,909	17.8	26.6	1.2
Asian Indian, Pakistani	237	0.6	29.8	4.2	168	0.6	34.0	4.5	69	0.6	19.0	8.8
Chinese	1,513	3.8	33.1	1.5	1,073	3.7	34.3	1.8	440	4.1	30.2	2.6
Filipino	955	2.4	22.9	1.7	659	2.3	21.3	1.9	296	2.8	26.9	3.2
Japanese	552	1.4	25.7	2.3	277	1.0	31.1	3.3	275	2.6	21.2	3.0
Korean	720	1.8	28.3	1.9	487	1.7	29.1	2.4	233	2.2	26.7	3.2
Vietnamese	1,321	3.3	28.7	1.6	1,008	3.5	28.0	1.8	313	2.9	31.8	3.3
Other Asian ^d	906	2.3	20.9	1.8	623	2.2	20.9	2.2	283	2.6	21.4	3.3
Pacific Islander	358	0.9	16.6	2.8	284	1.0	16.0	3.1	74	0.7	18.6	6.1
Hawaiian	186	0.5	13.4	3.4	146	0.5	14.5	4.1	40	0.4	9.2	5.8
Other Pacific Islander ^e	172	0.4	22.2	4.2	138	0.5	18.6	4.5	34	0.3	39.3	9.8
Hispanic ^f	7,371	18.7	20.7	0.6	5,373	18.7	20.2	0.7	1,998	18.6	22.1	1.2

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2005-2011 with follow-up through 2012.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorran, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.7

Cancer of the Lung and Bronchus

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2005-2011

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	264,559	100.0	19.7	0.1	140,132	100.0	17.0	0.1	124,427	100.0	22.6	0.1
White	217,451	82.2	20.0	0.1	113,654	81.1	17.4	0.1	103,797	83.4	22.8	0.2
White Non-Hispanic	203,195	76.8	20.0	0.1	105,881	75.6	17.5	0.1	97,314	78.2	22.7	0.2
White Hispanic	14,256	5.4	19.2	0.4	7,773	5.5	15.0	0.5	6,483	5.2	24.1	0.7
Black	29,769	11.3	16.6	0.3	16,629	11.9	14.1	0.3	13,140	10.6	19.8	0.4
American Indian/Alaska Native ^c	976	0.4	14.5	1.4	504	0.4	13.3	1.9	472	0.4	15.8	2.1
Asian	14,054	5.3	21.2	0.4	8,037	5.7	18.4	0.5	6,017	4.8	24.9	0.7
Asian Indian, Pakistani	641	0.2	26.7	2.3	432	0.3	23.9	2.6	209	0.2	32.6	4.5
Chinese	3,593	1.4	21.5	0.9	2,013	1.4	19.2	1.1	1,580	1.3	24.3	1.4
Filipino	3,571	1.3	22.5	0.9	2,102	1.5	18.5	1.1	1,469	1.2	28.2	1.5
Japanese	2,055	0.8	17.9	1.0	1,048	0.7	15.0	1.4	1,007	0.8	20.7	1.6
Korean	1,122	0.4	19.0	1.4	609	0.4	16.3	1.8	513	0.4	22.2	2.4
Vietnamese	1,620	0.6	21.2	1.3	1,064	0.8	18.9	1.5	556	0.4	25.8	2.4
Other Asian ^d	1,452	0.5	22.1	1.4	769	0.5	19.4	1.9	683	0.5	25.2	2.2
Pacific Islander	1,317	0.5	17.5	1.3	762	0.5	13.8	1.6	555	0.4	22.5	2.2
Hawaiian	870	0.3	17.8	1.6	493	0.4	14.0	2.1	377	0.3	22.4	2.6
Other Pacific Islander ^e	447	0.2	17.1	2.3	269	0.2	13.5	2.7	178	0.1	22.6	4.0
Hispanic ^f	14,800	5.6	19.1	0.4	8,057	5.7	15.0	0.5	6,743	5.4	23.9	0.6

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2005-2011 with follow-up through 2012.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorran, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.8

Cancer of the Prostate

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2005-2011

Race/Ethnicity	All Ages				Ages <65				Ages 65+			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	367,040	100.0	94.0	0.0	161,844	100.0	96.8	0.1	205,196	100.0	91.8	0.1
White	283,819	77.3	94.1	0.1	122,010	75.4	97.0	0.1	161,809	78.9	91.9	0.1
White Non-Hispanic	253,314	69.0	94.3	0.1	108,923	67.3	97.2	0.1	144,391	70.4	92.0	0.1
White Hispanic	30,505	8.3	93.2	0.2	13,087	8.1	95.7	0.2	17,418	8.5	91.3	0.3
Black	53,401	14.5	92.7	0.1	29,131	18.0	95.4	0.1	24,270	11.8	89.4	0.2
American Indian/Alaska Native ^c	938	0.3	89.2	1.2	389	0.2	92.5	1.6	549	0.3	86.8	1.8
Asian	15,792	4.3	94.6	0.2	5,337	3.3	96.9	0.3	10,455	5.1	93.5	0.3
Asian Indian, Pakistani	1,603	0.4	95.0	0.7	675	0.4	97.1	0.8	928	0.5	93.3	1.0
Chinese	3,274	0.9	95.0	0.5	981	0.6	98.6	0.5	2,293	1.1	93.5	0.6
Filipino	4,450	1.2	93.5	0.5	1,578	1.0	95.4	0.7	2,872	1.4	92.4	0.6
Japanese	2,481	0.7	95.1	0.5	684	0.4	97.7	0.7	1,797	0.9	94.1	0.6
Korean	781	0.2	94.2	1.0	186	0.1	94.1	2.1	595	0.3	94.2	1.2
Vietnamese	873	0.2	94.5	0.9	288	0.2	95.3	1.4	585	0.3	94.0	1.1
Other Asian ^d	2,330	0.6	95.9	0.5	945	0.6	98.2	0.5	1,385	0.7	94.2	0.8
Pacific Islander	1,254	0.3	90.0	1.0	492	0.3	92.9	1.4	762	0.4	88.2	1.4
Hawaiian	734	0.2	89.5	1.3	293	0.2	92.5	1.8	441	0.2	87.7	1.8
Other Pacific Islander ^e	520	0.1	90.8	1.6	199	0.1	93.7	2.2	321	0.2	89.1	2.1
Hispanic ^f	32,497	8.9	93.2	0.2	13,834	8.5	95.7	0.2	18,663	9.1	91.3	0.3

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2005-2011 with follow-up through 2012.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorroan, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.