

Table 19.14

Non-Hodgkin Lymphoma

Number of Cases

By SEER Location and Year of Diagnosis, 1975-2016

All Races, Males, Ages 20-54

Year of Diagnosis

SEER Location	1975- 1979	1980- 1981	1982- 1983	1984- 1985	1986- 1987	1988- 1989	1990- 1991	1992- 1993	1994- 1995	1996- 1997	1998- 1999	2000- 2001	2002- 2003	2004- 2005	2006- 2007	2008- 2009	2010- 2011	2012- 2013	2014- 2016
SEER 9 ^b	1,463	687	723	958	1,211	1,413	1,766	1,830	2,089	1,770	1,608	1,696	1,623	1,722	1,655	1,693	1,674	1,596	2,242
SEER 9 except SF-Oakland	1,208	563	588	730	827	923	1,195	1,294	1,483	1,325	1,269	1,327	1,308	1,410	1,383	1,401	1,402	1,328	1,848
San Francisco-Oakland SMSA ^c	255	124	135	228	384	490	571	536	606	445	339	369	315	312	272	292	272	268	394
San Francisco City/County	58	29	34	105	215	288	333	303	317	225	105	115	111	111	86	75	73	71	98

Age-Adjusted Cancer Incidence Rates^a

By SEER Location and Year of Diagnosis, 1975-2016

All Races, Males, Ages 20-54

Year of Diagnosis

SEER Location	1975- 1979	1980- 1981	1982- 1983	1984- 1985	1986- 1987	1988- 1989	1990- 1991	1992- 1993	1994- 1995	1996- 1997	1998- 1999	2000- 2001	2002- 2003	2004- 2005	2006- 2007	2008- 2009	2010- 2011	2012- 2013	2014- 2016
SEER 9 ^b	6.9	7.9	8.2	10.4	12.4	13.6	15.9	15.6	16.9	13.8	12.1	12.4	11.7	12.3	11.6	11.8	11.6	11.1	10.4
SEER 9 except SF-Oakland	6.9	7.8	8.1	9.7	10.3	10.9	13.1	13.4	14.4	12.3	11.5	11.6	11.2	11.9	11.5	11.6	11.5	11.0	10.3
San Francisco-Oakland SMSA ^c	6.9	8.4	8.9	13.9	22.6	26.4	29.5	26.7	29.8	21.3	15.6	16.4	13.9	14.1	12.3	13.0	11.8	11.4	11.0
San Francisco City/County	6.7	8.8	9.4	27.2	56.5	71.5	81.3	72.4	74.4	51.0	23.0	25.2	23.9	24.8	19.4	16.2	15.2	14.5	12.9

^a Rates are per 100,000 males aged 20-54 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco-Oakland SMSA, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

^c The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Non-Hodgkin Lymphoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.