

Table X-1

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1973-94

All Races, Males, All Ages

Year of Diagnosis

SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
All Areas [⊕]	174	34	44	109	217	367	517	723	853	1015	1110	1098	1131	1067	964	855
All Areas Excluding SF-Oakland SMSA	130	23	21	35	50	103	153	197	233	353	417	424	411	485	469	396
San Francisco-Oakland SMSA	44	11	23	74	167	264	364	526	620	662	693	674	720	582	495	459
San Francisco City/County	17	6	14	61	140	227	313	441	512	546	532	525	570	426	360	329

AGE-ADJUSTED CANCER INCIDENCE RATES^S

By SEER Location and Year of Diagnosis, 1973-94

All Races, Males, All Ages

Year of Diagnosis

SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
All Areas [⊕]	0.3	0.4	0.4	0.9	1.7	2.8	3.8	5.1	6.0	6.9	7.5	7.2	7.4	6.8	6.1	5.4
All Areas Excluding SF-Oakland SMSA	0.3	0.3	0.3	0.4	0.5	1.0	1.4	1.7	2.0	3.0	3.4	3.4	3.2	3.7	3.5	2.9
San Francisco-Oakland SMSA	0.5	0.7	1.3	3.5	7.3	11.5	15.7	21.6	25.4	26.1	27.2	25.7	27.8	22.1	18.4	17.4
San Francisco City/County	0.6	1.5	3.3	12.4	26.7	43.2	59.4	79.7	92.6	97.8	97.1	94.2	103.9	77.8	64.5	60.8

^S SEER Program. Rates are per 100,000 males and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

[⊕] All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

Table X-2

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1973-94

All Races, Males, Ages 20-54

SEER Location	Year of Diagnosis															
	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
All Areas [⊕]	18	5	20	75	178	328	475	682	799	944	1043	1037	1063	988	908	808
All Areas Excluding SF-Oakland SMSA	14	<4	4	9	25	77	128	174	205	311	378	388	376	452	436	376
San Francisco-Oakland SMSA	4	4	16	66	153	251	347	508	594	633	665	649	687	536	472	432
San Francisco City/County	<4	4	13	59	134	220	300	428	494	527	515	509	548	396	347	311

AGE-ADJUSTED CANCER INCIDENCE RATES^S

By SEER Location and Year of Diagnosis, 1973-94

All Races, Males, Ages 20-54

SEER Location	Year of Diagnosis															
	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
All Areas [⊕]	0.1	0.1	0.4	1.3	3.0	5.6	7.9	11.0	12.7	14.5	15.8	15.4	15.6	14.0	12.9	11.4
All Areas Excluding SF-Oakland SMSA	0.1	-	0.1	0.2	0.5	1.6	2.5	3.3	3.8	5.7	6.8	6.9	6.5	7.7	7.4	6.3
San Francisco-Oakland SMSA	0.1	0.5	1.9	6.9	14.9	24.9	34.0	47.5	55.3	56.3	59.0	56.1	59.7	44.6	39.6	36.4
San Francisco City/County	-	2.4	7.2	27.7	58.3	96.4	129.6	176.5	203.4	214.8	213.2	207.8	226.5	159.3	141.4	128.7

^S SEER Program. Rates are per 100,000 males aged 20-54 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

[⊕] All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

Table X-3

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1973-94

All Races, Males, Ages 55+

Year of Diagnosis

SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
All Areas [⊕]	156	29	24	34	39	38	42	39	54	70	66	61	68	78	56	47
All Areas Excluding SF-Oakland SMSA	116	22	17	26	25	26	25	22	28	42	39	36	35	32	33	20
San Francisco-Oakland SMSA	40	7	7	8	14	12	17	17	26	28	27	25	33	46	23	27
San Francisco City/County	16	<4	<4	<4	6	7	13	13	18	19	16	16	22	30	13	18

AGE-ADJUSTED CANCER INCIDENCE RATES^S

By SEER Location and Year of Diagnosis, 1973-94

All Races, Males, Ages 55+

Year of Diagnosis


SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
All Areas [⊕]	1.5	1.7	1.4	1.9	2.1	2.0	2.2	2.0	2.7	3.5	3.4	3.0	3.4	4.0	2.7	2.3
All Areas Excluding SF-Oakland SMSA	1.3	1.6	1.2	1.7	1.6	1.6	1.6	1.3	1.7	2.5	2.4	2.0	2.1	1.9	1.8	1.1
San Francisco-Oakland SMSA	2.3	2.7	2.4	2.7	4.8	4.0	5.4	5.5	8.3	9.0	8.9	8.1	10.8	15.3	7.2	8.9
San Francisco City/County	2.9	-	-	-	8.4	9.0	18.7	19.0	25.7	27.5	24.7	24.4	32.9	48.0	18.8	28.1

^S SEER Program. Rates are per 100,000 males aged 55+ and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

[⊕] All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

Figure X-1



All Cancer Sites vs. Kaposi's Sarcoma SEER Incidence Rates, 1973-1994 Males, Ages 20-54



Age-adjusted to 1970 Standard

San Francisco/Oakland
Ages 20-54, All Males, 1975-94

All SEER excluding San Francisco/Oakland
Ages 20-54, All Males, 1975-94



*Kaposi's excluded from Oral, Colorec., Lung, Melanoma, Testis, and NHL.