

Table XXVII-1

AGE-ADJUSTED CANCER SEER INCIDENCE RATES§ AND 23-YEAR TRENDS, 1973-95

By Primary Cancer Site

All Races, Males and Females

| Site | Ages 0-14 | | | | Ages 0-19 | | | |
|-------------------------|--------------------|----------------|-------------------|---------------|--------------------|----------------|-------------------|---------------|
| | Average 1973-74 | Rate⊕ 94-95 | % Change 73-95 | EAPC 73-95 | Average 1973-74 | Rate⊕ 94-95 | % Change 73-95 | EAPC 73-95 |
| All Sites | | | | | | | | |
| All Races | 12.8 | 13.6 | 5.8 | 0.7★ | 13.8 | 15.4 | 11.4 | 0.8★ |
| Whites | 13.3 | 13.9 | 4.7 | 0.6★ | 14.2 | 15.8 | 11.3 | 0.8★ |
| Blacks | 10.3 | 12.1 | 16.6 | 1.3★ | 11.1 | 12.6 | 14.0 | 0.9★ |
| Bone & Joint | 0.7 | 0.8 | 10.2 | 0.8 | 0.8 | 1.0 | 17.4 | 0.9★ |
| Brain & Other nervous | 2.4 | 3.2 | 30.4 | 1.7★ | 2.2 | 2.8 | 26.3 | 1.6★ |
| Hodgkin's disease | 0.7 | 0.4 | -40.3 | -1.8★ | 1.3 | 1.1 | -12.6 | -0.5 |
| Kidney & Renal pelvis | 0.7 | 0.8 | 19.4 | 0.4 | 0.6 | 0.7 | 18.2 | 0.5 |
| Leukemias | 4.3 | 3.8 | -11.0 | 0.4 | 3.8 | 3.5 | -8.1 | 0.4 |
| Acute lymphocytic | 2.7 | 3.1 | 12.4 | 1.0★ | 2.3 | 2.6 | 13.5 | 1.0★ |
| Non-Hodgkin's lymphomas | 0.8 | 1.0 | 25.6 | 0.5 | 0.8 | 1.2 | 53.2 | 1.3★ |
| Soft tissue | 0.7 | 0.9 | 31.5 | 0.7 | 0.8 | 0.9 | 20.2 | 0.6 |

The EAPC is the Estimated Annual Percent Change over the time interval.

§ SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

⊕ The Average Rate is the Average Annual Rate over the specified two-year period.

★ The EAPC is significantly different from zero ($p<.05$).

Table XXVII-2

AGE-ADJUSTED CANCER U.S. MORTALITY RATES* AND 23-YEAR TRENDS, 1973-95

By Race and Primary Cancer Site

Males and Females

| Race/Site | Ages 0-14 | | | | Ages 0-19 | | | |
|-------------------------|--------------------|----------------------------|-------------------|---------------|--------------------|----------------------------|-------------------|---------------|
| | Average 1973-74 | Rate [⊕] 94-95 | % Change 73-95 | EAPC 73-95 | Average 1973-74 | Rate [⊕] 94-95 | % Change 73-95 | EAPC 73-95 |
| <u>All Races</u> | | | | | | | | |
| All Sites | 5.4 | 2.8 | -48.1 | -3.0★ | 5.6 | 3.1 | -45.3 | -2.7★ |
| Bone & Joint | 0.2 | 0.1 | -41.4 | -2.8★ | 0.4 | 0.2 | -39.9 | -2.6★ |
| Brain & Other nervous | 1.0 | 0.8 | -25.6 | -1.2★ | 1.0 | 0.7 | -25.2 | -1.2★ |
| Hodgkin's disease | 0.1 | 0.0 | -83.2 | -7.3★ | 0.2 | 0.0 | -69.5 | -5.0★ |
| Kidney & Renal pelvis | 0.2 | 0.1 | -57.8 | -3.9★ | 0.2 | 0.1 | -54.7 | -3.6★ |
| Leukemias | 2.3 | 1.0 | -57.8 | -4.0★ | 2.3 | 1.0 | -53.5 | -3.5★ |
| Acute lymphocytic | 1.3 | 0.5 | -63.0 | -5.0★ | 1.1 | 0.5 | -57.8 | -4.3★ |
| Non-Hodgkin's lymphomas | 0.4 | 0.1 | -67.5 | -5.0★ | 0.5 | 0.2 | -61.4 | -4.0★ |
| Soft tissue | 0.6 | 0.1 | -75.5 | -6.6★ | 0.5 | 0.2 | -61.8 | -4.6★ |
| <u>Whites</u> | | | | | | | | |
| All Sites | 5.5 | 2.8 | -49.9 | -3.0★ | 5.7 | 3.1 | -46.8 | -2.8★ |
| Bone & Joint | 0.3 | 0.1 | -44.8 | -2.7★ | 0.4 | 0.2 | -40.4 | -2.4★ |
| Brain & Other nervous | 1.0 | 0.8 | -25.2 | -1.1★ | 0.9 | 0.7 | -24.0 | -1.1★ |
| Hodgkin's disease | 0.1 | 0.0 | -89.8 | -7.6★ | 0.2 | 0.0 | -74.7 | -5.4★ |
| Kidney & Renal pelvis | 0.2 | 0.1 | -55.9 | -3.8★ | 0.2 | 0.1 | -53.8 | -3.6★ |
| Leukemias | 2.5 | 1.0 | -60.1 | -4.2★ | 2.4 | 1.0 | -56.0 | -3.7★ |
| Acute lymphocytic | 1.4 | 0.5 | -63.9 | -5.1★ | 1.2 | 0.5 | -59.4 | -4.4★ |
| Non-Hodgkin's lymphomas | 0.4 | 0.1 | -69.0 | -4.9★ | 0.5 | 0.2 | -62.2 | -3.9★ |
| Soft tissue | 0.6 | 0.1 | -78.8 | -7.0★ | 0.5 | 0.2 | -65.4 | -4.9★ |
| <u>Blacks</u> | | | | | | | | |
| All Sites | 4.8 | 3.0 | -38.3 | -2.5★ | 5.2 | 3.3 | -36.8 | -2.3★ |
| Bone & Joint | 0.2 | 0.2 | -12.0 | -2.7★ | 0.3 | 0.2 | -31.1 | -3.1★ |
| Brain & Other nervous | 1.1 | 0.8 | -27.3 | -1.5★ | 1.1 | 0.7 | -29.6 | -1.4★ |
| Hodgkin's disease | 0.1 | 0.0 | -60.7 | - | 0.1 | 0.1 | -36.8 | -2.9★ |
| Kidney & Renal pelvis | 0.3 | 0.1 | -67.4 | -4.3★ | 0.2 | 0.1 | -59.6 | -3.8★ |
| Leukemias | 1.7 | 0.9 | -44.2 | -3.3★ | 1.7 | 1.1 | -36.0 | -2.5★ |
| Acute lymphocytic | 0.7 | 0.3 | -51.2 | -4.0★ | 0.7 | 0.4 | -37.2 | -3.0★ |
| Non-Hodgkin's lymphomas | 0.3 | 0.1 | -60.5 | -5.7★ | 0.4 | 0.2 | -56.6 | -4.5★ |
| Soft tissue | 0.5 | 0.2 | -60.5 | -5.0★ | 0.5 | 0.2 | -47.4 | -3.5★ |

The EAPC is the Estimated Annual Percent Change over the time interval.

* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

⊕ The Average Rate is the Average Annual Rate over the specified two-year period.

★ The EAPC is significantly different from zero ($p<.05$).

Table XXVII-3

AGE-ADJUSTED CANCER SEER INCIDENCE RATES\$, 1973-95

By Primary Cancer Site

All Races, Males and Females

Year of Diagnosis

| Age/Site | 1973 | 1974 | 1975 | 1976 | 1977 | 1978 | 1979 | 1980 | 1981 | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | |
|-----------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| <u>Ages 0-14</u> | | | | | | | | | | | | | | | | | | | | | | | | |
| All Sites | | | | | | | | | | | | | | | | | | | | | | | | |
| All Races | 12.7 | 13.0 | 11.3 | 12.9 | 12.7 | 13.0 | 12.9 | 12.7 | 12.3 | 12.8 | 12.8 | 13.7 | 14.3 | 14.0 | 14.0 | 13.2 | 14.8 | 14.0 | 15.0 | 13.5 | 14.7 | 14.0 | 13.2 | |
| Whites | 13.0 | 13.5 | 11.8 | 13.3 | 13.3 | 13.5 | 13.5 | 12.8 | 12.3 | 12.9 | 13.4 | 14.2 | 15.3 | 14.6 | 14.4 | 13.7 | 15.3 | 14.0 | 15.8 | 13.8 | 14.7 | 14.3 | 13.5 | |
| Blacks | 10.5 | 10.2 | 8.5 | 11.1 | 8.3 | 9.8 | 10.4 | 11.0 | 13.0 | 11.8 | 9.9 | 10.3 | 9.9 | 10.5 | 11.4 | 11.8 | 13.6 | 12.3 | 11.8 | 12.5 | 13.0 | 13.0 | 11.1 | |
| Bone & Joint | 0.8 | 0.6 | 0.5 | 0.5 | 0.6 | 0.7 | 0.8 | 0.7 | 0.6 | 0.6 | 0.7 | 0.7 | 0.8 | 0.7 | 0.7 | 0.6 | 0.6 | 0.7 | 0.9 | 0.6 | 0.6 | 0.8 | 0.8 | |
| Brain & Other nervous | 2.3 | 2.6 | 2.3 | 2.4 | 3.0 | 2.7 | 2.8 | 2.8 | 2.5 | 2.4 | 2.2 | 2.8 | 2.9 | 3.4 | 3.1 | 3.3 | 3.2 | 3.5 | 3.5 | 3.2 | 3.4 | 3.3 | 3.1 | |
| Hodgkin's disease | 0.9 | 0.6 | 0.7 | 0.8 | 0.7 | 0.7 | 0.9 | 0.5 | 0.6 | 0.5 | 0.7 | 0.7 | 0.7 | 0.5 | 0.7 | 0.6 | 0.8 | 0.7 | 0.6 | 0.5 | 0.6 | 0.6 | 0.3 | |
| Kidney & Renal pelvis | 0.7 | 0.7 | 0.6 | 0.9 | 0.7 | 0.9 | 0.9 | 0.8 | 0.8 | 0.9 | 1.1 | 0.7 | 0.8 | 0.9 | 0.8 | 0.7 | 1.1 | 0.7 | 0.9 | 0.7 | 0.9 | 1.1 | 0.5 | |
| Leukemias | 3.8 | 4.8 | 3.2 | 4.0 | 3.8 | 3.7 | 3.6 | 3.9 | 3.9 | 3.7 | 3.6 | 4.4 | 4.5 | 4.0 | 4.3 | 4.0 | 4.6 | 4.3 | 4.5 | 4.0 | 4.1 | 3.6 | 4.1 | |
| Acute lymphocytic | 2.4 | 3.1 | 2.1 | 2.9 | 3.0 | 2.9 | 2.7 | 3.0 | 3.1 | 2.9 | 2.9 | 3.4 | 3.4 | 3.2 | 3.3 | 3.2 | 3.8 | 3.3 | 3.4 | 3.0 | 3.0 | 2.9 | 3.3 | |
| Non-Hodgkin's lymph | 0.6 | 0.9 | 1.0 | 0.8 | 0.8 | 1.0 | 0.8 | 0.7 | 0.7 | 0.9 | 0.9 | 0.9 | 0.7 | 1.0 | 1.1 | 0.7 | 0.8 | 0.8 | 0.9 | 0.8 | 0.8 | 1.0 | 0.9 | |
| Soft tissue | 0.8 | 0.7 | 0.8 | 0.7 | 0.7 | 0.8 | 0.6 | 0.9 | 0.7 | 0.9 | 0.9 | 0.7 | 0.8 | 0.9 | 0.8 | 0.6 | 0.9 | 0.8 | 1.0 | 0.7 | 0.6 | 0.9 | 0.9 | |
| <u>Ages 0-19</u> | | | | | | | | | | | | | | | | | | | | | | | | |
| All Sites | | | | | | | | | | | | | | | | | | | | | | | | |
| All Races | 13.6 | 13.9 | 12.8 | 14.1 | 14.1 | 14.4 | 14.5 | 14.2 | 13.8 | 14.4 | 14.4 | 15.4 | 15.7 | 15.7 | 15.7 | 15.2 | 15.0 | 16.5 | 15.5 | 16.2 | 16.1 | 16.0 | 15.6 | 15.1 |
| Whites | 14.1 | 14.3 | 13.2 | 14.7 | 14.7 | 15.0 | 15.1 | 14.6 | 14.0 | 14.9 | 15.2 | 15.9 | 16.8 | 16.4 | 16.1 | 15.5 | 17.1 | 15.7 | 17.2 | 16.7 | 16.1 | 16.0 | 15.7 | |
| Blacks | 10.3 | 11.8 | 10.5 | 10.7 | 10.4 | 10.8 | 12.6 | 11.0 | 12.9 | 11.8 | 11.1 | 12.1 | 10.5 | 12.0 | 11.2 | 12.9 | 14.0 | 12.9 | 12.4 | 12.9 | 13.6 | 13.7 | 11.5 | |
| Bone & Joint | 0.9 | 0.8 | 0.7 | 0.6 | 0.7 | 0.9 | 0.9 | 0.9 | 1.0 | 0.8 | 0.8 | 0.9 | 1.0 | 0.9 | 1.0 | 0.8 | 0.9 | 0.8 | 1.0 | 0.9 | 0.9 | 1.0 | 0.9 | |
| Brain & Other nervous | 2.1 | 2.3 | 2.1 | 2.2 | 2.9 | 2.5 | 2.5 | 2.4 | 2.3 | 2.3 | 2.2 | 2.6 | 2.6 | 3.2 | 2.9 | 3.0 | 2.9 | 3.1 | 3.1 | 3.3 | 3.0 | 2.9 | 2.8 | |
| Hodgkin's disease | 1.4 | 1.1 | 1.5 | 1.4 | 1.4 | 1.5 | 1.6 | 1.2 | 1.2 | 1.5 | 1.4 | 1.5 | 1.4 | 1.3 | 1.4 | 1.3 | 1.7 | 1.3 | 1.4 | 1.4 | 1.3 | 1.3 | 1.0 | |
| Kidney & Renal pelvis | 0.6 | 0.5 | 0.5 | 0.7 | 0.6 | 0.7 | 0.7 | 0.6 | 0.6 | 0.7 | 0.8 | 0.6 | 0.7 | 0.7 | 0.6 | 0.8 | 0.6 | 0.7 | 0.6 | 0.8 | 0.9 | 0.4 | | |
| Leukemias | 3.5 | 4.2 | 3.0 | 3.6 | 3.4 | 3.4 | 3.2 | 3.4 | 3.6 | 3.4 | 3.4 | 3.9 | 4.0 | 3.6 | 3.6 | 3.5 | 4.1 | 3.7 | 4.0 | 3.6 | 3.6 | 3.3 | 3.7 | |
| Acute lymphocytic | 2.0 | 2.7 | 1.9 | 2.4 | 2.5 | 2.5 | 2.3 | 2.4 | 2.7 | 2.5 | 2.5 | 2.9 | 2.8 | 2.7 | 2.7 | 2.7 | 3.3 | 2.8 | 2.9 | 2.5 | 2.4 | 2.8 | | |
| Non-Hodgkin's lymph | 0.7 | 0.9 | 1.0 | 0.9 | 0.8 | 1.0 | 0.9 | 0.9 | 1.0 | 1.0 | 1.1 | 1.0 | 1.1 | 1.0 | 1.0 | 1.0 | 1.0 | 1.1 | 1.0 | 0.9 | 1.3 | 1.1 | | |
| Soft tissue | 0.7 | 0.8 | 0.8 | 0.7 | 0.7 | 0.9 | 0.7 | 0.9 | 0.7 | 1.0 | 1.0 | 0.7 | 0.9 | 0.9 | 0.7 | 0.9 | 0.8 | 1.0 | 0.7 | 0.7 | 0.8 | 1.0 | | |

§ SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-4

AGE-ADJUSTED CANCER U.S. MORTALITY RATES*, 1973-95

By Primary Cancer Site

All Races, Males and Females

Year of Diagnosis

| Age/Site | 1973 | 1974 | 1975 | 1976 | 1977 | 1978 | 1979 | 1980 | 1981 | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | |
|-----------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|-----|
| <u>Ages 0-14</u> | | | | | | | | | | | | | | | | | | | | | | | | |
| All Sites | 5.5 | 5.3 | 4.9 | 4.9 | 4.7 | 4.3 | 4.4 | 4.3 | 4.2 | 4.2 | 4.1 | 3.6 | 3.6 | 3.5 | 3.4 | 3.3 | 3.3 | 3.1 | 3.1 | 3.0 | 3.0 | 2.8 | 2.7 | |
| Bone & Joint | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.2 |
| Brain & Other nervous | 1.0 | 1.1 | 1.0 | 1.0 | 1.0 | 0.8 | 0.9 | 0.9 | 0.9 | 1.0 | 0.9 | 0.8 | 0.8 | 0.9 | 0.7 | 0.8 | 0.9 | 0.9 | 0.9 | 0.8 | 0.8 | 0.8 | 0.8 | 0.7 |
| Hodgkin's disease | 0.1 | 0.1 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Kidney & Renal pelvis | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Leukemias | 2.4 | 2.3 | 2.1 | 2.1 | 2.0 | 1.9 | 1.8 | 1.7 | 1.6 | 1.6 | 1.5 | 1.3 | 1.4 | 1.3 | 1.3 | 1.2 | 1.2 | 1.1 | 1.1 | 1.1 | 1.1 | 1.0 | 1.0 | 1.0 |
| Acute lymphocytic | 1.3 | 1.3 | 1.2 | 1.2 | 1.1 | 1.0 | 1.0 | 0.9 | 1.0 | 0.9 | 0.8 | 0.8 | 0.8 | 0.7 | 0.7 | 0.6 | 0.6 | 0.5 | 0.5 | 0.5 | 0.5 | 0.4 | 0.5 | 0.5 |
| Non-Hodgkin's lymph | 0.4 | 0.4 | 0.4 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 |
| Soft tissue | 0.6 | 0.5 | 0.5 | 0.6 | 0.6 | 0.5 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.1 | 0.1 |
| <u>Ages 0-19</u> | | | | | | | | | | | | | | | | | | | | | | | | |
| All Sites | 5.8 | 5.5 | 5.1 | 5.1 | 5.0 | 4.5 | 4.6 | 4.6 | 4.4 | 4.5 | 4.3 | 3.9 | 3.8 | 3.8 | 3.6 | 3.5 | 3.5 | 3.4 | 3.4 | 3.3 | 3.3 | 3.2 | 3.0 | |
| Bone & Joint | 0.4 | 0.4 | 0.3 | 0.4 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |
| Brain & Other nervous | 0.9 | 1.0 | 0.9 | 0.9 | 0.9 | 0.8 | 0.9 | 0.8 | 0.8 | 0.9 | 0.8 | 0.8 | 0.8 | 0.7 | 0.8 | 0.8 | 0.8 | 0.8 | 0.8 | 0.8 | 0.8 | 0.7 | 0.7 | 0.7 |
| Hodgkin's disease | 0.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 |
| Kidney & Renal pelvis | 0.2 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Leukemias | 2.3 | 2.2 | 2.0 | 2.1 | 2.0 | 1.8 | 1.8 | 1.8 | 1.6 | 1.6 | 1.5 | 1.4 | 1.4 | 1.3 | 1.4 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.1 | 1.1 | 1.1 | 1.0 |
| Acute lymphocytic | 1.2 | 1.1 | 1.0 | 1.1 | 1.0 | 0.9 | 1.0 | 0.9 | 0.9 | 0.9 | 0.8 | 0.7 | 0.7 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.5 | 0.6 | 0.5 | 0.5 | 0.5 | 0.5 |
| Non-Hodgkin's lymph | 0.5 | 0.5 | 0.4 | 0.3 | 0.4 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |
| Soft tissue | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 0.3 | 0.2 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |

* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-5

AGE-ADJUSTED CANCER U.S. MORTALITY RATES*, 1973-95

By Primary Cancer Site

Whites, Males and Females

Year of Diagnosis

| <u>Age/Site</u> | 1973 | 1974 | 1975 | 1976 | 1977 | 1978 | 1979 | 1980 | 1981 | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | |
|-----------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|-----|
| <u>Ages 0-14</u> | | | | | | | | | | | | | | | | | | | | | | | | |
| All Sites | 5.6 | 5.4 | 5.0 | 5.0 | 4.8 | 4.3 | 4.5 | 4.4 | 4.2 | 4.3 | 4.1 | 3.7 | 3.6 | 3.6 | 3.5 | 3.3 | 3.4 | 3.2 | 3.2 | 3.1 | 3.1 | 2.8 | 2.7 | |
| Bone & Joint | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 |
| Brain & Other nervous | 1.0 | 1.1 | 1.0 | 1.0 | 1.0 | 0.8 | 0.9 | 0.9 | 0.9 | 1.0 | 0.9 | 0.9 | 0.9 | 0.9 | 0.8 | 0.8 | 0.9 | 0.9 | 0.9 | 0.9 | 0.8 | 0.8 | 0.8 | 0.7 |
| Hodgkin's disease | 0.1 | 0.1 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Kidney & Renal pelvis | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Leukemias | 2.5 | 2.4 | 2.3 | 2.3 | 2.1 | 1.9 | 1.9 | 1.8 | 1.7 | 1.6 | 1.6 | 1.4 | 1.4 | 1.4 | 1.4 | 1.2 | 1.2 | 1.1 | 1.1 | 1.2 | 1.1 | 1.0 | 1.0 | 1.0 |
| Acute lymphocytic | 1.4 | 1.4 | 1.2 | 1.3 | 1.2 | 1.1 | 1.1 | 1.0 | 1.0 | 0.9 | 0.9 | 0.8 | 0.8 | 0.8 | 0.7 | 0.7 | 0.6 | 0.6 | 0.6 | 0.5 | 0.6 | 0.5 | 0.5 | 0.5 |
| Non-Hodgkin's lymph | 0.5 | 0.4 | 0.4 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.1 |
| Soft tissue | 0.6 | 0.5 | 0.5 | 0.6 | 0.6 | 0.5 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| <u>Ages 0-19</u> | | | | | | | | | | | | | | | | | | | | | | | | |
| All Sites | 5.9 | 5.6 | 5.2 | 5.2 | 5.1 | 4.6 | 4.7 | 4.7 | 4.4 | 4.5 | 4.4 | 4.0 | 3.9 | 3.9 | 3.7 | 3.6 | 3.6 | 3.4 | 3.5 | 3.4 | 3.3 | 3.2 | 3.0 | |
| Bone & Joint | 0.4 | 0.4 | 0.3 | 0.4 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 |
| Brain & Other nervous | 0.9 | 1.0 | 0.9 | 0.9 | 0.9 | 0.8 | 0.9 | 0.8 | 0.8 | 0.9 | 0.9 | 0.8 | 0.8 | 0.8 | 0.7 | 0.8 | 0.8 | 0.8 | 0.8 | 0.8 | 0.8 | 0.7 | 0.7 | 0.7 |
| Hodgkin's disease | 0.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 |
| Kidney & Renal pelvis | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Leukemias | 2.4 | 2.3 | 2.1 | 2.2 | 2.1 | 1.9 | 1.9 | 1.8 | 1.7 | 1.7 | 1.6 | 1.4 | 1.4 | 1.4 | 1.4 | 1.3 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.1 | 1.0 | 1.0 |
| Acute lymphocytic | 1.3 | 1.2 | 1.1 | 1.1 | 1.1 | 1.0 | 1.0 | 1.0 | 0.9 | 0.9 | 0.8 | 0.8 | 0.8 | 0.7 | 0.7 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 | 0.5 | 0.5 | 0.5 | 0.5 |
| Non-Hodgkin's lymph | 0.5 | 0.5 | 0.5 | 0.4 | 0.4 | 0.3 | 0.4 | 0.3 | 0.4 | 0.3 | 0.3 | 0.2 | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |
| Soft tissue | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |

* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-6

AGE-ADJUSTED CANCER U.S. MORTALITY RATES*, 1973-95

By Primary Cancer Site

Blacks, Males and Females

Year of Diagnosis

| <u>Age/Site</u> | 1973 | 1974 | 1975 | 1976 | 1977 | 1978 | 1979 | 1980 | 1981 | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
|-----------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| <u>Ages 0-14</u> | | | | | | | | | | | | | | | | | | | | | | | |
| All Sites | 5.2 | 4.4 | 4.5 | 4.2 | 4.2 | 4.1 | 4.0 | 3.8 | 4.2 | 4.0 | 3.9 | 3.5 | 3.3 | 3.4 | 3.0 | 3.2 | 3.2 | 3.0 | 3.1 | 2.7 | 2.7 | 2.9 | 3.0 |
| Bone & Joint | 0.2 | 0.2 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Brain & Other nervous | 1.2 | 1.1 | 0.9 | 1.0 | 1.0 | 0.8 | 0.9 | 0.9 | 1.0 | 0.8 | 0.8 | 0.7 | 0.6 | 0.9 | 0.7 | 1.0 | 0.8 | 0.7 | 0.8 | 0.7 | 0.6 | 0.8 | 0.8 |
| Hodgkin's disease | 0.1 | 0.1 | 0.1 | 0.0 | 0.1 | 0.1 | 0.1 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 |
| Kidney & Renal pelvis | 0.4 | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 |
| Leukemias | 1.8 | 1.6 | 1.6 | 1.6 | 1.4 | 1.5 | 1.3 | 1.2 | 1.6 | 1.5 | 1.2 | 1.3 | 1.2 | 1.1 | 1.0 | 0.9 | 1.0 | 0.9 | 1.0 | 0.8 | 0.9 | 0.8 | 1.1 |
| Acute lymphocytic | 0.7 | 0.7 | 0.8 | 0.8 | 0.7 | 0.6 | 0.6 | 0.6 | 0.8 | 0.7 | 0.7 | 0.7 | 0.7 | 0.7 | 0.6 | 0.5 | 0.4 | 0.5 | 0.5 | 0.4 | 0.4 | 0.3 | 0.4 |
| Non-Hodgkin's lymph | 0.3 | 0.3 | 0.4 | 0.2 | 0.3 | 0.4 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.3 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 |
| Soft tissue | 0.6 | 0.4 | 0.4 | 0.5 | 0.5 | 0.5 | 0.3 | 0.1 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.1 | 0.3 | 0.2 | 0.2 | 0.3 | 0.1 | 0.2 | 0.2 | 0.3 | 0.1 |
| <u>Ages 0-19</u> | | | | | | | | | | | | | | | | | | | | | | | |
| All Sites | 5.6 | 4.8 | 4.7 | 4.5 | 4.7 | 4.5 | 4.3 | 4.2 | 4.5 | 4.3 | 3.8 | 3.7 | 3.5 | 3.7 | 3.6 | 3.5 | 3.5 | 3.4 | 3.4 | 3.0 | 3.1 | 3.3 | 3.3 |
| Bone & Joint | 0.3 | 0.3 | 0.4 | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |
| Brain & Other nervous | 1.1 | 1.0 | 0.9 | 0.9 | 0.9 | 0.7 | 0.9 | 0.8 | 0.9 | 0.8 | 0.7 | 0.7 | 0.6 | 0.8 | 0.7 | 0.9 | 0.7 | 0.7 | 0.7 | 0.7 | 0.7 | 0.8 | 0.7 |
| Hodgkin's disease | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.0 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Kidney & Renal pelvis | 0.3 | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 |
| Leukemias | 1.8 | 1.5 | 1.6 | 1.5 | 1.4 | 1.5 | 1.4 | 1.4 | 1.5 | 1.5 | 1.1 | 1.3 | 1.1 | 1.1 | 1.2 | 1.0 | 1.1 | 1.0 | 1.1 | 0.9 | 1.0 | 1.0 | 1.1 |
| Acute lymphocytic | 0.7 | 0.6 | 0.7 | 0.7 | 0.7 | 0.6 | 0.6 | 0.6 | 0.7 | 0.7 | 0.6 | 0.6 | 0.6 | 0.6 | 0.5 | 0.5 | 0.5 | 0.5 | 0.4 | 0.3 | 0.3 | 0.3 | 0.5 |
| Non-Hodgkin's lymph | 0.5 | 0.3 | 0.4 | 0.3 | 0.4 | 0.4 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.3 | 0.2 | 0.1 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 |
| Soft tissue | 0.5 | 0.4 | 0.5 | 0.5 | 0.5 | 0.5 | 0.3 | 0.2 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 |

* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-7

AGE-ADJUSTED CANCER SEER INCIDENCES AND U.S. MORTALITY* RATES, 1991-95

By Primary Cancer Site

All Races, Males and Females

| Site | Ages 0-14 | | | | | | Ages 0-19 | | | | | |
|-------------------------|-----------|-------|---------|-----------|-------|---------|-----------|-------|---------|-----------|-------|---------|
| | Incidence | | | Mortality | | | Incidence | | | Mortality | | |
| | Total | Males | Females |
| All Sites | | | | | | | | | | | | |
| All Races | 14.1 | 14.9 | 13.1 | 2.9 | 3.3 | 2.6 | 15.8 | 16.5 | 15.0 | 3.2 | 3.7 | 2.8 |
| Whites | 14.4 | 15.3 | 13.5 | 3.0 | 3.3 | 2.6 | 16.3 | 17.1 | 15.5 | 3.3 | 3.7 | 2.8 |
| Blacks | 12.3 | 12.7 | 11.8 | 2.9 | 3.0 | 2.7 | 12.8 | 13.0 | 12.6 | 3.2 | 3.5 | 3.0 |
| Bone & Joint | 0.7 | 0.7 | 0.8 | 0.1 | 0.1 | 0.1 | 1.0 | 1.1 | 0.8 | 0.2 | 0.3 | 0.2 |
| Brain & Other nervous | 3.3 | 3.5 | 3.0 | 0.8 | 0.8 | 0.7 | 3.0 | 3.3 | 2.7 | 0.7 | 0.8 | 0.7 |
| Hodgkin's disease | 0.5 | 0.5 | 0.5 | 0.0 | 0.0 | 0.0 | 1.3 | 1.2 | 1.4 | 0.1 | 0.1 | 0.0 |
| Kidney & Renal pelvis | 0.8 | 0.9 | 0.8 | 0.1 | 0.1 | 0.1 | 0.7 | 0.7 | 0.7 | 0.1 | 0.1 | 0.1 |
| Leukemias | 4.0 | 4.4 | 3.7 | 1.1 | 1.2 | 0.9 | 3.7 | 3.9 | 3.4 | 1.1 | 1.3 | 0.9 |
| Acute lymphocytic | 3.1 | 3.4 | 2.8 | 0.5 | 0.6 | 0.4 | 2.6 | 2.9 | 2.3 | 0.5 | 0.6 | 0.4 |
| Non-Hodgkin's lymphomas | 0.9 | 1.3 | 0.5 | 0.1 | 0.2 | 0.1 | 1.1 | 1.5 | 0.6 | 0.2 | 0.3 | 0.1 |
| Soft tissue | 0.9 | 0.9 | 0.8 | 0.1 | 0.2 | 0.1 | 0.9 | 0.9 | 0.8 | 0.2 | 0.2 | 0.2 |

^s SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-8

5-YEAR RELATIVE SURVIVAL RATES (Percent)

By Selected Primary Cancer Site and Year Of Diagnosis

All Races, Males and Females

| Site | Ages 0-14 | | | | | | Ages 0-19 | | | | | |
|-------------------------|-----------|---------|---------|---------|---------|---------|-----------|---------|---------|---------|---------|---------|
| | 1974-76 | 1977-79 | 1980-82 | 1983-85 | 1986-88 | 1989-94 | 1974-76 | 1977-79 | 1980-82 | 1983-85 | 1986-88 | 1989-94 |
| All Sites | | | | | | | | | | | | |
| All Races | 55.6 | 61.5 | 64.8 | 67.8 | 70.0 | 73.8S | 58.4 | 64.0 | 66.5 | 69.8 | 72.8 | 74.7S |
| Whites | 55.7 | 61.9 | 65.3 | 69.3 | 71.2 | 74.6S | 58.7 | 64.4 | 67.4 | 71.4 | 74.3 | 75.9S |
| Blacks | 56.8 | 57.7 | 61.3 | 57.4 | 62.1 | 69.5S | 56.6 | 60.3 | 61.2 | 59.0 | 62.5 | 67.5S |
| Bone & Joint | 53.7↑ | 52.8↑ | 54.1↑ | 58.5↑ | 62.4↑ | 64.0S | 50.3 | 51.4 | 49.8 | 53.7 | 60.9 | 64.2S |
| Brain & Other nervous | 54.7 | 56.3 | 55.3 | 61.9 | 62.3 | 63.0S | 55.5 | 56.7 | 56.8 | 63.4 | 65.3 | 65.0S |
| Hodgkin's disease | 79.0 | 83.2 | 91.0 | 89.6 | 89.6 | 92.4S | 85.3 | 87.9 | 87.2 | 88.2 | 89.2 | 91.8S |
| Leukemias | 44.9 | 58.8 | 61.9 | 61.3 | 70.3 | 73.0S | 40.7 | 53.1 | 57.3 | 58.3 | 66.2 | 69.0S |
| Acute lymphocytic | 52.9 | 66.7 | 70.3 | 69.5 | 78.4 | 80.2S | 49.0 | 62.6 | 66.6 | 67.0 | 75.5 | 77.0S |
| Acute myeloid | 13.9 | 26.2↑ | 21.1↑ | 32.0↑ | 27.8↑ | 43.4S | 16.1 | 25.4 | 21.4 | 33.3↑ | 29.6↑ | 42.9S |
| Neuroblastoma | 52.2 | 53.6 | 52.9 | 54.0 | 59.9 | 69.2S | 51.7 | 53.6 | 53.1 | 53.3 | 58.9 | 67.8S |
| Non-Hodgkin's lymphomas | 44.6 | 50.7 | 61.8 | 70.4 | 69.2 | 77.6S | 44.0 | 49.7 | 59.4 | 66.5 | 73.1 | 74.5S |
| Soft tissue | 60.7 | 68.6 | 65.0 | 76.4 | 65.9 | 76.0S | 62.0 | 64.6 | 64.6 | 74.0 | 67.8 | 69.0 |
| Wilms' tumor | 74.3 | 77.3 | 86.8 | 86.4 | 91.3 | 92.9S | 73.6 | 76.8 | 86.2 | 86.7 | 91.4 | 92.3S |

Rates are from the SEER Program. They are based on data from population-based registries in Connecticut, New Mexico, Utah, Iowa, Hawaii, Atlanta, Detroit, Seattle-Puget Sound and San Francisco-Oakland. Rates are based on follow-up of patients through 1995.

S The difference in rates between 1974-76 and 1989-94 is statistically significant ($p<.05$).

↑ The standard error of the survival rate is between 5 and 10 percentage points.

↑↑ The standard error of the survival rate is greater than 10 percentage points.

Table XXVII-9

CHILDHOOD CANCER
 SEER Incidence, U.S. Mortality and 5-Year Relative Survival Rates
 For Top Cancer Sites[⊕] By 5-Year Age Groups
 All Races, Males and Females

| Age/Site | SEER Incidences | | | | US Mortality* | | | | Survivals§ | |
|---|-----------------|-------|---------|-------|---------------|-------|---------|-------|------------|---------|
| | 1991-95 | | 1973-95 | | 1991-95 | | 1973-95 | | 1974-76 | 1989-94 |
| | Num | Cases | Rate | PC | EAPC | Num | Cases | Rate | PC | EAPC |
| <u>Ages 0-4</u> | | | | | | | | | | |
| All Sites | 1,890 | 20.4 | 10.2 | 0.9★ | | 2,922 | 3.0 | -49.0 | -3.0★ | 56.2 |
| Soft tissue | 131 | 1.4 | 37.0 | 0.9 | | 141 | 0.1 | -86.4 | -9.9★ | 57.7↑ |
| Kidney and Renal pelvis | 188 | 2.0 | 31.8 | 0.5 | | 98 | 0.1 | -62.1 | -5.4★ | 74.2 |
| Eye and Orbit | 125 | 1.3 | -8.4 | 0.8 | | 54 | 0.1 | -62.6 | -3.5★ | 89.2 |
| Brain and Other nervous | 358 | 3.9 | 53.5 | 2.5★ | | 733 | 0.8 | -28.4 | -1.3★ | 51.2 |
| Other endocrine | 130 | 1.4 | - | - | | 532 | 0.5 | - | - | 32.8↑ |
| Leukemias | 639 | 6.9 | -8.2 | 0.3 | | 957 | 1.0 | -55.0 | -3.7★ | 50.8 |
| Acute lymphocytic | 511 | 5.5 | 18.3 | 0.9★ | | 311 | 0.3 | -69.9 | -5.5★ | 58.1 |
| <u>Ages 5-9</u> | | | | | | | | | | |
| All Sites | 896 | 10.2 | -0.7 | 0.4 | | 2,785 | 3.0 | -52.3 | -3.3★ | 55.4 |
| Brain and Other nervous | 275 | 3.1 | 15.0 | 1.2★ | | 872 | 0.9 | -27.0 | -0.9★ | 54.4 |
| Leukemias | 288 | 3.3 | -19.0 | 0.5 | | 1,017 | 1.1 | -66.9 | -5.1★ | 46.4 |
| Acute lymphocytic | 232 | 2.7 | 9.9 | 1.2★ | | 544 | 0.6 | -69.8 | -6.0★ | 53.4 |
| <u>Ages 10-14</u> | | | | | | | | | | |
| All Sites | 1,082 | 12.5 | 5.5 | 0.7★ | | 2,594 | 2.8 | -42.1 | -2.5★ | 55.1 |
| Bone and Joint | 114 | 1.3 | -8.9 | 0.5 | | 261 | 0.3 | -43.7 | -2.8★ | 52.8↑ |
| Brain and Other nervous | 257 | 3.0 | 25.1 | 1.5★ | | 635 | 0.7 | -21.1 | -1.4★ | 58.7 |
| Hodgkin's disease | 99 | 1.1 | -37.5 | -1.4★ | | 34 | 0.0 | -80.2 | -6.9★ | 75.9↑ |
| Leukemias | 207 | 2.4 | -7.2 | 0.5 | | 988 | 1.1 | -46.6 | -3.0★ | 29.0 |
| Acute lymphocytic | 139 | 1.6 | 0.4 | 1.2 | | 487 | 0.5 | -45.4 | -3.3★ | 37.2↑ |
| <u>Ages 15-19</u> | | | | | | | | | | |
| All Sites | 1,718 | 21.1 | 24.4 | 1.0★ | | 3,614 | 4.1 | -38.2 | -2.2★ | 63.8 |
| Bone and Joint | 133 | 1.6 | 29.5 | 1.1 | | 454 | 0.5 | -38.6 | -2.4★ | 46.6↑ |
| Melanomas of skin | 118 | 1.4 | 57.0 | 2.3★ | | 43 | 0.0 | -52.2 | -3.7★ | 76.7↑ |
| Ovary(females) | 100 | 2.5 | 78.4 | 3.5★ | | 38 | 0.1 | -76.4 | -6.5★ | 69.0↑ |
| Testis(males) | 142 | 3.4 | 65.4 | 2.6★ | | 52 | 0.1 | -75.6 | -7.8★ | 55.4↑ |
| Brain and Other nervous | 165 | 2.0 | 7.2 | 0.9 | | 536 | 0.6 | -23.6 | -1.1★ | 58.4↑ |
| Thyroid | 122 | 1.5 | 28.9 | 0.8 | | - | - | - | - | 100.0 |
| Hodgkin's disease | 288 | 3.5 | 7.9 | 0.1 | | 136 | 0.2 | -62.2 | -4.4★ | 88.8 |
| Non-Hodgkin's lymphomas | 137 | 1.7 | 128.2 | 2.9★ | | 324 | 0.4 | -48.8 | -2.5★ | 42.6↑ |
| Leukemias | 207 | 2.5 | 7.5 | 0.3 | | 1,139 | 1.3 | -38.2 | -2.1★ | 21.5 |
| Acute lymphocytic | 98 | 1.2 | 22.3 | 1.0 | | 480 | 0.5 | -31.8 | -1.7★ | 21.2↑ |
| <u>Note:</u> | | | | | | | | | | |
| The PC is the Percent Change over the time interval. | | | | | | | | | | |
| The EAPC is the Estimated Annual Percent Change over the time interval. | | | | | | | | | | |
| Incidence and mortality rates are per 100,000. Survival rates are expressed as percents. | | | | | | | | | | |
| SEER Program. | | | | | | | | | | |
| NCHS public use tape. | | | | | | | | | | |
| The EAPC is significantly different from zero ($p < .05$). | | | | | | | | | | |
| The difference in rates between 1974-76 and 1989-94 is statistically significant ($p < .05$). | | | | | | | | | | |
| The standard error of the survival rate is between 5 and 10 percentage points. | | | | | | | | | | |
| The standard error of the survival rate is greater than 10 percentage points. | | | | | | | | | | |
| Sites were chosen based on more than 100 SEER incident cases between 1991-95. | | | | | | | | | | |
| Statistic could not be calculated. | | | | | | | | | | |

The PC is the Percent Change over the time interval.

The EAPC is the Estimated Annual Percent Change over the time interval.

Incidence and mortality rates are per 100,000. Survival rates are expressed as percents.

SEER Program.

NCHS public use tape.

The EAPC is significantly different from zero ($p < .05$).

The difference in rates between 1974-76 and 1989-94 is statistically significant ($p < .05$).

The standard error of the survival rate is between 5 and 10 percentage points.

The standard error of the survival rate is greater than 10 percentage points.

Sites were chosen based on more than 100 SEER incident cases between 1991-95.

Statistic could not be calculated.

★ S ↑ ⊕ -

Table XXVII-10
ALL SITES (Invasive)

INCIDENCE, MORTALITY, AND SURVIVAL RATES

By Year of Diagnosis/Death

All Races, Males and Females

Ages 0-14

Year of Diagnosis/Death

| | 1975 | 1976 | 1977 | 1978 | 1979 | 1980 | 1981 | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
|---------------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Incidence | | | | | | | | | | | | | | | | | | | | | |
| SEER | 11.3 | 12.9 | 12.7 | 13.0 | 12.9 | 12.7 | 12.3 | 12.8 | 12.8 | 13.7 | 14.3 | 14.0 | 14.0 | 13.2 | 14.8 | 14.0 | 15.0 | 13.5 | 14.7 | 14.0 | 13.2 |
| Mortality | | | | | | | | | | | | | | | | | | | | | |
| U.S. | 4.9 | 4.9 | 4.7 | 4.3 | 4.4 | 4.3 | 4.2 | 4.2 | 4.1 | 3.6 | 3.6 | 3.5 | 3.4 | 3.3 | 3.3 | 3.1 | 3.1 | 3.0 | 3.0 | 2.8 | 2.7 |
| SEER | 5.0 | 4.7 | 4.9 | 4.3 | 4.6 | 4.2 | 4.1 | 4.3 | 4.5 | 3.6 | 4.0 | 4.1 | 3.8 | 3.6 | 2.8 | 3.2 | 3.4 | 2.9 | 3.1 | 2.6 | 2.7 |
| Relative Survival Rates (SEER) | | | | | | | | | | | | | | | | | | | | | |
| 1-year | 79.9 | 80.5 | 82.5 | 81.9 | 81.7 | 84.0 | 84.0 | 85.9 | 84.9 | 86.1 | 87.1 | 85.8 | 85.6 | 88.5 | 88.8 | 89.3 | 91.0 | 91.2 | 88.6 | 90.1 | |
| 2-year | 66.9 | 70.2 | 71.3 | 73.7 | 72.8 | 74.4 | 74.5 | 76.4 | 76.7 | 76.3 | 78.0 | 76.4 | 76.5 | 80.1 | 80.1 | 81.6 | 82.2 | 84.4 | 81.7 | | |
| 3-year | 61.2 | 63.7 | 65.4 | 69.3 | 66.7 | 68.2 | 70.0 | 71.2 | 72.3 | 71.3 | 73.8 | 72.5 | 71.8 | 76.4 | 75.8 | 77.3 | 78.6 | 80.8 | | | |
| 4-year | 58.7 | 60.1 | 62.1 | 65.6 | 62.9 | 64.5 | 65.6 | 69.8 | 69.7 | 67.1 | 70.6 | 71.0 | 69.8 | 73.2 | 73.1 | 74.8 | 75.9 | | | | |
| 5-year | 55.7 | 57.9 | 59.8 | 63.4 | 61.1 | 62.1 | 64.2 | 67.9 | 67.5 | 66.3 | 69.7 | 68.6 | 69.2 | 72.3 | 71.6 | 73.7 | | | | | |
| 6-year | 53.3 | 56.5 | 58.5 | 61.6 | 59.5 | 60.6 | 63.2 | 66.4 | 66.5 | 65.5 | 68.3 | 67.6 | 67.9 | 71.8 | 70.4 | | | | | | |
| 7-year | 51.5 | 55.2 | 57.5 | 61.0 | 58.3 | 59.8 | 61.7 | 65.8 | 66.2 | 64.4 | 67.2 | 67.3 | 66.6 | 70.8 | | | | | | | |
| 8-year | 50.5 | 54.8 | 57.0 | 60.3 | 57.1 | 58.9 | 61.0 | 65.3 | 65.5 | 63.3 | 65.9 | 67.0 | 66.6 | | | | | | | | |
| 9-year | 50.1 | 54.5 | 56.7 | 59.7 | 57.0 | 58.1 | 60.2 | 65.0 | 65.4 | 62.6 | 65.3 | 66.3 | | | | | | | | | |
| 10-year | 49.6 | 54.2 | 56.5 | 59.4 | 56.7 | 57.6 | 59.9 | 64.3 | 64.9 | 62.1 | 64.4 | | | | | | | | | | |
| 11-year | 49.6 | 54.2 | 56.0 | 58.9 | 56.0 | 57.4 | 59.1 | 64.0 | 64.6 | 61.8 | | | | | | | | | | | |
| 12-year | 49.1 | 53.6 | 55.7 | 58.6 | 55.7 | 57.1 | 58.3 | 63.4 | 63.8 | | | | | | | | | | | | |
| 13-year | 49.1 | 53.3 | 55.6 | 58.6 | 55.3 | 56.9 | 58.3 | 63.4 | | | | | | | | | | | | | |
| 14-year | 49.1 | 53.2 | 54.9 | 57.8 | 54.8 | 56.2 | 58.3 | | | | | | | | | | | | | | |
| 15-year | 49.1 | 52.9 | 54.8 | 57.5 | 54.0 | 56.2 | | | | | | | | | | | | | | | |
| 16-year | 49.1 | 52.0 | 54.1 | 56.9 | 53.9 | | | | | | | | | | | | | | | | |
| 17-year | 48.7 | 51.5 | 53.9 | 56.8 | | | | | | | | | | | | | | | | | |
| 18-year | 48.3 | 51.2 | 53.4 | | | | | | | | | | | | | | | | | | |
| 19-year | 48.3 | 50.4 | | | | | | | | | | | | | | | | | | | |
| 20-year | 48.3 | | | | | | | | | | | | | | | | | | | | |

Note: Incidence and mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Survival rates are relative rates expressed as percents.

Table XXVII-11
ALL SITES (Invasive)

INCIDENCE, MORTALITY, AND SURVIVAL RATES

By Year of Diagnosis/Death

All Races, Males and Females

Ages 0-19

Year of Diagnosis/Death

| | 1975 | 1976 | 1977 | 1978 | 1979 | 1980 | 1981 | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
|---------------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Incidence | | | | | | | | | | | | | | | | | | | | | |
| SEER | 12.8 | 14.1 | 14.1 | 14.4 | 14.5 | 14.2 | 13.8 | 14.4 | 14.4 | 15.4 | 15.7 | 15.7 | 15.2 | 15.0 | 16.5 | 15.5 | 16.2 | 16.1 | 16.0 | 15.6 | 15.1 |
| Mortality | | | | | | | | | | | | | | | | | | | | | |
| U.S. | 5.1 | 5.1 | 5.0 | 4.5 | 4.6 | 4.6 | 4.4 | 4.5 | 4.3 | 3.9 | 3.8 | 3.8 | 3.6 | 3.5 | 3.5 | 3.4 | 3.4 | 3.3 | 3.3 | 3.2 | 3.0 |
| SEER | 5.4 | 5.2 | 4.9 | 4.4 | 5.0 | 4.4 | 4.4 | 4.7 | 4.8 | 3.8 | 4.2 | 4.3 | 3.9 | 3.8 | 3.1 | 3.6 | 3.6 | 3.4 | 3.5 | 3.1 | 3.0 |
| Relative Survival Rates (SEER) | | | | | | | | | | | | | | | | | | | | | |
| 1-year | 81.4 | 82.4 | 84.0 | 83.2 | 84.0 | 86.3 | 84.9 | 86.0 | 87.0 | 87.4 | 87.8 | 88.2 | 87.9 | 90.0 | 89.7 | 90.4 | 91.6 | 90.6 | 89.8 | 90.4 | |
| 2-year | 69.4 | 73.1 | 74.0 | 73.9 | 74.3 | 76.6 | 74.8 | 77.9 | 79.3 | 77.9 | 78.6 | 79.2 | 79.5 | 82.6 | 81.4 | 82.8 | 83.6 | 83.5 | 82.1 | | |
| 3-year | 64.3 | 67.5 | 68.7 | 69.5 | 69.2 | 70.3 | 70.1 | 72.7 | 75.0 | 73.4 | 75.1 | 75.7 | 74.9 | 78.4 | 77.5 | 78.3 | 79.7 | 80.1 | | | |
| 4-year | 61.4 | 63.9 | 65.2 | 66.5 | 66.0 | 67.9 | 66.3 | 70.6 | 72.1 | 69.2 | 72.4 | 73.8 | 72.5 | 75.7 | 74.2 | 75.9 | 77.6 | | | | |
| 5-year | 58.6 | 62.2 | 63.2 | 64.5 | 64.2 | 65.9 | 64.9 | 68.8 | 69.7 | 68.2 | 71.5 | 71.8 | 74.8 | 72.8 | 74.5 | | | | | | |
| 6-year | 56.2 | 60.6 | 61.9 | 63.0 | 62.2 | 64.5 | 63.5 | 67.6 | 68.4 | 66.7 | 70.3 | 71.1 | 70.7 | 74.2 | 71.5 | | | | | | |
| 7-year | 54.9 | 59.3 | 61.2 | 62.4 | 61.1 | 63.9 | 62.6 | 66.9 | 67.8 | 65.8 | 69.1 | 70.6 | 69.5 | 73.2 | | | | | | | |
| 8-year | 54.1 | 58.7 | 60.6 | 61.8 | 59.6 | 63.3 | 61.9 | 66.3 | 66.7 | 64.5 | 68.2 | 70.2 | 69.1 | | | | | | | | |
| 9-year | 53.5 | 58.3 | 59.9 | 61.0 | 59.6 | 62.7 | 61.4 | 65.4 | 66.2 | 63.8 | 67.5 | 69.5 | | | | | | | | | |
| 10-year | 52.9 | 57.8 | 59.4 | 60.8 | 59.4 | 62.3 | 61.0 | 64.8 | 65.8 | 63.3 | 66.7 | | | | | | | | | | |
| 11-year | 52.9 | 57.6 | 59.0 | 60.3 | 58.8 | 62.0 | 60.3 | 64.4 | 65.6 | 63.1 | | | | | | | | | | | |
| 12-year | 52.4 | 57.1 | 58.8 | 60.0 | 58.3 | 61.8 | 59.6 | 64.1 | 64.8 | | | | | | | | | | | | |
| 13-year | 52.4 | 56.9 | 58.7 | 59.8 | 58.0 | 61.5 | 59.6 | 63.9 | | | | | | | | | | | | | |
| 14-year | 52.3 | 56.7 | 58.1 | 59.3 | 57.7 | 60.7 | 59.4 | | | | | | | | | | | | | | |
| 15-year | 52.2 | 56.6 | 57.7 | 59.1 | 57.1 | 60.3 | | | | | | | | | | | | | | | |
| 16-year | 52.2 | 55.9 | 57.1 | 58.6 | 56.8 | | | | | | | | | | | | | | | | |
| 17-year | 51.9 | 55.5 | 57.1 | 58.2 | | | | | | | | | | | | | | | | | |
| 18-year | 51.5 | 55.2 | 56.5 | | | | | | | | | | | | | | | | | | |
| 19-year | 51.5 | 54.6 | | | | | | | | | | | | | | | | | | | |
| 20-year | 51.3 | | | | | | | | | | | | | | | | | | | | |

Note: Incidence and mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.
Survival rates are relative rates expressed as percents.