

Section XXVII
Childhood Cancer by Site
Incidence, Survival and Mortality

The site groupings used in this section correspond to those used in the rest of the Cancer Statistics Review. While these groupings are not as appropriate for children as they are for adults, they are necessary in order to be able to compare incidence to mortality.

Childhood cancer is better categorized by groupings based on histologic type and site rather than primary site alone (see section XXVIII for Childhood Cancer by the International Classification of Childhood Cancer). For the leukemias and lymphomas, the categories are similar, but for solid tumors the categories can be quite different and the different terminology confusing. For example, in this section 'soft tissue' refers to any type of tumor that arises in connective or soft tissue in contrast to 'soft tissue sarcomas' in the other section, which refers to soft tissue sarcomas in any primary site.

Rates are presented per 100,000 children in this section. In contrast in the other childhood cancer section rates are presented per million children.

Table XXVII-1

AGE-ADJUSTED CANCER SEER INCIDENCE TRENDS, 1973-97, 1973-84, 1985-97§

By Primary Cancer Site

All Races, Males and Females

Site	Ages 0-14				Ages 0-19			
	% Change 1973-97	EAPC 73-97	EAPC 73-84	EAPC 85-97	% Change 1973-97	EAPC 73-97	EAPC 73-84	EAPC 85-97
All Sites								
All Races	10.0	0.7★	0.5	0.0	12.9	0.7★	0.8★	0.0♦
Whites	7.5	0.6★	0.3	-0.3	12.0	0.7★	0.9★	-0.2♦
Blacks	14.1	1.1★	1.2	1.1	10.0	0.8★	1.1	0.8
Bone & Joint	1.0	0.6	1.8	-0.4	11.8	0.8	1.8	0.0
Brain & Other nervous	17.9	1.4★	0.4	-0.4	21.4	1.3★	0.5	-0.4
Hodgkin's disease	-10.8	-1.3★	-1.6	-1.9	6.9	-0.4	0.8	-1.2
Kidney & Renal pelvis	33.0	0.6	2.3	0.5	30.8	0.7	2.1	0.6
Leukemias	0.0	0.4	0.0	-0.4	-2.9	0.4	0.1	-0.2
Acute lymphocytic	24.4	1.0★	2.1	-0.1◊	20.7	1.0★	2.3★	-0.2♦
Non-Hodgkin's lymphomas	0.3	0.2	1.0	-0.2	30.0	1.1★	2.3★	0.6◊
Soft tissue	41.0	0.9	1.1	1.5	32.6	0.9★	1.4	0.8

The EAPC is the Estimated Annual Percent Change over the time interval.
SEER Program.

The EAPC is significantly different from zero ($p<.05$).

The EAPC for 1985-97 is significantly different from the EAPC for 1973-84 ($p<.05$).

The EAPC for 1985-97 is significantly different from the EAPC for 1973-84 ($p<.10$).

Statistic could not be calculated.

§♦◊-

Table XXVII-2

AGE-ADJUSTED CANCER U.S. MORTALITY TRENDS, 1973-97, 1973-84, 1985-97

By Race and Primary Cancer Site

Males and Females

Race/Site	Ages 0-14			Ages 0-19				
	% Change 1973-97	EAPC 73-97	EAPC 73-84	EAPC 85-97	% Change 1973-97	EAPC 73-97	EAPC 73-84	EAPC 85-97
<u>All Races</u>								
All Sites	-50.4	-2.9★	-3.1★	-2.5★◊	-48.2	-2.7★	-2.9★	-2.3★♦
Bone & Joint	-58.5	-3.1★	-3.2★	-2.2	-46.7	-2.5★	-3.1★	-1.1♦
Brain & Other nervous	-23.9	-1.1★	-1.5★	-0.7	-24.1	-1.1★	-1.5★	-0.9★
Hodgkin's disease	-74.0	-6.3★	-9.1★	-5.7	-71.7	-5.0★	-4.4★	-5.6★
Kidney & Renal pelvis	-58.6	-3.7★	-4.5★	-3.3★♦	-53.8	-3.4★	-4.1★	-2.5★
Leukemias	-60.4	-3.9★	-4.8★	-3.3★♦	-57.1	-3.5★	-4.1★	-3.0★♦
Acute lymphocytic	-66.1	-4.9★	-4.5★	-4.2★	-62.0	-4.3★	-3.7★	-3.8★
Non-Hodgkin's lymphomas	-71.6	-5.0★	-5.9★	-5.2★	-64.9	-4.1★	-4.9★	-4.3★
Soft tissue	-7.6	-0.7	3.0★	-1.8♦	3.0	0.0	2.9★	-0.8♦
<u>Whites</u>								
All Sites	-50.9	-3.0★	-3.1★	-2.7★	-48.9	-2.8★	-2.9★	-2.5★
Bone & Joint	-55.4	-2.9★	-2.9★	-2.0	-45.0	-2.3★	-2.8★	-1.0♦
Brain & Other nervous	-20.8	-1.0★	-1.2★	-0.9	-21.5	-1.1★	-1.2★	-1.0★
Hodgkin's disease	-76.8	-6.7★	-10.0★	-6.5	-76.1	-5.5★	-5.1★	-6.9★
Kidney & Renal pelvis	-60.4	-3.7★	-4.6★	-3.1★	-55.2	-3.4★	-4.0★	-2.4★
Leukemias	-61.6	-4.1★	-5.0★	-3.5★♦	-58.0	-3.7★	-4.3★	-3.1★♦
Acute lymphocytic	-67.4	-5.0★	-4.8★	-4.0★	-63.0	-4.4★	-3.9★	-3.7★
Non-Hodgkin's lymphomas	-74.0	-5.1★	-5.7★	-5.8★	-65.5	-4.0★	-4.7★	-4.6★
Soft tissue	-11.7	-0.9	3.0★	-2.5★♦	-1.0	-0.1	3.0★	-1.2♦
<u>Blacks</u>								
All Sites	-45.4	-2.5★	-2.3★	-1.8★	-42.2	-2.2★	-2.6★	-1.5★♦
Bone & Joint	-65.6	-3.5★	-3.9★	-1.7	-43.4	-3.0★	-4.8★	-0.4♦
Brain & Other nervous	-35.7	-1.5★	-3.1★	-0.3♦	-33.4	-1.3★	-3.3★	-0.2♦
Hodgkin's disease	-51.3	-	-7.4	-	-41.2	-2.5★	-0.8	-2.1
Kidney & Renal pelvis	-49.3	-3.6★	-4.5★	-3.0	-46.3	-3.2★	-5.2★	-1.8
Leukemias	-52.5	-3.2★	-2.7★	-2.4★	-49.2	-2.6★	-2.3★	-2.1★
Acute lymphocytic	-52.0	-4.0★	-0.6	-5.9★♦	-48.6	-3.2★	-0.3	-4.9★♦
Non-Hodgkin's lymphomas	-51.1	-4.8★	-6.8★	-3.0	-55.8	-4.1★	-5.1★	-2.3
Soft tissue	14.5	0.0	3.9	-0.8	20.9	0.2	2.6	-0.5

The EAPC is the Estimated Annual Percent Change over the time interval.

The EAPC is significantly different from zero ($p<.05$).The EAPC for 1985-97 is significantly different from the EAPC for 1973-84 ($p<.05$).The EAPC for 1985-97 is significantly different from the EAPC for 1973-84 ($p<.10$).

Statistic could not be calculated.

★

♦

◊

-

Table XXVII-3

AGE-ADJUSTED CANCER SEER INCIDENCE RATES\$, 1975-97

By Primary Cancer Site

All Races, Males and Females

Year of Diagnosis

Age/Site	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
<u>Ages 0-14</u>																							
All Sites																							
All Races	11.3	12.9	12.7	13.0	13.0	12.7	12.3	12.8	12.8	13.8	14.3	14.0	14.0	13.2	14.8	14.0	14.9	13.4	14.7	14.0	13.8	14.5	13.7
Whites	11.8	13.3	13.4	13.5	13.5	12.8	12.3	12.9	13.4	14.2	15.3	14.7	14.4	13.7	15.3	14.0	15.8	13.8	14.8	14.3	14.3	14.4	14.1
Blacks	8.5	11.1	8.3	9.8	10.6	11.0	13.0	11.8	9.9	10.3	9.9	10.6	11.4	11.8	13.6	12.2	11.7	12.4	12.7	12.9	11.8	11.7	12.0
Bone & Joint	0.5	0.5	0.6	0.7	0.8	0.7	0.6	0.7	0.7	0.8	0.7	0.7	0.7	0.6	0.6	0.7	0.9	0.6	0.6	0.8	0.8	0.5	0.9
Brain & Other nervous	2.3	2.4	3.0	2.7	2.8	2.8	2.5	2.4	2.2	2.8	2.9	3.4	3.1	3.3	3.2	3.4	3.4	3.2	3.4	3.3	3.3	3.0	2.7
Hodgkin's disease	0.7	0.8	0.7	0.7	0.9	0.5	0.6	0.5	0.7	0.7	0.7	0.5	0.7	0.6	0.8	0.7	0.6	0.5	0.6	0.6	0.3	0.7	0.6
Kidney & Renal pelvis	0.6	0.9	0.7	0.9	0.9	0.8	0.9	0.9	1.1	0.7	0.8	0.9	0.8	0.7	1.1	0.7	0.9	0.7	0.9	1.1	0.6	0.9	0.9
Leukemias	3.2	4.0	3.9	3.7	3.6	3.9	3.9	3.7	3.6	4.4	4.5	4.0	4.3	4.0	4.6	4.3	4.5	3.9	4.1	3.5	4.3	4.5	4.1
Acute lymphocytic	2.1	2.9	3.0	2.9	2.7	3.0	3.1	2.9	2.9	3.4	3.4	3.2	3.3	3.2	3.8	3.3	3.4	3.0	3.0	2.8	3.5	3.4	3.4
Non-Hodgkin's lymph	1.0	0.8	0.7	1.0	0.8	0.7	0.7	0.9	0.9	0.9	0.7	1.0	1.1	0.7	0.8	0.8	0.9	0.8	0.8	1.0	0.9	0.7	0.9
Soft tissue	0.8	0.7	0.7	0.8	0.6	0.9	0.7	0.9	0.9	0.7	0.8	0.9	0.8	0.6	0.9	0.8	1.0	0.7	0.6	0.9	1.0	1.1	0.9
<u>Ages 0-19</u>																							
All Sites																							
All Races	12.8	14.1	14.1	14.4	14.5	14.2	13.8	14.4	14.4	15.4	15.7	15.7	15.2	15.0	16.5	15.5	16.2	16.0	16.0	15.6	15.6	15.9	15.2
Whites	13.2	14.7	14.7	15.0	15.0	14.6	14.0	14.9	15.2	15.9	16.8	16.5	16.0	15.5	17.1	15.7	17.2	16.7	16.2	16.0	16.3	16.0	16.0
Blacks	10.7	10.7	10.4	10.9	12.7	11.0	12.9	11.8	11.1	12.1	10.5	12.1	11.2	12.9	14.0	12.9	12.4	12.8	13.3	13.5	12.5	12.8	11.6
Bone & Joint	0.7	0.6	0.7	0.9	0.9	1.0	0.8	0.8	0.9	1.0	0.9	1.0	0.8	0.9	0.8	1.0	0.9	0.9	1.0	0.9	0.7	1.2	
Brain & Other nervous	2.1	2.2	2.9	2.5	2.5	2.4	2.3	2.3	2.2	2.6	2.6	3.2	2.9	3.0	2.9	3.1	3.0	3.2	3.0	2.9	2.9	2.9	2.5
Hodgkin's disease	1.5	1.4	1.4	1.5	1.6	1.2	1.3	1.5	1.5	1.4	1.3	1.4	1.4	1.7	1.3	1.4	1.4	1.3	1.3	1.0	1.4	1.3	
Kidney & Renal pelvis	0.5	0.7	0.6	0.7	0.7	0.6	0.7	0.7	0.8	0.6	0.7	0.7	0.7	0.6	0.8	0.6	0.7	0.6	0.8	0.9	0.5	0.7	0.7
Leukemias	3.0	3.6	3.4	3.4	3.3	3.4	3.6	3.4	3.4	3.9	4.0	3.6	3.6	3.5	4.1	3.7	4.0	3.6	3.6	3.3	3.9	3.9	3.6
Acute lymphocytic	1.9	2.4	2.5	2.5	2.3	2.4	2.7	2.5	2.5	2.9	2.8	2.7	2.7	2.7	3.3	2.8	2.9	2.5	2.5	2.4	3.0	2.8	2.8
Non-Hodgkin's lymph	1.0	0.9	0.8	1.0	0.9	0.9	1.0	1.0	1.1	1.0	1.0	1.1	1.0	1.0	1.0	1.1	1.0	1.0	0.9	1.3	1.2	1.0	1.1
Soft tissue	0.8	0.7	0.7	0.9	0.7	0.9	0.7	1.0	1.0	0.7	0.9	0.9	0.7	0.9	0.8	1.0	0.7	0.7	0.8	1.1	1.1	1.0	

§ SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-4

AGE-ADJUSTED CANCER U.S. MORTALITY RATES*, 1975-97

By Primary Cancer Site

All Races, Males and Females

Year of Diagnosis

<u>Age/Site</u>	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
<u>Ages 0-14</u>																							
All Sites																							
All Sites	4.9	4.9	4.7	4.3	4.4	4.3	4.2	4.2	4.1	3.6	3.6	3.5	3.4	3.3	3.3	3.1	3.1	3.0	3.0	2.8	2.7	2.7	2.7
Bone & Joint	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.2	0.1	0.1
Brain & Other nervous	1.0	1.0	1.0	0.8	0.9	0.9	0.9	1.0	0.9	0.8	0.8	0.9	0.7	0.8	0.9	0.9	0.9	0.8	0.8	0.8	0.7	0.8	0.7
Hodgkin's disease	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Kidney & Renal pelvis	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Leukemias	2.1	2.1	2.0	1.9	1.8	1.7	1.6	1.6	1.5	1.3	1.4	1.3	1.3	1.2	1.2	1.1	1.1	1.1	1.1	1.0	1.0	0.9	0.9
Acute lymphocytic	1.2	1.2	1.1	1.0	1.0	0.9	1.0	0.9	0.8	0.8	0.8	0.7	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.5	0.4	0.4
Non-Hodgkin's lymph	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.1
Soft tissue	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.2
<u>Ages 0-19</u>																							
All Sites	5.1	5.1	5.0	4.5	4.6	4.6	4.4	4.5	4.3	3.9	3.8	3.8	3.6	3.5	3.5	3.4	3.4	3.3	3.3	3.2	3.0	2.9	2.9
Bone & Joint	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Brain & Other nervous	0.9	0.9	0.9	0.8	0.9	0.8	0.8	0.9	0.8	0.8	0.8	0.7	0.8	0.8	0.8	0.8	0.7	0.8	0.7	0.7	0.8	0.7	0.7
Hodgkin's disease	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.1	0.0	0.0	0.0
Kidney & Renal pelvis	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Leukemias	2.0	2.1	2.0	1.8	1.8	1.8	1.6	1.6	1.5	1.4	1.4	1.3	1.4	1.2	1.2	1.2	1.2	1.2	1.1	1.1	1.0	1.0	1.0
Acute lymphocytic	1.0	1.1	1.0	0.9	1.0	0.9	0.9	0.9	0.8	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.5	0.6	0.5	0.5	0.4	0.4	0.4
Non-Hodgkin's lymph	0.4	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Soft tissue	0.2	0.2	0.2	0.2	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2

* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-5

AGE-ADJUSTED CANCER U.S. MORTALITY RATES*, 1975-97

By Primary Cancer Site

Whites, Males and Females

Year of Diagnosis

<u>Age/Site</u>	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
<u>Ages 0-14</u>																							
All Sites																							
All Sites	5.0	5.0	4.8	4.3	4.5	4.4	4.2	4.3	4.1	3.7	3.6	3.6	3.5	3.3	3.4	3.2	3.2	3.1	3.1	2.8	2.7	2.7	2.7
Bone & Joint	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1
Brain & Other nervous	1.0	1.0	1.0	0.8	0.9	0.9	0.9	1.0	0.9	0.9	0.9	0.9	0.8	0.8	0.9	0.9	0.9	0.8	0.8	0.8	0.7	0.8	0.8
Hodgkin's disease	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Kidney & Renal pelvis	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Leukemias	2.3	2.3	2.1	1.9	1.9	1.8	1.7	1.6	1.6	1.4	1.4	1.4	1.4	1.2	1.2	1.1	1.1	1.2	1.1	1.0	1.0	0.9	0.9
Acute lymphocytic	1.2	1.3	1.2	1.1	1.1	1.0	1.0	0.9	0.9	0.8	0.8	0.7	0.7	0.6	0.6	0.6	0.5	0.6	0.5	0.5	0.5	0.4	0.5
Non-Hodgkin's lymph	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1
Soft tissue	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.2
<u>Ages 0-19</u>																							
All Sites	5.2	5.2	5.1	4.6	4.7	4.7	4.4	4.5	4.4	4.0	3.9	3.9	3.7	3.6	3.6	3.4	3.5	3.4	3.3	3.2	3.0	2.9	2.9
Bone & Joint	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.2
Brain & Other nervous	0.9	0.9	0.9	0.8	0.9	0.8	0.8	0.9	0.9	0.8	0.8	0.7	0.8	0.8	0.8	0.8	0.8	0.7	0.7	0.8	0.7	0.8	0.7
Hodgkin's disease	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0
Kidney & Renal pelvis	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Leukemias	2.1	2.2	2.1	1.9	1.9	1.8	1.7	1.7	1.6	1.4	1.4	1.4	1.4	1.3	1.2	1.2	1.2	1.2	1.2	1.1	1.0	1.0	1.0
Acute lymphocytic	1.1	1.1	1.1	1.0	1.0	1.0	0.9	0.9	0.8	0.8	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.5
Non-Hodgkin's lymph	0.5	0.4	0.4	0.3	0.4	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Soft tissue	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2

* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-6

AGE-ADJUSTED CANCER U.S. MORTALITY RATES*, 1975-97

By Primary Cancer Site

Blacks, Males and Females

Year of Diagnosis

<u>Age/Site</u>	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
<u>Ages 0-14</u>																							
All Sites																							
All Sites	4.5	4.2	4.2	4.1	4.0	3.8	4.2	4.0	3.9	3.5	3.3	3.4	3.0	3.2	3.2	3.0	3.1	2.7	2.7	2.9	3.0	2.6	2.6
Bone & Joint	0.3	0.3	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1
Brain & Other nervous	0.9	1.0	1.0	0.8	0.9	0.9	1.0	0.8	0.8	0.7	0.6	0.9	0.7	1.0	0.8	0.7	0.8	0.7	0.6	0.8	0.8	0.8	0.7
Hodgkin's disease	0.1	0.0	0.1	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0
Kidney & Renal pelvis	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.2
Leukemias	1.6	1.6	1.4	1.5	1.3	1.2	1.6	1.5	1.2	1.3	1.2	1.1	1.0	0.9	1.0	0.9	1.0	0.8	0.9	0.8	1.1	0.8	0.8
Acute lymphocytic	0.8	0.8	0.7	0.6	0.6	0.6	0.8	0.7	0.7	0.7	0.7	0.6	0.5	0.4	0.5	0.5	0.4	0.4	0.3	0.3	0.4	0.4	0.3
Non-Hodgkin's lymph	0.4	0.2	0.3	0.4	0.3	0.2	0.2	0.2	0.2	0.1	0.2	0.3	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1
Soft tissue	0.2	0.2	0.2	0.2	0.3	0.1	0.3	0.3	0.3	0.2	0.2	0.1	0.3	0.2	0.2	0.3	0.1	0.2	0.2	0.3	0.1	0.2	0.2
<u>Ages 0-19</u>																							
All Sites	4.7	4.5	4.7	4.5	4.3	4.2	4.5	4.3	3.8	3.7	3.5	3.7	3.6	3.5	3.5	3.4	3.4	3.0	3.1	3.3	3.3	3.0	3.0
Bone & Joint	0.4	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.1
Brain & Other nervous	0.9	0.9	0.9	0.7	0.9	0.8	0.9	0.8	0.7	0.7	0.6	0.8	0.7	0.9	0.7	0.7	0.7	0.7	0.7	0.8	0.7	0.7	0.7
Hodgkin's disease	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Kidney & Renal pelvis	0.2	0.3	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1
Leukemias	1.6	1.5	1.4	1.5	1.4	1.4	1.5	1.5	1.1	1.3	1.1	1.1	1.2	1.0	1.1	1.1	1.0	1.1	0.9	1.0	1.0	1.1	0.8
Acute lymphocytic	0.7	0.7	0.7	0.6	0.6	0.6	0.7	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.3	0.3	0.3	0.5	0.4	0.3
Non-Hodgkin's lymph	0.4	0.3	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.3	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2
Soft tissue	0.2	0.2	0.3	0.2	0.3	0.2	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.3	0.2	0.2	0.3

* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-7

AGE-ADJUSTED CANCER SEER INCIDENCES AND U.S. MORTALITY* RATES, 1993-97

By Primary Cancer Site

All Races, Males and Females

Site	Ages 0-14						Ages 0-19					
	Incidence			Mortality			Incidence			Mortality		
	Total	Males	Females									
All Sites												
All Races	14.2	15.2	13.0	2.8	3.1	2.5	15.7	16.6	14.7	3.1	3.4	2.7
Whites	14.4	15.3	13.4	2.8	3.1	2.5	16.1	17.0	15.2	3.1	3.5	2.6
Blacks	12.2	13.4	11.0	2.8	2.9	2.7	12.7	13.5	11.9	3.1	3.3	3.0
Bone & Joint	0.7	0.9	0.6	0.1	0.1	0.1	0.9	1.2	0.7	0.2	0.2	0.2
Brain & Other nervous	3.2	3.5	2.8	0.8	0.8	0.7	2.9	3.2	2.5	0.7	0.8	0.7
Hodgkin's disease	0.6	0.5	0.6	0.0	0.0	0.0	1.2	1.2	1.4	0.0	0.0	0.0
Kidney & Renal pelvis	0.9	0.9	1.0	0.1	0.1	0.1	0.7	0.7	0.8	0.1	0.1	0.1
Leukemias	4.1	4.5	3.7	1.0	1.1	0.8	3.7	4.0	3.3	1.0	1.2	0.9
Acute lymphocytic	3.2	3.5	2.9	0.5	0.5	0.4	2.7	3.0	2.4	0.5	0.5	0.4
Non-Hodgkin's lymphomas	0.9	1.2	0.5	0.1	0.2	0.1	1.1	1.5	0.6	0.2	0.2	0.1
Soft tissue	0.9	0.9	0.9	0.1	0.2	0.1	0.9	0.9	0.9	0.2	0.2	0.2

§ SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-8

5-YEAR RELATIVE SURVIVAL RATES (Percent)

By Selected Primary Cancer Site and Year Of Diagnosis

All Races, Males and Females

Site	Ages 0-14						Ages 0-19					
	1974-76	1977-79	1980-82	1983-85	1986-88	1989-96	1974-76	1977-79	1980-82	1983-85	1986-88	1989-96
All Sites												
All Races	55.7	61.6	65.1	67.7	70.1	75.3S	58.5	64.0	66.6	69.7	72.9	75.7S
Whites	55.8	62.0	65.5	69.4	71.4	76.5S	58.8	64.5	67.4	71.4	74.5	77.1S
Blacks	56.8	58.0	61.3	57.4	62.4	69.6S	56.9	60.6	61.2	58.8	62.7	68.5S
Bone & Joint	53.3↑	53.3↑	54.6↑	56.6↑	63.2↑	68.7S	50.2	50.6	50.1	52.4	62.1	66.4S
Brain & Other nervous	54.7	56.1	55.4	61.7	62.8	65.1S	55.6	56.5	56.8	63.2	65.6	66.4S
Hodgkin's disease	78.3	83.5	90.9	89.6	89.9	94.2S	84.8	88.3	87.0	88.2	89.3	93.4S
Leukemias	45.3	59.3	62.7	61.1	70.2	74.9S	41.0	53.7	57.9	58.2	66.2	70.5S
Acute lymphocytic	53.2	67.0	71.1	69.3	78.3	81.9S	49.3	62.8	67.3	66.7	75.4	78.7S
Acute myeloid	14.7	29.5↑	21.1↑	31.5↑	32.2↑	44.3S	16.6	28.3	22.1	34.4↑	32.1↑	42.0S
Neuroblastoma	52.6	53.8	52.9	55.0	59.4	70.1S	52.2	53.8	53.1	54.2	58.7	69.2S
Non-Hodgkin's lymphomas	43.9	50.2	61.5	71.2	69.6	78.3S	43.3	49.5	59.0	67.0	73.1	74.0S
Soft tissue	60.7	68.9	64.6	75.8	66.3	76.4S	62.0	65.2	64.3	74.1	68.2	70.7S
Wilms' tumor	74.4	77.5	87.1	86.5	90.8	93.2S	73.6	77.1	86.5	86.8	90.9	92.9S

Rates are from the SEER Program. They are based on data from population-based registries in Connecticut, New Mexico, Utah, Iowa, Hawaii, Atlanta, Detroit, Seattle-Puget Sound and San Francisco-Oakland. Rates are based on follow-up of patients through 1997.

S The difference in rates between 1974-76 and 1989-96 is statistically significant ($p<.05$).

↑ The standard error of the survival rate is between 5 and 10 percentage points.

↑↑ The standard error of the survival rate is greater than 10 percentage points.

Table XXVII-9

CHILDHOOD CANCER
SEER Incidence, U.S. Mortality and 5-Year Relative Survival Rates
For Top Cancer Sites[⊕] By 5-Year Age Groups
All Races, Males and Females

Age/Site	SEER Incidences					US Mortality*					Survivals	
	1993-97 Num	1993-97 Cases	73-97 Rate	73-84 EAPC	85-97 EAPC	1993-97 Num	1993-97 Cases	73-97 Rate	73-84 EAPC	85-97 EAPC	1974-76	1989-96
<u>Ages 0-4</u>												
All Sites	1,899	20.6	0.8★	0.5	0.3	2,785	2.9	-3.0★♦	-2.9★♦	-2.7★	56.5	74.7\$
Soft tissue	124	1.3	0.8	3.1	-0.6	132	0.1	-2.2★♦	0.5♦	-3.6	57.7↑	82.1\$
Kidney and Renal pelvis	203	2.2	0.7	2.4	2.0	106	0.1	-4.9★♦	-6.5★♦	-1.8♦	74.2	91.9\$
Eye and Orbit	123	1.3	0.7	1.7	-0.8	32	0.0	-5.4★	-6.8★	-10.7★	89.6	94.5
Brain and Other nervous	352	3.8	2.1★	2.9	-0.5♦	749	0.8	-1.0★	-1.3★	0.1△	51.4	56.2
Other endocrine	132	1.4	-	-	-	494	0.5	-	-	-	32.8↑	61.4\$
Leukemias	646	7.0	0.3	-1.3	-0.2	899	0.9	-3.5★	-4.7★	-2.8★♦	51.2	77.1\$
Acute lymphocytic	521	5.7	0.9★	0.9	-0.1	294	0.3	-5.3★	-6.1★	-4.4★	58.4	84.2\$
<u>Ages 5-9</u>												
All Sites	944	10.5	0.4	0.4	-1.2★△	2,718	2.8	-3.2★♦	-3.6★♦	-2.7★△	55.4	75.6\$
Brain and Other nervous	267	3.0	0.7	-0.3	-2.0★	865	0.9	-1.0★	-1.6	-1.1	54.4	67.5\$
Leukemias	305	3.4	0.6	0.2	-0.5	921	1.0	-5.1★	-6.0★	-4.4★♦	47.0	77.5\$
Acute lymphocytic	258	2.9	1.3★	2.8	-0.1	491	0.5	-6.0★	-5.7★	-4.8★	53.8	83.0\$
<u>Ages 10-14</u>												
All Sites	1,096	12.3	0.7★	0.4	0.6	2,508	2.7	-2.5★♦	-2.6★♦	-2.2★	54.9	76.1\$
Bone and Joint	120	1.3	0.3	0.3	-1.2	236	0.3	-3.1★♦	-3.2★♦	-2.0	51.9↑	68.2\$
Brain and Other nervous	248	2.8	1.4★	-1.7	1.8△	624	0.7	-1.3★	-1.4	-0.9	58.7	75.8\$
Hodgkin's disease	104	1.2	-1.0★	-1.3	-1.5	32	0.0	-5.8★	-7.3★	-5.4	74.9↑	95.0\$
Leukemias	211	2.4	0.5	2.7	-0.3	964	1.0	-2.9★	-3.3★	-2.8★	28.8	63.6\$
Acute lymphocytic	139	1.6	1.1	4.2	0.1	486	0.5	-3.2★	-1.9★	-3.4★	37.2↑	71.8\$
<u>Ages 15-19</u>												
All Sites	1,745	20.3	0.9★	1.6★	0.1♦	3,522	3.9	-2.2★♦	-2.4★♦	-1.7★♦	63.8	76.5\$
Bone and Joint	142	1.7	1.0★	1.9	0.5	476	0.5	-2.1★♦	-3.1★♦	-0.4♦	46.6↑	63.3\$
Melanomas of skin	120	1.4	1.8★	0.6	-1.3	44	0.0	-4.2★	-4.9★	-5.3	76.7↑	93.6\$
Ovary(females)	98	2.3	2.7★	-3.5	0.9	32	0.1	-6.9★	-5.7★	-8.8★	69.7↑	90.8\$
Testis(males)	133	3.0	2.2★	7.3★	2.1	49	0.1	-7.7★	-9.6★	-3.0♦	55.3↑	92.1\$
Brain and Other nervous	165	1.9	0.9	0.9	-0.7	533	0.6	-1.1★	-1.6★	-1.4★	58.9↑	73.5\$
Thyroid	129	1.5	0.5	1.5	-0.5	-	-	-	-	-	100.0	99.5
Hodgkin's disease	289	3.4	0.1	2.2	-0.9♦	121	0.1	-4.8★	-2.7★	-6.2★△	88.4	93.0
Non-Hodgkin's lymphomas	151	1.8	2.8★♦	5.4★♦	2.2	293	0.3	-2.6★△	-3.3★△	-3.0★	41.7↑	66.9\$
Leukemias	198	2.3	0.0	0.7	0.9	1,089	1.2	-2.2★	-2.0★	-2.1★	21.5	45.0\$
Acute lymphocytic	95	1.1	0.5	4.0★	-0.6♦	441	0.5	-2.0★	0.1	-2.6★♦	21.2↑	49.8\$

Note: The EAPC is the Estimated Annual Percent Change over the time interval.

Incidence and mortality rates are per 100,000. Survival rates are expressed as percents.

SEER Program.

NCHS public use tape.

The EAPC is significantly different from zero ($p<.05$).

The EAPC for 1985-97 is significantly different from the EAPC for 1973-84 ($p<.05$).

The EAPC for 1985-97 is significantly different from the EAPC for 1973-84 ($p<.10$).

The difference in rates between 1974-76 and 1989-96 is statistically significant ($p<.05$).

The standard error of the survival rate is between 5 and 10 percentage points.

The standard error of the survival rate is greater than 10 percentage points.

Sites were chosen based on more than 100 SEER incident cases between 1993-97.

Statistic could not be calculated.

\$

*

♦

△

+

-

Table XXVII-10
ALL SITES (Invasive)

INCIDENCE, MORTALITY, AND SURVIVAL RATES

By Year of Diagnosis/Death

All Races, Males and Females

Ages 0-14

Year of Diagnosis/Death

	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	
Incidence																							
SEER	12.9	12.7	13.0	13.0	12.7	12.3	12.8	12.8	13.8	14.3	14.0	14.0	14.0	13.2	14.8	14.0	14.9	13.4	14.7	14.0	13.8	14.5	13.7
Mortality																							
U.S.	4.9	4.7	4.3	4.4	4.3	4.2	4.2	4.1	3.6	3.6	3.5	3.4	3.3	3.3	3.1	3.1	3.0	3.0	2.8	2.7	2.7	2.7	2.7
SEER	4.7	4.9	4.3	4.6	4.2	4.1	4.3	4.5	3.6	4.0	4.1	3.8	3.6	2.8	3.2	3.4	2.8	3.1	2.6	2.7	2.5	2.5	2.5
Relative Survival Rates (SEER)																							
1-year	80.5	82.3	81.9	81.9	84.1	84.2	85.8	84.6	86.2	87.1	85.7	85.8	88.6	88.6	89.2	90.9	90.9	88.4	90.6	90.0	92.8		
2-year	70.4	71.2	73.7	73.2	75.0	74.8	76.3	76.5	75.7	78.2	76.3	76.6	80.1	80.0	81.5	82.1	83.8	81.7	84.6	84.0			
3-year	64.0	65.0	69.4	67.2	68.6	70.3	71.1	72.2	70.7	74.0	72.3	72.3	76.5	75.5	77.1	78.5	80.4	78.0	81.3				
4-year	60.4	61.8	65.5	63.5	65.0	66.0	69.7	69.6	66.8	70.5	70.8	70.2	73.2	73.0	74.7	75.6	78.2	77.2					
5-year	58.2	59.5	63.5	61.8	62.5	64.5	68.1	67.4	66.0	69.7	68.6	69.6	72.3	71.5	73.6	74.4	76.9						
6-year	56.7	58.2	61.7	60.2	60.7	63.5	66.5	66.4	65.3	68.3	67.6	68.2	71.9	70.4	72.6	73.8							
7-year	55.4	57.0	61.0	59.0	59.6	61.9	65.8	66.1	64.1	67.3	67.4	66.9	71.0	68.8	71.4								
8-year	55.0	56.6	60.4	57.9	58.7	61.2	65.4	65.5	63.0	66.0	67.1	66.9	70.3	67.7									
9-year	54.7	56.3	59.6	57.8	57.7	60.3	65.1	65.3	62.4	65.4	66.5	66.6	70.0										
10-year	54.4	56.1	59.3	57.5	57.3	60.0	64.4	64.9	61.8	64.7	66.3	65.9											
11-year	54.4	55.6	58.8	56.8	57.1	59.2	64.1	64.4	61.5	63.8	66.3												
12-year	53.8	55.3	58.5	56.5	56.8	58.3	63.5	63.7	61.2	63.5													
13-year	53.5	55.2	58.5	56.2	56.7	58.3	63.3	63.2	60.2														
14-year	53.3	54.5	57.7	55.7	56.0	58.3	63.2	62.6															
15-year	53.0	54.4	57.4	55.0	56.0	58.3	63.0																
16-year	52.2	53.7	56.9	54.8	55.5	58.3																	
17-year	51.7	53.6	56.7	54.3	55.1																		
18-year	51.4	53.0	56.7	54.1																			
19-year	50.7	52.7	56.3																				
20-year	50.3	52.5																					
21-year	49.9																						

Note: Incidence and mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Survival rates are relative rates expressed as percents.

Table XXVII-11
ALL SITES (Invasive)

INCIDENCE, MORTALITY, AND SURVIVAL RATES

By Year of Diagnosis/Death

All Races, Males and Females

Ages 0-19

Year of Diagnosis/Death

	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
Incidence																						
SEER	14.1	14.1	14.4	14.5	14.2	13.8	14.4	14.4	15.4	15.7	15.7	15.2	15.0	16.5	15.5	16.2	16.0	16.0	15.6	15.6	15.9	15.2
Mortality																						
U.S.	5.1	5.0	4.5	4.6	4.6	4.4	4.5	4.3	3.9	3.8	3.8	3.6	3.5	3.5	3.4	3.4	3.3	3.3	3.2	3.0	2.9	2.9
SEER	5.2	4.9	4.4	5.0	4.4	4.4	4.7	4.8	3.8	4.2	4.3	3.9	3.8	3.1	3.6	3.5	3.3	3.5	3.1	2.9	2.7	2.7
Relative Survival Rates (SEER)																						
1-year	82.3	83.9	83.2	83.9	86.2	85.0	85.7	86.7	87.4	87.9	88.0	87.9	90.1	89.6	90.4	91.3	90.3	89.7	90.8	89.9	93.2	
2-year	73.1	73.9	73.9	74.6	76.8	75.0	77.6	79.2	77.5	78.8	79.0	79.6	82.4	81.3	82.8	83.3	83.1	82.3	84.3	84.1		
3-year	67.6	68.4	69.6	69.5	70.4	70.1	72.4	75.0	72.9	75.3	75.5	75.2	78.4	77.3	78.3	79.4	80.0	78.9	80.6			
4-year	64.1	65.0	66.6	66.2	68.1	66.4	70.5	72.1	68.9	72.4	73.7	72.6	75.6	74.1	75.8	77.1	77.8	77.4				
5-year	62.4	63.0	64.7	64.4	66.0	64.9	68.7	69.7	71.4	71.8	72.0	74.7	72.8	74.5	75.7	76.4						
6-year	60.6	61.8	63.2	62.4	64.3	63.5	67.4	68.5	66.5	70.2	71.1	70.9	74.1	71.6	73.3	75.0						
7-year	59.4	61.0	62.6	61.3	63.5	62.5	66.8	67.9	65.6	69.1	70.7	69.7	73.2	70.5	72.2							
8-year	58.8	60.4	62.0	59.8	62.9	61.9	66.2	66.8	64.3	68.2	70.3	69.3	72.1	69.8								
9-year	58.4	59.6	61.1	59.7	62.2	61.2	65.3	66.3	63.6	67.6	69.7	68.7	71.9									
10-year	58.0	59.1	60.9	59.6	61.8	60.9	64.7	65.9	63.0	66.9	69.3	68.2										
11-year	57.7	58.7	60.4	59.0	61.5	60.2	64.3	65.6	62.9	66.1	69.1											
12-year	57.2	58.6	60.1	58.5	61.3	59.4	63.9	64.9	62.6	65.7												
13-year	57.0	58.5	59.9	58.2	61.0	59.4	63.6	64.5	61.8													
14-year	56.8	57.9	59.4	57.9	60.3	59.2	63.1	64.1														
15-year	56.7	57.5	59.3	57.4	60.0	59.0	63.1															
16-year	56.0	56.9	58.7	57.0	59.0	59.0																
17-year	55.6	56.8	58.4	56.6	58.7																	
18-year	55.3	56.3	58.0	56.6																		
19-year	54.8	56.1	57.6																			
20-year	54.2	55.9																				
21-year	53.6																					

Note: Incidence and mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Survival rates are relative rates expressed as percents.