

Table VII-1
CORPUS AND UTERUS, NOS CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
WITH UP TO THREE JOINPOINTS BY RACE AND AGE

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	All Ages	1975-1979	-6.0 ^c	1979-1988	-1.7 ^c	1988-1998	0.7 ^c	1998-2000	-2.4
All Races	Under 50	1975-1984	-3.6 ^c	1984-2000	0.5				
All Races	50 and Over	1975-1979	-6.1 ^c	1979-1988	-1.7 ^c	1988-1998	0.6 ^c	1998-2000	-3.2
White	All Ages	1975-1979	-6.2 ^c	1979-1988	-1.6 ^c	1988-2000	0.4 ^c		
White	Under 50	1975-1983	-3.9 ^c	1983-2000	0.1				
White	50 and Over	1975-1979	-6.2 ^c	1979-1988	-1.7 ^c	1988-1998	0.7 ^c	1998-2000	-3.1
Black	All Ages	1975-2000	-0.1						
Black	Under 50	1975-2000	-0.5						
Black	50 and Over	1975-2000	0.0						
<u>U.S. Cancer Mortality^b</u>									
All Races	All Ages	1975-1991	-1.5 ^c	1991-2000	-0.3				
All Races	Under 50	1975-1988	-4.0 ^c	1988-2000	0.4				
All Races	50 and Over	1975-1991	-1.4 ^c	1991-2000	-0.4 ^c				
White	All Ages	1975-1991	-1.6 ^c	1991-2000	-0.3				
White	Under 50	1975-1987	-4.0 ^c	1987-2000	0.4				
White	50 and Over	1975-1991	-1.6 ^c	1991-2000	-0.4 ^c				
Black	All Ages	1975-2000	-0.7 ^c						
Black	Under 50	1975-2000	-3.7 ^c						
Black	50 and Over	1975-2000	-0.5 ^c						

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

Table VII-2
CORPUS AND UTERUS, NOS CANCER (Invasive)

SEER INCIDENCE^a AND U.S. DEATH^b RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	Incidence ^a			US Death ^b		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<u>AGE-SPECIFIC RATES, 1996-2000</u>						
AGE AT DIAGNOSIS/DEATH:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	1.3	1.2	-	0.1	-	-
30-34	2.8	2.3	-	0.2	0.2	-
35-39	6.1	6.0	3.8	0.4	0.4	0.5
40-44	12.5	12.7	6.7	0.8	0.7	1.0
45-49	24.0	24.3	12.7	1.7	1.7	1.7
50-54	47.9	51.3	22.4	3.3	3.1	4.4
55-59	68.4	73.2	38.9	6.3	6.0	9.7
60-64	86.5	91.9	69.4	10.8	9.9	20.4
65-69	94.4	100.5	83.3	15.4	13.8	32.8
70-74	101.9	108.1	81.1	20.5	18.9	40.3
75-79	99.0	105.0	68.7	24.4	23.0	43.1
80-84	95.2	100.6	73.8	30.0	28.9	46.5
85+	71.3	71.9	73.6	35.5	34.6	49.3
<u>AGE-ADJUSTED RATES, 1996-2000</u>						
AGE AT DIAGNOSIS/DEATH:						
All ages	25.2	26.5	17.8	4.1	3.8	7.0
Under 65	15.1	15.9	9.2	1.3	1.3	2.1
65 and over	94.5	99.9	77.1	23.1	21.7	40.8
All ages (world std.) ^c	18.1	19.1	12.7	2.5	2.3	4.3

^a SEER 9 areas. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups, unless noted.

^b NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups, unless noted.

^c Rates are per 100,000 and are age-adjusted to the world standard population.

- Statistic not shown. Rate based on less than 25 cases for the time interval.

Table VII-3
CORPUS AND UTERUS, NOS CANCER (Invasive)

SEER INCIDENCE^a AND U.S. DEATH^b RATES, AGE-ADJUSTED RATES, BY RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES</u>									
YEAR OF DIAGNOSIS:									
1975	35.5	6.5	111.3	37.3	6.8	117.2	19.9	-	62.6
1976	34.4	6.2	108.1	36.2	6.3	114.7	17.6	-	55.1
1977	31.6	5.7	99.6	33.1	5.9	104.5	20.1	-	63.4
1978	30.1	6.1	92.9	31.7	6.3	98.2	18.6	-	57.5
1979	27.9	5.3	87.0	29.2	5.3	91.7	16.8	-	50.5
1980	27.3	5.0	85.9	28.5	5.0	90.2	16.4	-	52.8
1981	27.3	5.6	84.0	28.3	5.5	88.2	17.4	-	55.5
1982	26.8	5.0	83.8	28.2	5.2	88.4	17.2	-	55.8
1983	26.6	4.9	83.4	27.8	4.9	87.8	18.9	-	59.6
1984	25.7	4.6	80.9	27.1	4.8	85.5	17.4	-	58.4
1985	25.2	4.6	79.2	26.5	4.8	83.4	17.5	-	57.5
1986	24.4	4.8	75.6	25.6	5.0	79.6	16.6	-	53.4
1987	24.7	5.0	76.4	26.1	5.3	80.7	15.5	-	51.7
1988	23.6	4.5	73.4	24.6	4.4	77.3	16.9	3.0	53.3
1989	24.4	4.5	76.6	25.5	4.8	79.7	19.4	-	65.5
1990	25.1	4.6	78.7	26.6	4.8	83.5	16.8	-	55.0
1991	24.6	4.4	77.5	25.9	4.4	82.2	17.6	3.3	55.0
1992	24.8	4.8	77.2	26.2	4.8	82.1	17.0	2.8	54.4
1993	24.3	5.2	74.5	25.7	5.1	79.6	17.0	3.6	52.3
1994	25.1	5.0	77.8	26.2	5.0	81.8	17.8	2.7	57.6
1995	25.3	4.7	79.3	26.4	4.6	83.3	18.6	2.5	60.6
1996	24.8	4.5	78.1	25.8	4.3	82.2	19.2	3.2	61.2
1997	25.8	5.0	80.4	27.3	5.0	85.6	16.2	2.4	52.5
1998	25.6	4.8	80.2	27.0	4.8	85.1	19.1	3.0	61.3
1999	25.2	5.2	77.6	26.6	5.0	82.9	17.1	2.7	54.6
2000	24.4	5.1	75.0	25.7	5.2	79.4	17.5	3.0	55.3
1996-2000	25.2	4.9	78.2	26.5	4.9	83.1	17.8	2.9	56.9
<u>U.S. DEATH RATES</u>									
YEAR OF DEATH:									
1975	5.3	0.5	17.7	5.1	0.5	17.1	7.8	0.9	25.8
1976	5.3	0.5	17.9	5.1	0.5	17.2	8.1	0.9	27.0
1977	5.2	0.5	17.5	5.0	0.5	16.7	8.4	0.8	28.0
1978	5.1	0.5	17.1	4.8	0.5	16.2	8.2	0.8	27.3
1979	5.0	0.4	17.0	4.7	0.4	16.2	7.9	0.7	26.7
1980	5.1	0.4	17.4	4.8	0.4	16.5	8.4	0.9	28.1
1981	4.9	0.4	16.6	4.7	0.4	15.9	7.3	0.6	25.1
1982	4.9	0.4	16.5	4.6	0.4	15.7	7.6	0.4	26.4
1983	4.8	0.4	16.4	4.6	0.3	15.8	7.3	0.6	24.8
1984	4.7	0.4	16.0	4.4	0.3	15.2	7.7	0.7	26.1
1985	4.6	0.4	15.7	4.4	0.3	15.0	7.4	0.5	25.3
1986	4.4	0.3	15.2	4.2	0.3	14.4	7.3	0.6	24.9
1987	4.4	0.3	15.1	4.2	0.3	14.3	7.3	0.5	25.1
1988	4.3	0.3	14.9	4.1	0.3	14.0	7.4	0.5	25.6
1989	4.3	0.3	14.5	4.0	0.3	13.6	7.3	0.5	25.0
1990	4.3	0.3	14.8	4.1	0.3	13.9	7.4	0.4	25.6
1991	4.2	0.3	14.2	3.9	0.3	13.3	7.3	0.5	25.1
1992	4.2	0.3	14.4	4.0	0.3	13.6	7.0	0.4	24.1
1993	4.2	0.3	14.3	3.9	0.3	13.3	7.3	0.4	25.2
1994	4.2	0.3	14.3	3.9	0.3	13.4	7.1	0.3	24.7
1995	4.1	0.3	14.1	4.0	0.3	13.5	6.5	0.3	22.5
1996	4.1	0.3	14.1	3.9	0.3	13.2	7.2	0.4	25.0
1997	4.0	0.3	13.7	3.8	0.3	12.9	6.7	0.3	23.6
1998	4.1	0.3	14.0	3.9	0.3	13.2	6.8	0.4	23.6
1999	4.1	0.3	13.9	3.9	0.3	13.1	6.8	0.4	23.7
2000	4.1	0.4	14.0	3.8	0.3	13.0	7.3	0.4	25.2
1996-2000	4.1	0.3	13.9	3.8	0.3	13.1	7.0	0.4	24.2

^a SEER 9 areas. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups, unless noted.

^b NCHS public use data file. Rates are per 100,000 and are age-adjusted to the world standard population.

- Statistic not shown. Rate based on less than 25 cases for the time interval.

Table VII-4
CORPUS AND UTERUS, NOS CANCER (Invasive)

SURVIVAL RATES, BY RACE, DIAGNOSIS YEAR, STAGE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>5-YR RELATIVE SURVIVAL RATES</u>									
YEAR OF DIAGNOSIS:									
1960-1963 ^a	-	-	-	73	-	-	31	-	-
1970-1973 ^a	-	-	-	81	-	-	44	-	-
1974-1976 ^b	88.0	92.2	87.4	88.9	92.9	88.4	61.5	84.4 ^d	56.5
1977-1979 ^b	85.1	92.1	84.1	86.4	92.9	85.5	57.9	81.1 ^d	52.6
1980-1982 ^b	81.6	91.1	80.4	82.8	92.4	81.7	55.1	85.6 ^d	50.2
1983-1985 ^b	82.9	92.5	81.6	84.5	93.3	83.5	54.5	81.4 ^d	50.5
1986-1988 ^b	82.9	92.3	81.4	84.5	92.5	83.3	57.2	86.5	52.0
1989-1991 ^b	83.7	88.4	83.0	85.6	90.8	84.8	57.7	72.1 ^d	55.2
1992-1999 ^b	84.4 ^c	90.0	83.4 ^c	86.3 ^c	91.5	85.5 ^c	60.0	80.9	55.3
1992-1999 ^b									
ALL STAGES	84.4	90.0	83.4	86.3	91.5	85.5	60.0	80.9	55.3
LOCALIZED	96.2	96.4	96.2	97.0	97.5	97.0	83.5	90.5	81.4
REGIONAL	64.7	80.1	61.9	67.3	82.0	65.0	42.5	71.7 ^d	37.1
DISTANT	26.0	44.2	23.2	28.0	46.7	25.2	13.4	-	10.2
UNSTAGED	53.9	79.6	50.3	53.4	80.8 ^d	50.1	48.5 ^d	-	46.7 ^d
<u>STAGE DISTRIBUTION (%) 1992-1999^b</u>									
ALL STAGES									
Number of cases	22,437	3,252	19,185	19,535	2,529	17,006	1,390	227	1,163
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
LOCALIZED	73	76	72	74	76	74	53	68	50
REGIONAL	15	14	15	14	14	14	22	17	23
DISTANT	8	7	8	8	7	8	17	9	19
UNSTAGED	4	4	5	4	3	4	8	7	8
<u>5-YR RELATIVE SURVIVAL RATES, 1992-1999^b</u>									
AGE AT DIAGNOSIS:									
<45	90.0	-	-	90.9	-	-	83.7	-	-
45-54	90.4	-	-	91.9	-	-	73.7	-	-
55-64	87.6	-	-	90.1	-	-	57.7	-	-
65-74	82.9	-	-	85.2	-	-	54.8	-	-
75+	74.7	-	-	76.4	-	-	42.9	-	-
Under 65	88.9	-	-	90.8	-	-	67.9	-	-
65 and over	79.8	-	-	81.9	-	-	50.8	-	-

- ^a Rates are based on End Results data from a series of hospital registries and one population-based registry.
- ^b Rates are from the SEER 9 areas. They are based on data from population-based registries in Connecticut, New Mexico, Utah, Iowa, Hawaii, Atlanta, Detroit, Seattle-Puget Sound and San Francisco-Oakland. Rates are based on follow-up of patients into 2000.
- ^c The difference in rates between 1974-1976 and 1992-1999 is statistically significant (p<.05).
- ^d The standard error of the survival rate is between 5 and 10 percentage points.
- ^e The standard error of the survival rate is greater than 10 percentage points.
- Statistic could not be calculated.

Table VII-5
CORPUS AND UTERUS, NOS CANCER (Invasive)

INCIDENCE, DEATH, AND SURVIVAL RATES^c

By Year of Diagnosis/Death

All Races, Females

Year of Diagnosis/Death

	1975- 1979	1980- 1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	
Incidence																			
SEER ^a	31.8	26.7	25.2	24.4	24.7	23.6	24.4	25.1	24.6	24.8	24.3	25.1	25.3	24.8	25.8	25.6	25.2	24.4	
Mortality																			
U.S. ^b	5.2	4.9	4.6	4.4	4.4	4.3	4.3	4.3	4.2	4.2	4.2	4.2	4.1	4.1	4.0	4.1	4.1	4.1	
SEER ^a	5.0	5.1	5.0	4.2	4.1	4.1	4.5	4.2	4.2	4.3	4.0	3.9	4.1	4.1	3.7	4.2	3.9	4.2	
Relative Survival Rates (SEER ^a)																			
1-year	93.9	91.7	93.2	93.1	92.1	92.3	92.3	92.7	92.0	93.1	92.5	92.0	93.2	92.0	93.3	93.7	94.0		
2-year	90.3	86.7	88.2	87.1	87.4	88.4	88.0	88.2	88.1	89.7	88.7	87.4	88.7	88.5	89.3	89.7			
3-year	88.1	84.1	85.5	85.4	84.4	85.5	85.9	85.3	85.8	87.4	85.9	85.2	86.5	86.4	86.3				
4-year	86.7	82.9	83.8	84.0	83.5	84.0	84.9	84.0	84.2	86.2	84.7	83.7	85.2	85.4					
5-year	86.4	82.0	83.4	82.9	82.5	83.2	84.1	83.3	83.7	86.1	83.9	83.0	84.8						
6-year	86.0	81.6	82.9	82.6	81.7	82.9	83.7	82.2	82.6	85.4	83.2	82.0							
7-year	85.8	81.3	82.3	82.5	81.7	82.6	83.7	82.2	82.5	85.4	83.1								
8-year	85.8	80.9	82.1	82.5	81.5	82.6	82.7	82.0	82.0	85.0									
9-year	85.8	80.7	81.7	82.2	81.5	82.3	82.7	82.0	81.9										
10-year	85.8	80.7	81.7	82.2	81.5	81.8	82.7	81.3											
11-year	85.8	80.5	81.7	82.2	81.4	81.8	82.6												
12-year	85.8	80.5	81.7	82.1	81.4	81.6													
13-year	85.8	80.3	81.2	82.1	81.4														
14-year	85.8	80.1	80.9	82.1															
15-year	85.8	80.1	80.9																
16-year	85.8	79.5																	
17-year	85.8																		
18-year	85.8																		
19-year	85.8																		
20-year	85.8																		

^a Rates are from the SEER 9 areas.

^b Rates are from the NCHS public use data file.

^c Incidence and death rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups. Survival rates are relative rates expressed as percents.

Table VII-6
CORPUS AND UTERUS, NOS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	2.62
10	0.00	0.01	0.06	2.66
20	0.01	0.06	0.23	2.67
30	0.05	0.22	0.73	2.67
40	0.18	0.69	1.47	2.65
50	0.53	1.32	2.02	2.52
60	0.83	1.57	2.00	2.09

Lifetime Risk of Being Diagnosed = 2.62%
Lifetime Risk of Dying = 0.50%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	2.82
10	0.00	0.01	0.06	2.86
20	0.01	0.06	0.23	2.87
30	0.05	0.23	0.78	2.87
40	0.18	0.74	1.59	2.84
50	0.57	1.44	2.19	2.71
60	0.90	1.69	2.15	2.24

Lifetime Risk of Being Diagnosed = 2.82%
Lifetime Risk of Dying = 0.48%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	1.67
10	0.00	0.01	0.04	1.72
20	0.01	0.04	0.13	1.72
30	0.03	0.12	0.39	1.73
40	0.09	0.37	0.97	1.73
50	0.29	0.92	1.40	1.70
60	0.69	1.22	1.48	1.55

Lifetime Risk of Being Diagnosed = 1.67%
Lifetime Risk of Dying = 0.72%

Table VII-7
CORPUS AND UTERUS, NOS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1996-2000	Trend 1992-2000
	Rate per 100,000 persons	APC (%)
	<u>Females</u>	<u>Females</u>
All Races	24.5	-0.1
White	26.1	0.0
White Hispanic ^b	17.2	0.5
White Non-Hispanic ^b	27.0	-0.1
Black	17.9	0.2
Asian/Pacific Islander	17.1	1.4
Amer Ind/Alaska Nat	10.6	1.0
Hispanic ^b	16.5	0.6

Mortality^c

RACE/ETHNICITY	Rate 1996-2000	Trend 1992-2000
	Rate per 100,000 persons	APC (%)
	<u>Females</u>	<u>Females</u>
All Races	4.1	-0.3
White	3.8	-0.4
White Hispanic ^b	3.2	0.2
White Non-Hispanic ^b	3.8	-0.3
Black	7.0	-0.1
Asian/Pacific Islander	2.3	1.4
Amer Ind/Alaska Nat	2.2	-3.8
Hispanic ^b	3.1	0.2

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Oklahoma, New York, and New Hampshire.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table VII-8
CORPUS AND UTERUS, NOS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE RATES^a AND SEER DEATH RATES^b
By Registry, Race and Age

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES^a, 1996-2000</u>									
REGISTRY									
12 registries	24.5	4.8	76.1	26.1	4.9	81.7	17.9	2.8	57.4
11 registries	24.5	4.8	76.1	26.1	4.9	81.7	17.9	2.8	57.4
9 registries	25.2	4.9	78.2	26.5	4.9	83.1	17.8	2.9	56.9
San Francisco-Oakland	25.2	4.5	79.3	27.8	4.5	88.6	17.7	-	56.9
Connecticut	28.2	5.1	88.7	28.8	5.2	90.6	19.0	-	60.4
Metropolitan Detroit	25.9	5.0	80.6	28.1	5.5	87.3	19.2	3.7	59.7
Hawaii	24.7	10.3	62.7	22.6	4.9	68.8	-	-	-
Iowa	28.1	5.4	87.4	28.3	5.4	88.1	-	-	-
New Mexico	18.2	4.2	54.8	18.5	3.9	56.5	-	-	-
Seattle-Puget Sound	26.2	4.6	82.8	27.3	4.6	86.6	19.0	-	65.3
Utah	23.3	4.9	71.6	23.5	4.8	72.4	-	-	-
Metropolitan Atlanta	20.5	3.5	65.1	22.6	4.4	70.4	16.2	2.1	53.3
San Jose-Monterey	23.9	3.6	76.9	26.2	3.7	85.0	-	-	-
Los Angeles	22.6	4.8	69.1	24.7	5.1	76.1	18.6	2.9	59.8
<u>SEER DEATH RATES^b, 1996-2000</u>									
REGISTRY									
12 registries	4.1	0.3	13.8	4.0	0.3	13.7	6.2	0.5	21.3
11 registries	4.1	0.4	13.9	4.0	0.3	13.7	6.3	0.5	21.3
9 registries	4.0	0.3	13.7	3.9	0.3	13.4	6.0	0.4	20.6
San Francisco-Oakland	4.0	0.4	13.4	4.1	-	13.8	5.7	-	20.3
Connecticut	4.0	0.4	13.5	3.9	-	13.2	6.0	-	20.0
Metropolitan Detroit	4.5	0.5	14.8	3.9	-	13.0	6.8	-	22.7
Hawaii	3.6	-	11.0	-	-	-	-	-	-
Iowa	4.4	-	15.0	4.4	-	14.9	-	-	-
New Mexico	3.5	-	11.6	3.5	-	11.9	-	-	-
Seattle-Puget Sound	4.0	-	13.7	4.0	-	14.0	-	-	-
Utah	3.7	-	12.8	3.7	-	12.9	-	-	-
Metropolitan Atlanta	3.8	-	13.5	3.3	-	11.7	5.5	-	19.0
San Jose-Monterey	4.0	-	14.0	4.2	-	14.9	-	-	-
Los Angeles	4.3	0.4	14.5	4.3	0.4	14.7	6.9	-	23.1

Note: San Jose-Monterey and Los Angeles are new SEER areas as of 1992 and their data are only used in tables where indicated.

^a SEER 9 areas. Incidence rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

^b NCHS public use data file. Death rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

- Statistic could not be calculated or populations are inadequate.

Table VII-9
CORPUS AND UTERUS, NOS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, 1996-2000

All Races, Females

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
TOTAL U.S.	4.1	0.02							
High Five States									
District of Columbia	7.4 ^{bc}	0.68	(01)	80.5					
New Jersey	5.0 ^{bc}	0.14	(02)	22.0					
New York	4.8 ^{bc}	0.09	(03)	17.1					
Vermont	4.8 ^b	0.53	(04)	17.1					
Pennsylvania	4.7 ^c	0.11	(05)	14.6					
Low Five States									
Colorado	3.2 ^{bc}	0.18	(47)	-22.0					
Alaska	3.2 ^b	0.69	(48)	-22.0					
Nevada	3.1 ^{bc}	0.27	(49)	-24.4					
Arizona	3.0 ^{bc}	0.15	(50)	-26.8					
North Dakota	2.8 ^{bc}	0.36	(51)	-31.7					
Alabama	3.6	0.17	(40)	-12.2	Montana	4.1	0.39	(18)	0.0
Alaska	3.2 ^b	0.69	(48)	-22.0	Nebraska	4.4	0.30	(11)	7.3
Arizona	3.0 ^{bc}	0.15	(50)	-26.8	Nevada	3.1 ^{bc}	0.27	(49)	-24.4
Arkansas	3.5	0.21	(43)	-14.6	New Hampshire	4.0	0.35	(23)	-2.4
California	3.9	0.07	(29)	-4.9	New Jersey	5.0 ^{bc}	0.14	(02)	22.0
Colorado	3.2 ^{bc}	0.18	(47)	-22.0	New Mexico	3.5	0.28	(44)	-14.6
Connecticut	4.0	0.19	(25)	-2.4	New York	4.8 ^{bc}	0.09	(03)	17.1
Delaware	4.5	0.45	(10)	9.8	North Carolina	4.1	0.14	(17)	0.0
District of Columbia	7.4 ^{bc}	0.68	(01)	80.5	North Dakota	2.8 ^{bc}	0.36	(51)	-31.7
Florida	3.5 ^c	0.08	(42)	-14.6	Ohio	4.6 ^c	0.12	(06)	12.2
Georgia	3.8	0.14	(35)	-7.3	Oklahoma	3.7	0.19	(37)	-9.8
Hawaii	3.6	0.34	(39)	-12.2	Oregon	4.1	0.21	(21)	0.0
Idaho	3.8	0.35	(34)	-7.3	Pennsylvania	4.7 ^c	0.11	(05)	14.6
Illinois	4.5 ^c	0.11	(08)	9.8	Rhode Island	3.6	0.33	(41)	-12.2
Indiana	4.5	0.16	(07)	9.8	South Carolina	4.0	0.19	(24)	-2.4
Iowa	4.4	0.22	(12)	7.3	South Dakota	3.5	0.40	(45)	-14.6
Kansas	3.9	0.22	(28)	-4.9	Tennessee	3.8	0.16	(30)	-7.3
Kentucky	3.7	0.18	(38)	-9.8	Texas	3.8	0.09	(33)	-7.3
Louisiana	4.0	0.18	(26)	-2.4	Utah	3.7	0.30	(36)	-9.8
Maine	4.2	0.33	(15)	2.4	Vermont	4.8 ^b	0.53	(04)	17.1
Maryland	4.3	0.18	(13)	4.9	Virginia	4.2	0.15	(16)	2.4
Massachusetts	4.1	0.15	(19)	0.0	Washington	3.8	0.16	(32)	-7.3
Michigan	4.3	0.12	(14)	4.9	West Virginia	4.5	0.27	(09)	9.8
Minnesota	4.0	0.17	(27)	-2.4	Wisconsin	4.1	0.16	(20)	0.0
Mississippi	3.5	0.21	(46)	-14.6	Wyoming	3.8	0.56	(31)	-7.3
Missouri	4.1	0.16	(22)	0.0					

^a NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

^b Absolute percent difference between state rate and total U.S. rate is 15% or more.

^c Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

Table VII-10
CORPUS AND UTERUS, NOS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race</u>									
All Races ^b	139,001	104,062	81,837	68,466	71,469	243,063	464,835	91,805	556,640
White ^b	125,672	96,898	76,756	65,068	68,327	222,570	432,721	84,323	517,044
Black ^b	7,941	4,098	3,128	1,897	1,691	12,039	18,755	1,430	20,185
Asian/Pacific Islander ^c	3,092	1,783	+	+	+	4,875	+	+	+
Hispanic ^d	5,704	3,726	+	+	+	9,430	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^e	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>												
All Races ^c	0.1498%	-	-	0.0023%	0.0178%	0.0797%	0.2850%	0.5841%	0.7167%	0.6118%	0.1525%	
White ^c	0.1722%	-	-	0.0021%	0.0170%	0.0795%	0.3105%	0.6509%	0.7923%	0.6643%	0.1662%	
Black ^c	0.0610%	-	-	-	0.0129%	0.0471%	0.1235%	0.3303%	0.4035%	0.2910%	0.0813%	
Asian/Pacific Islander ^c	0.0923%	-	-	-	0.0240%	0.1040%	0.2401%	0.3289%	0.3306%	0.2635%	0.0991%	
Hispanic ^d	0.0547%	-	-	-	0.0204%	0.0646%	0.2043%	0.3765%	0.4079%	0.3212%	0.0991%	

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

SEER Incidence and US Death Rates+ Corpus Uteri Cancer, by Race





Figure VII-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines are calculated using the Joinpoint Regression Program.

SEER Incidence and US Death Rates+ Corpus Uteri Cancer, by Age and Race




Figure VII-2

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines are calculated using the Joinpoint Regression Program.