

Table II-14
ALL SITES (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	3,685,031	2,355,137	1,352,174	794,296	535,422	6,040,168	8,722,060	833,252	9,555,312
Males	1,841,190	1,171,670	533,649	279,217	154,814	3,012,860	3,980,540	261,159	4,241,699
Females	1,843,841	1,183,467	818,525	515,079	380,608	3,027,308	4,741,520	572,093	5,313,613
White ^b	3,212,365	2,100,734	1,227,413	726,380	490,382	5,313,099	7,757,274	754,307	8,511,581
Males	1,592,187	1,041,352	486,147	257,079	142,532	2,633,539	3,519,297	242,731	3,762,028
Females	1,620,178	1,059,382	741,266	469,301	347,850	2,679,560	4,237,977	511,576	4,749,553
Black ^b	334,267	181,271	85,607	47,316	30,371	515,538	678,832	45,780	724,612
Males	180,538	96,491	33,090	15,402	7,953	277,029	333,474	14,469	347,943
Females	153,729	84,780	52,517	31,914	22,418	238,509	345,358	31,311	376,669
Asian/Pacific Islander ^c	76,199	39,771	+	+	+	115,970	+	+	+
Males	33,025	16,066	+	+	+	49,091	+	+	+
Females	43,174	23,705	+	+	+	66,879	+	+	+
Hispanic ^d	166,028	94,199	+	+	+	260,227	+	+	+
Males	79,075	42,040	+	+	+	121,115	+	+	+
Females	86,953	52,159	+	+	+	139,112	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	1.9111%	0.0748%	0.0993%	0.2076%	0.5325%	1.2379%	2.9800%	6.4966%	10.5383%	10.6130%	2.1055%
Males	1.9139%	0.0801%	0.1037%	0.1876%	0.4086%	0.8068%	2.3870%	7.4372%	14.4841%	15.9608%	2.4352%
Females	1.9083%	0.0693%	0.0946%	0.2285%	0.6596%	1.6613%	3.5440%	5.6667%	7.5967%	7.8301%	1.9259%
White ^c	2.0821%	0.0808%	0.1064%	0.2286%	0.5746%	1.2864%	3.0774%	6.7927%	10.8803%	10.7544%	2.1801%
Males	2.0585%	0.0867%	0.1104%	0.2098%	0.4480%	0.8440%	2.4332%	7.6148%	14.6828%	15.9405%	2.4787%
Females	2.1054%	0.0745%	0.1021%	0.2488%	0.7089%	1.7345%	3.7053%	6.0493%	8.0202%	8.1040%	2.0281%
Black ^c	1.3706%	0.0488%	0.0742%	0.1358%	0.3684%	1.0616%	2.9355%	6.6802%	10.5336%	9.3292%	2.0092%
Males	1.5439%	0.0510%	0.0797%	0.1059%	0.2441%	0.7367%	2.9184%	9.3119%	17.1580%	16.7625%	2.7691%
Females	1.2137%	0.0466%	0.0686%	0.1628%	0.4789%	1.3448%	2.9498%	4.5986%	6.1660%	5.9977%	1.5498%
Asian/Pacific Islander ^c	1.1831%	0.0662%	0.0793%	0.1250%	0.3426%	0.9577%	2.0406%	3.7762%	6.7261%	8.1208%	1.4109%
Males	1.0844%	0.0697%	0.0842%	0.1037%	0.2178%	0.4891%	1.2481%	3.7580%	9.0719%	11.7285%	1.4915%
Females	1.2757%	0.0625%	0.0740%	0.1457%	0.4597%	1.3769%	2.7444%	3.7910%	4.9490%	5.5891%	1.3824%
Hispanic ^d	0.7301%	0.0704%	0.0889%	0.1413%	0.3699%	0.8641%	1.9564%	4.2461%	7.6889%	8.0374%	1.4911%
Males	0.6513%	0.0757%	0.0928%	0.1265%	0.2583%	0.4514%	1.3952%	4.7963%	11.0101%	12.2605%	1.7297%
Females	0.8128%	0.0649%	0.0848%	0.1584%	0.4948%	1.2773%	2.4666%	3.7969%	5.3269%	5.7843%	1.3812%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas (d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table III-14
BRAIN AND OTHER NERVOUS SYSTEM (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	33,993	20,589	14,073	8,437	6,459	54,582	83,551	14,131	97,682
Males	19,234	11,447	7,270	4,481	3,220	30,681	45,652	7,051 ^h	52,703
Females	14,759	9,142	6,803	3,956	3,239	23,901	37,899	7,080 ^h	44,979
White ^b	30,155	18,062	12,531	7,525	5,830	48,217	74,103	11,167	85,270
Males	17,145	10,056	6,315	4,004	2,925	27,201	40,445	5,795 ^h	46,240
Females	13,010	8,006	6,216	3,521	2,905	21,016	33,658	5,372 ^h	39,030
Black ^b	2,530	1,719	1,168	696	458	4,249	6,571	1,488	8,059
Males	1,452	886	711	369	-	2,338	3,622	718	4,340
Females	1,078	833	457	327	-	1,911	2,949	770	3,719
Asian/Pacific Islander ^c	816	431	+	+	+	1,247	+	+	+
Males	440	256	+	+	+	696	+	+	+
Females	376	175	+	+	+	551	+	+	+
Hispanic ^d	2,640	1,546	+	+	+	4,186	+	+	+
Males	1,433	801	+	+	+	2,234	+	+	+
Females	1,207	745	+	+	+	1,952	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0181%	0.0119%	0.0195%	0.0141%	0.0195%	0.0218%	0.0225%	0.0221%	0.0176%	0.0103%	0.0182%
Males	0.0204%	0.0137%	0.0205%	0.0159%	0.0212%	0.0252%	0.0259%	0.0262%	0.0210%	0.0115%	0.0205%
Females	0.0158%	0.0101%	0.0184%	0.0122%	0.0176%	0.0186%	0.0193%	0.0186%	0.0151%	0.0097%	0.0159%
White ^c	0.0200%	0.0134%	0.0205%	0.0157%	0.0219%	0.0247%	0.0249%	0.0243%	0.0191%	0.0094%	0.0200%
Males	0.0224%	0.0153%	0.0214%	0.0175%	0.0234%	0.0283%	0.0283%	0.0287%	0.0233%	0.0111%	0.0225%
Females	0.0175%	0.0113%	0.0195%	0.0137%	0.0204%	0.0210%	0.0216%	0.0203%	0.0160%	0.0085%	0.0177%
Black ^c	0.0118%	0.0079%	0.0172%	0.0097%	0.0113%	0.0122%	0.0151%	0.0115%	-	-	0.0119%
Males	0.0132%	0.0093%	0.0174%	0.0106%	0.0132%	0.0151%	0.0194%	-	-	-	0.0131%
Females	0.0106%	-	0.0170%	0.0089%	0.0096%	0.0096%	-	-	-	-	0.0108%
Asian/Pacific Islander ^c	0.0108%	0.0068%	0.0148%	0.0078%	0.0104%	0.0115%	0.0117%	0.0161%	-	-	0.0110%
Males	0.0124%	-	0.0163%	0.0104%	0.0118%	0.0119%	0.0136%	0.0209%	-	-	0.0126%
Females	0.0093%	-	0.0133%	-	0.0091%	0.0111%	-	-	-	-	0.0096%
Hispanic ^d	0.0122%	0.0101%	0.0122%	0.0076%	0.0136%	0.0163%	0.0160%	0.0215%	0.0166%	-	0.0133%
Males	0.0127%	0.0110%	0.0121%	0.0084%	0.0133%	0.0169%	0.0165%	0.0248%	-	-	0.0142%
Females	0.0117%	0.0091%	0.0123%	0.0067%	0.0140%	0.0157%	0.0156%	0.0188%	-	-	0.0125%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g h} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25. (h) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table IV-17
BREAST CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	774,287	529,022	357,116	192,249	127,139	1,303,309	1,979,813	+	+
Males	4,591	2,707	1,714	930	487	7,298	10,429	+	+
Females	769,696	526,315	355,402	191,319	126,652	1,296,011	1,969,384	228,120	2,197,504
White ^b	682,617	475,623	325,741	175,439	117,057	1,158,240	1,776,477	+	+
Males	3,985	2,382	1,529	863	458	6,367	9,217	+	+
Females	678,632	473,241	324,212	174,576	116,599	1,151,873	1,767,260	212,145	1,979,405
Black ^b	64,797	38,575	23,047	12,512	7,251	103,372	146,182	+	+
Males	476	-	-	-	-	713	960	+	+
Females	64,321	38,338	22,887	12,451	7,225	102,659	145,222	14,350	159,572
Asian/Pacific Islander ^c	18,403	10,246	+	+	+	28,649	+	+	+
Males	70	55	+	+	+	125	+	+	+
Females	18,333	10,191	+	+	+	28,524	+	+	+
Hispanic ^d	32,491	19,602	+	+	+	52,093	+	+	+
Males	143	-	+	+	+	212	+	+	+
Females	32,348	19,533	+	+	+	51,881	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.4150%	-	-	0.0055%	0.0744%	0.3810%	0.9432%	1.3844%	1.8618%	2.0733%	0.4559%
Males	0.0048%	-	-	-	-	0.0022%	0.0085%	0.0205%	0.0300%	0.0444%	0.0061%
Females	0.8150%	-	-	0.0113%	0.1503%	0.7530%	1.8320%	2.5879%	3.2275%	3.1290%	0.8278%
White ^c	0.4564%	-	-	0.0050%	0.0735%	0.3863%	0.9790%	1.4735%	1.9851%	2.1902%	0.4778%
Males	0.0050%	-	-	-	-	0.0023%	0.0076%	0.0211%	0.0305%	0.0452%	0.0061%
Females	0.9037%	-	-	0.0104%	0.1510%	0.7753%	1.9257%	2.7867%	3.4552%	3.2864%	0.8744%
Black ^c	0.2787%	-	-	0.0095%	0.0892%	0.3613%	0.8436%	1.1468%	1.4813%	1.6075%	0.3905%
Males	0.0046%	-	-	-	-	-	0.0178%	-	-	-	0.0082%
Females	0.5270%	-	-	0.0180%	0.1683%	0.6750%	1.5367%	2.0354%	2.4327%	2.3020%	0.6740%
Asian/Pacific Islander ^c	0.2803%	-	-	0.0054%	0.0650%	0.3582%	0.7525%	0.9437%	1.1177%	1.0866%	0.3206%
Males	0.0028%	-	-	-	-	-	-	-	-	-	0.0038%
Females	0.5408%	-	-	0.0106%	0.1260%	0.6770%	1.4169%	1.7012%	1.9495%	1.8276%	0.5826%
Hispanic ^d	0.1478%	-	-	0.0043%	0.0495%	0.2642%	0.6028%	0.8906%	1.2280%	1.3950%	0.3024%
Males	0.0011%	-	-	-	-	-	-	-	-	-	0.0035%
Females	0.3015%	-	-	0.0093%	0.1048%	0.5272%	1.1475%	1.6094%	2.0937%	2.1181%	0.5416%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table V-10
CERVIX UTERI CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race</u>									
All Races ^b	46,867	38,414	32,786	26,196	25,729	85,281	169,992	61,072	231,064
White ^b	36,314	31,576	27,065	21,433	20,969	67,890	137,357	51,303	188,660
Black ^b	7,211	4,611	3,990	3,688	3,818	11,822	23,318	+	+
Asian/Pacific Islander ^c	2,045	1,485	+	+	+	3,530	+	+	+
Hispanic ^d	7,383	6,009	+	+	+	13,392	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	0-9	10-19	20-29	Age Specific (Crude)						Age-Adjusted ^e
					30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	0.0645%	-	-	0.0131%	0.0814%	0.1305%	0.1283%	0.1193%	0.0999%	0.0641%	0.0653%
White ^c	0.0643%	-	-	0.0141%	0.0904%	0.1328%	0.1207%	0.1074%	0.0851%	0.0516%	0.0641%
Black ^c	0.0609%	-	-	0.0132%	0.0635%	0.1291%	0.1334%	0.1400%	0.1623%	0.1437%	0.0709%
Asian/Pacific Islander ^c	0.0648%	-	-	-	0.0375%	0.1056%	0.1628%	0.1756%	0.1663%	0.1372%	0.0678%
Hispanic ^d	0.0793%	-	-	0.0109%	0.1057%	0.2116%	0.2445%	0.2440%	0.2570%	0.1712%	0.1165%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table VI-20
COLON AND RECTUM CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	393,820	236,523	164,128	97,076	57,130	630,343	948,677	92,822	1,041,499
Males	196,538	116,985	80,538	43,332	23,287	313,523	460,680	38,338	499,018
Females	197,282	119,538	83,590	53,744	33,843	316,820	487,997	54,484	542,481
White ^b	342,690	209,535	146,732	87,741	51,606	552,225	838,304	84,701	923,005
Males	172,471	104,237	72,559	39,555	21,096	276,708	409,918	35,074	444,992
Females	170,219	105,298	74,173	48,186	30,510	275,517	428,386	49,627	478,013
Black ^b	36,608	19,469	12,589	6,880	3,861	56,077	79,407	5,143	84,550
Males	16,366	8,777	5,389	2,525	1,380	25,143	34,437	2,032	36,469
Females	20,242	10,692	7,200	4,355	2,481	30,934	44,970	3,111	48,081
Asian/Pacific Islander ^c	10,196	5,217	+	+	+	15,413	+	+	+
Males	5,414	2,695	+	+	+	8,109	+	+	+
Females	4,782	2,522	+	+	+	7,304	+	+	+
Hispanic ^d	16,066	8,077	+	+	+	24,143	+	+	+
Males	8,829	4,142	+	+	+	12,971	+	+	+
Females	7,237	3,935	+	+	+	11,172	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.1986%	-	-	0.0030%	0.0175%	0.0674%	0.2398%	0.6765%	1.2631%	1.7659%	0.2230%
Males	0.2022%	-	-	0.0029%	0.0177%	0.0708%	0.2733%	0.8188%	1.5359%	2.0505%	0.2642%
Females	0.1951%	-	-	0.0031%	0.0173%	0.0642%	0.2079%	0.5510%	1.0597%	1.6178%	0.1921%
White ^c	0.2115%	-	-	0.0028%	0.0168%	0.0641%	0.2299%	0.6772%	1.2758%	1.7922%	0.2227%
Males	0.2141%	-	-	0.0028%	0.0170%	0.0668%	0.2664%	0.8227%	1.5550%	2.0731%	0.2647%
Females	0.2089%	-	-	0.0029%	0.0165%	0.0613%	0.1944%	0.5457%	1.0657%	1.6486%	0.1907%
Black ^c	0.1508%	-	-	-	0.0186%	0.0813%	0.3115%	0.7534%	1.2790%	1.5555%	0.2338%
Males	0.1457%	-	-	-	0.0160%	0.0885%	0.3415%	0.8567%	1.4122%	1.8668%	0.2647%
Females	0.1554%	-	-	-	0.0208%	0.0751%	0.2864%	0.6716%	1.1912%	1.4159%	0.2142%
Asian/Pacific Islander ^c	0.1650%	-	-	-	0.0201%	0.0747%	0.2403%	0.6000%	1.1210%	1.6286%	0.2070%
Males	0.1797%	-	-	-	0.0222%	0.0787%	0.2600%	0.7363%	1.4297%	1.9281%	0.2487%
Females	0.1512%	-	-	-	0.0181%	0.0712%	0.2228%	0.4893%	0.8872%	1.4183%	0.1752%
Hispanic ^d	0.0667%	-	-	0.0027%	0.0132%	0.0451%	0.1776%	0.4783%	0.9238%	1.2154%	0.1603%
Males	0.0691%	-	-	-	0.0135%	0.0462%	0.2059%	0.6167%	1.2189%	1.3973%	0.1979%
Females	0.0642%	-	-	-	0.0130%	0.0441%	0.1519%	0.3653%	0.7140%	1.1184%	0.1330%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table VII-10
CORPUS AND UTERUS, NOS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race</u>									
All Races ^b	139,001	104,062	81,837	68,466	71,469	243,063	464,835	91,805	556,640
White ^b	125,672	96,898	76,756	65,068	68,327	222,570	432,721	84,323	517,044
Black ^b	7,941	4,098	3,128	1,897	1,691	12,039	18,755	1,430	20,185
Asian/Pacific Islander ^c	3,092	1,783	+	+	+	4,875	+	+	+
Hispanic ^d	5,704	3,726	+	+	+	9,430	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^e	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>												
All Races ^c	0.1498%	-	-	0.0023%	0.0178%	0.0797%	0.2850%	0.5841%	0.7167%	0.6118%	0.1525%	
White ^c	0.1722%	-	-	0.0021%	0.0170%	0.0795%	0.3105%	0.6509%	0.7923%	0.6643%	0.1662%	
Black ^c	0.0610%	-	-	-	0.0129%	0.0471%	0.1235%	0.3303%	0.4035%	0.2910%	0.0813%	
Asian/Pacific Islander ^c	0.0923%	-	-	-	0.0240%	0.1040%	0.2401%	0.3289%	0.3306%	0.2635%	0.0991%	
Hispanic ^d	0.0547%	-	-	-	0.0204%	0.0646%	0.2043%	0.3765%	0.4079%	0.3212%	0.0991%	

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.
 - Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.
 + Not available.

Table VIII-14
ESOPHAGUS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	14,718	3,754	1,414	624	-	18,472	20,766	762	21,528
Males	11,243	2,684	1,028	422	-	13,927	15,568	470	16,038
Females	3,475	1,070	386	-	-	4,545	5,198	292	5,490
White ^b	12,720	3,139	1,120	502	-	15,859	17,692	444	18,136
Males	9,905	2,237	846	326	-	12,142	13,489	267	13,756
Females	2,815	902	-	-	-	3,717	4,203	177	4,380
Black ^b	1,626	536	-	-	-	2,162	2,585	196	2,781
Males	1,049	381	-	-	-	1,430	1,703	110	1,813
Females	577	-	-	-	-	732	882	86	968
Asian/Pacific Islander ^c	252	64	+	+	+	316	+	+	+
Males	201	-	+	+	+	251	+	+	+
Females	-	-	+	+	+	65	+	+	+
Hispanic ^d	601	157	+	+	+	758	+	+	+
Males	463	133	+	+	+	596	+	+	+
Females	138	-	+	+	+	162	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0055%	-	-	-	0.0004%	0.0024%	0.0105%	0.0258%	0.0308%	0.0244%	0.0061%
Males	0.0084%	-	-	-	-	0.0042%	0.0183%	0.0424%	0.0521%	0.0407%	0.0103%
Females	0.0027%	-	-	-	-	-	0.0030%	0.0112%	0.0150%	0.0160%	0.0027%
White ^c	0.0059%	-	-	-	-	0.0025%	0.0107%	0.0263%	0.0316%	0.0248%	0.0063%
Males	0.0091%	-	-	-	-	0.0044%	0.0189%	0.0436%	0.0532%	0.0423%	0.0106%
Females	0.0028%	-	-	-	-	-	0.0026%	0.0107%	0.0154%	0.0159%	0.0026%
Black ^c	0.0052%	-	-	-	-	-	0.0154%	0.0329%	0.0332%	-	0.0075%
Males	0.0068%	-	-	-	-	-	0.0253%	0.0487%	0.0489%	-	0.0108%
Females	0.0036%	-	-	-	-	-	-	0.0204%	-	-	0.0049%
Asian/Pacific Islander ^c	0.0033%	-	-	-	-	-	-	0.0163%	0.0241%	-	0.0040%
Males	0.0055%	-	-	-	-	-	-	0.0290%	0.0476%	-	0.0073%
Females	0.0013%	-	-	-	-	-	-	-	-	-	0.0015%
Hispanic ^d	0.0021%	-	-	-	-	-	0.0075%	0.0157%	0.0244%	-	0.0049%
Males	0.0032%	-	-	-	-	-	0.0138%	0.0274%	0.0419%	-	0.0087%
Females	0.0009%	-	-	-	-	-	-	-	-	-	0.0020%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.
 - Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.
 + Not available.

Table IX-14
HODGKIN LYMPHOMA (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	32,937	27,817	23,536	18,177	13,527	60,754	115,994	15,285	131,279
Males	17,233	14,316	11,868	9,823	6,881	31,549	60,121	6,804 ^h	66,925
Females	15,704	13,501	11,668	8,354	6,646	29,205	55,873	8,481 ^h	64,354
White ^b	28,864	24,806	21,686	16,929	12,585	53,670	104,870	13,287	118,157
Males	15,099	12,914	10,869	9,053	6,242	28,013	54,177	5,664 ^h	59,841
Females	13,765	11,892	10,817	7,876	6,343	25,657	50,693	7,623 ^h	58,316
Black ^b	3,252	2,514	1,436	1,016	789	5,766	9,007	+	+
Males	1,659	1,159	769	650	545	2,818	4,782	+	+
Females	1,593	1,355	667	366	-	2,948	4,225	+	+
Asian/Pacific Islander ^c	401	273	+	+	+	674	+	+	+
Males	222	116	+	+	+	338	+	+	+
Females	179	157	+	+	+	336	+	+	+
Hispanic ^d	2,352	1,632	+	+	+	3,984	+	+	+
Males	1,296	912	+	+	+	2,208	+	+	+
Females	1,056	720	+	+	+	1,776	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0199%	0.0007%	0.0085%	0.0308%	0.0365%	0.0242%	0.0199%	0.0188%	0.0173%	0.0120%	0.0196%
Males	0.0213%	0.0012%	0.0082%	0.0294%	0.0369%	0.0269%	0.0238%	0.0250%	0.0206%	0.0143%	0.0211%
Females	0.0186%	-	0.0088%	0.0324%	0.0360%	0.0215%	0.0160%	0.0133%	0.0149%	0.0108%	0.0183%
White ^c	0.0226%	0.0009%	0.0099%	0.0359%	0.0414%	0.0271%	0.0218%	0.0206%	0.0188%	0.0131%	0.0221%
Males	0.0240%	0.0015%	0.0093%	0.0339%	0.0414%	0.0301%	0.0260%	0.0278%	0.0218%	0.0157%	0.0236%
Females	0.0211%	-	0.0106%	0.0380%	0.0414%	0.0240%	0.0178%	0.0141%	0.0165%	0.0118%	0.0209%
Black ^c	0.0162%	-	0.0057%	0.0246%	0.0307%	0.0214%	0.0202%	0.0192%	-	-	0.0165%
Males	0.0168%	-	-	0.0225%	0.0327%	0.0245%	0.0241%	-	-	-	0.0177%
Females	0.0155%	-	-	0.0265%	0.0289%	0.0187%	0.0168%	-	-	-	0.0155%
Asian/Pacific Islander ^c	0.0058%	-	-	0.0077%	0.0105%	0.0065%	-	-	-	-	0.0057%
Males	0.0061%	-	-	-	0.0100%	-	-	-	-	-	0.0063%
Females	0.0056%	-	-	0.0080%	0.0110%	0.0077%	-	-	-	-	0.0052%
Hispanic ^d	0.0116%	-	0.0088%	0.0163%	0.0172%	0.0121%	0.0167%	0.0189%	0.0204%	-	0.0128%
Males	0.0125%	-	0.0097%	0.0153%	0.0173%	0.0147%	0.0202%	-	-	-	0.0145%
Females	0.0107%	-	0.0079%	0.0176%	0.0170%	0.0095%	0.0135%	0.0186%	-	-	0.0114%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g h} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25. (h) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XI-14
KIDNEY AND RENAL PELVIS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	80,552	50,321	28,809	17,196	9,893	130,873	186,771	15,302	202,073
Males	48,830	29,757	16,502	10,211	5,478	78,587	110,778	8,864	119,642
Females	31,722	20,564	12,307	6,985	4,415	52,286	75,993	6,438	82,431
White ^b	69,345	44,447	25,877	15,340	8,913	113,792	163,922	14,958	178,880
Males	42,530	26,471	14,974	9,372	4,983	69,001	98,330	7,682	106,012
Females	26,815	17,976	10,903	5,968	3,930	44,791	65,592	7,276	72,868
Black ^b	8,758	4,639	2,450	1,533	807	13,397	18,187	1,860	20,047
Males	4,774	2,533	1,249	668	400	7,307	9,624	743	10,367
Females	3,984	2,106	1,201	865	407	6,090	8,563	1,117	9,680
Asian/Pacific Islander ^c	1,419	594	+	+	+	2,013	+	+	+
Males	894	378	+	+	+	1,272	+	+	+
Females	525	216	+	+	+	741	+	+	+
Hispanic ^d	4,987	2,714	+	+	+	7,701	+	+	+
Males	2,838	1,538	+	+	+	4,376	+	+	+
Females	2,149	1,176	+	+	+	3,325	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0406%	0.0074%	0.0039%	0.0013%	0.0068%	0.0275%	0.0775%	0.1517%	0.2044%	0.1736%	0.0446%
Males	0.0497%	0.0066%	0.0037%	0.0012%	0.0071%	0.0347%	0.1052%	0.2060%	0.2856%	0.2635%	0.0604%
Females	0.0316%	0.0083%	0.0041%	0.0015%	0.0064%	0.0204%	0.0511%	0.1037%	0.1439%	0.1268%	0.0321%
White ^c	0.0444%	0.0081%	0.0041%	0.0011%	0.0068%	0.0286%	0.0798%	0.1603%	0.2170%	0.1782%	0.0466%
Males	0.0543%	0.0075%	0.0039%	-	0.0072%	0.0365%	0.1080%	0.2142%	0.3026%	0.2746%	0.0630%
Females	0.0346%	0.0087%	0.0043%	0.0013%	0.0063%	0.0206%	0.0523%	0.1115%	0.1526%	0.1289%	0.0334%
Black ^c	0.0346%	0.0072%	0.0050%	-	0.0081%	0.0324%	0.0995%	0.1595%	0.1878%	0.1669%	0.0478%
Males	0.0406%	-	-	-	0.0076%	0.0381%	0.1368%	0.2238%	0.2360%	0.2548%	0.0631%
Females	0.0292%	0.0093%	-	-	0.0086%	0.0275%	0.0681%	0.1087%	0.1561%	0.1275%	0.0369%
Asian/Pacific Islander ^c	0.0204%	-	-	-	0.0050%	0.0151%	0.0381%	0.0780%	0.1076%	0.1209%	0.0241%
Males	0.0267%	-	-	-	-	0.0191%	0.0539%	0.1148%	0.1682%	0.1619%	0.0342%
Females	0.0145%	-	-	-	-	0.0116%	0.0240%	0.0481%	0.0617%	0.0922%	0.0162%
Hispanic ^d	0.0215%	0.0056%	0.0031%	-	0.0056%	0.0249%	0.0645%	0.1450%	0.2308%	0.1897%	0.0441%
Males	0.0236%	0.0039%	-	-	0.0061%	0.0265%	0.0896%	0.1950%	0.3036%	0.2676%	0.0575%
Females	0.0193%	0.0072%	0.0044%	-	0.0049%	0.0233%	0.0418%	0.1042%	0.1789%	0.1481%	0.0340%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XII-14
LARYNX CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	34,446	23,797	16,659	11,280	6,620	58,243	92,802	6,029	98,831
Males	27,153	19,083	13,632	9,018	5,410	46,236	74,296	4,885 ^h	79,181
Females	7,293	4,714	3,027	2,262	1,210	12,007	18,506	1,144 ^h	19,650
White ^b	28,877	20,640	14,610	10,126	6,016	49,517	80,269	5,109	85,378
Males	23,010	16,507	12,088	8,090	4,908	39,517	64,603	4,300 ^h	68,903
Females	5,867	4,133	2,522	2,036	1,108	10,000	15,666	809 ^h	16,475
Black ^b	4,795	2,712	1,756	1,009	531	7,507	10,803	+	+
Males	3,530	2,206	1,364	806	436	5,736	8,342	+	+
Females	1,265	506	392	-	-	1,771	2,461	+	+
Asian/Pacific Islander ^c	484	211	+	+	+	695	+	+	+
Males	400	182	+	+	+	582	+	+	+
Females	84	-	+	+	+	113	+	+	+
Hispanic ^d	1,272	864	+	+	+	2,136	+	+	+
Males	1,095	720	+	+	+	1,815	+	+	+
Females	177	144	+	+	+	321	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0176%	-	-	-	0.0010%	0.0072%	0.0314%	0.0862%	0.1043%	0.0710%	0.0196%
Males	0.0284%	-	-	-	0.0011%	0.0108%	0.0530%	0.1470%	0.1979%	0.1627%	0.0356%
Females	0.0070%	-	-	-	0.0009%	0.0037%	0.0109%	0.0324%	0.0346%	0.0233%	0.0071%
White ^c	0.0188%	-	-	-	0.0011%	0.0072%	0.0313%	0.0897%	0.1062%	0.0713%	0.0200%
Males	0.0303%	-	-	-	0.0011%	0.0104%	0.0530%	0.1521%	0.2006%	0.1646%	0.0361%
Females	0.0074%	-	-	-	-	0.0040%	0.0101%	0.0332%	0.0352%	0.0236%	0.0072%
Black ^c	0.0195%	-	-	-	-	0.0130%	0.0553%	0.1252%	0.1405%	0.0842%	0.0287%
Males	0.0314%	-	-	-	-	0.0226%	0.0920%	0.2156%	0.2753%	0.2004%	0.0527%
Females	0.0088%	-	-	-	-	-	0.0245%	0.0536%	0.0515%	-	0.0117%
Asian/Pacific Islander ^c	0.0075%	-	-	-	-	-	0.0104%	0.0304%	0.0634%	0.0563%	0.0094%
Males	0.0131%	-	-	-	-	-	0.0178%	0.0543%	0.1275%	0.1199%	0.0182%
Females	0.0023%	-	-	-	-	-	-	-	-	-	0.0026%
Hispanic ^d	0.0058%	-	-	-	-	0.0035%	0.0178%	0.0590%	0.0703%	0.0571%	0.0131%
Males	0.0097%	-	-	-	-	-	0.0343%	0.1117%	0.1430%	0.1481%	0.0265%
Females	0.0018%	-	-	-	-	-	-	-	-	-	0.0033%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g h} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25. (h) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XIII-20
LEUKEMIA (Invasive)Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	75,378	43,008	25,059	14,578	8,315	118,386	166,338	12,068	178,406
Males	43,595	24,187	14,236	7,980	4,088	67,782	94,086	5,870	99,956
Females	31,783	18,821	10,823	6,598	4,227	50,604	72,252	6,198	78,450
White ^b	67,138	38,946	22,845	13,340	7,749	106,084	150,018	11,773	161,791
Males	38,725	22,203	13,063	7,333	3,874	60,928	85,198	5,748	90,946
Females	28,413	16,743	9,782	6,007	3,875	45,156	64,820	6,025	70,845
Black ^b	4,988	2,528	1,207	786	418	7,516	9,927	912	10,839
Males	2,813	1,164	618	431	-	3,977	5,185	420	5,605
Females	2,175	1,364	589	355	-	3,539	4,742	492	5,234
Asian/Pacific Islander ^c	1,590	828	+	+	+	2,418	+	+	+
Males	926	427	+	+	+	1,353	+	+	+
Females	664	401	+	+	+	1,065	+	+	+
Hispanic ^d	5,615	3,121	+	+	+	8,736	+	+	+
Males	3,241	1,837	+	+	+	5,078	+	+	+
Females	2,374	1,284	+	+	+	3,658	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0387%	0.0284%	0.0273%	0.0119%	0.0131%	0.0208%	0.0482%	0.0959%	0.1502%	0.1582%	0.0412%
Males	0.0452%	0.0303%	0.0299%	0.0136%	0.0155%	0.0253%	0.0616%	0.1282%	0.2058%	0.2133%	0.0522%
Females	0.0323%	0.0265%	0.0246%	0.0101%	0.0106%	0.0164%	0.0355%	0.0675%	0.1087%	0.1295%	0.0323%
White ^c	0.0433%	0.0315%	0.0297%	0.0134%	0.0133%	0.0216%	0.0518%	0.1060%	0.1653%	0.1700%	0.0448%
Males	0.0505%	0.0333%	0.0327%	0.0155%	0.0156%	0.0264%	0.0664%	0.1404%	0.2261%	0.2318%	0.0568%
Females	0.0362%	0.0297%	0.0265%	0.0112%	0.0109%	0.0168%	0.0376%	0.0749%	0.1195%	0.1385%	0.0350%
Black ^c	0.0206%	0.0113%	0.0168%	0.0052%	0.0113%	0.0162%	0.0364%	0.0581%	0.0917%	0.0878%	0.0255%
Males	0.0236%	0.0130%	0.0180%	-	0.0144%	0.0166%	0.0482%	0.0816%	0.1164%	-	0.0316%
Females	0.0180%	0.0096%	0.0156%	-	0.0086%	0.0159%	0.0264%	0.0396%	0.0755%	0.0781%	0.0210%
Asian/Pacific Islander ^c	0.0217%	0.0306%	0.0239%	0.0096%	0.0114%	0.0164%	0.0241%	0.0327%	0.0520%	0.0540%	0.0233%
Males	0.0253%	0.0342%	0.0252%	0.0106%	0.0138%	0.0204%	0.0234%	0.0401%	0.0794%	0.0744%	0.0278%
Females	0.0184%	0.0268%	0.0226%	0.0086%	0.0092%	0.0127%	0.0247%	0.0267%	0.0312%	-	0.0195%
Hispanic ^d	0.0254%	0.0312%	0.0319%	0.0127%	0.0135%	0.0174%	0.0308%	0.0485%	0.0848%	0.0934%	0.0307%
Males	0.0287%	0.0329%	0.0363%	0.0159%	0.0153%	0.0205%	0.0364%	0.0670%	0.1139%	0.1099%	0.0371%
Females	0.0220%	0.0295%	0.0271%	0.0090%	0.0115%	0.0143%	0.0256%	0.0334%	0.0641%	0.0847%	0.0252%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XIII-21
ACUTE LYMPHOCYTIC LEUKEMIA (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	13,335	9,125	7,885	5,839	3,771	22,460	39,955	+	+
Males	7,637	5,043	4,474	2,973	1,888	12,680	22,015	+	+
Females	5,698	4,082	3,411	2,866	1,883	9,780	17,940	+	+
White ^b	11,467	7,970	6,914	5,268	3,516	19,437	35,135	+	+
Males	6,507	4,517	3,966	2,698	1,791	11,024	19,479	+	+
Females	4,960	3,453	2,948	2,570	1,725	8,413	15,656	+	+
Black ^b	1,035	641	497	319	-	1,676	2,648	+	+
Males	641	-	-	-	-	926	1,414	+	+
Females	394	356	-	-	-	750	1,234	+	+
Asian/Pacific Islander ^c	550	371	+	+	+	921	+	+	+
Males	340	170	+	+	+	510	+	+	+
Females	210	201	+	+	+	411	+	+	+
Hispanic ^d	2,791	1,681	+	+	+	4,472	+	+	+
Males	1,597	998	+	+	+	2,595	+	+	+
Females	1,194	683	+	+	+	1,877	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0088%	0.0250%	0.0230%	0.0055%	0.0026%	0.0015%	0.0019%	0.0021%	0.0013%	-	0.0087%
Males	0.0101%	0.0269%	0.0251%	0.0071%	0.0030%	0.0016%	0.0021%	0.0025%	-	-	0.0096%
Females	0.0076%	0.0230%	0.0207%	0.0037%	0.0022%	0.0014%	0.0016%	-	-	-	0.0077%
White ^c	0.0097%	0.0281%	0.0257%	0.0064%	0.0029%	0.0016%	0.0020%	0.0021%	-	-	0.0097%
Males	0.0110%	0.0299%	0.0282%	0.0083%	0.0033%	0.0016%	0.0022%	0.0028%	-	-	0.0108%
Females	0.0083%	0.0261%	0.0230%	0.0042%	0.0025%	0.0016%	0.0017%	-	-	-	0.0086%
Black ^c	0.0045%	0.0093%	0.0124%	-	-	-	-	-	-	-	0.0038%
Males	0.0053%	0.0109%	0.0135%	-	-	-	-	-	-	-	0.0043%
Females	0.0037%	0.0078%	0.0113%	-	-	-	-	-	-	-	0.0033%
Asian/Pacific Islander ^c	0.0077%	0.0253%	0.0179%	0.0039%	-	-	-	-	-	-	0.0078%
Males	0.0087%	0.0289%	0.0190%	-	-	-	-	-	-	-	0.0085%
Females	0.0066%	0.0216%	0.0168%	-	-	-	-	-	-	-	0.0071%
Hispanic ^d	0.0133%	0.0286%	0.0273%	0.0059%	0.0031%	-	-	-	-	-	0.0104%
Males	0.0150%	0.0306%	0.0306%	0.0081%	-	-	-	-	-	-	0.0117%
Females	0.0115%	0.0265%	0.0238%	-	-	-	-	-	-	-	0.0090%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XIV-14
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	10,269	1,498	901	396	-	11,767	13,266	+	+
Males	6,904	896	354	-	-	7,800	8,391	+	+
Females	3,365	602	547	-	-	3,967	4,875	+	+
White ^b	7,625	1,198	737	344	-	8,823	10,036	+	+
Males	5,043	698	-	-	-	5,741	6,171	+	+
Females	2,582	500	494	-	-	3,082	3,865	+	+
Black ^b	1,285	-	-	-	-	1,406	1,584	+	+
Males	917	-	-	-	-	991	1,087	+	+
Females	368	-	-	-	-	415	497	+	+
Asian/Pacific Islander ^c	1,183	174	+	+	+	1,357	+	+	+
Males	842	109	+	+	+	951	+	+	+
Females	341	65	+	+	+	406	+	+	+
Hispanic ^d	1,163	228	+	+	+	1,391	+	+	+
Males	710	142	+	+	+	852	+	+	+
Females	453	-	+	+	+	539	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0050%	0.0015%	0.0007%	-	0.0007%	0.0037%	0.0095%	0.0194%	0.0232%	0.0179%	0.0055%
Males	0.0066%	0.0016%	-	-	0.0009%	0.0052%	0.0147%	0.0285%	0.0337%	0.0257%	0.0079%
Females	0.0034%	0.0013%	-	-	-	0.0022%	0.0046%	0.0114%	0.0154%	0.0138%	0.0035%
White ^c	0.0040%	0.0014%	0.0007%	-	0.0005%	0.0027%	0.0069%	0.0137%	0.0172%	0.0156%	0.0041%
Males	0.0050%	0.0015%	-	-	-	0.0036%	0.0105%	0.0198%	0.0240%	0.0203%	0.0057%
Females	0.0029%	0.0012%	-	-	-	0.0018%	0.0034%	0.0082%	0.0120%	0.0132%	0.0028%
Black ^c	0.0037%	-	-	-	-	-	0.0120%	0.0160%	0.0190%	-	0.0049%
Males	0.0052%	-	-	-	-	-	0.0210%	-	-	-	0.0075%
Females	0.0023%	-	-	-	-	-	-	-	-	-	0.0029%
Asian/Pacific Islander ^c	0.0136%	-	-	-	-	0.0108%	0.0261%	0.0627%	0.0734%	0.0479%	0.0159%
Males	0.0198%	-	-	-	-	0.0164%	0.0418%	0.0989%	0.1152%	0.0803%	0.0248%
Females	0.0079%	-	-	-	-	-	0.0121%	0.0334%	0.0417%	-	0.0086%
Hispanic ^d	0.0039%	-	-	-	-	0.0031%	0.0129%	0.0272%	0.0395%	-	0.0078%
Males	0.0046%	-	-	-	-	-	0.0209%	0.0417%	0.0415%	-	0.0098%
Females	0.0031%	-	-	-	-	-	-	-	0.0380%	-	0.0060%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XV-19
LUNG AND BRONCHUS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	202,049	64,492	33,754	18,075	9,871	266,541	328,241	14,216	342,457
Males	102,908	31,191	17,500	9,081	5,180	134,099	165,860	8,687	174,547
Females	99,141	33,301	16,254	8,994	4,691	132,442	162,381	5,529	167,910
White ^b	175,244	57,029	29,944	16,104	8,738	232,273	287,059	7,605	294,664
Males	87,928	27,199	15,212	7,978	4,568	115,127	142,885	4,114	146,999
Females	87,316	29,830	14,732	8,126	4,170	117,146	144,174	3,491	147,665
Black ^b	21,428	5,961	3,094	1,588	906	27,389	32,977	885	33,862
Males	11,703	3,143	1,835	880	520	14,846	18,081	504	18,585
Females	9,725	2,818	1,259	708	386	12,543	14,896	381	15,277
Asian/Pacific Islander ^c	3,925	954	+	+	+	4,879	+	+	+
Males	2,398	566	+	+	+	2,964	+	+	+
Females	1,527	388	+	+	+	1,915	+	+	+
Hispanic ^d	4,917	1,584	+	+	+	6,501	+	+	+
Males	2,687	803	+	+	+	3,490	+	+	+
Females	2,230	781	+	+	+	3,011	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0801%	-	-	0.0011%	0.0044%	0.0262%	0.1207%	0.3629%	0.5526%	0.3722%	0.0899%
Males	0.0824%	-	-	0.0009%	0.0037%	0.0270%	0.1287%	0.4061%	0.6610%	0.4959%	0.1042%
Females	0.0779%	-	-	0.0013%	0.0051%	0.0255%	0.1131%	0.3248%	0.4718%	0.3079%	0.0794%
White ^c	0.0863%	-	-	0.0012%	0.0043%	0.0254%	0.1195%	0.3786%	0.5669%	0.3761%	0.0917%
Males	0.0857%	-	-	-	0.0034%	0.0246%	0.1204%	0.4112%	0.6548%	0.4997%	0.1030%
Females	0.0869%	-	-	0.0015%	0.0051%	0.0262%	0.1187%	0.3492%	0.5008%	0.3130%	0.0839%
Black ^c	0.0711%	-	-	-	0.0063%	0.0416%	0.1776%	0.4320%	0.6057%	0.3352%	0.1061%
Males	0.0816%	-	-	-	-	0.0508%	0.2276%	0.5404%	0.8168%	0.4669%	0.1377%
Females	0.0616%	-	-	-	-	0.0336%	0.1356%	0.3463%	0.4665%	0.2761%	0.0837%
Asian/Pacific Islander ^c	0.0522%	-	-	-	0.0036%	0.0186%	0.0807%	0.2094%	0.4076%	0.3660%	0.0642%
Males	0.0657%	-	-	-	-	0.0240%	0.1061%	0.2728%	0.6199%	0.4894%	0.0892%
Females	0.0394%	-	-	-	-	0.0138%	0.0582%	0.1579%	0.2467%	0.2793%	0.0447%
Hispanic ^d	0.0179%	-	-	-	-	0.0116%	0.0447%	0.1515%	0.2724%	0.2304%	0.0423%
Males	0.0185%	-	-	-	-	0.0119%	0.0498%	0.1796%	0.3573%	0.3140%	0.0528%
Females	0.0172%	-	-	-	-	0.0114%	0.0400%	0.1286%	0.2120%	0.1858%	0.0349%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b ^c ^d Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^f ^g (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XVI-14
MELANOMA OF THE SKIN (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	186,458	127,203	95,775	63,264	39,655	313,661	512,355	38,505	550,860
Males	98,180	64,958	44,976	27,690	16,325	163,138	252,129	15,303 ^h	267,432
Females	88,278	62,245	50,799	35,574	23,330	150,523	260,226	23,202 ^h	283,428
White ^b	177,406	120,797	89,989	60,841	37,821	298,203	486,854	34,500	521,354
Males	93,419	61,948	42,405	26,690	15,532	155,367	239,994	13,784 ^h	253,778
Females	83,987	58,849	47,584	34,151	22,289	142,836	246,860	20,716 ^h	267,576
Black ^b	953	391	401	-	-	1,344	2,041	+	+
Males	478	-	-	-	-	657	848	+	+
Females	475	-	-	-	-	687	1,193	+	+
Asian/Pacific Islander ^c	529	244	+	+	+	773	+	+	+
Males	266	99	+	+	+	365	+	+	+
Females	263	145	+	+	+	408	+	+	+
Hispanic ^d	2,923	1,750	+	+	+	4,673	+	+	+
Males	1,178	687	+	+	+	1,865	+	+	+
Females	1,745	1,063	+	+	+	2,808	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0956%	-	0.0021%	0.0237%	0.0691%	0.1292%	0.1957%	0.2552%	0.3017%	0.2920%	0.1018%
Males	0.1013%	-	0.0015%	0.0170%	0.0524%	0.1229%	0.2193%	0.3335%	0.4228%	0.4530%	0.1182%
Females	0.0901%	-	0.0028%	0.0307%	0.0862%	0.1354%	0.1731%	0.1861%	0.2115%	0.2082%	0.0906%
White ^c	0.1157%	-	0.0026%	0.0282%	0.0830%	0.1555%	0.2289%	0.2999%	0.3451%	0.3206%	0.1190%
Males	0.1220%	-	0.0017%	0.0197%	0.0614%	0.1459%	0.2540%	0.3882%	0.4819%	0.5064%	0.1365%
Females	0.1096%	-	0.0035%	0.0374%	0.1060%	0.1653%	0.2045%	0.2201%	0.2421%	0.2257%	0.1075%
Black ^c	0.0036%	-	-	-	-	-	0.0086%	0.0139%	0.0178%	-	0.0050%
Males	0.0037%	-	-	-	-	-	-	-	-	-	0.0062%
Females	0.0035%	-	-	-	-	-	-	-	-	-	0.0043%
Asian/Pacific Islander ^c	0.0075%	-	-	-	0.0050%	0.0072%	0.0153%	0.0175%	0.0302%	0.0360%	0.0084%
Males	0.0074%	-	-	-	-	-	0.0170%	0.0205%	0.0366%	-	0.0092%
Females	0.0075%	-	-	-	0.0068%	0.0086%	0.0138%	-	0.0254%	-	0.0079%
Hispanic ^d	0.0133%	-	-	0.0038%	0.0138%	0.0267%	0.0416%	0.0507%	0.0824%	0.1004%	0.0235%
Males	0.0102%	-	-	-	0.0078%	0.0202%	0.0327%	0.0505%	0.1074%	0.1209%	0.0225%
Females	0.0166%	-	-	0.0059%	0.0206%	0.0333%	0.0496%	0.0509%	0.0646%	0.0894%	0.0252%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g h} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25. (h) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XVII-14
MYELOMA (Invasive)Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
All Races ^b	33,366	9,302	2,686	1,127	384	42,668	46,865	+	+
Males	17,836	5,051	1,494	557	-	22,887	25,088	+	+
Females	15,530	4,251	1,192	570	-	19,781	21,777	+	+
White ^b	26,249	7,026	2,048	765	-	33,275	36,329	+	+
Males	14,426	3,835	1,142	347	-	18,261	19,844	+	+
Females	11,823	3,191	906	418	-	15,014	16,485	+	+
Black ^b	6,144	1,976	536	340	-	8,120	9,124	+	+
Males	2,881	1,041	321	-	-	3,922	4,495	+	+
Females	3,263	935	-	-	-	4,198	4,629	+	+
Asian/Pacific Islander ^c	537	161	+	+	+	698	+	+	+
Males	303	88	+	+	+	391	+	+	+
Females	234	73	+	+	+	307	+	+	+
Hispanic ^d	1,833	437	+	+	+	2,270	+	+	+
Males	981	251	+	+	+	1,232	+	+	+
Females	852	186	+	+	+	1,038	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.0134%	-	-	-	0.0013%	0.0065%	0.0239%	0.0526%	0.0773%	0.0780%	0.0149%
Males	0.0146%	-	-	-	0.0018%	0.0082%	0.0287%	0.0643%	0.0933%	0.0978%	0.0182%
Females	0.0122%	-	-	-	0.0009%	0.0049%	0.0194%	0.0424%	0.0653%	0.0677%	0.0123%
White ^c	0.0131%	-	-	-	0.0014%	0.0056%	0.0217%	0.0491%	0.0728%	0.0719%	0.0138%
Males	0.0144%	-	-	-	0.0019%	0.0072%	0.0272%	0.0594%	0.0892%	0.0907%	0.0171%
Females	0.0118%	-	-	-	-	0.0039%	0.0163%	0.0398%	0.0605%	0.0623%	0.0111%
Black ^c	0.0210%	-	-	-	-	0.0164%	0.0528%	0.1138%	0.1566%	0.1553%	0.0315%
Males	0.0219%	-	-	-	-	0.0205%	0.0568%	0.1405%	0.1771%	0.2057%	0.0372%
Females	0.0202%	-	-	-	-	0.0127%	0.0494%	0.0928%	0.1430%	0.1327%	0.0275%
Asian/Pacific Islander ^c	0.0073%	-	-	-	-	0.0042%	0.0149%	0.0225%	0.0486%	0.0598%	0.0090%
Males	0.0084%	-	-	-	-	-	0.0162%	0.0333%	0.0593%	0.0754%	0.0113%
Females	0.0063%	-	-	-	-	-	0.0138%	-	0.0404%	-	0.0072%
Hispanic ^d	0.0063%	-	-	-	0.0022%	0.0054%	0.0217%	0.0441%	0.0736%	0.0829%	0.0139%
Males	0.0066%	-	-	-	-	0.0065%	0.0248%	0.0489%	0.0794%	0.1451%	0.0173%
Females	0.0060%	-	-	-	-	-	0.0189%	0.0401%	0.0695%	-	0.0116%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b ^c ^d Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^f ^g (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XVIII-19
NON-HODGKIN LYMPHOMA (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	144,844	77,015	43,437	23,792	12,350	221,859	301,438	12,962	314,400
Males	75,865	39,258	22,045	12,036	5,922	115,123	155,126	6,980	162,106
Females	68,979	37,757	21,392	11,756	6,428	106,736	146,312	5,982	152,294
White ^b	129,635	69,533	39,728	21,814	11,464	199,168	272,174	13,846	286,020
Males	67,559	35,232	20,212	11,076	5,551	102,791	139,630	7,966	147,596
Females	62,076	34,301	19,516	10,738	5,913	96,377	132,544	5,880	138,424
Black ^b	9,837	4,870	2,446	1,276	677	14,707	19,106	612	19,718
Males	5,449	2,623	1,215	617	-	8,072	10,172	318	10,490
Females	4,388	2,247	1,231	659	409	6,635	8,934	294	9,228
Asian/Pacific Islander ^c	3,020	1,477	+	+	+	4,497	+	+	+
Males	1,613	733	+	+	+	2,346	+	+	+
Females	1,407	744	+	+	+	2,151	+	+	+
Hispanic ^d	7,080	3,461	+	+	+	10,541	+	+	+
Males	3,814	1,694	+	+	+	5,508	+	+	+
Females	3,266	1,767	+	+	+	5,033	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0706%	0.0021%	0.0079%	0.0126%	0.0285%	0.0582%	0.1234%	0.2217%	0.3263%	0.3379%	0.0768%
Males	0.0746%	0.0034%	0.0110%	0.0150%	0.0340%	0.0688%	0.1429%	0.2496%	0.3644%	0.3854%	0.0883%
Females	0.0667%	-	0.0046%	0.0102%	0.0229%	0.0478%	0.1048%	0.1972%	0.2979%	0.3131%	0.0670%
White ^c	0.0782%	0.0020%	0.0084%	0.0131%	0.0294%	0.0609%	0.1308%	0.2391%	0.3508%	0.3516%	0.0815%
Males	0.0818%	0.0031%	0.0116%	0.0155%	0.0349%	0.0712%	0.1508%	0.2678%	0.3903%	0.4006%	0.0931%
Females	0.0746%	-	0.0049%	0.0105%	0.0235%	0.0506%	0.1112%	0.2131%	0.3211%	0.3265%	0.0713%
Black ^c	0.0397%	-	0.0053%	0.0123%	0.0273%	0.0552%	0.1020%	0.1369%	0.1533%	0.1670%	0.0515%
Males	0.0452%	-	0.0071%	0.0164%	0.0332%	0.0709%	0.1262%	0.1546%	0.1712%	0.2032%	0.0621%
Females	0.0347%	-	-	0.0087%	0.0220%	0.0414%	0.0817%	0.1228%	0.1415%	0.1508%	0.0429%
Asian/Pacific Islander ^c	0.0448%	-	0.0081%	0.0098%	0.0202%	0.0366%	0.0712%	0.1491%	0.2181%	0.2700%	0.0521%
Males	0.0479%	-	0.0115%	0.0105%	0.0230%	0.0418%	0.0785%	0.1681%	0.2509%	0.3071%	0.0596%
Females	0.0420%	-	-	0.0092%	0.0175%	0.0319%	0.0646%	0.1336%	0.1933%	0.2440%	0.0460%
Hispanic ^d	0.0297%	0.0024%	0.0062%	0.0088%	0.0184%	0.0399%	0.0837%	0.1553%	0.2634%	0.2533%	0.0562%
Males	0.0301%	0.0038%	0.0085%	0.0102%	0.0214%	0.0435%	0.0960%	0.1611%	0.2827%	0.2502%	0.0604%
Females	0.0293%	-	-	0.0073%	0.0151%	0.0364%	0.0726%	0.1506%	0.2496%	0.2549%	0.0522%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XIX-14
ORAL CAVITY AND PHARYNX CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	82,193	50,625	33,074	21,741	14,560	132,818	202,193	19,215	221,408
Males	54,722	32,620	20,158	13,452	8,912	87,342	129,864	13,071	142,935
Females	27,471	18,005	12,916	8,289	5,648	45,476	72,329	6,144	78,473
White ^b	70,799	44,761	29,625	19,704	13,016	115,560	177,905	15,772	193,677
Males	47,443	29,030	18,245	12,421	8,158	76,473	115,297	10,628	125,925
Females	23,356	15,731	11,380	7,283	4,858	39,087	62,608	5,144	67,752
Black ^b	7,237	3,704	2,238	1,246	914	10,941	15,339	+	+
Males	4,752	2,214	1,231	604	361	6,966	9,162	380	9,542
Females	2,485	1,490	1,007	642	553	3,975	6,177	+	+
Asian/Pacific Islander ^c	2,573	1,345	+	+	+	3,918	+	+	+
Males	1,658	863	+	+	+	2,521	+	+	+
Females	915	482	+	+	+	1,397	+	+	+
Hispanic ^d	3,096	1,737	+	+	+	4,833	+	+	+
Males	2,060	996	+	+	+	3,056	+	+	+
Females	1,036	741	+	+	+	1,777	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0427%	-	0.0016%	0.0048%	0.0124%	0.0369%	0.0905%	0.1570%	0.1857%	0.1713%	0.0467%
Males	0.0567%	-	0.0016%	0.0046%	0.0143%	0.0515%	0.1330%	0.2293%	0.2770%	0.2672%	0.0679%
Females	0.0291%	-	0.0015%	0.0049%	0.0105%	0.0225%	0.0501%	0.0932%	0.1177%	0.1214%	0.0293%
White ^c	0.0451%	-	0.0016%	0.0044%	0.0112%	0.0339%	0.0903%	0.1648%	0.1938%	0.1783%	0.0473%
Males	0.0598%	-	0.0016%	0.0043%	0.0130%	0.0474%	0.1331%	0.2391%	0.2876%	0.2841%	0.0689%
Females	0.0306%	-	0.0017%	0.0045%	0.0093%	0.0202%	0.0486%	0.0977%	0.1233%	0.1242%	0.0294%
Black ^c	0.0296%	-	-	0.0052%	0.0101%	0.0401%	0.0898%	0.1380%	0.1309%	0.0845%	0.0394%
Males	0.0397%	-	-	-	0.0118%	0.0541%	0.1336%	0.2129%	0.2114%	0.1164%	0.0582%
Females	0.0204%	-	-	-	0.0086%	0.0278%	0.0530%	0.0787%	0.0779%	0.0701%	0.0253%
Asian/Pacific Islander ^c	0.0378%	-	-	0.0062%	0.0196%	0.0517%	0.0849%	0.1100%	0.1444%	0.1345%	0.0424%
Males	0.0502%	-	-	-	0.0221%	0.0754%	0.1244%	0.1584%	0.2141%	0.1714%	0.0598%
Females	0.0263%	-	-	-	0.0173%	0.0305%	0.0500%	0.0707%	0.0916%	0.1086%	0.0279%
Hispanic ^d	0.0135%	-	-	0.0029%	0.0058%	0.0178%	0.0483%	0.0840%	0.1185%	0.1245%	0.0271%
Males	0.0165%	-	-	-	0.0060%	0.0223%	0.0710%	0.1249%	0.1797%	0.1808%	0.0389%
Females	0.0104%	-	-	-	0.0057%	0.0134%	0.0278%	0.0507%	0.0750%	0.0944%	0.0181%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XX-10
OVARY CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race</u>									
All Races ^b	70,400	42,213	25,933	17,685	13,578	112,613	169,809	33,140 ^h	202,949
White ^b	62,914	37,690	23,540	16,020	12,177	100,604	152,341	27,175 ^h	179,516
Black ^b	4,610	2,827	1,624	1,107	1,021	7,437	11,189	+	+
Asian/Pacific Islander ^c	2,009	1,145	+	+	+	3,154	+	+	+
Hispanic ^d	4,435	2,684	+	+	+	7,119	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	0-9	Age Specific (Crude)							Age-Adjusted ^e All Ages	
			10-19	20-29	30-39	40-49	50-59	60-69	70-79		80+
<u>Race</u>											
All Races ^c	0.0741%	-	0.0055%	0.0225%	0.0479%	0.0905%	0.1594%	0.2063%	0.2142%	0.1460%	0.0753%
White ^c	0.0821%	-	0.0050%	0.0227%	0.0514%	0.0970%	0.1722%	0.2278%	0.2301%	0.1525%	0.0810%
Black ^c	0.0395%	-	-	0.0180%	0.0302%	0.0605%	0.0872%	0.1116%	0.1508%	0.0982%	0.0470%
Asian/Pacific Islander ^c	0.0567%	-	0.0103%	0.0236%	0.0419%	0.0736%	0.1332%	0.1363%	0.1325%	0.1062%	0.0584%
Hispanic ^d	0.0420%	-	0.0055%	0.0238%	0.0394%	0.0679%	0.1320%	0.1644%	0.1952%	0.1278%	0.0632%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g h} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25. (h) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XXI-14
PANCREAS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	17,407	3,126	1,438	773	390	20,533	23,134	400	23,534
Males	8,680	1,412	555	319	-	10,092	11,166	198	11,364
Females	8,727	1,714	883	454	-	10,441	11,968	202	12,170
White ^b	14,636	2,656	939	651	350	17,292	19,232	311	19,543
Males	7,437	1,182	357	295	-	8,619	9,456	159	9,615
Females	7,199	1,474	582	356	-	8,673	9,776	152	9,928
Black ^b	2,026	336	401	-	-	2,362	2,895	78	2,973
Males	947	-	-	-	-	1,114	1,305	43	1,348
Females	1,079	-	-	-	-	1,248	1,590	35	1,625
Asian/Pacific Islander ^c	477	82	+	+	+	559	+	+	+
Males	202	-	+	+	+	239	+	+	+
Females	275	-	+	+	+	320	+	+	+
Hispanic ^d	964	248	+	+	+	1,212	+	+	+
Males	406	-	+	+	+	511	+	+	+
Females	558	143	+	+	+	701	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0066%	-	-	-	0.0012%	0.0035%	0.0113%	0.0274%	0.0356%	0.0357%	0.0073%
Males	0.0065%	-	-	-	0.0010%	0.0036%	0.0131%	0.0327%	0.0371%	0.0385%	0.0081%
Females	0.0067%	-	-	-	0.0014%	0.0035%	0.0095%	0.0228%	0.0345%	0.0342%	0.0067%
White ^c	0.0069%	-	-	-	0.0011%	0.0034%	0.0111%	0.0281%	0.0339%	0.0349%	0.0072%
Males	0.0068%	-	-	-	-	0.0036%	0.0131%	0.0340%	0.0355%	0.0380%	0.0081%
Females	0.0069%	-	-	-	0.0013%	0.0032%	0.0093%	0.0228%	0.0327%	0.0334%	0.0065%
Black ^c	0.0058%	-	-	-	-	0.0045%	0.0133%	0.0276%	0.0495%	0.0406%	0.0087%
Males	0.0059%	-	-	-	-	-	0.0178%	0.0305%	0.0522%	-	0.0098%
Females	0.0058%	-	-	-	-	-	-	0.0253%	0.0477%	-	0.0079%
Asian/Pacific Islander ^c	0.0058%	-	-	-	-	-	0.0099%	0.0228%	0.0373%	0.0396%	0.0071%
Males	0.0052%	-	-	-	-	-	-	0.0262%	0.0371%	-	0.0070%
Females	0.0064%	-	-	-	-	-	-	0.0200%	0.0375%	-	0.0071%
Hispanic ^d	0.0034%	-	-	-	-	0.0039%	0.0088%	0.0239%	0.0368%	0.0494%	0.0074%
Males	0.0027%	-	-	-	-	-	-	0.0307%	-	-	0.0072%
Females	0.0040%	-	-	-	-	-	-	0.0184%	0.0390%	-	0.0076%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.
 - Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.
 + Not available.

Table XXII-10
PROSTATE CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race</u>									
All Races ^b	819,511	588,141	152,551	46,239	15,756	1,407,652	1,622,198	15,010	1,637,208
White ^b	685,949	513,474	138,206	41,698	14,155	1,199,423	1,393,482	15,919	1,409,401
Black ^b	104,419	60,678	12,057	3,995	1,431	165,097	182,580	1,326	183,906
Asian/Pacific Islander ^c	11,264	6,146	+	+	+	17,410	+	+	+
Hispanic ^d	32,992	17,951	+	+	+	50,943	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	0.8761%	-	-	-	-	0.0427%	0.7034%	3.7619%	8.5564%	9.2620%	1.1740%
White ^c	0.9205%	-	-	-	-	0.0376%	0.6873%	3.7467%	8.5122%	9.0022%	1.1577%
Black ^c	0.9056%	-	-	-	-	0.1044%	1.3088%	6.1735%	12.5219%	12.0319%	1.7584%
Asian/Pacific Islander ^c	0.4170%	-	-	-	-	0.0135%	0.2073%	1.4279%	4.6443%	6.5568%	0.6266%
Hispanic ^d	0.2682%	-	-	-	-	0.0187%	0.3938%	2.4883%	6.9089%	7.5599%	0.8991%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XXIII-14
STOMACH CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	29,179	11,579	7,614	4,212	2,495	40,758	55,079	4,448	59,527
Males	17,369	6,351	4,031	2,349	1,379	23,720	31,479	2,426	33,905
Females	11,810	5,228	3,583	1,863	1,116	17,038	23,600	2,022	25,622
White ^b	22,513	8,768	5,968	3,213	2,067	31,281	42,529	3,457	45,986
Males	13,688	4,884	3,213	1,851	1,126	18,572	24,762	1,939	26,701
Females	8,825	3,884	2,755	1,362	941	12,709	17,767	1,518	19,285
Black ^b	4,191	1,675	1,039	649	-	5,866	7,815	431	8,246
Males	2,235	911	515	330	-	3,146	4,154	213	4,367
Females	1,956	764	524	319	-	2,720	3,661	218	3,879
Asian/Pacific Islander ^c	2,211	972	+	+	+	3,183	+	+	+
Males	1,294	539	+	+	+	1,833	+	+	+
Females	917	433	+	+	+	1,350	+	+	+
Hispanic ^d	2,933	1,056	+	+	+	3,989	+	+	+
Males	1,625	585	+	+	+	2,210	+	+	+
Females	1,308	471	+	+	+	1,779	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0159%	-	-	0.0005%	0.0029%	0.0088%	0.0205%	0.0519%	0.0981%	0.1196%	0.0177%
Males	0.0186%	-	-	-	0.0028%	0.0104%	0.0266%	0.0709%	0.1352%	0.1643%	0.0238%
Females	0.0133%	-	-	-	0.0030%	0.0071%	0.0146%	0.0351%	0.0704%	0.0964%	0.0131%
White ^c	0.0133%	-	-	-	0.0024%	0.0064%	0.0167%	0.0426%	0.0747%	0.0947%	0.0140%
Males	0.0157%	-	-	-	0.0026%	0.0076%	0.0224%	0.0596%	0.1035%	0.1300%	0.0190%
Females	0.0109%	-	-	-	0.0023%	0.0052%	0.0112%	0.0272%	0.0530%	0.0766%	0.0101%
Black ^c	0.0157%	-	-	-	-	0.0153%	0.0273%	0.0650%	0.1280%	0.1645%	0.0240%
Males	0.0173%	-	-	-	-	0.0189%	0.0334%	0.0903%	0.1717%	0.1744%	0.0301%
Females	0.0143%	-	-	-	-	0.0121%	0.0222%	0.0450%	0.0992%	0.1601%	0.0194%
Asian/Pacific Islander ^c	0.0344%	-	-	-	0.0051%	0.0190%	0.0407%	0.1049%	0.2611%	0.3506%	0.0432%
Males	0.0410%	-	-	-	-	0.0227%	0.0519%	0.1406%	0.3628%	0.4566%	0.0570%
Females	0.0282%	-	-	-	-	0.0157%	0.0308%	0.0759%	0.1840%	0.2763%	0.0326%
Hispanic ^d	0.0111%	-	-	-	0.0050%	0.0116%	0.0274%	0.0645%	0.1397%	0.1971%	0.0254%
Males	0.0118%	-	-	-	0.0051%	0.0111%	0.0317%	0.0932%	0.1823%	0.2582%	0.0325%
Females	0.0103%	-	-	-	0.0049%	0.0121%	0.0234%	0.0411%	0.1094%	0.1645%	0.0204%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XXIV-10
TESTIS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race</u>									
All Races ^b	34,721	32,106	27,686	21,014	14,087	66,827	129,614	21,237 ^h	150,851
White ^b	32,680	30,633	26,610	20,278	13,525	63,313	123,726	19,583 ^h	143,309
Black ^b	962	496	469	342	-	1,458	2,506	+	+
Asian/Pacific Islander ^c	531	467	+	+	+	998	+	+	+
Hispanic ^d	2,928	2,271	+	+	+	5,199	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	0.0465%	0.0022%	0.0041%	0.0481%	0.1083%	0.0938%	0.0420%	0.0182%	0.0103%	0.0065%	0.0448%
White ^c	0.0558%	0.0022%	0.0045%	0.0584%	0.1305%	0.1107%	0.0493%	0.0211%	0.0121%	-	0.0533%
Black ^c	0.0093%	-	-	-	0.0193%	0.0241%	-	-	-	-	0.0094%
Asian/Pacific Islander ^c	0.0175%	-	-	0.0180%	0.0361%	0.0367%	0.0141%	-	-	-	0.0166%
Hispanic ^d	0.0297%	-	0.0041%	0.0388%	0.0698%	0.0566%	0.0221%	-	-	-	0.0289%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XXV-14
 THYROID CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	78,174	57,487	44,328	33,818	31,359	135,661	245,166	47,389	292,555
Males	18,056	13,957	9,935	7,230	7,218	32,013	56,396	8,626 ^h	65,022
Females	60,118	43,530	34,393	26,588	24,141	103,648	188,770	38,763 ^h	227,533
White ^b	68,647	51,024	39,765	30,601	28,239	119,671	218,276	39,915	258,191
Males	16,396	12,668	9,164	6,474	6,702	29,064	51,404	7,449 ^h	58,853
Females	52,251	38,356	30,601	24,127	21,537	90,607	166,872	32,466 ^h	199,338
Black ^b	4,501	3,298	1,994	1,596	1,588	7,799	12,977	+	+
Males	686	719	336	356	-	1,405	2,284	+	+
Females	3,815	2,579	1,658	1,240	1,401	6,394	10,693	+	+
Asian/Pacific Islander ^c	3,412	2,331	+	+	+	5,743	+	+	+
Males	626	430	+	+	+	1,056	+	+	+
Females	2,786	1,901	+	+	+	4,687	+	+	+
Hispanic ^d	5,908	4,379	+	+	+	10,287	+	+	+
Males	1,057	867	+	+	+	1,924	+	+	+
Females	4,851	3,512	+	+	+	8,363	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0469%	-	0.0033%	0.0266%	0.0607%	0.0815%	0.0905%	0.0858%	0.0749%	0.0497%	0.0481%
Males	0.0219%	-	0.0012%	0.0078%	0.0207%	0.0355%	0.0488%	0.0570%	0.0543%	0.0350%	0.0237%
Females	0.0714%	-	0.0055%	0.0461%	0.1017%	0.1267%	0.1302%	0.1111%	0.0903%	0.0573%	0.0716%
White ^c	0.0499%	-	0.0036%	0.0292%	0.0662%	0.0862%	0.0929%	0.0869%	0.0726%	0.0472%	0.0502%
Males	0.0240%	-	-	0.0086%	0.0226%	0.0389%	0.0520%	0.0603%	0.0547%	0.0324%	0.0252%
Females	0.0756%	-	0.0061%	0.0513%	0.1125%	0.1342%	0.1327%	0.1110%	0.0861%	0.0547%	0.0751%
Black ^c	0.0208%	-	-	0.0103%	0.0232%	0.0369%	0.0555%	0.0484%	0.0590%	-	0.0249%
Males	0.0085%	-	-	-	0.0082%	0.0135%	0.0269%	0.0254%	0.0452%	-	0.0115%
Females	0.0320%	-	-	0.0181%	0.0366%	0.0573%	0.0795%	0.0666%	0.0682%	-	0.0360%
Asian/Pacific Islander ^c	0.0522%	-	-	0.0253%	0.0585%	0.0893%	0.1017%	0.1037%	0.1017%	0.0917%	0.0544%
Males	0.0203%	-	-	-	0.0188%	0.0316%	0.0414%	0.0537%	0.0566%	0.0704%	0.0231%
Females	0.0822%	-	-	0.0430%	0.0957%	0.1409%	0.1553%	0.1444%	0.1359%	0.1067%	0.0820%
Hispanic ^d	0.0299%	-	0.0031%	0.0183%	0.0438%	0.0702%	0.0802%	0.0745%	0.0847%	0.0699%	0.0422%
Males	0.0106%	-	-	0.0052%	0.0108%	0.0244%	0.0347%	0.0481%	0.0568%	-	0.0181%
Females	0.0500%	-	0.0054%	0.0335%	0.0807%	0.1161%	0.1216%	0.0961%	0.1045%	0.0897%	0.0653%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g h} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25. (h) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XXVI-14
URINARY BLADDER CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	180,332	118,309	77,702	47,214	26,184	298,641	449,741	29,395	479,136
Males	134,833	88,472	57,731	34,455	18,022	223,305	333,513	20,020	353,533
Females	45,499	29,837	19,971	12,759	8,162	75,336	116,228	9,375	125,603
White ^b	168,728	111,753	74,003	45,243	24,876	280,481	424,603	23,179	447,782
Males	126,775	83,817	54,991	33,040	17,154	210,592	315,777	15,751	331,528
Females	41,953	27,936	19,012	12,203	7,722	69,889	108,826	7,428	116,254
Black ^b	7,124	4,231	2,527	1,261	726	11,355	15,869	+	+
Males	4,690	2,876	1,882	878	536	7,566	10,862	+	+
Females	2,434	1,355	645	383	-	3,789	5,007	113	5,120
Asian/Pacific Islander ^c	2,043	1,186	+	+	+	3,229	+	+	+
Males	1,513	910	+	+	+	2,423	+	+	+
Females	530	276	+	+	+	806	+	+	+
Hispanic ^d	4,422	2,829	+	+	+	7,251	+	+	+
Males	3,382	2,143	+	+	+	5,525	+	+	+
Females	1,040	686	+	+	+	1,726	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0901%	-	-	0.0017%	0.0064%	0.0271%	0.1061%	0.3191%	0.6225%	0.7205%	0.1013%
Males	0.1373%	-	-	0.0021%	0.0094%	0.0417%	0.1691%	0.5296%	1.1155%	1.4651%	0.1811%
Females	0.0440%	-	-	0.0013%	0.0034%	0.0128%	0.0463%	0.1335%	0.2550%	0.3329%	0.0436%
White ^c	0.1069%	-	-	0.0021%	0.0073%	0.0309%	0.1207%	0.3665%	0.6915%	0.7723%	0.1129%
Males	0.1622%	-	-	0.0027%	0.0102%	0.0467%	0.1899%	0.6014%	1.2411%	1.5966%	0.2013%
Females	0.0521%	-	-	0.0015%	0.0041%	0.0149%	0.0533%	0.1540%	0.2781%	0.3510%	0.0483%
Black ^c	0.0306%	-	-	-	-	0.0152%	0.0545%	0.1349%	0.2918%	0.3605%	0.0484%
Males	0.0440%	-	-	-	-	0.0257%	0.0885%	0.2285%	0.4906%	0.6462%	0.0821%
Females	0.0184%	-	-	-	-	0.0260%	0.0610%	0.1608%	0.2325%	0.2325%	0.0259%
Asian/Pacific Islander ^c	0.0357%	-	-	-	-	0.0109%	0.0391%	0.1198%	0.2856%	0.4002%	0.0459%
Males	0.0551%	-	-	-	-	0.0183%	0.0640%	0.2017%	0.4855%	0.7179%	0.0790%
Females	0.0174%	-	-	-	-	0.0169%	0.0532%	0.1342%	0.1773%	0.1773%	0.0205%
Hispanic ^d	0.0199%	-	-	-	0.0034%	0.0107%	0.0451%	0.1396%	0.2953%	0.4252%	0.0496%
Males	0.0294%	-	-	-	0.0044%	0.0142%	0.0767%	0.2497%	0.5465%	0.8270%	0.0908%
Females	0.0100%	-	-	-	-	0.0073%	0.0164%	0.0496%	0.1167%	0.2108%	0.0210%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b ^c ^d Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^f ^g (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.

Table XXVII-12
 CHILDHOOD CANCER (Invasive, <20 Years old at diagnosis)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	52,974	44,110	37,906	31,324	27,777	97,084	194,091	+	+
Males	28,094	23,182	19,331	15,713	12,813	51,276	99,133	+	+
Females	24,880	20,928	18,575	15,611	14,964	45,808	94,958	+	+
White ^b	44,348	36,920	32,634	27,374	24,481	81,268	165,757	+	+
Males	23,439	19,714	16,675	13,877	11,396	43,153	85,101	+	+
Females	20,909	17,206	15,959	13,497	13,085	38,115	80,656	+	+
Black ^b	5,524	4,765	3,555	2,775	2,372	10,289	18,991	+	+
Males	2,973	2,284	1,735	1,280	1,004	5,257	9,276	+	+
Females	2,551	2,481	1,820	1,495	1,368	5,032	9,715	+	+
Asian/Pacific Islander ^c	1,873	1,395	+	+	+	3,268	+	+	+
Males	1,018	667	+	+	+	1,685	+	+	+
Females	855	728	+	+	+	1,583	+	+	+
Hispanic ^d	7,557	5,538	+	+	+	13,095	+	+	+
Males	3,990	3,089	+	+	+	7,079	+	+	+
Females	3,567	2,449	+	+	+	6,016	+	+	+

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{f g} (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.

- Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.

+ Not available.