

Table X-1

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1976-2001

All Races, Males, All Ages

Year of Diagnosis

SEER Location	1976-1977	1978-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001
All Areas <sup>b</sup>	62	43	79	328	888	1578	2145	2247	2065	1802	838	386	386
All Areas but SF-Oakland	45	33	44	87	259	428	775	843	975	872	462	242	255
San Francisco-Oakland SMSA	17	10	35	241	629	1150	1370	1404	1090	930	376	144	131
San Francisco City/County	9	<4	21	201	540	956	1085	1099	799	690	253	85	78

AGE-ADJUSTED CANCER INCIDENCE RATES<sup>a</sup>

By SEER Location and Year of Diagnosis, 1976-2001

All Races, Males, All Ages

Year of Diagnosis

SEER Location	1976-1977	1978-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001
All Areas <sup>b</sup>	0.5	0.3	0.5	1.7	4.2	7.1	9.2	9.4	8.2	7.0	3.2	1.5	1.5
All Areas but SF-Oakland	0.4	0.3	0.4	0.6	1.5	2.3	4.0	4.2	4.6	4.0	2.1	1.1	1.2
San Francisco-Oakland SMSA	0.9	0.4	1.3	7.0	17.2	30.1	34.3	34.3	25.8	22.0	8.9	3.4	3.0
San Francisco City/County	1.5	-	3.1	25.1	65.3	111.4	125.4	126.7	89.8	77.5	28.1	9.3	8.4

<sup>a</sup> SEER 9 areas. Rates are per 100,000 males and are age-adjusted to the 2000 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

<sup>b</sup> All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.

Table X-2

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1976-2001

All Races, Males, Ages 20-54

## Year of Diagnosis

SEER Location	1976-1977	1978-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001
All Areas <sup>b</sup>	6	4	26	254	807	1484	2007	2117	1930	1676	754	313	333
All Areas but SF-Oakland	5	<4	5	35	207	378	694	773	909	806	409	195	216
San Francisco-Oakland SMSA	<4	<4	21	219	600	1106	1313	1344	1021	870	345	118	117
San Francisco City/County	<4	0	18	193	520	925	1049	1061	755	649	236	75	72

AGE-ADJUSTED CANCER INCIDENCE RATES<sup>a</sup>

By SEER Location and Year of Diagnosis, 1976-2001

All Races, Males, Ages 20-54

## Year of Diagnosis

SEER Location	1976-1977	1978-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001
All Areas <sup>b</sup>	0.1	0.0	0.3	2.4	7.4	13.1	17.0	17.2	15.0	12.7	5.6	2.3	2.4
All Areas but SF-Oakland	0.1	-	0.1	0.4	2.2	3.9	6.9	7.4	8.4	7.2	3.6	1.7	1.9
San Francisco-Oakland SMSA	-	-	1.3	12.1	32.4	57.5	65.0	64.5	47.2	40.3	15.9	5.2	5.1
San Francisco City/County	-	0.0	5.3	47.8	125.1	214.2	240.7	242.3	166.8	142.7	51.4	15.8	15.2

<sup>a</sup> SEER 9 areas. Rates are per 100,000 males aged 20-54 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

<sup>b</sup> All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.

Table X-3

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1976-2001

All Races, Males, Ages 55+

## Year of Diagnosis

SEER Location	1976-1977	1978-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001
All Areas <sup>b</sup>	56	39	53	73	80	93	135	130	134	126	84	73	53
All Areas but SF-Oakland	40	30	39	51	52	50	81	70	65	66	53	47	39
San Francisco-Oakland SMSA	16	9	14	22	28	43	54	60	69	60	31	26	14
San Francisco City/County	8	<4	<4	8	20	31	35	38	44	41	17	10	6

AGE-ADJUSTED CANCER INCIDENCE RATES<sup>a</sup>

By SEER Location and Year of Diagnosis, 1976-2001

All Races, Males, Ages 55+

## Year of Diagnosis

SEER Location	1976-1977	1978-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001
All Areas <sup>b</sup>	2.2	1.4	1.9	2.4	2.4	2.6	3.6	3.6	3.3	3.0	2.1	1.7	1.3
All Areas but SF-Oakland	1.8	1.3	1.7	2.0	2.0	1.8	2.7	2.5	2.0	1.9	1.6	1.3	1.1
San Francisco-Oakland SMSA	4.1	1.8	3.1	4.6	4.7	6.4	8.1	9.6	10.4	8.8	4.6	3.7	2.0
San Francisco City/County	6.0	-	-	5.6	12.9	20.5	23.5	26.4	30.5	29.0	11.5	6.7	3.7


<sup>a</sup> SEER 9 areas. Rates are per 100,000 males aged 55+ and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

<sup>b</sup> All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.


Figure X-1

# All Cancer Sites vs Kaposi's Sarcoma SEER Incidence Rates, 1975-2001 Males, Ages 20-54




Source: SEER 9 areas. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups.  
Regression lines are calculated using the Joinpoint Regression Program Version 2.7, September 2003, National Cancer Institute.

All SEER excluding  
San Francisco/Oakland  
Males, Ages 20-54  
1975-2001



San Francisco/Oakland  
Males, Ages 20-54  
1975-2001



San Francisco County/City  
Males, Ages 20-54  
1975-2001




Figure X-2

Source: SEER 9 areas. Rates are age-adjusted to the 2000 U.S. standard million population by 5-year age groups.

\* Kaposi's sarcoma excluded from Colorectal, Lung, and NHL<sup>^</sup>.

<sup>^</sup> Non-Hodgkin Lymphoma.