

Table VIII-1
ESOPHAGUS CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2003
WITH UP TO THREE JOINPOINTS BY RACE AND SEX

| | | Joinpoint Segment 1 | | Joinpoint Segment 2 | | Joinpoint Segment 3 | | Joinpoint Segment 4 | |
|--|-----------------|---------------------|-------|---------------------|-------|---------------------|-------|---------------------|-----|
| | | Year | APC | Year | APC | Year | APC | Year | APC |
| <u>SEER Cancer Incidence^a</u> | | | | | | | | | |
| All Races | Male and Female | 1975-2003 | 0.5* | | | | | | |
| All Races | Male | 1975-2003 | 0.7* | | | | | | |
| All Races | Female | 1975-2003 | -0.4* | | | | | | |
| White | Male and Female | 1975-2003 | 1.3* | | | | | | |
| White | Male | 1975-2003 | 1.6* | | | | | | |
| White | Female | 1975-2003 | 0.0 | | | | | | |
| Black | Male and Female | 1975-1986 | 1.0 | 1986-2003 | -4.4* | | | | |
| Black | Male | 1975-1986 | 1.1 | 1986-2003 | -4.8* | | | | |
| Black | Female | 1975-1980 | 9.9 | 1980-2003 | -2.6* | | | | |
| <u>SEER Delay-Adjusted Incidence^a</u> | | | | | | | | | |
| All Races | Male and Female | 1975-2003 | 0.5* | | | | | | |
| All Races | Male | 1975-2003 | 0.7* | | | | | | |
| All Races | Female | 1975-2003 | -0.3* | | | | | | |
| White | Male and Female | 1975-1981 | -0.4 | 1981-2003 | 1.6* | | | | |
| White | Male | 1975-2003 | 1.7* | | | | | | |
| White | Female | 1975-2003 | 0.0 | | | | | | |
| Black | Male and Female | 1975-1986 | 1.0 | 1986-2003 | -4.4* | | | | |
| Black | Male | 1975-1986 | 1.1 | 1986-2003 | -4.8* | | | | |
| Black | Female | 1975-1980 | 9.8 | 1980-2003 | -2.5* | | | | |
| <u>U.S. Cancer Mortality^b</u> | | | | | | | | | |
| All Races | Male and Female | 1975-2001 | 0.7* | 2001-2003 | -0.6 | | | | |
| All Races | Male | 1975-1985 | 0.7* | 1985-1994 | 1.2* | 1994-2003 | 0.5* | | |
| All Races | Female | 1975-2003 | 0.0 | | | | | | |
| White | Male and Female | 1975-1983 | 0.4* | 1983-2001 | 1.7* | 2001-2003 | 0.1 | | |
| White | Male | 1975-1982 | 0.4 | 1982-2000 | 1.9* | 2000-2003 | 0.6 | | |
| White | Female | 1975-2003 | 0.4* | | | | | | |
| Black | Male and Female | 1975-1984 | 0.6 | 1984-1993 | -1.9* | 1993-2003 | -4.3* | | |
| Black | Male | 1975-1984 | 0.8 | 1984-1993 | -1.9* | 1993-2003 | -4.7* | | |
| Black | Female | 1975-1990 | 0.0 | 1990-2003 | -3.2* | | | | |

Joinpoint Regression Program Version 3.1, April 2006, National Cancer Institute. (<http://srab.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

^b Trends are from the NCHS public use data file for the total US.

* The APC is significantly different from zero (p<.05).

Table VIII-2
ESOPHAGUS CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE^a RATES BY YEAR, RACE AND SEX

| | All Races | | | Whites | | | Blacks | | |
|---------------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| <u>AGE-ADJUSTED RATES</u> | | | | | | | | | |
| YEAR OF DIAGNOSIS: | | | | | | | | | |
| 1975 | 4.1 | 6.8 | 2.1 | 3.5 | 5.6 | 2.0 | 10.9 | 19.9 | 3.7 |
| 1976 | 4.3 | 6.9 | 2.3 | 3.7 | 5.9 | 2.1 | 10.7 | 16.9 | 5.6 |
| 1977 | 4.1 | 6.6 | 2.2 | 3.3 | 5.3 | 1.9 | 12.0 | 20.9 | 5.4 |
| 1978 | 4.1 | 7.0 | 1.9 | 3.4 | 5.7 | 1.7 | 13.0 | 24.4 | 4.3 |
| 1979 | 4.4 | 7.4 | 2.2 | 3.7 | 6.2 | 2.0 | 12.9 | 23.0 | 5.2 |
| 1980 | 4.3 | 6.9 | 2.3 | 3.5 | 5.7 | 1.9 | 12.3 | 18.8 | 7.4 |
| 1981 | 4.1 | 6.5 | 2.4 | 3.3 | 5.2 | 1.9 | 13.5 | 21.3 | 7.5 |
| 1982 | 4.3 | 6.8 | 2.3 | 3.5 | 5.5 | 2.0 | 13.6 | 22.7 | 6.4 |
| 1983 | 4.3 | 7.4 | 2.0 | 3.6 | 6.1 | 1.8 | 12.2 | 22.2 | 4.3 |
| 1984 | 4.2 | 6.7 | 2.2 | 3.5 | 5.6 | 1.9 | 12.0 | 19.9 | 6.0 |
| 1985 | 4.4 | 7.4 | 2.2 | 3.7 | 6.3 | 1.9 | 12.7 | 22.1 | 5.8 |
| 1986 | 4.6 | 7.5 | 2.5 | 3.9 | 6.2 | 2.2 | 14.1 | 24.0 | 6.3 |
| 1987 | 4.5 | 7.5 | 2.3 | 3.9 | 6.4 | 2.0 | 11.4 | 20.1 | 4.8 |
| 1988 | 4.4 | 7.3 | 2.2 | 3.9 | 6.3 | 2.0 | 11.2 | 19.5 | 5.0 |
| 1989 | 4.4 | 7.3 | 2.2 | 3.8 | 6.2 | 2.0 | 10.5 | 17.3 | 5.4 |
| 1990 | 4.8 | 8.3 | 2.2 | 4.2 | 7.2 | 1.9 | 12.5 | 21.4 | 5.8 |
| 1991 | 4.6 | 7.4 | 2.4 | 4.2 | 6.7 | 2.2 | 10.2 | 17.8 | 4.9 |
| 1992 | 4.6 | 8.0 | 2.0 | 4.2 | 7.4 | 1.8 | 10.6 | 18.0 | 5.3 |
| 1993 | 4.6 | 7.7 | 2.2 | 4.2 | 6.9 | 2.1 | 10.0 | 17.5 | 4.6 |
| 1994 | 4.5 | 7.7 | 2.1 | 4.2 | 7.2 | 1.9 | 9.2 | 14.7 | 5.1 |
| 1995 | 4.4 | 7.1 | 2.2 | 4.1 | 6.6 | 2.1 | 8.3 | 13.7 | 4.3 |
| 1996 | 4.8 | 8.1 | 2.2 | 4.4 | 7.5 | 2.0 | 9.7 | 16.4 | 4.8 |
| 1997 | 4.7 | 7.8 | 2.2 | 4.6 | 7.7 | 2.0 | 7.1 | 10.8 | 4.5 |
| 1998 | 4.7 | 8.2 | 2.0 | 4.5 | 7.9 | 1.9 | 7.7 | 12.4 | 4.2 |
| 1999 | 4.9 | 8.4 | 2.1 | 4.9 | 8.4 | 2.0 | 7.2 | 11.5 | 4.1 |
| 2000 | 4.8 | 8.0 | 2.2 | 4.7 | 8.0 | 2.0 | 7.0 | 10.6 | 4.4 |
| 2001 | 4.8 | 8.4 | 1.9 | 4.8 | 8.4 | 1.9 | 6.3 | 11.4 | 2.6 |
| 2002 | 4.6 | 7.8 | 2.0 | 4.6 | 8.0 | 1.8 | 6.3 | 10.0 | 3.6 |
| 2003 | 4.6 | 7.9 | 1.9 | 4.7 | 8.1 | 1.9 | 6.4 | 10.1 | 3.8 |
| 1975-2003 | 4.5 | 7.5 | 2.2 | 4.1 | 6.8 | 2.0 | 9.9 | 16.7 | 4.9 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VIII-3
ESOPHAGUS CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH^a RATES BY YEAR, RACE AND SEX

| | All Races | | | Whites | | | Blacks | | |
|---------------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| <u>AGE-ADJUSTED RATES</u> | | | | | | | | | |
| YEAR OF DEATH: | | | | | | | | | |
| 1975 | 3.7 | 6.4 | 1.7 | 3.1 | 5.4 | 1.5 | 9.7 | 17.0 | 4.1 |
| 1976 | 3.7 | 6.4 | 1.8 | 3.1 | 5.3 | 1.5 | 10.2 | 17.7 | 4.3 |
| 1977 | 3.7 | 6.3 | 1.8 | 3.1 | 5.2 | 1.5 | 10.1 | 17.6 | 4.4 |
| 1978 | 3.8 | 6.4 | 1.8 | 3.2 | 5.4 | 1.6 | 10.0 | 17.6 | 4.3 |
| 1979 | 3.8 | 6.4 | 1.8 | 3.1 | 5.3 | 1.5 | 10.3 | 18.6 | 4.2 |
| 1980 | 3.8 | 6.6 | 1.8 | 3.2 | 5.5 | 1.5 | 10.6 | 18.3 | 4.9 |
| 1981 | 3.8 | 6.6 | 1.8 | 3.2 | 5.4 | 1.5 | 10.4 | 18.7 | 4.3 |
| 1982 | 3.8 | 6.5 | 1.8 | 3.2 | 5.4 | 1.6 | 10.5 | 18.6 | 4.5 |
| 1983 | 3.9 | 6.6 | 1.8 | 3.2 | 5.6 | 1.5 | 10.2 | 17.9 | 4.5 |
| 1984 | 3.9 | 6.7 | 1.9 | 3.3 | 5.6 | 1.6 | 10.6 | 18.8 | 4.6 |
| 1985 | 3.9 | 6.7 | 1.7 | 3.3 | 5.7 | 1.5 | 9.9 | 17.7 | 4.3 |
| 1986 | 3.9 | 6.8 | 1.8 | 3.3 | 5.8 | 1.6 | 10.2 | 18.0 | 4.5 |
| 1987 | 4.0 | 6.9 | 1.8 | 3.4 | 6.0 | 1.5 | 9.7 | 17.1 | 4.4 |
| 1988 | 4.0 | 7.0 | 1.8 | 3.5 | 6.0 | 1.6 | 10.0 | 18.4 | 3.9 |
| 1989 | 4.1 | 7.1 | 1.8 | 3.6 | 6.2 | 1.6 | 9.7 | 17.3 | 4.3 |
| 1990 | 4.1 | 7.2 | 1.8 | 3.6 | 6.3 | 1.5 | 9.3 | 16.3 | 4.4 |
| 1991 | 4.1 | 7.2 | 1.8 | 3.6 | 6.4 | 1.6 | 9.2 | 16.5 | 4.1 |
| 1992 | 4.2 | 7.3 | 1.8 | 3.7 | 6.6 | 1.6 | 8.6 | 15.1 | 4.1 |
| 1993 | 4.2 | 7.3 | 1.8 | 3.7 | 6.6 | 1.6 | 9.0 | 16.0 | 4.1 |
| 1994 | 4.3 | 7.6 | 1.8 | 3.9 | 6.9 | 1.6 | 8.6 | 15.3 | 3.9 |
| 1995 | 4.3 | 7.6 | 1.8 | 3.9 | 7.0 | 1.6 | 8.0 | 14.4 | 3.5 |
| 1996 | 4.3 | 7.5 | 1.9 | 4.0 | 7.1 | 1.7 | 7.8 | 13.6 | 3.7 |
| 1997 | 4.3 | 7.5 | 1.8 | 4.0 | 7.1 | 1.6 | 7.3 | 12.8 | 3.4 |
| 1998 | 4.4 | 7.7 | 1.8 | 4.1 | 7.3 | 1.7 | 7.1 | 12.2 | 3.6 |
| 1999 | 4.3 | 7.7 | 1.8 | 4.2 | 7.4 | 1.6 | 6.5 | 11.6 | 3.1 |
| 2000 | 4.4 | 7.7 | 1.8 | 4.3 | 7.6 | 1.7 | 6.4 | 10.9 | 3.3 |
| 2001 | 4.4 | 7.8 | 1.8 | 4.3 | 7.7 | 1.7 | 6.4 | 11.3 | 3.0 |
| 2002 | 4.4 | 7.8 | 1.7 | 4.3 | 7.8 | 1.6 | 5.9 | 10.2 | 2.9 |
| 2003 | 4.4 | 7.7 | 1.7 | 4.3 | 7.7 | 1.7 | 5.7 | 9.7 | 2.9 |
| 1975-2003 | 4.1 | 7.2 | 1.8 | 3.7 | 6.5 | 1.6 | 8.6 | 15.2 | 3.9 |

^a NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VIII-4
ESOPHAGUS CANCER (Invasive)

SEER INCIDENCE^a AND U.S. DEATH^b RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE AND SEX

| | All Races | | | Whites | | | Blacks | | |
|---------------------------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| <u>SEER INCIDENCE</u> | | | | | | | | | |
| <u>AGE AT DIAGNOSIS</u> | | | | | | | | | |
| AGE-ADJUSTED RATES, 2000-2003 | | | | | | | | | |
| All ages | 4.5 | 7.8 | 2.0 | 4.5 | 7.8 | 1.9 | 6.4 | 10.8 | 3.3 |
| Under 65 | 1.9 | 3.1 | 0.7 | 1.8 | 3.1 | 0.6 | 3.2 | 5.2 | 1.6 |
| 65 and over | 22.9 | 39.6 | 11.1 | 23.2 | 40.2 | 11.0 | 28.5 | 49.6 | 15.0 |
| All ages(IARC world std) ^c | 3.0 | 5.1 | 1.2 | 3.0 | 5.1 | 1.2 | 4.5 | 7.5 | 2.2 |
| AGE-SPECIFIC RATES, 2000-2003 | | | | | | | | | |
| <1 | - | - | - | - | - | - | - | - | - |
| 1-4 | - | - | - | - | - | - | - | - | - |
| 5-9 | - | - | - | - | - | - | - | - | - |
| 10-14 | - | - | - | - | - | - | - | - | - |
| 15-19 | - | - | - | - | - | - | - | - | - |
| 20-24 | - | - | - | - | - | - | - | - | - |
| 25-29 | - | - | - | - | - | - | - | - | - |
| 30-34 | 0.2 | 0.3 | - | 0.2 | 0.3 | - | - | - | - |
| 35-39 | 0.4 | 0.6 | 0.2 | 0.4 | 0.7 | - | - | - | - |
| 40-44 | 1.0 | 1.7 | 0.3 | 1.0 | 1.7 | 0.3 | 1.8 | 2.9 | - |
| 45-49 | 2.7 | 4.3 | 1.1 | 2.6 | 4.4 | 0.9 | 4.3 | 5.7 | 3.0 |
| 50-54 | 5.2 | 9.2 | 1.4 | 5.1 | 9.1 | 1.2 | 9.4 | 15.5 | 4.2 |
| 55-59 | 9.3 | 15.7 | 3.3 | 9.0 | 15.3 | 2.9 | 17.0 | 28.1 | 7.7 |
| 60-64 | 13.8 | 23.0 | 5.3 | 13.5 | 22.7 | 5.1 | 23.1 | 38.3 | 10.9 |
| 65-69 | 19.3 | 32.6 | 7.9 | 19.1 | 32.1 | 7.7 | 29.3 | 50.3 | 13.2 |
| 70-74 | 23.8 | 39.9 | 10.9 | 24.1 | 40.7 | 10.7 | 31.8 | 52.1 | 17.7 |
| 75-79 | 25.2 | 44.6 | 11.6 | 25.9 | 46.1 | 11.6 | 31.2 | 55.3 | 16.3 |
| 80-84 | 24.9 | 43.9 | 13.2 | 25.3 | 45.0 | 13.1 | 25.7 | 43.3 | 16.2 |
| 85+ | 23.0 | 40.8 | 15.0 | 23.7 | 41.7 | 15.7 | 18.6 | 40.4 | 10.0 |
| <u>U.S. MORTALITY</u> | | | | | | | | | |
| <u>AGE AT DEATH</u> | | | | | | | | | |
| AGE-ADJUSTED RATES, 2000-2003 | | | | | | | | | |
| All ages | 4.4 | 7.8 | 1.8 | 4.3 | 7.7 | 1.7 | 6.1 | 10.5 | 3.0 |
| Under 65 | 1.7 | 3.0 | 0.5 | 1.6 | 2.9 | 0.4 | 2.9 | 5.0 | 1.2 |
| 65 and over | 22.9 | 40.7 | 10.4 | 22.8 | 40.7 | 10.1 | 28.3 | 49.1 | 15.5 |
| All ages(IARC world std) ^c | 2.8 | 5.0 | 1.0 | 2.8 | 4.9 | 1.0 | 4.2 | 7.3 | 2.0 |
| AGE-SPECIFIC RATES, 2000-2003 | | | | | | | | | |
| <1 | - | - | - | - | - | - | - | - | - |
| 1-4 | - | - | - | - | - | - | - | - | - |
| 5-9 | - | - | - | - | - | - | - | - | - |
| 10-14 | - | - | - | - | - | - | - | - | - |
| 15-19 | - | - | - | - | - | - | - | - | - |
| 20-24 | - | - | - | - | - | - | - | - | - |
| 25-29 | 0.0 | 0.1 | - | 0.0 | 0.1 | - | - | - | - |
| 30-34 | 0.1 | 0.2 | 0.0 | 0.1 | 0.2 | - | - | - | - |
| 35-39 | 0.3 | 0.6 | 0.1 | 0.4 | 0.7 | 0.1 | 0.3 | 0.4 | - |
| 40-44 | 1.0 | 1.7 | 0.3 | 1.0 | 1.7 | 0.3 | 1.5 | 2.4 | 0.7 |
| 45-49 | 2.4 | 4.1 | 0.7 | 2.3 | 4.0 | 0.6 | 3.9 | 6.3 | 1.8 |
| 50-54 | 4.8 | 8.4 | 1.3 | 4.5 | 7.9 | 1.1 | 8.3 | 14.4 | 3.2 |
| 55-59 | 8.5 | 14.9 | 2.6 | 8.0 | 14.1 | 2.2 | 15.4 | 27.0 | 6.0 |
| 60-64 | 12.9 | 22.6 | 4.1 | 12.3 | 21.8 | 3.6 | 21.6 | 37.0 | 9.5 |
| 65-69 | 18.1 | 31.3 | 6.6 | 17.4 | 30.4 | 6.0 | 27.5 | 47.1 | 13.0 |
| 70-74 | 22.4 | 39.1 | 9.1 | 22.2 | 38.9 | 8.6 | 30.0 | 51.3 | 15.7 |
| 75-79 | 25.4 | 45.6 | 11.2 | 25.4 | 45.8 | 10.8 | 31.2 | 55.3 | 16.8 |
| 80-84 | 27.3 | 49.8 | 13.7 | 27.5 | 50.6 | 13.4 | 28.3 | 47.7 | 18.3 |
| 85+ | 25.3 | 46.0 | 16.5 | 25.9 | 47.3 | 16.8 | 21.9 | 39.8 | 15.1 |

^a SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^c Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VIII-5
ESOPHAGUS CANCER (Invasive)

SURVIVAL RATES, BY RACE, SEX, DIAGNOSIS YEAR, STAGE AND AGE

| | All Races | | | Whites | | | Blacks | | |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 5-YR RELATIVE SURVIVAL RATES | | | | | | | | | |
| YEAR OF DIAGNOSIS: | | | | | | | | | |
| 1960-1963 ^a | - | - | - | 4 | 4 | 6 | 1 | 0 | 2 |
| 1970-1973 ^a | - | - | - | 4 | 4 | 4 | 4 | 4 | 3 |
| 1975-1977 ^b | 5.1 | 4.2 | 7.4 | 5.7 | 5.1 | 6.9 | 3.1 | 1.6 | 7.1 |
| 1978-1980 ^b | 5.1 | 4.9 | 5.6 | 5.5 | 5.5 | 5.2 | 4.3 | 3.3 | 7.0 |
| 1981-1983 ^b | 6.8 | 6.2 | 8.3 | 7.5 | 6.8 | 9.1 | 4.3 | 3.6 | 6.4 |
| 1984-1986 ^b | 9.6 | 8.5 | 12.3 | 10.5 | 9.2 | 13.3 | 8.2 | 7.6 | 9.9 |
| 1987-1989 ^b | 9.7 | 9.6 | 10.0 | 11.0 | 11.4 | 9.7 | 6.4 | 5.0 | 10.3 |
| 1990-1992 ^b | 12.6 | 12.2 | 13.8 | 13.5 | 12.8 | 15.4 | 9.4 | 9.8 | 8.4 |
| 1993-1995 ^b | 12.9 | 13.1 | 12.4 | 14.6 | 14.8 | 13.9 | 7.6 | 8.1 | 6.3 |
| 1996-2002 ^b | 16.2 ^e | 15.9 ^e | 16.9 ^e | 17.1 ^e | 16.9 ^e | 17.8 ^e | 11.9 ^e | 10.8 ^e | 14.1 |
| 5-YR PERIOD SURVIVAL RATES^{cd} | | | | | | | | | |
| YEAR OF DIAGNOSIS: | | | | | | | | | |
| 1996-2002 | 16.3 | 16.0 | 17.5 | 17.0 | 16.5 | 18.9 | 12.5 | 13.0 | 11.9 |
| STAGE DISTRIBUTION (%) 1996-2002^c | | | | | | | | | |
| All Stages | | | | | | | | | |
| Number of cases | 12,553 | 9,484 | 3,069 | 10,181 | 7,781 | 2,400 | 1,620 | 1,130 | 490 |
| Percent | 100% | 100% | 100% | 100% | 100% | 100% | 100% | 100% | 100% |
| Localized | 24 | 24 | 25 | 25 | 24 | 27 | 20 | 20 | 20 |
| Regional | 29 | 30 | 27 | 29 | 30 | 26 | 28 | 29 | 28 |
| Distant | 30 | 31 | 25 | 29 | 31 | 24 | 33 | 34 | 31 |
| Unstaged | 17 | 15 | 23 | 16 | 14 | 23 | 19 | 17 | 21 |
| 5-YR RELATIVE SURVIVAL RATES, 1996-2002^c | | | | | | | | | |
| AGE AT DIAGNOSIS: | | | | | | | | | |
| <45 | 22.4 | 21.7 | 25.8 ^f | 21.6 | 20.7 | 29.2 ^f | 21.8 | 22.7 ^f | 19.5 ^f |
| 45-54 | 18.3 | 18.0 | 19.7 | 20.1 | 20.1 | 18.7 | 11.8 | 10.1 | 18.4 ^f |
| 55-64 | 16.1 | 15.1 | 20.5 | 16.8 | 15.8 | 22.2 | 11.9 | 10.5 | 14.9 |
| 65-74 | 15.5 | 15.3 | 15.9 | 16.5 | 16.1 | 18.0 | 6.4 | 7.3 | 6.0 |
| 75+ | 11.0 | 10.7 | 11.1 | 11.5 | 11.2 | 11.6 | 7.9 | 10.6 ^f | 4.7 |
| Under 65 | 17.4 | 16.7 | 20.9 | 18.4 | 17.8 | 21.9 | 13.1 | 11.8 | 16.5 |
| 65 and over | 13.8 | 13.9 | 13.6 | 14.6 | 14.5 | 14.8 | 6.9 | 8.1 | 5.8 |
| STAGE: | | | | | | | | | |
| All Stages | 15.6 | 15.4 | 16.0 | 16.3 | 16.2 | 16.8 | 11.1 | 10.8 | 11.8 |
| Localized | 33.6 | 34.3 | 30.8 | 35.8 | 37.1 | 31.0 | 19.5 | 15.5 | 26.4 ^f |
| Regional | 16.8 | 16.0 | 19.3 | 17.4 | 16.7 | 20.1 | 14.3 | 14.4 | 14.0 |
| Distant | 2.6 | 2.7 | 2.2 | 2.8 | 2.7 | 2.9 | 1.2 | 1.3 | - |
| Unstaged | 10.8 | 10.8 | 10.7 | 9.4 | 8.7 | 10.6 | 14.5 | 16.8 | 8.9 |

^a Rates are based on End Results data from a series of hospital registries and one population-based registry.

^b Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

^c Rates are based on follow-up of patients into 2003.

^d Rates are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2002. The remaining 13 SEER Areas contribute cases for the entire period 1996-2002. Rates are based on follow-up of patients into 2003.

^e Period survival provides a more up-to-date estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three calendar year blocks (2000-2002: 0-1 year survival), (1999-2001: 1-2 year survival), (1998-2000: 2-3 year survival), (1997-1999: 3-4 year survival), (1996-1998: 4-5 years survival).

^f The difference in rates between 1975-1977 and 1996-2002 is statistically significant ($p < .05$).

^g The standard error of the survival rate is between 5 and 10 percentage points.

⁻ Statistic could not be calculated due to fewer than 16 cases during the time period.

Table VIII-6
ESOPHAGUS CANCER (Invasive)

SURVIVAL RATES^a

By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

| | 1975- | 1980- | | | | | | | | | | | | | | | | | | | | |
|---|-------|-------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|--|--|
| | 1979 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | | |
| Relative Survival Rates (SEER^b) | | | | | | | | | | | | | | | | | | | | | | |
| 1-year | 27.7 | 31.4 | 36.0 | 35.0 | 35.6 | 36.1 | 38.7 | 39.2 | 41.8 | 41.8 | 38.2 | 39.0 | 41.6 | 43.8 | 41.7 | 42.0 | 44.9 | 47.8 | 47.8 | 42.9 | | |
| 2-year | 11.7 | 14.6 | 20.6 | 19.1 | 18.1 | 20.4 | 20.1 | 20.5 | 21.4 | 24.6 | 22.5 | 23.5 | 24.0 | 24.6 | 24.3 | 24.7 | 27.2 | 29.1 | 29.9 | | | |
| 3-year | 7.8 | 10.2 | 13.2 | 14.4 | 11.9 | 13.9 | 14.4 | 14.7 | 15.4 | 19.6 | 16.4 | 18.6 | 17.7 | 19.0 | 18.0 | 19.1 | 21.1 | 24.6 | | | | |
| 4-year | 6.0 | 8.5 | 10.0 | 11.8 | 9.5 | 12.0 | 11.6 | 11.8 | 13.2 | 17.0 | 14.2 | 15.6 | 13.8 | 16.4 | 14.8 | 14.9 | 18.7 | | | | | |
| 5-year | 5.0 | 7.1 | 8.8 | 10.5 | 8.9 | 10.0 | 10.2 | 10.3 | 12.2 | 15.3 | 12.7 | 13.8 | 12.2 | 15.7 | 13.0 | 13.3 | | | | | | |
| 6-year | 4.4 | 6.4 | 7.3 | 9.4 | 7.9 | 9.2 | 9.1 | 10.0 | 10.8 | 14.4 | 11.5 | 12.5 | 11.5 | 14.7 | 12.0 | | | | | | | |
| 7-year | 4.0 | 5.8 | 6.9 | 8.7 | 7.1 | 8.7 | 8.4 | 9.0 | 10.3 | 13.0 | 10.8 | 10.8 | 11.6 | 11.0 | 11.0 | 13.8 | | | | | | |
| 8-year | 3.6 | 5.2 | 6.0 | 7.9 | 6.2 | 8.3 | 8.3 | 8.6 | 9.2 | 12.5 | 10.5 | 10.5 | 11.0 | 11.0 | | | | | | | | |
| 9-year | 3.3 | 4.8 | 5.5 | 7.6 | 5.5 | 7.0 | 7.7 | 7.9 | 9.1 | 11.9 | 9.6 | 9.6 | 10.9 | | | | | | | | | |
| 10-year | 3.0 | 4.6 | 5.5 | 6.8 | 5.4 | 6.5 | 7.0 | 7.6 | 8.8 | 11.0 | 9.2 | | | | | | | | | | | |
| 11-year | 2.9 | 4.3 | 5.0 | 6.4 | 5.4 | 6.1 | 6.0 | 7.6 | 8.8 | 9.8 | | | | | | | | | | | | |
| 12-year | 2.8 | 4.2 | 4.3 | 6.1 | 5.4 | 5.7 | 4.8 | 5.8 | 8.5 | | | | | | | | | | | | | |
| 13-year | 2.6 | 4.0 | 4.3 | 5.8 | 4.2 | 5.7 | 4.6 | 5.8 | | | | | | | | | | | | | | |
| 14-year | 2.6 | 3.9 | 4.3 | 5.8 | 4.0 | 5.2 | 4.0 | | | | | | | | | | | | | | | |
| 15-year | 2.6 | 3.9 | 4.3 | 5.7 | 3.9 | 4.8 | | | | | | | | | | | | | | | | |
| 16-year | 2.5 | 3.8 | 4.3 | 4.4 | 3.7 | | | | | | | | | | | | | | | | | |
| 17-year | 2.5 | 3.6 | 3.5 | 3.9 | | | | | | | | | | | | | | | | | | |
| 18-year | 2.5 | 3.2 | 3.5 | | | | | | | | | | | | | | | | | | | |
| 19-year | 2.5 | 3.1 | | | | | | | | | | | | | | | | | | | | |
| 20-year | 2.4 | | | | | | | | | | | | | | | | | | | | | |

^a Survival rates are relative rates expressed as percents.

^b Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table VIII-7
ESOPHAGUS CANCER (Invasive)

SURVIVAL RATES^a

By Year of Diagnosis

All Races, Males

Year of Diagnosis

| | 1975- | 1980- | | | | | | | | | | | | | | | | | | | | |
|---|-------|-------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|--|--|
| | 1979 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | | |
| Relative Survival Rates (SEER^b) | | | | | | | | | | | | | | | | | | | | | | |
| 1-year | 25.4 | 30.7 | 35.0 | 34.2 | 34.0 | 34.1 | 38.3 | 41.3 | 41.4 | 41.2 | 39.4 | 38.4 | 41.3 | 45.1 | 42.1 | 42.2 | 44.6 | 49.1 | 49.0 | 44.2 | | |
| 2-year | 10.3 | 14.1 | 19.6 | 19.7 | 17.2 | 19.1 | 19.6 | 21.4 | 21.1 | 24.4 | 22.2 | 22.0 | 24.5 | 25.2 | 25.1 | 25.0 | 26.9 | 29.5 | 30.4 | | | |
| 3-year | 6.9 | 9.5 | 11.1 | 15.6 | 12.3 | 12.1 | 14.4 | 14.8 | 15.5 | 19.5 | 16.8 | 18.3 | 18.0 | 18.8 | 19.0 | 19.2 | 21.2 | 24.3 | | | | |
| 4-year | 5.2 | 7.7 | 7.2 | 12.5 | 9.7 | 10.7 | 11.5 | 11.8 | 12.9 | 16.0 | 14.4 | 15.1 | 13.9 | 15.7 | 15.8 | 14.6 | 18.5 | | | | | |
| 5-year | 4.3 | 6.2 | 7.0 | 11.2 | 9.1 | 9.4 | 10.0 | 10.1 | 12.2 | 14.2 | 12.8 | 13.7 | 12.6 | 14.7 | 13.9 | 12.5 | | | | | | |
| 6-year | 3.8 | 5.4 | 6.2 | 10.3 | 8.4 | 8.8 | 8.5 | 9.7 | 10.6 | 13.2 | 11.8 | 12.3 | 11.9 | 13.9 | 12.7 | | | | | | | |
| 7-year | 3.5 | 4.7 | 5.7 | 9.4 | 7.1 | 8.5 | 7.8 | 8.8 | 10.6 | 11.9 | 11.5 | 11.6 | 11.6 | 11.6 | 13.0 | | | | | | | |
| 8-year | 3.2 | 4.3 | 5.0 | 8.2 | 6.4 | 8.1 | 7.6 | 8.1 | 9.1 | 11.3 | 11.2 | 10.7 | 10.7 | 11.4 | | | | | | | | |
| 9-year | 3.0 | 4.0 | 4.8 | 8.1 | 6.0 | 6.8 | 7.5 | 7.4 | 9.0 | 10.9 | 10.1 | 10.1 | 10.4 | | | | | | | | | |
| 10-year | 2.8 | 3.8 | 4.8 | 7.7 | 6.0 | 6.3 | 7.3 | 7.1 | 8.5 | 9.9 | 9.4 | | | | | | | | | | | |
| 11-year | 2.7 | 3.7 | 4.8 | 7.0 | 6.0 | 5.8 | 6.5 | 7.1 | 8.5 | 8.3 | | | | | | | | | | | | |
| 12-year | 2.5 | 3.6 | 3.6 | 6.5 | 6.0 | 5.2 | 5.5 | 5.6 | 8.1 | | | | | | | | | | | | | |
| 13-year | 2.5 | 3.6 | 3.6 | 6.0 | 4.5 | 5.1 | 5.1 | 5.6 | | | | | | | | | | | | | | |
| 14-year | 2.4 | 3.5 | 3.6 | 6.0 | 4.1 | 4.5 | 4.2 | | | | | | | | | | | | | | | |
| 15-year | 2.4 | 3.5 | 3.6 | 5.9 | 3.9 | 3.7 | | | | | | | | | | | | | | | | |
| 16-year | 2.4 | 3.5 | 3.6 | 4.8 | 3.5 | | | | | | | | | | | | | | | | | |
| 17-year | 2.4 | 3.5 | 2.6 | 3.8 | | | | | | | | | | | | | | | | | | |
| 18-year | 2.4 | 3.3 | 2.6 | | | | | | | | | | | | | | | | | | | |
| 19-year | 2.4 | 3.2 | | | | | | | | | | | | | | | | | | | | |
| 20-year | 2.4 | | | | | | | | | | | | | | | | | | | | | |

^a Survival rates are relative rates expressed as percents.

^b Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table VIII-8
ESOPHAGUS CANCER (Invasive)

SURVIVAL RATES^a

By Year of Diagnosis

All Races, Females

Year of Diagnosis

| | 1975- | 1980- | | | | | | | | | | | | | | | | | | | | |
|---|-------|-------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|--|--|
| | 1979 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | | |
| Relative Survival Rates (SEER^b) | | | | | | | | | | | | | | | | | | | | | | |
| 1-year | 33.5 | 33.2 | 38.8 | 36.9 | 39.8 | 41.3 | 40.0 | 33.0 | 42.9 | 43.8 | 35.2 | 41.0 | 42.4 | 40.1 | 40.3 | 41.5 | 46.0 | 44.0 | 43.3 | 39.0 | | |
| 2-year | 15.1 | 16.0 | 23.5 | 17.7 | 20.4 | 23.6 | 21.5 | 17.6 | 22.1 | 25.4 | 23.2 | 28.1 | 22.4 | 22.6 | 21.9 | 23.4 | 28.3 | 27.9 | 28.0 | | | |
| 3-year | 10.2 | 12.1 | 18.7 | 11.6 | 10.9 | 18.5 | 14.5 | 14.5 | 14.9 | 20.0 | 15.4 | 19.5 | 16.8 | 19.7 | 15.1 | 18.7 | 20.8 | 25.5 | | | | |
| 4-year | 8.0 | 10.6 | 17.1 | 10.0 | 8.7 | 15.4 | 11.8 | 11.8 | 13.7 | 19.6 | 13.4 | 16.5 | 13.7 | 18.1 | 11.7 | 15.7 | 19.0 | | | | | |
| 5-year | 6.5 | 9.4 | 13.4 | 8.8 | 7.9 | 11.4 | 10.7 | 11.0 | 12.2 | 18.5 | 12.2 | 14.2 | 10.6 | 18.1 | 10.6 | 15.6 | | | | | | |
| 6-year | 6.0 | 8.7 | 10.2 | 7.1 | 6.6 | 10.2 | 10.4 | 10.6 | 11.1 | 18.3 | 10.7 | 13.1 | 10.1 | 16.7 | 9.8 | | | | | | | |
| 7-year | 5.2 | 8.4 | 9.8 | 6.8 | 6.6 | 9.3 | 9.6 | 9.7 | 9.5 | 16.4 | 8.7 | 11.6 | 9.1 | 16.1 | | | | | | | | |
| 8-year | 4.5 | 7.5 | 8.2 | 6.8 | 5.3 | 8.5 | 9.6 | 9.7 | 9.2 | 16.3 | 8.5 | 11.3 | 9.1 | | | | | | | | | |
| 9-year | 4.1 | 6.6 | 7.2 | 6.6 | 4.2 | 7.4 | 8.1 | 9.2 | 9.2 | 14.7 | 8.3 | 11.3 | | | | | | | | | | |
| 10-year | 3.6 | 6.6 | 6.8 | 4.5 | 3.5 | 7.2 | 6.2 | 8.8 | 9.2 | 14.4 | 8.3 | | | | | | | | | | | |
| 11-year | 3.4 | 5.6 | 5.5 | 4.5 | 3.5 | 6.8 | 4.6 | 8.8 | 9.2 | 14.4 | | | | | | | | | | | | |
| 12-year | 3.2 | 5.3 | 5.5 | 4.5 | 3.5 | 6.8 | 3.1 | 6.3 | 9.1 | | | | | | | | | | | | | |
| 13-year | 2.9 | 4.9 | 5.5 | 4.5 | 3.5 | 6.8 | 3.1 | 5.8 | | | | | | | | | | | | | | |
| 14-year | 2.9 | 4.7 | 5.5 | 4.5 | 3.5 | 6.8 | 3.1 | | | | | | | | | | | | | | | |
| 15-year | 2.9 | 4.7 | 5.5 | 4.5 | 3.5 | 6.8 | | | | | | | | | | | | | | | | |
| 16-year | 2.8 | 4.0 | 5.2 | 3.6 | 3.5 | | | | | | | | | | | | | | | | | |
| 17-year | 2.5 | 3.2 | 5.2 | 3.6 | | | | | | | | | | | | | | | | | | |
| 18-year | 2.3 | 3.1 | 5.2 | | | | | | | | | | | | | | | | | | | |
| 19-year | 2.3 | 2.8 | | | | | | | | | | | | | | | | | | | | |
| 20-year | 2.3 | | | | | | | | | | | | | | | | | | | | | |

^a Survival rates are relative rates expressed as percents.

^b Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table VIII-9
ESOPHAGUS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

| Race/ Ethnicity | Current Age | Males | | | | Females | | | |
|-------------------------------|----------------|---------|---------|---------|------------|---------|---------|---------|------------|
| | | +10 yrs | +20 yrs | +30 yrs | Eventually | +10 yrs | +20 yrs | +30 yrs | Eventually |
| All Races | 0 | 0.00 | 0.00 | 0.00 | 0.76 | 0.00 | 0.00 | 0.00 | 0.25 |
| | 10 | 0.00 | 0.00 | 0.01 | 0.77 | 0.00 | 0.00 | 0.00 | 0.25 |
| | 20 | 0.00 | 0.01 | 0.04 | 0.78 | 0.00 | 0.00 | 0.01 | 0.25 |
| | 30 | 0.00 | 0.04 | 0.15 | 0.79 | 0.00 | 0.01 | 0.03 | 0.25 |
| | 40 | 0.03 | 0.15 | 0.37 | 0.80 | 0.01 | 0.03 | 0.09 | 0.25 |
| | 50 | 0.12 | 0.35 | 0.62 | 0.79 | 0.02 | 0.08 | 0.17 | 0.25 |
| | 60 | 0.25 | 0.54 | 0.70 | 0.73 | 0.06 | 0.15 | 0.22 | 0.24 |
| White | 0 | 0.00 | 0.00 | 0.00 | 0.78 | 0.00 | 0.00 | 0.00 | 0.24 |
| | 10 | 0.00 | 0.00 | 0.01 | 0.78 | 0.00 | 0.00 | 0.00 | 0.25 |
| | 20 | 0.00 | 0.01 | 0.04 | 0.79 | 0.00 | 0.00 | 0.01 | 0.25 |
| | 30 | 0.00 | 0.04 | 0.15 | 0.80 | 0.00 | 0.01 | 0.03 | 0.25 |
| | 40 | 0.03 | 0.14 | 0.37 | 0.81 | 0.01 | 0.03 | 0.08 | 0.25 |
| | 50 | 0.12 | 0.35 | 0.63 | 0.80 | 0.02 | 0.08 | 0.16 | 0.25 |
| | 60 | 0.25 | 0.55 | 0.71 | 0.74 | 0.06 | 0.15 | 0.22 | 0.24 |
| Black | 0 | 0.00 | 0.00 | 0.00 | 0.82 | 0.00 | 0.00 | 0.00 | 0.30 |
| | 10 | 0.00 | 0.00 | 0.00 | 0.84 | 0.00 | 0.00 | 0.00 | 0.30 |
| | 20 | 0.00 | 0.00 | 0.05 | 0.85 | 0.00 | 0.00 | 0.02 | 0.30 |
| | 30 | 0.00 | 0.05 | 0.23 | 0.87 | 0.00 | 0.02 | 0.07 | 0.31 |
| | 40 | 0.05 | 0.24 | 0.54 | 0.89 | 0.02 | 0.07 | 0.16 | 0.31 |
| | 50 | 0.20 | 0.53 | 0.78 | 0.91 | 0.05 | 0.15 | 0.25 | 0.30 |
| | 60 | 0.38 | 0.68 | 0.80 | 0.82 | 0.10 | 0.22 | 0.27 | 0.27 |
| Asian/ Pacific Islander | 0 | 0.00 | 0.00 | 0.00 | 0.51 | 0.00 | 0.00 | 0.00 | 0.16 |
| | 10 | 0.00 | 0.00 | 0.00 | 0.52 | 0.00 | 0.00 | 0.00 | 0.16 |
| | 20 | 0.00 | 0.00 | 0.02 | 0.52 | 0.00 | 0.00 | 0.01 | 0.16 |
| | 30 | 0.00 | 0.02 | 0.07 | 0.52 | 0.00 | 0.01 | 0.02 | 0.16 |
| | 40 | 0.01 | 0.06 | 0.19 | 0.52 | 0.01 | 0.02 | 0.05 | 0.16 |
| | 50 | 0.05 | 0.18 | 0.33 | 0.52 | 0.01 | 0.04 | 0.09 | 0.16 |
| | 60 | 0.14 | 0.29 | 0.45 | 0.48 | 0.03 | 0.08 | 0.13 | 0.15 |
| Hispanic ^a | 0 | 0.00 | 0.00 | 0.00 | 0.62 | 0.00 | 0.00 | 0.00 | 0.22 |
| | 10 | 0.00 | 0.00 | 0.00 | 0.63 | 0.00 | 0.00 | 0.00 | 0.22 |
| | 20 | 0.00 | 0.00 | 0.02 | 0.63 | 0.00 | 0.00 | 0.00 | 0.22 |
| | 30 | 0.00 | 0.02 | 0.09 | 0.64 | 0.00 | 0.00 | 0.01 | 0.23 |
| | 40 | 0.02 | 0.08 | 0.23 | 0.65 | 0.00 | 0.01 | 0.04 | 0.23 |
| | 50 | 0.07 | 0.22 | 0.40 | 0.65 | 0.01 | 0.03 | 0.09 | 0.23 |
| | 60 | 0.16 | 0.35 | 0.52 | 0.62 | 0.03 | 0.09 | 0.14 | 0.23 |

| Race/Ethnicity | Lifetime Risk of Developing | | Lifetime Risk of Dying | |
|------------------------|-----------------------------|---------|------------------------|---------|
| | Males | Females | Males | Females |
| All Races | 0.76% | 0.25% | 0.75% | 0.22% |
| White | 0.78% | 0.24% | 0.75% | 0.21% |
| Black | 0.82% | 0.30% | 0.79% | 0.30% |
| Asian/Pacific Islander | 0.51% | 0.16% | 0.42% | 0.13% |
| Hispanic ^a | 0.62% | 0.22% | 0.48% | 0.16% |

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table VIII-10
ESOPHAGUS CANCER (Invasive)

SEER INCIDENCE AND U.S. MORTALITY
AGE-ADJUSTED RATES AND TRENDS^a
By Race/Ethnicity and Sex

SEER Incidence

| RACE/ETHNICITY | Rate 2000-2003 ^b Rate per 100,000 persons | | | Trend 1994-2003 ^c APC (%) | | |
|----------------------------------|---|-------|---------|---|-------|---------|
| | Total | Males | Females | Total | Males | Females |
| | All Races | 4.5 | 7.8 | 2.0 | 0.0 | 0.3 |
| White | 4.5 | 7.8 | 1.9 | 1.1 | 1.4* | -0.6 |
| White Hispanic ^d | 2.9 | 5.2 | 1.2 | -2.4 | -1.7 | -4.0 |
| White Non-Hispanic ^d | 4.7 | 8.1 | 2.1 | 1.8* | 2.1* | 0.1 |
| Black | 6.4 | 10.8 | 3.3 | -4.3* | -4.5* | -3.6* |
| Asian/Pacific Islander | 2.4 | 4.0 | 1.2 | -3.0* | -3.1* | -0.9 |
| Amer Ind/Alaska Nat ^e | 4.7 | 9.5 | - | - | - | - |
| Hispanic ^d | 2.9 | 5.2 | 1.2 | -2.2 | -1.6 | -3.5 |

U.S. Mortality^f

| RACE/ETHNICITY | Rate 2000-2003 Rate per 100,000 persons | | | Trend 1994-2003 APC (%) | | |
|---------------------------------|--|-------|---------|----------------------------|-------|---------|
| | Total | Males | Females | Total | Males | Females |
| | All Races | 4.4 | 7.8 | 1.8 | 0.4* | 0.4* |
| White | 4.3 | 7.7 | 1.7 | 1.4* | 1.4* | 0.5 |
| White Hispanic ^d | 2.5 | 4.4 | 1.0 | -1.5* | -2.0* | -0.5 |
| White Non-Hispanic ^d | 4.4 | 7.9 | 1.7 | 1.6* | 1.6* | 0.6* |
| Black | 6.1 | 10.5 | 3.0 | -4.3* | -4.7* | -3.2* |
| Asian/Pacific Islander | 1.8 | 3.0 | 0.8 | -3.7* | -3.8* | -3.2 |
| Amer Ind/Alaska Nat | 2.9 | 5.0 | 1.2 | 1.1 | 0.1 | - |
| Hispanic ^d | 2.4 | 4.2 | 1.0 | -1.5* | -2.0* | -0.7 |

The APC is the Annual Percent Change over the time interval.

— Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Incidence data used in calculating the rates are from the 17 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^c Incidence data used in calculating the trends are from the 13 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).

^d Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. The 2000-2003 Hispanic and Non-Hispanic death rates exclude deaths from Minnesota, New Hampshire and North Dakota. The 1994-2003 Hispanic and Non-Hispanic mortality trends exclude deaths from Maine, Minnesota, New Hampshire, North Dakota, and Oklahoma.

^e Incidence data for American Indians/Alaska Natives include cases from Connecticut, Detroit, Iowa, New Mexico, Seattle, Utah, Atlanta, and the Alaska Native Registry. Incidence rates are for diagnosis years 1999-2002. Incidence trends are for diagnosis years 1994-2002.

^f Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

* The APC is significantly different from zero ($p < .05$).

Table VIII-11
ESOPHAGUS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE RATES^a
By Registry, Race/Ethnicity and Sex

| | All Races | | | Whites | | Blacks | | Asian/Pacific Islanders | | Hispanics ^b | |
|--|-----------|-------|---------|--------|---------|--------|---------|-------------------------|---------|------------------------|---------|
| | Total | Males | Females | Males | Females | Males | Females | Males | Females | Males | Females |
| <u>SEER INCIDENCE RATES^a, 2000-2003</u> | | | | | | | | | | | |
| REGISTRY | | | | | | | | | | | |
| 17 SEER Areas ^c | 4.5 | 7.8 | 2.0 | 7.8 | 1.9 | 10.8 | 3.3 | 4.0 | 1.2 | 5.2 | 1.2 |
| 13 SEER Areas ^c | 4.4 | 7.6 | 2.0 | 7.7 | 1.9 | 10.1 | 3.5 | 4.4 | 1.1 | 5.4 | 0.9 |
| 11 SEER Areas ^c | 4.4 | 7.5 | 1.9 | 7.7 | 1.9 | 9.9 | 3.5 | 4.4 | 1.1 | 5.4 | 0.9 |
| 9 SEER Areas ^c | 4.7 | 8.0 | 2.0 | 8.1 | 1.9 | 10.5 | 3.6 | 5.0 | 1.0 | 5.3 | 1.0 |
| San Francisco-Oakland | 3.9 | 6.5 | 1.9 | 7.2 | 1.9 | 7.3 | 3.0 | 4.0 | 1.1 | 3.8 | - |
| Connecticut | 5.5 | 9.3 | 2.6 | 9.2 | 2.5 | 10.6 | 4.4 | - | - | 15.3 | - |
| Metropolitan Detroit | 5.5 | 9.6 | 2.4 | 8.9 | 2.0 | 12.4 | 4.0 | - | - | - | - |
| Hawaii | 3.4 | 6.1 | 1.0 | 5.8 | - | - | - | 6.2 | 0.9 | - | - |
| Iowa | 5.1 | 9.2 | 1.8 | 9.2 | 1.8 | - | - | - | - | - | - |
| New Mexico | 3.6 | 6.5 | 1.3 | 6.6 | 1.4 | - | - | - | - | 4.8 | - |
| Seattle-Puget Sound | 5.0 | 8.5 | 2.2 | 8.7 | 2.2 | - | - | 5.5 | - | - | - |
| Utah | 3.2 | 5.7 | 1.1 | 5.9 | 1.1 | - | - | - | - | - | - |
| Metropolitan Atlanta | 4.7 | 8.1 | 2.1 | 7.1 | 1.6 | 11.6 | 3.0 | - | - | - | - |
| San Jose-Monterey | 3.9 | 6.7 | 1.8 | 7.5 | 1.8 | - | - | 3.3 | - | 5.4 | - |
| Los Angeles | 3.7 | 6.1 | 1.9 | 6.2 | 1.8 | 8.4 | 3.1 | 3.6 | 1.2 | 5.5 | 1.0 |
| Rural Georgia | 8.0 | 12.8 | - | - | - | 24.1 | - | - | - | - | - |
| California exclud. SF/SJM/LA | 4.3 | 7.2 | 1.9 | 7.5 | 2.0 | 9.0 | 2.0 | 3.1 | 1.3 | 4.6 | 1.2 |
| Kentucky | 4.9 | 9.0 | 1.6 | 8.7 | 1.6 | 16.5 | 3.1 | - | - | - | - |
| Louisiana | 5.0 | 8.8 | 1.9 | 8.3 | 1.6 | 10.6 | 2.7 | - | - | - | - |
| New Jersey | 5.1 | 8.5 | 2.4 | 8.2 | 2.3 | 13.3 | 3.7 | 3.1 | - | 5.6 | 2.6 |

- ^a Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130)
^b Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.
^c Incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky.
The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.
The SEER 11 areas comprise the SEER 9 areas plus San Jose-Monterey and Los Angeles.
The SEER 13 areas comprise the SEER 11 areas plus the Alaska Native Registry and Rural Georgia.
The SEER 17 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VIII-12
ESOPHAGUS CANCER (Invasive)

AGE-ADJUSTED SEER DEATH RATES^a
By Registry, Race/Ethnicity and Sex

| | All Races | | | Whites | | Blacks | | Asian/Pacific Islanders | | Hispanics ^b | |
|--|-----------|-------|---------|--------|---------|--------|---------|-------------------------|---------|------------------------|---------|
| | Total | Males | Females | Males | Females | Males | Females | Males | Females | Males | Females |
| <u>SEER DEATH RATES^a, 2000-2003</u> | | | | | | | | | | | |
| REGISTRY | | | | | | | | | | | |
| 17 SEER Areas ^c | 4.1 | 7.3 | 1.7 | 7.5 | 1.7 | 9.2 | 2.6 | 3.3 | 0.9 | 4.2 | 0.9 |
| 13 SEER Areas ^c | 4.0 | 7.0 | 1.7 | 7.3 | 1.6 | 8.6 | 2.8 | 3.6 | 0.9 | 4.5 | 0.8 |
| 11 SEER Areas ^c | 4.0 | 7.0 | 1.6 | 7.3 | 1.6 | 8.4 | 2.8 | 3.6 | 0.9 | 4.5 | 0.8 |
| 9 SEER Areas ^c | 4.3 | 7.5 | 1.7 | 7.7 | 1.7 | 9.2 | 2.8 | 4.0 | 0.9 | 5.0 | 0.8 |
| San Francisco-Oakland | 3.8 | 6.3 | 1.8 | 6.9 | 1.8 | 7.0 | 3.5 | 3.6 | 1.1 | 3.6 | - |
| Connecticut | 4.6 | 8.1 | 1.9 | 7.9 | 1.7 | 10.4 | 3.9 | - | - | 9.3 | - |
| Metropolitan Detroit | 5.0 | 9.3 | 1.8 | 9.5 | 1.7 | 9.3 | 2.5 | - | - | - | - |
| Hawaii | 2.8 | 5.2 | 0.8 | 6.1 | - | - | - | 4.8 | - | - | - |
| Iowa | 4.6 | 8.2 | 1.8 | 8.2 | 1.8 | - | - | - | - | - | - |
| New Mexico | 3.8 | 6.9 | 1.2 | 7.1 | 1.1 | - | - | - | - | 5.8 | - |
| Seattle-Puget Sound | 4.8 | 8.4 | 2.0 | 8.7 | 2.0 | - | - | - | - | - | - |
| Utah | 2.8 | 5.2 | 0.8 | 5.3 | 0.8 | - | - | - | - | - | - |
| Metropolitan Atlanta | 4.0 | 7.1 | 1.8 | 5.8 | 1.5 | 11.5 | 2.5 | - | - | - | - |
| San Jose-Monterey | 3.3 | 5.3 | 1.6 | 6.1 | 1.8 | - | - | - | - | 3.0 | - |
| Los Angeles | 3.3 | 5.7 | 1.5 | 6.0 | 1.4 | 6.7 | 2.6 | 3.3 | 1.1 | 4.3 | 0.8 |
| Rural Georgia | 7.7 | 11.5 | - | - | - | 21.2 | - | - | - | - | - |
| California exclud. SF/SJM/LA | 4.1 | 7.1 | 1.7 | 7.4 | 1.8 | 7.7 | 2.0 | 2.6 | 0.8 | 3.8 | 1.0 |
| Kentucky | 4.4 | 8.4 | 1.3 | 8.2 | 1.3 | 13.4 | - | - | - | - | - |
| Louisiana | 4.5 | 8.1 | 1.7 | 7.7 | 1.5 | 9.8 | 2.2 | - | - | - | - |
| New Jersey | 4.4 | 7.7 | 1.9 | 7.6 | 1.8 | 10.8 | 2.6 | - | - | 4.2 | - |
| Total U.S. | 4.4 | 7.8 | 1.8 | 7.7 | 1.7 | 10.5 | 3.0 | 3.0 | 0.8 | 4.2 | 1.0 |

^a NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. The 2000-2003 Hispanic death rates do not include deaths from Minnesota, New Hampshire and North Dakota.

^c The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 11 areas comprise the SEER 9 areas plus San Jose-Monterey and Los Angeles.

The SEER 13 areas comprise the SEER 11 areas plus the Alaska Native Registry and Rural Georgia.

The SEER 17 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VIII-13
ESOPHAGUS CANCER (INVASIVE)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2000-2003

Males and Females

| <u>State</u> | <u>Rate</u> | <u>SE</u> | <u>Rank</u> | <u>PD</u> | <u>State</u> | <u>Rate</u> | <u>SE</u> | <u>Rank</u> | <u>PD</u> |
|----------------------|-------------------|-----------|-------------|-----------|----------------|-------------------|-----------|-------------|-----------|
| TOTAL U.S. | 4.4 | 0.02 | | | | | | | |
| High Five States | | | | | | | | | |
| District of Columbia | 7.1 ^{bc} | 0.56 | (01) | 61.4 | | | | | |
| New Hampshire | 5.8 ^{bc} | 0.34 | (02) | 31.8 | | | | | |
| Maine | 5.5 ^{bc} | 0.31 | (03) | 25.0 | | | | | |
| Wyoming | 5.4 ^b | 0.53 | (04) | 22.7 | | | | | |
| Alaska | 5.3 ^b | 0.64 | (05) | 20.5 | | | | | |
| Low Five States | | | | | | | | | |
| New Mexico | 3.8 | 0.23 | (47) | -13.6 | | | | | |
| Kansas | 3.7 ^{bc} | 0.18 | (48) | -15.9 | | | | | |
| Arkansas | 3.6 ^{bc} | 0.18 | (49) | -18.2 | | | | | |
| Hawaii | 2.8 ^{bc} | 0.23 | (50) | -36.4 | | | | | |
| Utah | 2.8 ^{bc} | 0.21 | (51) | -36.4 | | | | | |
| State | Rate | SE | Rank | PD | State | Rate | SE | Rank | PD |
| Alabama | 4.1 | 0.15 | (40) | -6.8 | Montana | 4.9 | 0.35 | (13) | 11.4 |
| Alaska | 5.3 ^b | 0.64 | (05) | 20.5 | Nebraska | 4.6 | 0.25 | (24) | 4.5 |
| Arizona | 4.1 | 0.14 | (38) | -6.8 | Nevada | 4.7 | 0.25 | (21) | 6.8 |
| Arkansas | 3.6 ^{bc} | 0.18 | (49) | -18.2 | New Hampshire | 5.8 ^{bc} | 0.34 | (02) | 31.8 |
| California | 3.8 ^c | 0.06 | (46) | -13.6 | New Jersey | 4.4 | 0.11 | (31) | 0.0 |
| Colorado | 4.0 | 0.17 | (43) | -9.1 | New Mexico | 3.8 | 0.23 | (47) | -13.6 |
| Connecticut | 4.6 | 0.18 | (23) | 4.5 | New York | 4.4 | 0.07 | (32) | 0.0 |
| Delaware | 4.7 | 0.38 | (19) | 6.8 | North Carolina | 4.1 | 0.11 | (39) | -6.8 |
| District of Columbia | 7.1 ^{bc} | 0.56 | (01) | 61.4 | North Dakota | 4.0 | 0.38 | (42) | -9.1 |
| Florida | 4.2 | 0.07 | (37) | -4.5 | Ohio | 5.1 ^{bc} | 0.10 | (08) | 15.9 |
| Georgia | 4.5 | 0.13 | (26) | 2.3 | Oklahoma | 3.9 | 0.16 | (44) | -11.4 |
| Hawaii | 2.8 ^{bc} | 0.23 | (50) | -36.4 | Oregon | 5.0 | 0.19 | (11) | 13.6 |
| Idaho | 4.5 | 0.30 | (27) | 2.3 | Pennsylvania | 4.9 ^c | 0.09 | (15) | 11.4 |
| Illinois | 4.7 | 0.10 | (20) | 6.8 | Rhode Island | 4.9 | 0.33 | (16) | 11.4 |
| Indiana | 4.8 | 0.14 | (18) | 9.1 | South Carolina | 5.1 ^{bc} | 0.18 | (07) | 15.9 |
| Iowa | 4.6 | 0.19 | (22) | 4.5 | South Dakota | 4.2 | 0.36 | (36) | -4.5 |
| Kansas | 3.7 ^{bc} | 0.18 | (48) | -15.9 | Tennessee | 4.0 | 0.13 | (41) | -9.1 |
| Kentucky | 4.4 | 0.16 | (30) | 0.0 | Texas | 3.8 ^c | 0.07 | (45) | -13.6 |
| Louisiana | 4.5 | 0.16 | (25) | 2.3 | Utah | 2.8 ^{bc} | 0.21 | (51) | -36.4 |
| Maine | 5.5 ^{bc} | 0.31 | (03) | 25.0 | Vermont | 5.0 | 0.44 | (10) | 13.6 |
| Maryland | 4.9 | 0.16 | (14) | 11.4 | Virginia | 4.3 | 0.13 | (33) | -2.3 |
| Massachusetts | 5.3 ^{bc} | 0.14 | (06) | 20.5 | Washington | 4.8 | 0.15 | (17) | 9.1 |
| Michigan | 5.0 ^c | 0.11 | (12) | 13.6 | West Virginia | 4.5 | 0.23 | (29) | 2.3 |
| Minnesota | 4.5 | 0.15 | (28) | 2.3 | Wisconsin | 5.0 ^c | 0.15 | (09) | 13.6 |
| Mississippi | 4.3 | 0.20 | (34) | -2.3 | Wyoming | 5.4 ^b | 0.53 | (04) | 22.7 |
| Missouri | 4.3 | 0.13 | (35) | -2.3 | | | | | |

^a NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Absolute percent difference between state rate and total U.S. rate is 15% or more.

^c Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

Table VIII-14
ESOPHAGUS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2000-2003

Males

| <u>State</u> | <u>Rate</u> | <u>SE</u> | <u>Rank</u> | <u>PD</u> | <u>State</u> | <u>Rate</u> | <u>SE</u> | <u>Rank</u> | <u>PD</u> |
|----------------------|--------------------|-----------|-------------|-----------|----------------|--------------------|-----------|-------------|-----------|
| TOTAL U.S. | 7.8 | 0.04 | | | | | | | |
| High Five States | | | | | | | | | |
| District of Columbia | 12.6 ^{bc} | 1.17 | (01) | 61.5 | | | | | |
| New Hampshire | 10.5 ^{bc} | 0.70 | (02) | 34.6 | | | | | |
| Massachusetts | 9.9 ^{bc} | 0.30 | (03) | 26.9 | | | | | |
| Maine | 9.9 ^b | 0.62 | (04) | 26.9 | | | | | |
| Wyoming | 9.3 ^b | 1.08 | (05) | 19.2 | | | | | |
| Low Five States | | | | | | | | | |
| Texas | 6.7 ^c | 0.15 | (47) | -14.1 | | | | | |
| California | 6.5 ^{bc} | 0.11 | (48) | -16.7 | | | | | |
| Arkansas | 6.1 ^{bc} | 0.35 | (49) | -21.8 | | | | | |
| Utah | 5.2 ^{bc} | 0.42 | (50) | -33.3 | | | | | |
| Hawaii | 5.2 ^{bc} | 0.47 | (51) | -33.3 | | | | | |
| State | Rate | SE | Rank | PD | State | Rate | SE | Rank | PD |
| Alabama | 7.2 | 0.30 | (41) | -7.7 | Montana | 8.5 | 0.70 | (16) | 9.0 |
| Alaska | 8.9 | 1.26 | (11) | 14.1 | Nebraska | 8.2 | 0.51 | (22) | 5.1 |
| Arizona | 7.3 | 0.28 | (40) | -6.4 | Nevada | 8.1 | 0.48 | (27) | 3.8 |
| Arkansas | 6.1 ^{bc} | 0.35 | (49) | -21.8 | New Hampshire | 10.5 ^{bc} | 0.70 | (02) | 34.6 |
| California | 6.5 ^{bc} | 0.11 | (48) | -16.7 | New Jersey | 7.7 | 0.23 | (31) | -1.3 |
| Colorado | 7.3 | 0.35 | (39) | -6.4 | New Mexico | 6.9 | 0.48 | (45) | -11.5 |
| Connecticut | 8.1 | 0.36 | (26) | 3.8 | New York | 7.5 | 0.15 | (34) | -3.8 |
| Delaware | 8.7 | 0.79 | (13) | 11.5 | North Carolina | 7.1 | 0.23 | (44) | -9.0 |
| District of Columbia | 12.6 ^{bc} | 1.17 | (01) | 61.5 | North Dakota | 7.3 | 0.76 | (37) | -6.4 |
| Florida | 7.2 ^c | 0.14 | (43) | -7.7 | Ohio | 9.2 ^{bc} | 0.21 | (06) | 17.9 |
| Georgia | 7.9 | 0.26 | (29) | 1.3 | Oklahoma | 7.2 | 0.34 | (42) | -7.7 |
| Hawaii | 5.2 ^{bc} | 0.47 | (51) | -33.3 | Oregon | 8.9 | 0.37 | (10) | 14.1 |
| Idaho | 7.6 | 0.58 | (33) | -2.6 | Pennsylvania | 8.7 ^c | 0.19 | (12) | 11.5 |
| Illinois | 8.4 | 0.20 | (18) | 7.7 | Rhode Island | 8.2 | 0.64 | (23) | 5.1 |
| Indiana | 8.7 | 0.29 | (14) | 11.5 | South Carolina | 9.1 ^{bc} | 0.36 | (07) | 16.7 |
| Iowa | 8.2 | 0.38 | (24) | 5.1 | South Dakota | 7.5 | 0.72 | (35) | -3.8 |
| Kansas | 6.8 | 0.38 | (46) | -12.8 | Tennessee | 7.3 | 0.27 | (38) | -6.4 |
| Kentucky | 8.4 | 0.35 | (19) | 7.7 | Texas | 6.7 ^c | 0.15 | (47) | -14.1 |
| Louisiana | 8.1 | 0.34 | (25) | 3.8 | Utah | 5.2 ^{bc} | 0.42 | (50) | -33.3 |
| Maine | 9.9 ^b | 0.62 | (04) | 26.9 | Vermont | 8.4 | 0.87 | (17) | 7.7 |
| Maryland | 8.5 | 0.32 | (15) | 9.0 | Virginia | 7.5 | 0.26 | (36) | -3.8 |
| Massachusetts | 9.9 ^{bc} | 0.30 | (03) | 26.9 | Washington | 8.3 | 0.29 | (20) | 6.4 |
| Michigan | 9.0 ^{bc} | 0.23 | (08) | 15.4 | West Virginia | 7.9 | 0.47 | (28) | 1.3 |
| Minnesota | 7.8 | 0.30 | (30) | 0.0 | Wisconsin | 9.0 ^{bc} | 0.31 | (09) | 15.4 |
| Mississippi | 8.3 | 0.42 | (21) | 6.4 | Wyoming | 9.3 ^b | 1.08 | (05) | 19.2 |
| Missouri | 7.6 | 0.27 | (32) | -2.6 | | | | | |

^a NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Absolute percent difference between state rate and total U.S. rate is 15% or more.

^c Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

Table VIII-15
ESOPHAGUS CANCER (INVASIVE)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2000-2003

Females

| <u>State</u> | <u>Rate</u> | <u>SE</u> | <u>Rank</u> | <u>PD</u> | <u>State</u> | <u>Rate</u> | <u>SE</u> | <u>Rank</u> | <u>PD</u> |
|----------------------|-------------------|-----------|-------------|-----------|----------------|-------------------|-----------|-------------|-----------|
| TOTAL U.S. | 1.8 | 0.02 | | | Montana | 2.0 | 0.30 | (07) | 11.1 |
| High Five States | | | | | Nebraska | 1.8 | 0.21 | (27) | 0.0 |
| District of Columbia | 3.0 ^b | 0.49 | (01) | 66.7 | Nevada | 1.7 | 0.21 | (32) | -5.6 |
| Wyoming | 2.5 ^b | 0.48 | (02) | 38.9 | New Hampshire | 2.0 | 0.26 | (14) | 11.1 |
| Vermont | 2.4 ^b | 0.41 | (03) | 33.3 | New Jersey | 1.9 | 0.09 | (21) | 5.6 |
| Rhode Island | 2.2 ^b | 0.28 | (04) | 22.2 | New Mexico | 1.2 ^{bc} | 0.17 | (49) | -33.3 |
| Maryland | 2.2 ^b | 0.14 | (05) | 22.2 | New York | 2.0 | 0.07 | (10) | 11.1 |
| Low Five States | | | | | North Carolina | 1.8 | 0.10 | (29) | 0.0 |
| Mississippi | 1.2 ^{bc} | 0.14 | (47) | -33.3 | North Dakota | 1.4 ^b | 0.29 | (44) | -22.2 |
| Oklahoma | 1.2 ^{bc} | 0.12 | (48) | -33.3 | Ohio | 2.0 | 0.09 | (08) | 11.1 |
| New Mexico | 1.2 ^{bc} | 0.17 | (49) | -33.3 | Oklahoma | 1.2 ^{bc} | 0.12 | (48) | -33.3 |
| Hawaii | 0.8 ^{bc} | 0.17 | (50) | -55.6 | Oregon | 1.9 | 0.15 | (20) | 5.6 |
| Utah | 0.8 ^{bc} | 0.15 | (51) | -55.6 | Pennsylvania | 2.0 | 0.08 | (15) | 11.1 |
| Alabama | 1.6 | 0.12 | (40) | -11.1 | Rhode Island | 2.2 ^b | 0.28 | (04) | 22.2 |
| Alaska | 2.1 ^b | 0.56 | (06) | 16.7 | South Carolina | 2.0 | 0.15 | (09) | 11.1 |
| Arizona | 1.6 | 0.12 | (38) | -11.1 | South Dakota | 1.4 ^b | 0.28 | (43) | -22.2 |
| Arkansas | 1.6 | 0.16 | (39) | -11.1 | Tennessee | 1.5 ^b | 0.11 | (41) | -16.7 |
| California | 1.7 | 0.05 | (35) | -5.6 | Texas | 1.6 | 0.06 | (37) | -11.1 |
| Colorado | 1.5 ^b | 0.14 | (42) | -16.7 | Utah | 0.8 ^{bc} | 0.15 | (51) | -55.6 |
| Connecticut | 1.9 | 0.15 | (18) | 5.6 | Vermont | 2.4 ^b | 0.41 | (03) | 33.3 |
| Delaware | 1.7 | 0.30 | (34) | -5.6 | Virginia | 1.8 | 0.11 | (23) | 0.0 |
| District of Columbia | 3.0 ^b | 0.49 | (01) | 66.7 | Washington | 2.0 | 0.13 | (11) | 11.1 |
| Florida | 1.7 | 0.06 | (31) | -5.6 | West Virginia | 1.7 | 0.19 | (30) | -5.6 |
| Georgia | 1.9 | 0.11 | (16) | 5.6 | Wisconsin | 1.9 | 0.12 | (17) | 5.6 |
| Hawaii | 0.8 ^{bc} | 0.17 | (50) | -55.6 | Wyoming | 2.5 ^b | 0.48 | (02) | 38.9 |
| Idaho | 1.8 | 0.26 | (25) | 0.0 | | | | | |
| Illinois | 2.0 | 0.08 | (13) | 11.1 | | | | | |
| Indiana | 1.8 | 0.11 | (28) | 0.0 | | | | | |
| Iowa | 1.8 | 0.15 | (26) | 0.0 | | | | | |
| Kansas | 1.4 ^b | 0.15 | (45) | -22.2 | | | | | |
| Kentucky | 1.3 ^{bc} | 0.12 | (46) | -27.8 | | | | | |
| Louisiana | 1.7 | 0.13 | (33) | -5.6 | | | | | |
| Maine | 2.0 | 0.24 | (12) | 11.1 | | | | | |
| Maryland | 2.2 ^b | 0.14 | (05) | 22.2 | | | | | |
| Massachusetts | 1.9 | 0.11 | (19) | 5.6 | | | | | |
| Michigan | 1.8 | 0.09 | (22) | 0.0 | | | | | |
| Minnesota | 1.8 | 0.13 | (24) | 0.0 | | | | | |
| Mississippi | 1.2 ^{bc} | 0.14 | (47) | -33.3 | | | | | |
| Missouri | 1.6 | 0.11 | (36) | -11.1 | | | | | |

^a NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Absolute percent difference between state rate and total U.S. rate is 15% or more.

^c Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

Table VIII-16
ESOPHAGUS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2003
By Race/Ethnicity, Sex and Years Since Diagnosis

| Years Since Diagnosis | 0 to <5 | 5 to <10 | 10 to <15 | 15 to <20 | 20 to <25 | 0 to <13 ^e | 0 to <28 ^e | >=28 ^g | Complete ^h |
|-------------------------------------|---------|----------|-----------|-----------|-----------|-----------------------|-----------------------|-------------------|-----------------------|
| <u>Race/Sex</u> | | | | | | | | | |
| All Races ^b | 16,590 | 4,363 | 1,804 | 674 | 271 | 22,336 | 23,825 | 240 | 24,065 |
| Males | 12,747 | 3,273 | 1,236 | 425 | 210 | 16,955 | 17,985 | 119 | 18,104 |
| Females | 3,843 | 1,090 | 568 | 249 | 61 | 5,381 | 5,840 | 121 | 5,961 |
| White ^b | 14,345 | 3,867 | 1,427 | 594 | 222 | 19,327 | 20,552 | 199 | 20,751 |
| Males | 11,238 | 2,932 | 1,005 | 373 | 188 | 14,922 | 15,818 | 112 | 15,930 |
| Females | 3,107 | 935 | 422 | 221 | 34 | 4,405 | 4,734 | 87 | 4,821 |
| Black ^b | 1,751 | 438 | 314 | 77 | 40 | 2,417 | 2,648 | 63 | 2,711 |
| Males | 1,162 | 290 | 187 | 52 | 14 | 1,604 | 1,719 | 19 | 1,738 |
| Females | 589 | 148 | 127 | 25 | 26 | 813 | 929 | 44 | 973 |
| Asian/Pacific Islander ^c | 276 | 66 | + | + | + | 374 | + | + | + |
| Males | 199 | 51 | + | + | + | 275 | + | + | + |
| Females | 77 | 15 | + | + | + | 99 | + | + | + |
| Hispanic ^d | 566 | 153 | + | + | + | 776 | + | + | + |
| Males | 442 | 114 | + | + | + | 608 | + | + | + |
| Females | 124 | 39 | + | + | + | 168 | + | + | + |

Estimated prevalence percent^a on January 1, 2003, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

| Age at Prevalence | Age Specific (Crude) | | | | | | | | Age-Adjusted ^f All Ages | |
|-------------------------------------|----------------------|-----|-------|---------|---------|---------|---------|---------|---------------------------------------|---------|
| | All Ages | 0-9 | 10-19 | 20-29 | 30-39 | 40-49 | 50-59 | 60-69 | 70-79 | |
| <u>Race/Sex</u> | | | | | | | | | | |
| All Races ^c | 0.0060% | - | - | 0.0001% | 0.0004% | 0.0025% | 0.0110% | 0.0250% | 0.0347% | 0.0267% |
| Males | 0.0092% | - | - | 0.0002% | 0.0007% | 0.0040% | 0.0187% | 0.0417% | 0.0596% | 0.0460% |
| Females | 0.0029% | - | - | - | 0.0011% | 0.0037% | 0.0101% | 0.0161% | 0.0161% | 0.0028% |
| White ^c | 0.0066% | - | - | 0.0001% | 0.0004% | 0.0026% | 0.0116% | 0.0260% | 0.0365% | 0.0281% |
| Males | 0.0101% | - | - | - | 0.0007% | 0.0041% | 0.0202% | 0.0436% | 0.0631% | 0.0479% |
| Females | 0.0030% | - | - | - | - | 0.0010% | 0.0031% | 0.0100% | 0.0161% | 0.0175% |
| Black ^c | 0.0053% | - | - | - | - | 0.0037% | 0.0134% | 0.0321% | 0.0363% | 0.0217% |
| Males | 0.0074% | - | - | - | - | 0.0054% | 0.0198% | 0.0484% | 0.0564% | 0.0411% |
| Females | 0.0035% | - | - | - | - | 0.0022% | 0.0081% | 0.0192% | 0.0231% | 0.0124% |
| Asian/Pacific Islander ^c | 0.0032% | - | - | - | - | 0.0010% | 0.0054% | 0.0123% | 0.0211% | 0.0184% |
| Males | 0.0048% | - | - | - | - | 0.0015% | 0.0074% | 0.0221% | 0.0362% | 0.0367% |
| Females | 0.0016% | - | - | - | - | - | 0.0035% | 0.0042% | 0.0102% | 0.0017% |
| Hispanic ^d | 0.0018% | - | - | - | - | 0.0013% | 0.0056% | 0.0129% | 0.0207% | 0.0219% |
| Males | 0.0027% | - | - | - | - | 0.0020% | 0.0101% | 0.0225% | 0.0379% | 0.0317% |
| Females | 0.0009% | - | - | - | - | - | - | 0.0049% | 0.0084% | 0.0164% |

^a US 2003 cancer prevalence counts are based on 2003 cancer prevalence proportions from the SEER registries and 1/1/2003 US population estimates based on the average of 2002 and 2003 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia excluding Hawaii and Seattle.

^e Maximum limited-duration prevalence: 28 years for 1975-2003 SEER 9 data; 13 years for 1990-2003 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^{f g h} Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^(g) Cases diagnosed more than 28 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). ^(h)Complete prevalence is obtained by summing 0 to <28 and >=28.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

SEER Incidence, Delay Adjusted Incidence and US Death Rates⁺ Esophagus Cancer, by Race and Sex




Figure VIII-1

+ Source: SEER 9 areas and NCHS public use data file for the total US. Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines and APCs are calculated using the Joinpoint Regression Program Version 3.1, April 2006, National Cancer Institute.


The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

SEER Incidence and US Death Rates*

Esophagus Cancer, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2003
and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2003



Source: Incidence data for Whites and Blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from NCHS public use data file for the total US.

* Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 3.1, April 2006, National Cancer Institute.

API = Asian/Pacific Islander.

~ AI/AN = American Indian/Alaska Native. Incidence data for AI/AN include cases from Connecticut, Detroit, Iowa, New Mexico, Seattle, Utah, Atlanta and the Alaska Native Registry for the time period 1992-2002. Mortality data are from the entire US for the time period 1992-2003.

^ Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, and the Alaska Native Registry. Mortality data for Hispanics exclude cases from Connecticut, Maine, Maryland, Minnesota, New Hampshire, New York, North Dakota, Oklahoma, and Vermont.

Figure VIII-2