



Trends in SEER Incidence & US Death Rates by Primary Cancer Site 1994-2003

Trends in SEER Incidence Rates



Trends in US Cancer Death Rates



Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia) and NCHS public use file for the total US. For sex-specific cancer sites, the population was limited to the population of the appropriate sex. Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

The APC is the Annual Percent Change over the time interval.


* The APC is significantly different from zero ($p<.05$).

[^] Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.


Figure I-4

Trends in SEER Incidence Rates by Primary Cancer Site 1994-2003

Ages Less Than 65



Ages 65 and Over



Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). For sex-specific cancer sites, the population was limited to the population of the appropriate sex.

Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

The APC is the Annual Percent Change over the time interval.

* The APC is significantly different from zero ($p<.05$).

[^] Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Figure I-5

Trends in US Death Rates by Primary Cancer Site 1994-2003





Figure I-6

Source: NCHS public use data file for the total US. For sex-specific cancer sites, the population was limited to the population of the appropriate sex. Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

The APC is the Annual Percent Change over the time interval.

* The APC is significantly different from zero ($p<.05$).

Trends in SEER Incidence Rates by Primary Cancer Site 1994-2003



Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).

Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

The APC is the Annual Percent Change over the time interval.

* The APC is significantly different from zero ($p<.05$).

^ Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Figure I-7

Trends in US Death Rates by Primary Cancer Site 1994-2003




Figure I-8

Source: NCHS public use data file for the total US.


Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

The APC is the Annual Percent Change over the time interval.

* The APC is significantly different from zero ($p<.05$).

Figure I-10

Trends in US Death Rates, 1994-2003 by Primary Cancer Site All Ages, by Race




Source: NCHS public use data file for the total US.

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

US Death Rates 1975-2003



Source: NCHS public use data file for the total US. Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).