

Table I-14

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site, Race/Ethnicity and Sex

17 SEER Areas, 2001-2003

Site	All Races		Whites		Blacks		Asian/Pacific Islanders		Hispanics ^a	
	Males	Females	Males	Females	Males	Females	Males	Females	Males	Females
All Sites	45.31	37.86	45.28	38.88	42.67	32.25	39.99	32.65	40.95	33.30
Invasive and In Situ	47.21	41.69	47.28	42.84	43.36	34.84	40.92	35.65	42.01	35.80
Oral Cavity and Pharynx	1.39	0.66	1.41	0.66	1.26	0.53	1.11	0.66	0.96	0.47
Esophagus	0.76	0.25	0.78	0.24	0.82	0.30	0.51	0.16	0.62	0.22
Stomach	1.12	0.68	1.01	0.57	1.31	0.99	2.44	1.72	1.80	1.36
Colon and Rectum	5.79	5.37	5.79	5.34	5.17	5.41	6.16	5.50	5.03	4.48
Invasive and In Situ	6.14	5.64	6.14	5.61	5.51	5.73	6.50	5.74	5.30	4.71
Liver and Intrahepatic Bile Duct	0.88	0.40	0.74	0.34	0.92	0.36	2.46	1.19	1.46	0.76
Pancreas	1.27	1.27	1.29	1.25	1.18	1.46	1.28	1.31	1.18	1.54
Larynx	0.63	0.15	0.62	0.15	0.86	0.18	0.43	0.04	0.60	0.09
Invasive and In Situ	0.68	0.16	0.67	0.17	0.90	0.19	0.43	0.04	0.62	0.09
Lung and Bronchus	8.02	6.15	8.09	6.47	8.26	5.21	7.07	4.18	5.21	3.36
Melanoma of the Skin	2.04	1.38	2.33	1.60	0.08	0.09	0.19	0.14	0.45	0.49
Invasive and In Situ	3.26	2.25	3.69	2.59	0.09	0.11	0.25	0.20	0.66	0.71
Breast	0.12	12.67	0.12	13.25	0.14	9.98	0.08	9.34	0.06	9.23
Invasive and In Situ	0.13	15.16	0.13	15.80	0.16	11.91	0.08	11.69	0.07	10.82
Cervix Uteri	-	0.73	-	0.69	-	0.94	-	0.82	-	1.22
Corpus and Uterus, NOS	-	2.49	-	2.61	-	1.93	-	1.72	-	1.89
Invasive and In Situ	-	2.52	-	2.64	-	1.96	-	1.74	-	1.91
Ovary ^b	-	1.44	-	1.53	-	0.95	-	1.12	-	1.35
Prostate	17.12	-	16.58	-	19.76	-	12.62	-	16.29	-
Testis	0.36	-	0.42	-	0.10	-	0.11	-	0.26	-
Urinary Bladder (Invasive and In Situ)	3.61	1.14	3.95	1.22	1.40	0.78	2.22	0.67	2.34	0.84
Kidney and Renal Pelvis	1.65	0.97	1.72	1.01	1.46	0.91	1.01	0.58	1.68	1.09
Brain and Other Nervous System	0.65	0.52	0.71	0.57	0.34	0.31	0.41	0.30	0.55	0.51
Thyroid	0.36	1.02	0.38	1.06	0.17	0.59	0.40	1.12	0.30	1.02
Hodgkin Lymphoma	0.24	0.20	0.26	0.21	0.19	0.15	0.13	0.10	0.27	0.15
Non-Hodgkin Lymphoma	2.14	1.83	2.24	1.92	1.25	1.07	1.91	1.60	1.96	1.77
Myeloma	0.67	0.53	0.63	0.49	1.06	0.89	0.45	0.43	0.77	0.63
Leukemia	1.49	1.05	1.56	1.10	0.94	0.76	1.01	0.72	1.19	0.93
Acute Lymphocytic Leukemia	0.13	0.11	0.14	0.11	0.06	0.05	0.12	0.13	0.18	0.16
Chronic Lymphocytic Leukemia	0.53	0.34	0.57	0.36	0.31	0.22	0.13	0.09	0.30	0.17
Acute Myeloid Leukemia	0.44	0.33	0.45	0.34	0.29	0.24	0.43	0.31	0.39	0.35
Chronic Myeloid Leukemia	0.18	0.13	0.18	0.13	0.14	0.11	0.15	0.09	0.15	0.11
Kaposi Sarcoma	0.10	0.02	0.08	0.01	0.16	0.01	0.06	0.00	0.14	0.04
Mesothelioma	0.20	0.05	0.23	0.06	0.08	0.02	0.09	0.02	0.20	0.07

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives.

Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky.

^b Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Note: Invasive cancer only unless specified otherwise.

Table I-15

Lifetime Risk (Percent) of Dying from Cancer by Site, Race/Ethnicity and Sex

Total U.S., 2001-2003

Site	All Races		Whites		Blacks		Asian/Pacific Islanders		Hispanics ^a	
	Males	Females	Males	Females	Males	Females	Males	Females	Males	Females
All Sites	23.42	19.82	23.50	20.04	23.89	19.46	20.68	16.42	19.96	15.67
Oral Cavity and Pharynx	0.39	0.19	0.37	0.19	0.50	0.17	0.45	0.24	0.32	0.14
Esophagus	0.75	0.22	0.75	0.21	0.79	0.30	0.42	0.13	0.48	0.16
Stomach	0.57	0.38	0.52	0.34	0.87	0.64	1.56	1.20	1.06	0.76
Colon and Rectum	2.33	2.18	2.32	2.15	2.43	2.50	2.37	2.16	2.14	1.82
Liver and Intrahepatic Bile Duct	0.67	0.38	0.63	0.36	0.72	0.41	1.97	1.13	1.18	0.76
Pancreas	1.18	1.18	1.19	1.17	1.14	1.33	1.08	1.19	1.03	1.18
Larynx	0.23	0.06	0.22	0.06	0.38	0.08	0.13	0.02	0.22	0.03
Lung and Bronchus	7.18	4.96	7.26	5.16	7.18	4.03	5.41	3.21	4.31	2.20
Melanoma of the Skin	0.36	0.20	0.41	0.22	0.03	0.04	0.07	0.06	0.11	0.10
Breast	0.03	2.92	0.03	2.91	0.04	3.27	0.01	1.69	0.02	2.06
Cervix Uteri	-	0.25	-	0.22	-	0.44	-	0.32	-	0.36
Corpus and Uterus, NOS	-	0.51	-	0.49	-	0.75	-	0.33	-	0.47
Ovary	-	1.05	-	1.11	-	0.76	-	0.69	-	0.81
Prostate	2.91	-	2.73	-	4.59	-	2.18	-	3.24	-
Testis	0.02	-	0.02	-	0.01	-	0.01	-	0.02	-
Urinary Bladder	0.76	0.32	0.81	0.32	0.39	0.33	0.52	0.25	0.59	0.26
Kidney and Renal Pelvis	0.59	0.34	0.61	0.35	0.45	0.29	0.35	0.20	0.60	0.35
Brain and Other Nervous System	0.48	0.38	0.52	0.42	0.24	0.20	0.31	0.17	0.35	0.29
Thyroid	0.04	0.06	0.04	0.06	0.03	0.05	0.05	0.12	0.06	0.09
Hodgkin Lymphoma	0.05	0.04	0.05	0.04	0.04	0.02	0.04	0.03	0.06	0.04
Non-Hodgkin Lymphoma	0.93	0.79	0.99	0.83	0.49	0.44	0.87	0.70	0.79	0.72
Myeloma	0.46	0.40	0.45	0.37	0.62	0.66	0.25	0.26	0.45	0.39
Leukemia	0.97	0.71	1.02	0.74	0.63	0.55	0.63	0.49	0.72	0.57
Acute Lymphocytic Leukemia	0.05	0.04	0.05	0.04	0.03	0.03	0.04	0.03	0.06	0.06
Chronic Lymphocytic Leukemia	0.23	0.15	0.24	0.16	0.15	0.12	0.06	0.03	0.11	0.06
Acute Myeloid Leukemia	0.34	0.25	0.36	0.27	0.20	0.18	0.26	0.24	0.25	0.19
Chronic Myeloid Leukemia	0.06	0.05	0.06	0.05	0.06	0.04	0.04	0.05	0.05	0.04

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table II-9
ALL SITES (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.17	0.35	0.71	45.31	0.15	0.31	0.78	37.86
	10	0.18	0.55	1.27	45.83	0.16	0.64	1.90	38.18
	20	0.37	1.09	3.09	45.97	0.48	1.74	4.79	38.17
	30	0.74	2.77	9.23	46.34	1.28	4.35	10.19	38.03
	40	2.08	8.69	22.54	46.67	3.14	9.09	18.29	37.49
	50	6.96	21.54	37.37	46.94	6.23	15.87	26.72	35.97
	60	16.58	34.60	43.67	45.49	10.57	22.48	30.60	32.63
White	0	0.18	0.38	0.77	45.28	0.16	0.33	0.83	38.88
	10	0.20	0.60	1.34	45.71	0.17	0.68	1.98	39.16
	20	0.40	1.15	3.14	45.83	0.51	1.81	4.92	39.14
	30	0.77	2.78	9.13	46.13	1.32	4.45	10.44	38.98
	40	2.06	8.55	22.42	46.39	3.19	9.30	18.86	38.40
	50	6.83	21.39	37.22	46.58	6.39	16.39	27.57	36.83
	60	16.50	34.42	43.29	45.02	10.96	23.22	31.41	33.37
Black	0	0.12	0.24	0.48	42.67	0.12	0.22	0.57	32.25
	10	0.13	0.37	1.01	43.75	0.11	0.47	1.56	32.87
	20	0.25	0.89	3.37	44.04	0.36	1.46	4.32	32.88
	30	0.66	3.21	11.94	44.98	1.11	4.01	9.51	32.89
	40	2.64	11.71	26.92	46.00	2.97	8.61	16.93	32.60
	50	9.87	26.42	40.28	47.18	6.01	14.86	24.04	31.53
	60	20.64	37.92	45.01	46.53	10.02	20.42	26.99	28.90
Asian/ Pacific Islander	0	0.15	0.28	0.51	39.99	0.12	0.25	0.57	32.65
	10	0.14	0.36	0.83	40.29	0.13	0.45	1.45	32.82
	20	0.23	0.69	1.96	40.33	0.32	1.32	3.88	32.79
	30	0.47	1.75	5.56	40.43	1.01	3.58	8.03	32.65
	40	1.29	5.14	14.48	40.42	2.61	7.12	13.52	32.05
	50	3.95	13.54	27.43	40.17	4.66	11.26	19.60	30.40
	60	10.27	25.16	35.74	38.81	7.02	15.89	23.85	27.38
Hispanic ^a	0	0.17	0.34	0.61	40.95	0.16	0.30	0.64	33.30
	10	0.17	0.45	0.96	41.34	0.14	0.49	1.49	33.51
	20	0.28	0.79	2.13	41.46	0.35	1.35	3.82	33.49
	30	0.52	1.87	6.32	41.74	1.01	3.50	7.97	33.35
	40	1.38	5.91	16.94	41.99	2.52	7.07	14.14	32.82
	50	4.72	16.21	30.84	42.29	4.71	12.04	21.01	31.39
	60	12.69	28.85	38.35	41.50	7.87	17.49	25.35	28.64

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	45.31%	37.86%	23.42%	19.82%
White	45.28%	38.88%	23.50%	20.04%
Black	42.67%	32.25%	23.89%	19.46%
Asian/Pacific Islander	39.99%	32.65%	20.68%	16.42%
Hispanic ^a	40.95%	33.30%	19.96%	15.67%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table III-9
BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.04	0.06	0.09	0.65	0.03	0.05	0.08	0.52
	10	0.03	0.05	0.09	0.63	0.02	0.04	0.08	0.49
	20	0.03	0.06	0.12	0.60	0.02	0.05	0.10	0.47
	30	0.04	0.10	0.20	0.59	0.03	0.07	0.14	0.45
	40	0.06	0.16	0.31	0.56	0.04	0.11	0.22	0.43
	50	0.10	0.25	0.42	0.51	0.07	0.18	0.31	0.39
	60	0.16	0.34	0.43	0.44	0.11	0.24	0.32	0.33
White	0	0.04	0.07	0.09	0.71	0.04	0.06	0.08	0.57
	10	0.03	0.06	0.10	0.69	0.02	0.05	0.08	0.54
	20	0.03	0.07	0.14	0.66	0.03	0.06	0.11	0.52
	30	0.04	0.11	0.22	0.64	0.04	0.08	0.16	0.50
	40	0.07	0.18	0.34	0.61	0.05	0.13	0.24	0.47
	50	0.11	0.28	0.46	0.56	0.08	0.20	0.34	0.43
	60	0.18	0.38	0.46	0.48	0.12	0.27	0.35	0.36
Black	0	0.03	0.05	0.06	0.34	0.03	0.04	0.05	0.31
	10	0.02	0.04	0.06	0.32	0.01	0.03	0.04	0.29
	20	0.02	0.04	0.07	0.30	0.01	0.03	0.06	0.28
	30	0.02	0.06	0.11	0.29	0.02	0.05	0.09	0.26
	40	0.04	0.10	0.17	0.28	0.03	0.07	0.13	0.25
	50	0.06	0.14	0.22	0.26	0.05	0.11	0.17	0.23
	60	0.09	0.19	0.23	0.23	0.07	0.14	0.19	0.20
Asian/ Pacific Islander	0	0.02	0.05	0.06	0.41	0.03	0.04	0.06	0.30
	10	0.02	0.04	0.06	0.39	0.02	0.03	0.05	0.28
	20	0.02	0.04	0.07	0.37	0.02	0.03	0.05	0.26
	30	0.02	0.06	0.11	0.35	0.02	0.04	0.07	0.24
	40	0.04	0.09	0.15	0.34	0.02	0.06	0.12	0.23
	50	0.06	0.12	0.21	0.31	0.04	0.10	0.15	0.21
	60	0.06	0.16	0.24	0.26	0.06	0.12	0.17	0.18
Hispanic ^a	0	0.03	0.05	0.07	0.55	0.04	0.05	0.07	0.51
	10	0.02	0.04	0.06	0.53	0.02	0.03	0.06	0.48
	20	0.02	0.04	0.09	0.51	0.02	0.05	0.08	0.47
	30	0.03	0.08	0.16	0.50	0.03	0.06	0.13	0.45
	40	0.05	0.13	0.25	0.48	0.03	0.10	0.20	0.42
	50	0.08	0.21	0.36	0.44	0.06	0.17	0.31	0.40
	60	0.14	0.30	0.37	0.39	0.11	0.25	0.32	0.34

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	0.65%	0.52%	0.48%	0.38%
White	0.71%	0.57%	0.52%	0.42%
Black	0.34%	0.31%	0.24%	0.20%
Asian/Pacific Islander	0.41%	0.30%	0.31%	0.17%
Hispanic ^a	0.55%	0.51%	0.35%	0.29%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table IV-14
BREAST CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.00	0.12	0.00	0.00	0.06	12.67
	10	0.00	0.00	0.00	0.12	0.00	0.06	0.48	12.81
	20	0.00	0.00	0.01	0.12	0.05	0.48	1.90	12.84
	30	0.00	0.01	0.03	0.12	0.43	1.85	4.36	12.86
	40	0.01	0.03	0.06	0.12	1.44	3.98	7.27	12.58
	50	0.02	0.05	0.09	0.12	2.63	6.03	9.26	11.52
	60	0.04	0.08	0.10	0.11	3.65	7.12	9.11	9.54
White	0	0.00	0.00	0.00	0.12	0.00	0.00	0.05	13.25
	10	0.00	0.00	0.00	0.12	0.00	0.06	0.48	13.38
	20	0.00	0.00	0.01	0.12	0.05	0.48	1.92	13.41
	30	0.00	0.01	0.03	0.12	0.43	1.88	4.48	13.42
	40	0.01	0.02	0.06	0.12	1.47	4.10	7.59	13.15
	50	0.02	0.05	0.09	0.12	2.72	6.33	9.74	12.06
	60	0.04	0.08	0.10	0.11	3.86	7.52	9.58	10.01
Black	0	0.00	0.00	0.00	0.14	0.00	0.00	0.07	9.98
	10	0.00	0.00	0.00	0.15	0.00	0.07	0.53	10.20
	20	0.00	0.00	0.01	0.15	0.07	0.53	1.87	10.23
	30	0.00	0.01	0.04	0.15	0.46	1.81	4.02	10.24
	40	0.01	0.04	0.08	0.16	1.37	3.63	6.20	9.98
	50	0.03	0.07	0.12	0.15	2.38	5.09	7.45	9.08
	60	0.05	0.10	0.13	0.14	3.01	5.64	7.07	7.45
Asian/ Pacific Islander	0	0.00	0.00	0.00	0.08	0.00	0.00	0.04	9.34
	10	0.00	0.00	0.00	0.08	0.00	0.04	0.44	9.41
	20	0.00	0.00	0.01	0.08	0.04	0.44	1.70	9.43
	30	0.00	0.00	0.02	0.08	0.40	1.66	3.62	9.42
	40	0.00	0.02	0.04	0.08	1.28	3.25	5.45	9.09
	50	0.02	0.03	0.05	0.07	2.02	4.27	6.32	7.99
	60	0.02	0.04	0.06	0.06	2.36	4.50	5.83	6.25
Hispanic ^a	0	0.00	0.00	0.00	0.06	0.00	0.00	0.05	9.23
	10	0.00	0.00	0.00	0.06	0.00	0.05	0.38	9.33
	20	0.00	0.00	0.00	0.06	0.05	0.38	1.47	9.35
	30	0.00	0.00	0.01	0.06	0.33	1.43	3.21	9.34
	40	0.00	0.01	0.03	0.06	1.11	2.91	5.12	9.09
	50	0.01	0.03	0.04	0.06	1.84	4.11	6.32	8.18
	60	0.02	0.03	0.05	0.06	2.38	4.71	6.12	6.67

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	0.12%	12.67%	0.03%	2.92%
White	0.12%	13.25%	0.03%	2.91%
Black	0.14%	9.98%	0.04%	3.27%
Asian/Pacific Islander	0.08%	9.34%	0.01%	1.69%
Hispanic ^a	0.06%	9.23%	0.02%	2.06%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table V-7
CERVIX UTERI CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/Ethnicity	Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.04	0.73
	10	0.00	0.04	0.16	0.73
	20	0.04	0.16	0.31	0.73
	30	0.12	0.27	0.41	0.70
	40	0.15	0.29	0.42	0.58
	50	0.14	0.28	0.37	0.44
	60	0.14	0.25	0.30	0.32
White	0	0.00	0.00	0.04	0.69
	10	0.00	0.04	0.17	0.70
	20	0.04	0.17	0.31	0.70
	30	0.13	0.27	0.40	0.66
	40	0.15	0.28	0.40	0.54
	50	0.13	0.25	0.34	0.40
	60	0.13	0.22	0.26	0.28
Black	0	0.00	0.00	0.04	0.94
	10	0.00	0.04	0.15	0.96
	20	0.04	0.15	0.31	0.96
	30	0.11	0.27	0.44	0.93
	40	0.17	0.34	0.55	0.84
	50	0.18	0.40	0.56	0.70
	60	0.24	0.41	0.53	0.57
Asian/Pacific Islander	0	0.00	0.00	0.02	0.82
	10	0.00	0.02	0.09	0.83
	20	0.02	0.09	0.23	0.83
	30	0.07	0.21	0.37	0.82
	40	0.14	0.30	0.48	0.75
	50	0.16	0.34	0.49	0.61
	60	0.19	0.33	0.43	0.46
Hispanic ^a	0	0.00	0.00	0.04	1.22
	10	0.00	0.04	0.19	1.23
	20	0.04	0.19	0.43	1.23
	30	0.15	0.40	0.64	1.20
	40	0.25	0.50	0.75	1.06
	50	0.25	0.51	0.70	0.82
	60	0.27	0.47	0.56	0.59

Race/Ethnicity	Lifetime Risk of Developing	Lifetime Risk of Dying
All Races	0.73%	0.25%
White	0.69%	0.22%
Black	0.94%	0.44%
Asian/Pacific Islander	0.82%	0.32%
Hispanic ^a	1.22%	0.36%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table VI-15
COLON AND RECTUM CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.02	5.79	0.00	0.00	0.01	5.37
	10	0.00	0.02	0.08	5.87	0.00	0.01	0.07	5.43
	20	0.01	0.07	0.29	5.90	0.01	0.07	0.26	5.44
	30	0.06	0.28	0.98	5.96	0.05	0.25	0.77	5.45
	40	0.22	0.93	2.41	6.01	0.20	0.73	1.81	5.45
	50	0.74	2.27	4.34	6.01	0.54	1.64	3.35	5.37
	60	1.67	3.91	5.39	5.73	1.16	2.96	4.61	5.09
White	0	0.00	0.00	0.02	5.79	0.00	0.00	0.01	5.34
	10	0.00	0.02	0.07	5.86	0.00	0.01	0.07	5.39
	20	0.01	0.07	0.28	5.89	0.01	0.07	0.25	5.41
	30	0.06	0.27	0.94	5.95	0.05	0.24	0.74	5.42
	40	0.21	0.90	2.37	5.99	0.19	0.69	1.75	5.41
	50	0.71	2.24	4.32	5.99	0.51	1.60	3.31	5.33
	60	1.65	3.90	5.39	5.71	1.14	2.94	4.60	5.06
Black	0	0.00	0.00	0.01	5.17	0.00	0.00	0.01	5.41
	10	0.00	0.01	0.08	5.31	0.00	0.01	0.07	5.53
	20	0.01	0.08	0.35	5.36	0.01	0.07	0.33	5.54
	30	0.07	0.35	1.23	5.49	0.06	0.32	1.05	5.58
	40	0.29	1.20	2.77	5.60	0.27	1.01	2.34	5.62
	50	0.98	2.66	4.44	5.69	0.78	2.17	3.89	5.59
	60	1.96	4.04	5.20	5.50	1.53	3.42	4.87	5.29
Asian/ Pacific Islander	0	0.00	0.00	0.02	6.16	0.00	0.00	0.01	5.50
	10	0.00	0.02	0.09	6.22	0.00	0.01	0.06	5.55
	20	0.02	0.08	0.28	6.24	0.01	0.06	0.26	5.56
	30	0.07	0.27	0.89	6.27	0.05	0.25	0.72	5.56
	40	0.20	0.83	2.15	6.25	0.19	0.67	1.53	5.54
	50	0.64	1.98	4.07	6.17	0.48	1.35	2.79	5.41
	60	1.41	3.59	5.31	5.80	0.90	2.39	4.10	5.09
Hispanic ^a	0	0.00	0.00	0.01	5.03	0.00	0.00	0.01	4.48
	10	0.00	0.01	0.06	5.09	0.00	0.01	0.06	4.53
	20	0.01	0.06	0.22	5.12	0.01	0.06	0.21	4.54
	30	0.05	0.21	0.76	5.17	0.05	0.20	0.60	4.54
	40	0.17	0.72	1.97	5.19	0.15	0.55	1.40	4.52
	50	0.57	1.86	3.63	5.19	0.41	1.27	2.54	4.45
	60	1.38	3.28	4.59	4.95	0.89	2.20	3.55	4.19

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	5.79%	5.37%	0.05%	0.10%
White	5.79%	5.34%	0.05%	0.10%
Black	5.17%	5.41%	0.04%	0.11%
Asian/Pacific Islander	6.16%	5.50%	0.11%	0.13%
Hispanic ^a	5.03%	4.48%	0.09%	0.19%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table VII-7
CORPUS AND UTERUS, NOS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/Ethnicity	Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.01	2.49
	10	0.00	0.01	0.06	2.52
	20	0.01	0.06	0.24	2.53
	30	0.05	0.24	0.75	2.53
	40	0.18	0.70	1.45	2.50
	50	0.53	1.30	1.95	2.37
	60	0.81	1.50	1.87	1.94
White	0	0.00	0.00	0.01	2.61
	10	0.00	0.01	0.06	2.64
	20	0.01	0.06	0.25	2.64
	30	0.05	0.24	0.78	2.65
	40	0.19	0.73	1.52	2.62
	50	0.56	1.36	2.05	2.48
	60	0.85	1.57	1.95	2.02
Black	0	0.00	0.00	0.01	1.93
	10	0.00	0.01	0.06	1.98
	20	0.01	0.06	0.17	1.98
	30	0.05	0.16	0.49	1.99
	40	0.11	0.45	1.13	1.98
	50	0.36	1.06	1.64	1.95
	60	0.77	1.40	1.69	1.74
Asian/Pacific Islander	0	0.00	0.00	0.01	1.72
	10	0.00	0.01	0.06	1.73
	20	0.01	0.06	0.28	1.73
	30	0.06	0.27	0.69	1.73
	40	0.22	0.64	1.08	1.68
	50	0.43	0.87	1.22	1.48
	60	0.46	0.82	1.04	1.09
Hispanic ^a	0	0.00	0.00	0.01	1.89
	10	0.00	0.01	0.06	1.91
	20	0.01	0.06	0.23	1.91
	30	0.05	0.22	0.60	1.91
	40	0.17	0.55	1.04	1.87
	50	0.39	0.88	1.34	1.73
	60	0.52	0.99	1.30	1.40

Race/Ethnicity	Lifetime Risk of Developing	Lifetime Risk of Dying
All Races	2.49%	0.51%
White	2.61%	0.49%
Black	1.93%	0.75%
Asian/Pacific Islander	1.72%	0.33%
Hispanic ^a	1.89%	0.47%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table VIII-9
ESOPHAGUS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.00	0.76	0.00	0.00	0.00	0.25
	10	0.00	0.00	0.01	0.77	0.00	0.00	0.00	0.25
	20	0.00	0.01	0.04	0.78	0.00	0.00	0.01	0.25
	30	0.00	0.04	0.15	0.79	0.00	0.01	0.03	0.25
	40	0.03	0.15	0.37	0.80	0.01	0.03	0.09	0.25
	50	0.12	0.35	0.62	0.79	0.02	0.08	0.17	0.25
	60	0.25	0.54	0.70	0.73	0.06	0.15	0.22	0.24
White	0	0.00	0.00	0.00	0.78	0.00	0.00	0.00	0.24
	10	0.00	0.00	0.01	0.78	0.00	0.00	0.00	0.25
	20	0.00	0.01	0.04	0.79	0.00	0.00	0.01	0.25
	30	0.00	0.04	0.15	0.80	0.00	0.01	0.03	0.25
	40	0.03	0.14	0.37	0.81	0.01	0.03	0.08	0.25
	50	0.12	0.35	0.63	0.80	0.02	0.08	0.16	0.25
	60	0.25	0.55	0.71	0.74	0.06	0.15	0.22	0.24
Black	0	0.00	0.00	0.00	0.82	0.00	0.00	0.00	0.30
	10	0.00	0.00	0.00	0.84	0.00	0.00	0.00	0.30
	20	0.00	0.00	0.05	0.85	0.00	0.00	0.02	0.30
	30	0.00	0.05	0.23	0.87	0.00	0.02	0.07	0.31
	40	0.05	0.24	0.54	0.89	0.02	0.07	0.16	0.31
	50	0.20	0.53	0.78	0.91	0.05	0.15	0.25	0.30
	60	0.38	0.68	0.80	0.82	0.10	0.22	0.27	0.27
Asian/ Pacific Islander	0	0.00	0.00	0.00	0.51	0.00	0.00	0.00	0.16
	10	0.00	0.00	0.00	0.52	0.00	0.00	0.00	0.16
	20	0.00	0.00	0.02	0.52	0.00	0.00	0.01	0.16
	30	0.00	0.02	0.07	0.52	0.00	0.01	0.02	0.16
	40	0.01	0.06	0.19	0.52	0.01	0.02	0.05	0.16
	50	0.05	0.18	0.33	0.52	0.01	0.04	0.09	0.16
	60	0.14	0.29	0.45	0.48	0.03	0.08	0.13	0.15
Hispanic ^a	0	0.00	0.00	0.00	0.62	0.00	0.00	0.00	0.22
	10	0.00	0.00	0.00	0.63	0.00	0.00	0.00	0.22
	20	0.00	0.00	0.02	0.63	0.00	0.00	0.00	0.22
	30	0.00	0.02	0.09	0.64	0.00	0.00	0.01	0.23
	40	0.02	0.08	0.23	0.65	0.00	0.01	0.04	0.23
	50	0.07	0.22	0.40	0.65	0.01	0.03	0.09	0.23
	60	0.16	0.35	0.52	0.62	0.03	0.09	0.14	0.23

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	0.76%	0.25%	0.75%	0.22%
White	0.78%	0.24%	0.75%	0.21%
Black	0.82%	0.30%	0.79%	0.30%
Asian/Pacific Islander	0.51%	0.16%	0.42%	0.13%
Hispanic ^a	0.62%	0.22%	0.48%	0.16%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table IX-9
HODGKIN LYMPHOMA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.02	0.06	0.24	0.00	0.02	0.06	0.20
	10	0.02	0.06	0.10	0.24	0.02	0.06	0.10	0.20
	20	0.04	0.08	0.11	0.22	0.04	0.08	0.10	0.18
	30	0.04	0.07	0.10	0.18	0.03	0.05	0.07	0.14
	40	0.03	0.06	0.10	0.15	0.02	0.04	0.06	0.10
	50	0.03	0.07	0.10	0.12	0.02	0.04	0.06	0.08
	60	0.04	0.08	0.09	0.10	0.02	0.05	0.07	0.07
White	0	0.00	0.03	0.07	0.26	0.00	0.02	0.07	0.21
	10	0.02	0.07	0.11	0.26	0.02	0.07	0.10	0.21
	20	0.04	0.08	0.12	0.23	0.05	0.08	0.11	0.19
	30	0.04	0.07	0.10	0.19	0.04	0.06	0.08	0.15
	40	0.03	0.06	0.10	0.15	0.02	0.04	0.07	0.11
	50	0.03	0.07	0.11	0.12	0.02	0.04	0.07	0.09
	60	0.05	0.08	0.10	0.10	0.03	0.05	0.07	0.07
Black	0	0.00	0.01	0.04	0.19	0.00	0.01	0.05	0.15
	10	0.01	0.04	0.07	0.19	0.01	0.05	0.08	0.15
	20	0.03	0.06	0.09	0.18	0.04	0.06	0.09	0.14
	30	0.03	0.06	0.10	0.15	0.03	0.05	0.07	0.11
	40	0.03	0.07	0.10	0.13	0.03	0.05	0.06	0.08
	50	0.04	0.07	0.09	0.10	0.02	0.04	0.05	0.05
	60	0.04	0.06	0.07	0.07	0.02	0.03	0.03	0.04
Asian/ Pacific Islander	0	0.00	0.01	0.03	0.13	0.00	0.01	0.03	0.10
	10	0.01	0.03	0.05	0.13	0.01	0.03	0.05	0.10
	20	0.02	0.03	0.04	0.11	0.02	0.04	0.04	0.09
	30	0.01	0.03	0.04	0.10	0.01	0.02	0.03	0.07
	40	0.01	0.03	0.05	0.08	0.01	0.01	0.02	0.05
	50	0.02	0.04	0.05	0.07	0.00	0.01	0.02	0.05
	60	0.02	0.04	0.06	0.06	0.01	0.02	0.04	0.04
Hispanic ^a	0	0.01	0.02	0.05	0.27	0.00	0.02	0.04	0.15
	10	0.02	0.04	0.06	0.27	0.01	0.04	0.05	0.15
	20	0.02	0.04	0.07	0.25	0.02	0.04	0.05	0.13
	30	0.02	0.05	0.08	0.23	0.02	0.03	0.04	0.11
	40	0.03	0.06	0.11	0.21	0.01	0.03	0.05	0.09
	50	0.03	0.09	0.15	0.19	0.02	0.04	0.06	0.08
	60	0.06	0.12	0.15	0.17	0.02	0.05	0.07	0.07

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	0.24%	0.20%	0.05%	0.04%
White	0.26%	0.21%	0.05%	0.04%
Black	0.19%	0.15%	0.04%	0.02%
Asian/Pacific Islander	0.13%	0.10%	0.04%	0.03%
Hispanic ^a	0.27%	0.15%	0.06%	0.04%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table X-10
KAPOSI SARCOMA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.01	0.10	0.00	0.00	0.00	0.02
	10	0.00	0.01	0.03	0.10	0.00	0.00	0.00	0.02
	20	0.01	0.03	0.05	0.10	0.00	0.00	0.00	0.02
	30	0.02	0.05	0.06	0.09	0.00	0.00	0.00	0.02
	40	0.02	0.03	0.04	0.07	0.00	0.00	0.00	0.01
	50	0.01	0.02	0.03	0.05	0.00	0.00	0.01	0.01
	60	0.01	0.02	0.03	0.04	0.00	0.01	0.01	0.01
White	0	0.00	0.00	0.00	0.08	0.00	0.00	0.00	0.01
	10	0.00	0.00	0.02	0.09	0.00	0.00	0.00	0.01
	20	0.00	0.02	0.04	0.09	0.00	0.00	0.00	0.01
	30	0.02	0.04	0.05	0.08	0.00	0.00	0.00	0.01
	40	0.02	0.03	0.04	0.06	0.00	0.00	0.00	0.01
	50	0.01	0.02	0.03	0.04	0.00	0.00	0.01	0.01
	60	0.01	0.02	0.03	0.04	0.00	0.01	0.01	0.02
Black	0	0.00	0.00	0.03	0.16	0.00	0.00	0.00	0.01
	10	0.00	0.03	0.09	0.17	0.00	0.00	0.00	0.01
	20	0.03	0.08	0.12	0.17	0.00	0.00	0.01	0.01
	30	0.06	0.10	0.12	0.15	0.00	0.00	0.01	0.01
	40	0.04	0.07	0.07	0.09	0.00	0.00	0.00	0.01
	50	0.02	0.03	0.04	0.05	0.00	0.00	0.00	0.01
	60	0.01	0.01	0.03	0.03	0.00	0.00	0.00	0.00
Asian/ Pacific Islander	0	0.00	0.00	0.00	0.06	0.00	0.00	0.00	0.00
	10	0.00	0.00	0.01	0.06	0.00	0.00	0.00	0.00
	20	0.00	0.01	0.02	0.06	0.00	0.00	0.00	0.00
	30	0.01	0.01	0.02	0.06	0.00	0.00	0.00	0.00
	40	0.01	0.01	0.02	0.05	0.00	0.00	0.00	0.00
	50	0.00	0.01	0.02	0.05	0.00	0.00	0.00	0.00
	60	0.01	0.01	0.03	0.04	0.00	0.00	0.00	0.00
Hispanic ^a	0	0.00	0.00	0.01	0.14	0.00	0.00	0.00	0.04
	10	0.00	0.01	0.03	0.15	0.00	0.00	0.00	0.04
	20	0.01	0.03	0.06	0.15	0.00	0.00	0.00	0.04
	30	0.03	0.06	0.07	0.14	0.00	0.00	0.00	0.04
	40	0.03	0.05	0.06	0.11	0.00	0.00	0.00	0.04
	50	0.02	0.04	0.06	0.09	0.00	0.00	0.01	0.04
	60	0.02	0.05	0.07	0.08	0.00	0.01	0.04	0.04

Race/Ethnicity	Lifetime Risk of Developing	
	Males	Females
All Races	0.10%	0.01%
White	0.08%	0.01%
Black	0.16%	0.02%
Asian/Pacific Islander	0.06%	0.00%
Hispanic ^a	0.14%	0.01%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky.

Table XI-9
KIDNEY AND RENAL PELVIS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.01	0.01	0.02	1.65	0.01	0.01	0.02	0.97
	10	0.00	0.01	0.04	1.66	0.00	0.01	0.03	0.97
	20	0.01	0.03	0.15	1.67	0.01	0.02	0.09	0.97
	30	0.03	0.14	0.42	1.69	0.02	0.08	0.23	0.97
	40	0.12	0.40	0.89	1.69	0.06	0.21	0.46	0.96
	50	0.29	0.80	1.33	1.63	0.15	0.41	0.70	0.92
	60	0.55	1.12	1.40	1.45	0.27	0.58	0.77	0.81
White	0	0.01	0.01	0.02	1.72	0.01	0.01	0.02	1.01
	10	0.00	0.01	0.04	1.73	0.00	0.01	0.03	1.01
	20	0.00	0.04	0.15	1.73	0.01	0.02	0.09	1.01
	30	0.03	0.15	0.43	1.75	0.02	0.08	0.23	1.01
	40	0.12	0.41	0.92	1.75	0.07	0.22	0.48	1.00
	50	0.30	0.83	1.38	1.69	0.15	0.43	0.73	0.95
	60	0.57	1.17	1.45	1.50	0.28	0.60	0.79	0.83
Black	0	0.01	0.01	0.02	1.46	0.01	0.02	0.03	0.91
	10	0.00	0.01	0.04	1.48	0.00	0.01	0.04	0.92
	20	0.01	0.04	0.16	1.49	0.01	0.03	0.11	0.92
	30	0.03	0.16	0.49	1.52	0.03	0.10	0.25	0.92
	40	0.13	0.47	0.96	1.54	0.07	0.23	0.47	0.91
	50	0.36	0.88	1.29	1.51	0.16	0.41	0.69	0.87
	60	0.60	1.09	1.30	1.34	0.27	0.58	0.74	0.78
Asian/ Pacific Islander	0	0.00	0.00	0.01	1.01	0.00	0.01	0.01	0.58
	10	0.00	0.00	0.02	1.02	0.00	0.00	0.01	0.58
	20	0.00	0.02	0.07	1.02	0.00	0.01	0.04	0.58
	30	0.01	0.07	0.22	1.03	0.01	0.04	0.11	0.58
	40	0.06	0.21	0.48	1.02	0.03	0.10	0.24	0.57
	50	0.15	0.43	0.73	0.98	0.07	0.21	0.40	0.55
	60	0.29	0.61	0.80	0.86	0.14	0.33	0.46	0.49
Hispanic ^a	0	0.01	0.01	0.02	1.68	0.01	0.01	0.01	1.09
	10	0.00	0.01	0.03	1.69	0.00	0.00	0.02	1.09
	20	0.00	0.03	0.13	1.70	0.00	0.02	0.09	1.10
	30	0.02	0.13	0.39	1.71	0.02	0.08	0.24	1.10
	40	0.10	0.37	0.83	1.71	0.07	0.22	0.50	1.09
	50	0.28	0.75	1.31	1.66	0.16	0.44	0.77	1.04
	60	0.50	1.10	1.42	1.48	0.29	0.63	0.85	0.91

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	1.65%	0.97%	0.59%	0.34%
White	1.72%	1.01%	0.61%	0.35%
Black	1.46%	0.91%	0.45%	0.29%
Asian/Pacific Islander	1.01%	0.58%	0.35%	0.20%
Hispanic ^a	1.68%	1.09%	0.60%	0.35%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XII-9
LARYNX CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.00	0.63	0.00	0.00	0.00	0.15
	10	0.00	0.00	0.00	0.64	0.00	0.00	0.00	0.15
	20	0.00	0.00	0.04	0.65	0.00	0.00	0.01	0.15
	30	0.00	0.04	0.15	0.65	0.00	0.01	0.04	0.15
	40	0.03	0.15	0.36	0.66	0.01	0.03	0.09	0.15
	50	0.12	0.34	0.54	0.65	0.03	0.08	0.12	0.14
	60	0.24	0.46	0.56	0.57	0.05	0.10	0.12	0.12
White	0	0.00	0.00	0.00	0.62	0.00	0.00	0.00	0.15
	10	0.00	0.00	0.01	0.63	0.00	0.00	0.00	0.16
	20	0.00	0.01	0.04	0.63	0.00	0.00	0.01	0.16
	30	0.00	0.04	0.15	0.64	0.00	0.01	0.04	0.16
	40	0.03	0.14	0.35	0.65	0.01	0.03	0.09	0.15
	50	0.11	0.33	0.53	0.64	0.03	0.08	0.13	0.15
	60	0.23	0.45	0.55	0.56	0.06	0.11	0.13	0.13
Black	0	0.00	0.00	0.00	0.86	0.00	0.00	0.00	0.18
	10	0.00	0.00	0.01	0.89	0.00	0.00	0.00	0.19
	20	0.00	0.01	0.08	0.89	0.00	0.00	0.03	0.19
	30	0.01	0.08	0.29	0.92	0.00	0.02	0.07	0.19
	40	0.07	0.30	0.61	0.94	0.02	0.07	0.13	0.19
	50	0.24	0.57	0.84	0.93	0.05	0.11	0.16	0.17
	60	0.39	0.70	0.79	0.80	0.07	0.13	0.14	0.14
Asian/ Pacific Islander	0	0.00	0.00	0.00	0.43	0.00	0.00	0.00	0.04
	10	0.00	0.00	0.00	0.43	0.00	0.00	0.00	0.04
	20	0.00	0.00	0.01	0.43	0.00	0.00	0.00	0.04
	30	0.00	0.01	0.05	0.44	0.00	0.00	0.01	0.04
	40	0.01	0.05	0.15	0.44	0.00	0.01	0.02	0.04
	50	0.04	0.14	0.26	0.43	0.00	0.02	0.03	0.04
	60	0.10	0.23	0.33	0.41	0.01	0.02	0.04	0.04
Hispanic ^a	0	0.00	0.00	0.00	0.60	0.00	0.00	0.00	0.09
	10	0.00	0.00	0.00	0.60	0.00	0.00	0.00	0.09
	20	0.00	0.00	0.02	0.61	0.00	0.00	0.00	0.09
	30	0.00	0.02	0.10	0.61	0.00	0.00	0.01	0.09
	40	0.02	0.10	0.24	0.62	0.00	0.01	0.03	0.09
	50	0.08	0.23	0.46	0.62	0.01	0.03	0.05	0.09
	60	0.16	0.40	0.54	0.57	0.02	0.04	0.06	0.08

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	0.63%	0.15%	0.23%	0.06%
White	0.62%	0.15%	0.22%	0.06%
Black	0.86%	0.18%	0.38%	0.08%
Asian/Pacific Islander	0.43%	0.04%	0.13%	0.02%
Hispanic ^a	0.60%	0.09%	0.22%	0.03%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XIII-17
LEUKEMIA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.06	0.09	0.12	1.49	0.05	0.08	0.09	1.05
	10	0.04	0.06	0.10	1.45	0.02	0.04	0.07	1.01
	20	0.03	0.07	0.13	1.42	0.02	0.05	0.09	0.99
	30	0.04	0.11	0.26	1.41	0.03	0.07	0.17	0.98
	40	0.07	0.22	0.53	1.40	0.05	0.14	0.33	0.96
	50	0.16	0.48	0.96	1.38	0.10	0.29	0.58	0.93
	60	0.35	0.86	1.24	1.32	0.20	0.50	0.79	0.87
White	0	0.06	0.10	0.13	1.56	0.05	0.08	0.10	1.10
	10	0.04	0.07	0.10	1.52	0.03	0.05	0.07	1.05
	20	0.03	0.07	0.13	1.49	0.02	0.05	0.10	1.03
	30	0.04	0.11	0.26	1.48	0.03	0.08	0.17	1.01
	40	0.07	0.23	0.55	1.46	0.05	0.15	0.35	0.99
	50	0.16	0.50	1.00	1.44	0.10	0.30	0.60	0.96
	60	0.36	0.90	1.29	1.38	0.21	0.52	0.81	0.90
Black	0	0.03	0.05	0.07	0.94	0.03	0.05	0.06	0.76
	10	0.02	0.05	0.09	0.94	0.02	0.03	0.06	0.75
	20	0.02	0.06	0.13	0.92	0.02	0.04	0.08	0.73
	30	0.04	0.11	0.24	0.92	0.02	0.07	0.14	0.72
	40	0.07	0.21	0.45	0.91	0.04	0.12	0.27	0.71
	50	0.15	0.41	0.70	0.90	0.08	0.24	0.46	0.70
	60	0.30	0.64	0.83	0.87	0.17	0.42	0.60	0.68
Asian/ Pacific Islander	0	0.06	0.08	0.11	1.01	0.05	0.08	0.09	0.72
	10	0.03	0.05	0.09	0.96	0.02	0.04	0.06	0.67
	20	0.03	0.06	0.12	0.94	0.02	0.04	0.08	0.65
	30	0.04	0.09	0.17	0.92	0.02	0.06	0.12	0.63
	40	0.05	0.13	0.32	0.89	0.04	0.10	0.20	0.61
	50	0.08	0.28	0.57	0.85	0.06	0.16	0.32	0.58
	60	0.21	0.51	0.76	0.81	0.10	0.27	0.45	0.53
Hispanic ^a	0	0.07	0.11	0.14	1.19	0.06	0.09	0.12	0.93
	10	0.05	0.08	0.12	1.14	0.03	0.05	0.08	0.88
	20	0.03	0.07	0.13	1.10	0.02	0.05	0.10	0.85
	30	0.04	0.09	0.18	1.08	0.03	0.07	0.15	0.83
	40	0.06	0.15	0.35	1.06	0.04	0.12	0.28	0.80
	50	0.09	0.31	0.63	1.03	0.08	0.24	0.45	0.77
	60	0.23	0.57	0.88	1.00	0.17	0.38	0.60	0.72

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	1.49%	1.05%	0.97%	0.71%
White	1.56%	1.10%	1.02%	0.74%
Black	0.94%	0.76%	0.63%	0.55%
Asian/Pacific Islander	1.01%	0.72%	0.63%	0.49%
Hispanic ^a	1.19%	0.93%	0.72%	0.57%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XIV-9
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.01	0.01	0.88	0.00	0.00	0.01	0.40
	10	0.00	0.00	0.01	0.88	0.00	0.00	0.01	0.40
	20	0.00	0.01	0.08	0.89	0.00	0.01	0.02	0.40
	30	0.01	0.08	0.25	0.90	0.00	0.02	0.06	0.40
	40	0.07	0.24	0.49	0.90	0.01	0.06	0.16	0.40
	50	0.18	0.43	0.70	0.86	0.04	0.15	0.28	0.40
	60	0.27	0.56	0.71	0.74	0.11	0.25	0.34	0.37
White	0	0.00	0.01	0.01	0.74	0.00	0.01	0.01	0.34
	10	0.00	0.00	0.01	0.74	0.00	0.00	0.01	0.34
	20	0.00	0.01	0.06	0.74	0.00	0.00	0.02	0.34
	30	0.01	0.06	0.20	0.75	0.00	0.02	0.05	0.34
	40	0.06	0.20	0.40	0.76	0.01	0.05	0.13	0.34
	50	0.15	0.35	0.58	0.72	0.04	0.12	0.23	0.33
	60	0.22	0.47	0.60	0.62	0.08	0.20	0.29	0.31
Black	0	0.00	0.00	0.01	0.92	0.00	0.00	0.01	0.36
	10	0.00	0.00	0.02	0.95	0.00	0.00	0.01	0.37
	20	0.00	0.02	0.11	0.95	0.00	0.01	0.03	0.37
	30	0.01	0.11	0.39	0.97	0.01	0.02	0.09	0.37
	40	0.10	0.39	0.66	0.99	0.02	0.08	0.17	0.37
	50	0.31	0.60	0.83	0.95	0.06	0.16	0.27	0.37
	60	0.34	0.61	0.72	0.75	0.11	0.22	0.31	0.33
Asian/ Pacific Islander	0	0.01	0.01	0.02	2.46	0.01	0.01	0.01	1.19
	10	0.00	0.01	0.05	2.48	0.00	0.00	0.01	1.20
	20	0.01	0.05	0.19	2.48	0.00	0.01	0.04	1.20
	30	0.04	0.18	0.53	2.49	0.01	0.03	0.12	1.20
	40	0.14	0.50	1.17	2.48	0.02	0.11	0.41	1.19
	50	0.36	1.04	1.79	2.38	0.08	0.39	0.77	1.18
	60	0.71	1.50	1.98	2.11	0.31	0.70	1.03	1.13
Hispanic ^a	0	0.00	0.01	0.01	1.46	0.00	0.00	0.01	0.76
	10	0.00	0.00	0.01	1.47	0.00	0.00	0.01	0.76
	20	0.00	0.01	0.11	1.48	0.00	0.01	0.02	0.76
	30	0.01	0.11	0.37	1.50	0.00	0.02	0.08	0.77
	40	0.11	0.37	0.78	1.51	0.01	0.08	0.24	0.77
	50	0.27	0.69	1.17	1.45	0.06	0.23	0.50	0.76
	60	0.45	0.96	1.20	1.26	0.17	0.45	0.67	0.73

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	0.88%	0.40%	0.67%	0.38%
White	0.74%	0.34%	0.63%	0.36%
Black	0.92%	0.36%	0.72%	0.41%
Asian/Pacific Islander	2.46%	1.19%	1.97%	1.13%
Hispanic ^a	1.46%	0.76%	1.18%	0.76%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XV-15
LUNG AND BRONCHUS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.01	8.02	0.00	0.00	0.01	6.15
	10	0.00	0.01	0.03	8.13	0.00	0.01	0.04	6.22
	20	0.00	0.03	0.23	8.17	0.00	0.04	0.21	6.23
	30	0.03	0.23	1.10	8.28	0.03	0.21	0.88	6.26
	40	0.21	1.09	3.41	8.39	0.18	0.85	2.57	6.28
	50	0.92	3.32	6.55	8.50	0.69	2.44	4.74	6.24
	60	2.61	6.10	7.93	8.21	1.84	4.26	5.61	5.83
White	0	0.00	0.00	0.01	8.09	0.00	0.00	0.01	6.47
	10	0.00	0.01	0.03	8.18	0.00	0.01	0.04	6.53
	20	0.00	0.03	0.22	8.23	0.01	0.04	0.21	6.55
	30	0.03	0.22	1.05	8.32	0.03	0.21	0.90	6.57
	40	0.19	1.04	3.39	8.43	0.18	0.88	2.70	6.59
	50	0.88	3.31	6.60	8.53	0.71	2.58	5.02	6.54
	60	2.63	6.18	7.99	8.26	1.95	4.51	5.90	6.11
Black	0	0.00	0.00	0.00	8.26	0.00	0.00	0.00	5.21
	10	0.00	0.00	0.04	8.49	0.00	0.00	0.04	5.32
	20	0.00	0.04	0.39	8.56	0.00	0.04	0.28	5.34
	30	0.03	0.40	1.87	8.77	0.03	0.28	1.07	5.38
	40	0.38	1.89	4.67	9.03	0.25	1.06	2.64	5.44
	50	1.62	4.59	7.66	9.26	0.84	2.49	4.32	5.41
	60	3.46	7.03	8.66	8.90	1.81	3.81	4.81	5.01
Asian/ Pacific Islander	0	0.00	0.00	0.00	7.07	0.00	0.00	0.01	4.18
	10	0.00	0.00	0.04	7.14	0.00	0.01	0.03	4.21
	20	0.00	0.04	0.17	7.16	0.00	0.03	0.12	4.22
	30	0.03	0.17	0.71	7.21	0.03	0.12	0.46	4.22
	40	0.14	0.68	2.18	7.24	0.10	0.44	1.25	4.22
	50	0.55	2.08	4.66	7.23	0.34	1.17	2.46	4.17
	60	1.60	4.30	6.48	6.99	0.85	2.18	3.50	3.93
Hispanic ^a	0	0.00	0.00	0.00	5.21	0.00	0.00	0.00	3.36
	10	0.00	0.00	0.02	5.28	0.00	0.00	0.02	3.39
	20	0.00	0.02	0.10	5.31	0.00	0.02	0.09	3.40
	30	0.01	0.09	0.46	5.37	0.01	0.09	0.36	3.41
	40	0.08	0.45	1.64	5.43	0.08	0.34	1.08	3.42
	50	0.39	1.60	3.68	5.52	0.27	1.02	2.18	3.39
	60	1.30	3.52	5.02	5.48	0.77	1.98	2.88	3.24

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	8.02%	6.15%	7.18%	4.96%
White	8.09%	6.47%	7.26%	5.16%
Black	8.26%	5.21%	7.18%	4.03%
Asian/Pacific Islander	7.07%	4.18%	5.41%	3.21%
Hispanic ^a	5.21%	3.36%	4.31%	2.20%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XVI-9
MELANOMA OF SKIN (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.01	0.05	2.04	0.00	0.02	0.09	1.38
	10	0.01	0.05	0.13	2.06	0.01	0.09	0.22	1.39
	20	0.04	0.12	0.30	2.07	0.07	0.20	0.39	1.38
	30	0.09	0.27	0.61	2.06	0.13	0.32	0.55	1.31
	40	0.19	0.53	1.03	2.01	0.19	0.42	0.69	1.20
	50	0.36	0.88	1.46	1.89	0.24	0.51	0.79	1.03
	60	0.56	1.19	1.58	1.66	0.29	0.58	0.78	0.83
White	0	0.00	0.01	0.06	2.33	0.00	0.02	0.11	1.60
	10	0.01	0.06	0.15	2.36	0.02	0.11	0.26	1.62
	20	0.04	0.14	0.36	2.36	0.09	0.24	0.47	1.60
	30	0.10	0.32	0.71	2.35	0.15	0.38	0.65	1.52
	40	0.22	0.62	1.19	2.29	0.23	0.50	0.82	1.38
	50	0.41	1.01	1.67	2.14	0.28	0.60	0.92	1.18
	60	0.64	1.36	1.79	1.87	0.34	0.68	0.89	0.94
Black	0	0.00	0.00	0.00	0.08	0.00	0.00	0.00	0.09
	10	0.00	0.00	0.01	0.08	0.00	0.00	0.01	0.09
	20	0.00	0.01	0.01	0.08	0.00	0.01	0.01	0.09
	30	0.00	0.01	0.02	0.08	0.00	0.01	0.02	0.08
	40	0.01	0.02	0.04	0.08	0.00	0.01	0.03	0.08
	50	0.01	0.04	0.06	0.08	0.01	0.03	0.05	0.08
	60	0.03	0.06	0.07	0.07	0.02	0.05	0.06	0.08
Asian/ Pacific Islander	0	0.00	0.00	0.00	0.19	0.00	0.00	0.00	0.14
	10	0.00	0.00	0.01	0.19	0.00	0.00	0.02	0.14
	20	0.00	0.01	0.02	0.19	0.00	0.02	0.03	0.14
	30	0.01	0.02	0.04	0.19	0.01	0.03	0.05	0.14
	40	0.01	0.03	0.06	0.18	0.01	0.03	0.05	0.13
	50	0.02	0.05	0.10	0.18	0.02	0.04	0.06	0.12
	60	0.03	0.09	0.12	0.16	0.02	0.04	0.08	0.10
Hispanic ^a	0	0.00	0.00	0.01	0.45	0.00	0.00	0.02	0.49
	10	0.00	0.01	0.02	0.46	0.00	0.02	0.06	0.49
	20	0.01	0.02	0.05	0.46	0.02	0.05	0.11	0.49
	30	0.01	0.04	0.09	0.46	0.04	0.10	0.16	0.47
	40	0.03	0.08	0.17	0.45	0.06	0.12	0.21	0.44
	50	0.05	0.14	0.28	0.43	0.06	0.15	0.26	0.39
	60	0.10	0.25	0.39	0.41	0.09	0.20	0.28	0.34

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	2.04%	1.38%	0.36%	0.20%
White	2.33%	1.60%	0.41%	0.22%
Black	0.08%	0.09%	0.03%	0.04%
Asian/Pacific Islander	0.19%	0.14%	0.07%	0.06%
Hispanic ^a	0.45%	0.49%	0.11%	0.10%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XVII-8
MESOTHELIOMA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.00	0.20	0.00	0.00	0.00	0.05
	10	0.00	0.00	0.00	0.21	0.00	0.00	0.00	0.05
	20	0.00	0.00	0.00	0.21	0.00	0.00	0.00	0.05
	30	0.00	0.00	0.02	0.21	0.00	0.00	0.01	0.05
	40	0.00	0.02	0.06	0.21	0.00	0.01	0.02	0.05
	50	0.02	0.06	0.15	0.22	0.01	0.02	0.03	0.05
	60	0.05	0.14	0.21	0.22	0.01	0.03	0.04	0.05
White	0	0.00	0.00	0.00	0.23	0.00	0.00	0.00	0.06
	10	0.00	0.00	0.00	0.23	0.00	0.00	0.00	0.06
	20	0.00	0.00	0.00	0.23	0.00	0.00	0.00	0.06
	30	0.00	0.00	0.02	0.23	0.00	0.00	0.01	0.06
	40	0.00	0.02	0.07	0.23	0.00	0.01	0.02	0.06
	50	0.02	0.07	0.17	0.24	0.01	0.02	0.04	0.05
	60	0.06	0.16	0.23	0.24	0.01	0.03	0.05	0.05
Black	0	0.00	0.00	0.00	0.08	0.00	0.00	0.00	0.02
	10	0.00	0.00	0.00	0.08	0.00	0.00	0.00	0.02
	20	0.00	0.00	0.00	0.08	0.00	0.00	0.00	0.02
	30	0.00	0.00	0.02	0.08	0.00	0.00	0.00	0.02
	40	0.00	0.02	0.03	0.08	0.00	0.00	0.01	0.02
	50	0.01	0.03	0.06	0.08	0.00	0.01	0.02	0.02
	60	0.01	0.05	0.08	0.08	0.01	0.01	0.02	0.02
Asian/ Pacific Islander	0	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.02
	10	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.02
	20	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.02
	30	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.02
	40	0.00	0.00	0.02	0.09	0.00	0.00	0.01	0.02
	50	0.00	0.02	0.06	0.09	0.00	0.01	0.01	0.02
	60	0.02	0.06	0.08	0.09	0.01	0.01	0.02	0.02
Hispanic ^a	0	0.00	0.00	0.00	0.20	0.00	0.00	0.00	0.07
	10	0.00	0.00	0.00	0.20	0.00	0.00	0.00	0.07
	20	0.00	0.00	0.01	0.21	0.00	0.00	0.00	0.07
	30	0.00	0.01	0.02	0.21	0.00	0.00	0.01	0.07
	40	0.00	0.02	0.07	0.21	0.00	0.01	0.02	0.07
	50	0.02	0.06	0.15	0.21	0.00	0.02	0.04	0.07
	60	0.05	0.14	0.20	0.21	0.01	0.04	0.06	0.06

Race/Ethnicity	Lifetime Risk of Developing	
	Males	Females
All Races	0.20%	0.16%
White	0.23%	0.18%
Black	0.08%	0.05%
Asian/Pacific Islander	0.09%	0.07%
Hispanic ^a	0.20%	0.11%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky.

Table XVIII-9
MULTIPLE MYELOMA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.00	0.67	0.00	0.00	0.00	0.53
	10	0.00	0.00	0.01	0.68	0.00	0.00	0.00	0.54
	20	0.00	0.01	0.03	0.68	0.00	0.00	0.02	0.54
	30	0.01	0.03	0.12	0.69	0.00	0.02	0.09	0.54
	40	0.03	0.11	0.29	0.69	0.02	0.08	0.20	0.54
	50	0.09	0.27	0.51	0.69	0.06	0.19	0.37	0.53
	60	0.20	0.46	0.62	0.66	0.13	0.32	0.46	0.49
White	0	0.00	0.00	0.00	0.63	0.00	0.00	0.00	0.49
	10	0.00	0.00	0.01	0.64	0.00	0.00	0.00	0.49
	20	0.00	0.01	0.03	0.64	0.00	0.00	0.02	0.50
	30	0.01	0.03	0.11	0.65	0.00	0.02	0.07	0.50
	40	0.02	0.10	0.26	0.65	0.02	0.07	0.18	0.50
	50	0.08	0.25	0.48	0.65	0.06	0.17	0.34	0.49
	60	0.18	0.43	0.59	0.62	0.12	0.30	0.43	0.45
Black	0	0.00	0.00	0.00	1.06	0.00	0.00	0.00	0.89
	10	0.00	0.00	0.01	1.08	0.00	0.00	0.01	0.91
	20	0.00	0.01	0.06	1.09	0.00	0.01	0.05	0.91
	30	0.01	0.07	0.23	1.12	0.01	0.05	0.18	0.91
	40	0.05	0.22	0.54	1.14	0.04	0.17	0.41	0.92
	50	0.18	0.52	0.90	1.17	0.13	0.38	0.68	0.92
	60	0.40	0.84	1.10	1.15	0.27	0.60	0.81	0.85
Asian/ Pacific Islander	0	0.00	0.00	0.00	0.45	0.00	0.00	0.00	0.43
	10	0.00	0.00	0.00	0.46	0.00	0.00	0.00	0.43
	20	0.00	0.00	0.02	0.46	0.00	0.00	0.02	0.43
	30	0.00	0.01	0.06	0.46	0.00	0.01	0.06	0.43
	40	0.01	0.06	0.17	0.46	0.01	0.05	0.13	0.44
	50	0.05	0.16	0.32	0.46	0.04	0.12	0.26	0.43
	60	0.12	0.29	0.40	0.43	0.08	0.23	0.35	0.40
Hispanic ^a	0	0.00	0.00	0.00	0.77	0.00	0.00	0.00	0.63
	10	0.00	0.00	0.01	0.78	0.00	0.00	0.00	0.63
	20	0.00	0.01	0.03	0.79	0.00	0.00	0.02	0.63
	30	0.01	0.03	0.10	0.79	0.00	0.02	0.09	0.64
	40	0.02	0.10	0.27	0.80	0.02	0.08	0.23	0.64
	50	0.08	0.26	0.50	0.80	0.07	0.22	0.42	0.63
	60	0.19	0.45	0.71	0.77	0.15	0.37	0.53	0.58

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	0.67%	0.53%	0.46%	0.40%
White	0.63%	0.49%	0.45%	0.37%
Black	1.06%	0.89%	0.62%	0.66%
Asian/Pacific Islander	0.45%	0.43%	0.25%	0.26%
Hispanic ^a	0.77%	0.63%	0.45%	0.39%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XIX-13
NON-HODGKIN LYMPHOMA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.01	0.03	0.06	2.14	0.00	0.01	0.04	1.83
	10	0.02	0.06	0.13	2.16	0.01	0.03	0.08	1.85
	20	0.04	0.11	0.26	2.15	0.02	0.07	0.17	1.84
	30	0.07	0.23	0.52	2.14	0.05	0.14	0.36	1.83
	40	0.16	0.45	0.96	2.11	0.10	0.32	0.73	1.80
	50	0.31	0.83	1.50	2.03	0.23	0.64	1.21	1.74
	60	0.57	1.29	1.78	1.86	0.44	1.03	1.49	1.59
White	0	0.01	0.03	0.06	2.24	0.00	0.01	0.04	1.92
	10	0.02	0.06	0.13	2.26	0.01	0.03	0.08	1.94
	20	0.04	0.11	0.26	2.25	0.02	0.07	0.17	1.93
	30	0.07	0.23	0.53	2.25	0.05	0.14	0.37	1.92
	40	0.16	0.46	1.00	2.21	0.10	0.33	0.77	1.89
	50	0.31	0.87	1.58	2.13	0.23	0.68	1.28	1.83
	60	0.60	1.37	1.87	1.96	0.47	1.09	1.57	1.67
Black	0	0.01	0.02	0.06	1.25	0.00	0.01	0.04	1.07
	10	0.02	0.06	0.16	1.27	0.01	0.04	0.10	1.08
	20	0.04	0.14	0.32	1.27	0.03	0.09	0.19	1.08
	30	0.10	0.29	0.54	1.26	0.06	0.16	0.34	1.06
	40	0.19	0.45	0.76	1.19	0.11	0.28	0.52	1.01
	50	0.29	0.62	0.92	1.08	0.19	0.44	0.73	0.95
	60	0.38	0.73	0.90	0.92	0.28	0.59	0.79	0.84
Asian/ Pacific Islander	0	0.01	0.02	0.05	1.91	0.00	0.02	0.04	1.60
	10	0.02	0.05	0.09	1.92	0.01	0.03	0.06	1.61
	20	0.03	0.07	0.17	1.91	0.02	0.05	0.11	1.60
	30	0.04	0.14	0.32	1.89	0.03	0.09	0.25	1.58
	40	0.10	0.28	0.63	1.86	0.07	0.23	0.48	1.56
	50	0.19	0.55	1.12	1.80	0.16	0.42	0.89	1.51
	60	0.37	0.97	1.49	1.69	0.27	0.75	1.24	1.39
Hispanic ^a	0	0.01	0.03	0.06	1.96	0.00	0.01	0.03	1.77
	10	0.02	0.05	0.11	1.98	0.01	0.02	0.06	1.79
	20	0.03	0.09	0.22	1.97	0.02	0.05	0.13	1.78
	30	0.06	0.19	0.42	1.97	0.04	0.11	0.30	1.77
	40	0.14	0.36	0.75	1.93	0.07	0.27	0.63	1.75
	50	0.24	0.64	1.26	1.86	0.20	0.57	1.12	1.70
	60	0.43	1.10	1.59	1.73	0.38	0.96	1.42	1.55

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	2.14%	1.83%	0.93%	0.79%
White	2.24%	1.92%	0.99%	0.83%
Black	1.25%	1.07%	0.49%	0.44%
Asian/Pacific Islander	1.91%	1.60%	0.87%	0.70%
Hispanic ^a	1.96%	1.77%	0.79%	0.72%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XX-9
ORAL CAVITY AND PHARYNX CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.01	1.39	0.00	0.00	0.01	0.66
	10	0.00	0.01	0.04	1.41	0.00	0.01	0.03	0.66
	20	0.01	0.04	0.16	1.41	0.01	0.03	0.07	0.66
	30	0.03	0.16	0.45	1.42	0.02	0.06	0.16	0.66
	40	0.13	0.43	0.83	1.42	0.05	0.14	0.29	0.64
	50	0.31	0.73	1.10	1.34	0.10	0.25	0.44	0.61
	60	0.45	0.85	1.06	1.11	0.16	0.36	0.50	0.54
White	0	0.00	0.00	0.01	1.41	0.00	0.00	0.01	0.66
	10	0.00	0.01	0.04	1.43	0.00	0.01	0.03	0.67
	20	0.01	0.03	0.16	1.44	0.01	0.03	0.07	0.67
	30	0.03	0.15	0.45	1.45	0.02	0.06	0.15	0.66
	40	0.13	0.43	0.84	1.44	0.04	0.14	0.29	0.65
	50	0.31	0.74	1.12	1.36	0.10	0.25	0.45	0.62
	60	0.46	0.87	1.08	1.13	0.17	0.37	0.51	0.55
Black	0	0.00	0.01	0.02	1.26	0.00	0.00	0.01	0.53
	10	0.01	0.01	0.05	1.30	0.00	0.01	0.03	0.54
	20	0.01	0.04	0.19	1.30	0.01	0.03	0.10	0.54
	30	0.03	0.18	0.55	1.33	0.02	0.09	0.19	0.54
	40	0.16	0.53	0.94	1.34	0.07	0.18	0.31	0.52
	50	0.41	0.84	1.14	1.27	0.12	0.26	0.37	0.48
	60	0.51	0.85	0.98	1.00	0.16	0.28	0.37	0.40
Asian/ Pacific Islander	0	0.00	0.00	0.01	1.11	0.00	0.00	0.02	0.66
	10	0.00	0.01	0.05	1.13	0.00	0.02	0.05	0.67
	20	0.01	0.04	0.17	1.13	0.01	0.04	0.09	0.67
	30	0.04	0.16	0.37	1.12	0.03	0.08	0.16	0.65
	40	0.12	0.34	0.60	1.10	0.05	0.13	0.26	0.63
	50	0.22	0.49	0.76	0.99	0.09	0.21	0.35	0.59
	60	0.29	0.56	0.75	0.81	0.13	0.27	0.45	0.52
Hispanic ^a	0	0.00	0.00	0.01	0.96	0.00	0.00	0.01	0.47
	10	0.00	0.01	0.02	0.97	0.00	0.01	0.02	0.47
	20	0.00	0.02	0.08	0.98	0.00	0.01	0.04	0.47
	30	0.02	0.08	0.24	0.98	0.01	0.04	0.09	0.47
	40	0.07	0.23	0.50	0.98	0.03	0.08	0.15	0.46
	50	0.17	0.45	0.72	0.94	0.05	0.13	0.25	0.44
	60	0.30	0.59	0.74	0.83	0.08	0.21	0.34	0.40

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	1.39%	0.66%	0.39%	0.19%
White	1.41%	0.66%	0.37%	0.19%
Black	1.26%	0.53%	0.50%	0.17%
Asian/Pacific Islander	1.11%	0.66%	0.45%	0.24%
Hispanic ^a	0.96%	0.47%	0.32%	0.14%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XXI-7
OVARY CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/Ethnicity	Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.01	0.03	1.44
	10	0.01	0.03	0.07	1.46
	20	0.02	0.06	0.19	1.45
	30	0.04	0.17	0.41	1.44
	40	0.12	0.37	0.73	1.41
	50	0.25	0.62	1.00	1.31
	60	0.38	0.78	1.05	1.11
White	0	0.00	0.01	0.03	1.53
	10	0.01	0.03	0.08	1.54
	20	0.02	0.06	0.19	1.54
	30	0.04	0.17	0.44	1.52
	40	0.13	0.40	0.78	1.49
	50	0.27	0.66	1.07	1.39
	60	0.41	0.83	1.11	1.17
Black	0	0.00	0.01	0.02	0.95
	10	0.01	0.02	0.06	0.97
	20	0.02	0.05	0.13	0.97
	30	0.04	0.12	0.26	0.96
	40	0.08	0.22	0.47	0.94
	50	0.15	0.40	0.65	0.90
	60	0.28	0.54	0.76	0.82
Asian/Pacific Islander	0	0.00	0.01	0.04	1.12
	10	0.01	0.04	0.08	1.13
	20	0.03	0.07	0.19	1.12
	30	0.04	0.16	0.34	1.10
	40	0.12	0.30	0.54	1.06
	50	0.19	0.42	0.69	0.95
	60	0.24	0.51	0.71	0.79
Hispanic ^a	0	0.00	0.02	0.04	1.35
	10	0.01	0.03	0.07	1.36
	20	0.02	0.06	0.16	1.35
	30	0.04	0.14	0.34	1.33
	40	0.11	0.31	0.60	1.30
	50	0.20	0.50	0.87	1.22
	60	0.31	0.69	0.98	1.05

Race/Ethnicity	Lifetime Risk of Developing	Lifetime Risk of Dying
All Races	1.44%	1.05%
White	1.53%	1.11%
Black	0.95%	0.76%
Asian/Pacific Islander	1.12%	0.69%
Hispanic ^a	1.35%	0.81%

Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XXII-9
PANCREAS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.00	1.27	0.00	0.00	0.00	1.27
	10	0.00	0.00	0.01	1.29	0.00	0.00	0.01	1.29
	20	0.00	0.01	0.05	1.30	0.00	0.01	0.04	1.29
	30	0.01	0.05	0.20	1.31	0.01	0.04	0.14	1.30
	40	0.04	0.20	0.53	1.33	0.03	0.13	0.40	1.30
	50	0.16	0.50	0.97	1.33	0.11	0.38	0.81	1.30
	60	0.37	0.88	1.20	1.27	0.29	0.74	1.13	1.25
White	0	0.00	0.00	0.00	1.29	0.00	0.00	0.00	1.25
	10	0.00	0.00	0.01	1.30	0.00	0.00	0.01	1.26
	20	0.00	0.01	0.05	1.31	0.00	0.01	0.04	1.27
	30	0.01	0.05	0.20	1.32	0.01	0.04	0.14	1.27
	40	0.04	0.20	0.53	1.34	0.03	0.13	0.39	1.27
	50	0.16	0.50	0.98	1.34	0.10	0.37	0.80	1.27
	60	0.36	0.88	1.20	1.27	0.28	0.73	1.11	1.22
Black	0	0.00	0.00	0.00	1.18	0.00	0.00	0.00	1.46
	10	0.00	0.00	0.01	1.21	0.00	0.00	0.01	1.49
	20	0.00	0.01	0.07	1.22	0.00	0.01	0.06	1.50
	30	0.01	0.07	0.27	1.25	0.01	0.06	0.20	1.51
	40	0.06	0.26	0.64	1.28	0.05	0.20	0.56	1.53
	50	0.22	0.62	1.06	1.31	0.15	0.53	1.02	1.54
	60	0.47	0.98	1.23	1.26	0.41	0.95	1.36	1.52
Asian/ Pacific Islander	0	0.00	0.00	0.00	1.28	0.00	0.00	0.00	1.31
	10	0.00	0.00	0.01	1.30	0.00	0.00	0.01	1.32
	20	0.00	0.01	0.03	1.30	0.00	0.01	0.03	1.32
	30	0.01	0.03	0.13	1.31	0.01	0.03	0.11	1.32
	40	0.03	0.13	0.39	1.31	0.03	0.11	0.33	1.32
	50	0.10	0.37	0.79	1.31	0.08	0.30	0.70	1.31
	60	0.28	0.71	1.13	1.26	0.23	0.64	1.07	1.26
Hispanic ^a	0	0.00	0.00	0.00	1.18	0.00	0.00	0.00	1.54
	10	0.00	0.00	0.01	1.20	0.00	0.00	0.01	1.55
	20	0.00	0.01	0.03	1.20	0.00	0.01	0.03	1.56
	30	0.01	0.03	0.15	1.22	0.01	0.03	0.12	1.56
	40	0.03	0.15	0.42	1.23	0.02	0.12	0.39	1.56
	50	0.13	0.40	0.86	1.24	0.10	0.37	0.84	1.57
	60	0.30	0.78	1.11	1.18	0.29	0.77	1.27	1.52

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	1.27%	1.27%	1.18%	1.18%
White	1.29%	1.25%	1.19%	1.17%
Black	1.18%	1.46%	1.14%	1.33%
Asian/Pacific Islander	1.28%	1.31%	1.08%	1.19%
Hispanic ^a	1.18%	1.54%	1.03%	1.18%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XXIII-7
PROSTATE CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/Ethnicity	Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.00	17.12
	10	0.00	0.00	0.01	17.36
	20	0.00	0.01	0.30	17.46
	30	0.01	0.30	2.56	17.69
	40	0.30	2.59	8.70	17.99
	50	2.39	8.74	15.25	18.40
	60	7.03	14.25	17.21	17.74
White	0	0.00	0.00	0.00	16.58
	10	0.00	0.00	0.01	16.78
	20	0.00	0.01	0.27	16.87
	30	0.01	0.27	2.43	17.07
	40	0.27	2.47	8.45	17.35
	50	2.28	8.49	14.79	17.72
	60	6.84	13.79	16.54	17.02
Black	0	0.00	0.00	0.00	19.76
	10	0.00	0.00	0.02	20.30
	20	0.00	0.02	0.61	20.48
	30	0.02	0.62	4.37	20.99
	40	0.62	4.50	12.45	21.67
	50	4.17	12.73	19.69	22.64
	60	10.37	18.81	21.82	22.39
Asian/Pacific Islander	0	0.00	0.00	0.00	12.62
	10	0.00	0.00	0.00	12.75
	20	0.00	0.00	0.09	12.79
	30	0.00	0.09	0.97	12.87
	40	0.09	0.97	4.29	12.97
	50	0.90	4.28	9.43	13.14
	60	3.56	8.99	12.08	12.90
Hispanic ^a	0	0.00	0.00	0.00	16.29
	10	0.00	0.00	0.00	16.49
	20	0.00	0.00	0.17	16.59
	30	0.00	0.17	1.68	16.78
	40	0.17	1.70	6.81	17.01
	50	1.57	6.86	13.22	17.40
	60	5.73	12.63	16.02	17.15

Race/Ethnicity	Lifetime Risk of Developing	Lifetime Risk of Dying
All Races	17.12%	2.91%
White	16.58%	2.73%
Black	19.76%	4.59%
Asian/Pacific Islander	12.62%	2.18%
Hispanic ^a	16.29%	3.24%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XXIV-9
STOMACH CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.00	1.12	0.00	0.00	0.00	0.68
	10	0.00	0.00	0.02	1.14	0.00	0.00	0.02	0.68
	20	0.00	0.02	0.06	1.14	0.00	0.01	0.04	0.68
	30	0.01	0.06	0.18	1.15	0.01	0.04	0.10	0.69
	40	0.05	0.17	0.45	1.16	0.03	0.09	0.21	0.68
	50	0.13	0.41	0.81	1.15	0.06	0.19	0.40	0.66
	60	0.31	0.74	1.04	1.11	0.13	0.35	0.56	0.63
White	0	0.00	0.00	0.00	1.01	0.00	0.00	0.00	0.57
	10	0.00	0.00	0.02	1.02	0.00	0.00	0.01	0.58
	20	0.00	0.02	0.06	1.02	0.00	0.01	0.04	0.58
	30	0.01	0.05	0.16	1.03	0.01	0.04	0.09	0.58
	40	0.04	0.15	0.40	1.04	0.02	0.08	0.18	0.57
	50	0.12	0.37	0.73	1.03	0.05	0.16	0.33	0.56
	60	0.28	0.66	0.93	0.98	0.11	0.29	0.47	0.53
Black	0	0.00	0.00	0.01	1.31	0.00	0.00	0.00	0.99
	10	0.00	0.01	0.02	1.34	0.00	0.00	0.02	1.01
	20	0.01	0.02	0.09	1.35	0.00	0.02	0.05	1.01
	30	0.02	0.09	0.27	1.38	0.01	0.05	0.14	1.02
	40	0.07	0.26	0.63	1.41	0.04	0.13	0.30	1.02
	50	0.20	0.59	1.07	1.43	0.09	0.27	0.59	1.02
	60	0.46	1.01	1.34	1.43	0.20	0.55	0.88	1.02
Asian/ Pacific Islander	0	0.00	0.00	0.01	2.44	0.00	0.00	0.00	1.72
	10	0.00	0.01	0.03	2.46	0.00	0.00	0.03	1.74
	20	0.01	0.03	0.10	2.47	0.00	0.02	0.08	1.74
	30	0.02	0.09	0.27	2.48	0.02	0.07	0.19	1.74
	40	0.07	0.25	0.66	2.48	0.05	0.17	0.39	1.73
	50	0.18	0.60	1.38	2.46	0.11	0.34	0.83	1.69
	60	0.44	1.26	2.11	2.38	0.23	0.74	1.31	1.62
Hispanic ^a	0	0.00	0.00	0.01	1.80	0.00	0.00	0.00	1.36
	10	0.00	0.01	0.03	1.82	0.00	0.00	0.03	1.38
	20	0.01	0.03	0.09	1.83	0.00	0.02	0.08	1.38
	30	0.02	0.09	0.26	1.84	0.02	0.07	0.20	1.38
	40	0.06	0.24	0.62	1.85	0.05	0.18	0.40	1.37
	50	0.18	0.57	1.21	1.84	0.13	0.35	0.73	1.34
	60	0.41	1.09	1.57	1.77	0.23	0.62	1.07	1.25

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	1.12%	0.68%	0.57%	0.38%
White	1.01%	0.57%	0.52%	0.34%
Black	1.31%	0.99%	0.87%	0.64%
Asian/Pacific Islander	2.44%	1.72%	1.56%	1.20%
Hispanic ^a	1.80%	1.36%	1.06%	0.76%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XXV-7
TESTIS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/Ethnicity	Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.02	0.12	0.36
	10	0.02	0.12	0.24	0.36
	20	0.10	0.22	0.29	0.34
	30	0.12	0.19	0.22	0.24
	40	0.07	0.11	0.12	0.13
	50	0.03	0.05	0.05	0.06
	60	0.01	0.02	0.02	0.03
White	0	0.00	0.03	0.15	0.42
	10	0.03	0.14	0.28	0.42
	20	0.12	0.26	0.34	0.40
	30	0.14	0.23	0.26	0.29
	40	0.09	0.12	0.14	0.15
	50	0.04	0.05	0.06	0.06
	60	0.02	0.02	0.03	0.03
Black	0	0.00	0.01	0.03	0.10
	10	0.01	0.03	0.06	0.10
	20	0.02	0.06	0.08	0.10
	30	0.03	0.05	0.06	0.07
	40	0.02	0.03	0.04	0.04
	50	0.01	0.02	0.02	0.02
	60	0.01	0.01	0.01	0.01
Asian/Pacific Islander	0	0.00	0.01	0.04	0.11
	10	0.01	0.04	0.07	0.11
	20	0.03	0.06	0.08	0.10
	30	0.03	0.05	0.06	0.07
	40	0.02	0.03	0.03	0.04
	50	0.01	0.01	0.02	0.02
	60	0.00	0.01	0.01	0.01
Hispanic ^a	0	0.00	0.03	0.11	0.26
	10	0.02	0.11	0.19	0.26
	20	0.09	0.17	0.21	0.24
	30	0.08	0.12	0.14	0.15
	40	0.04	0.06	0.06	0.07
	50	0.02	0.03	0.03	0.03
	60	0.01	0.01	0.01	0.01

Race/Ethnicity	Lifetime Risk of Developing	Lifetime Risk of Dying
All Races	0.36%	0.02%
White	0.42%	0.02%
Black	0.10%	0.01%
Asian/Pacific Islander	0.11%	0.01%
Hispanic ^a	0.26%	0.02%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XXVI-9
THYROID CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.02	0.36	0.00	0.02	0.12	1.02
	10	0.00	0.02	0.06	0.36	0.02	0.12	0.29	1.03
	20	0.02	0.05	0.11	0.36	0.10	0.27	0.47	1.01
	30	0.04	0.09	0.17	0.35	0.17	0.37	0.57	0.91
	40	0.05	0.13	0.22	0.32	0.20	0.40	0.57	0.75
	50	0.08	0.17	0.25	0.27	0.20	0.37	0.50	0.56
	60	0.10	0.18	0.21	0.21	0.18	0.31	0.37	0.38
White	0	0.00	0.01	0.02	0.38	0.00	0.03	0.13	1.06
	10	0.01	0.02	0.06	0.39	0.02	0.13	0.32	1.07
	20	0.02	0.06	0.12	0.38	0.11	0.29	0.51	1.05
	30	0.04	0.10	0.18	0.37	0.19	0.40	0.60	0.95
	40	0.06	0.14	0.24	0.34	0.22	0.42	0.59	0.77
	50	0.08	0.18	0.26	0.29	0.20	0.38	0.50	0.57
	60	0.11	0.19	0.22	0.22	0.19	0.32	0.37	0.38
Black	0	0.00	0.00	0.01	0.17	0.00	0.01	0.05	0.59
	10	0.00	0.01	0.02	0.17	0.01	0.05	0.13	0.60
	20	0.01	0.02	0.04	0.17	0.04	0.12	0.22	0.59
	30	0.02	0.04	0.08	0.17	0.08	0.18	0.31	0.56
	40	0.02	0.07	0.12	0.16	0.10	0.23	0.34	0.48
	50	0.05	0.10	0.13	0.14	0.13	0.25	0.35	0.40
	60	0.06	0.10	0.11	0.11	0.13	0.24	0.28	0.29
Asian/ Pacific Islander	0	0.00	0.00	0.02	0.40	0.00	0.02	0.11	1.12
	10	0.00	0.02	0.06	0.40	0.02	0.11	0.28	1.12
	20	0.02	0.06	0.10	0.40	0.09	0.26	0.46	1.11
	30	0.03	0.08	0.14	0.38	0.16	0.37	0.59	1.02
	40	0.05	0.10	0.19	0.35	0.20	0.43	0.62	0.86
	50	0.06	0.15	0.24	0.31	0.23	0.42	0.58	0.66
	60	0.10	0.19	0.24	0.26	0.20	0.37	0.44	0.45
Hispanic ^a	0	0.00	0.00	0.01	0.30	0.00	0.02	0.09	1.02
	10	0.00	0.01	0.04	0.30	0.02	0.09	0.23	1.03
	20	0.01	0.03	0.07	0.30	0.08	0.21	0.38	1.02
	30	0.02	0.06	0.11	0.29	0.14	0.31	0.50	0.94
	40	0.04	0.09	0.17	0.27	0.17	0.37	0.56	0.81
	50	0.05	0.14	0.21	0.24	0.20	0.39	0.54	0.65
	60	0.09	0.17	0.20	0.20	0.20	0.36	0.44	0.47

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	0.36%	1.02%	0.04%	0.06%
White	0.38%	1.06%	0.04%	0.06%
Black	0.17%	0.59%	0.03%	0.05%
Asian/Pacific Islander	0.40%	1.12%	0.05%	0.12%
Hispanic ^a	0.30%	1.02%	0.06%	0.09%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table XXVII-9
URINARY BLADDER CANCER (Invasive and In Situ)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (17 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2001-2003

Race/ Ethnicity	Current Age	Males				Females			
		+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
All Races	0	0.00	0.00	0.01	3.61	0.00	0.00	0.00	1.14
	10	0.00	0.00	0.02	3.66	0.00	0.00	0.01	1.16
	20	0.00	0.02	0.10	3.68	0.00	0.01	0.04	1.16
	30	0.02	0.10	0.43	3.73	0.01	0.04	0.13	1.16
	40	0.08	0.41	1.26	3.77	0.03	0.13	0.37	1.17
	50	0.34	1.22	2.63	3.83	0.10	0.35	0.74	1.16
	60	0.96	2.48	3.55	3.78	0.26	0.66	1.01	1.11
White	0	0.00	0.00	0.01	3.95	0.00	0.00	0.00	1.22
	10	0.00	0.01	0.02	4.00	0.00	0.00	0.01	1.23
	20	0.00	0.02	0.11	4.02	0.00	0.01	0.04	1.24
	30	0.02	0.11	0.47	4.07	0.01	0.04	0.15	1.24
	40	0.09	0.45	1.40	4.11	0.03	0.14	0.41	1.24
	50	0.38	1.36	2.89	4.17	0.11	0.39	0.80	1.23
	60	1.06	2.72	3.86	4.09	0.29	0.72	1.08	1.17
Black	0	0.00	0.00	0.00	1.40	0.00	0.00	0.00	0.78
	10	0.00	0.00	0.01	1.44	0.00	0.00	0.01	0.79
	20	0.00	0.01	0.07	1.45	0.00	0.01	0.03	0.80
	30	0.01	0.07	0.26	1.49	0.00	0.02	0.08	0.80
	40	0.06	0.26	0.63	1.52	0.02	0.08	0.24	0.81
	50	0.22	0.61	1.16	1.57	0.06	0.23	0.50	0.82
	60	0.46	1.10	1.49	1.57	0.18	0.48	0.74	0.83
Asian/ Pacific Islander	0	0.00	0.00	0.00	2.22	0.00	0.00	0.00	0.67
	10	0.00	0.00	0.01	2.24	0.00	0.00	0.00	0.67
	20	0.00	0.01	0.04	2.25	0.00	0.00	0.01	0.67
	30	0.01	0.04	0.16	2.26	0.00	0.01	0.04	0.67
	40	0.03	0.16	0.51	2.27	0.01	0.04	0.13	0.67
	50	0.13	0.49	1.22	2.28	0.03	0.12	0.31	0.67
	60	0.38	1.14	1.95	2.25	0.09	0.29	0.56	0.66
Hispanic ^a	0	0.00	0.00	0.00	2.34	0.00	0.00	0.00	0.84
	10	0.00	0.00	0.01	2.37	0.00	0.00	0.01	0.85
	20	0.00	0.01	0.05	2.39	0.00	0.01	0.02	0.85
	30	0.01	0.05	0.20	2.41	0.01	0.02	0.08	0.85
	40	0.04	0.20	0.64	2.44	0.02	0.07	0.20	0.85
	50	0.16	0.61	1.41	2.47	0.06	0.19	0.41	0.85
	60	0.48	1.33	2.14	2.47	0.14	0.37	0.64	0.82

Race/Ethnicity	Lifetime Risk of Developing		Lifetime Risk of Dying	
	Males	Females	Males	Females
All Races	3.61%	1.14%	0.76%	0.32%
White	3.95%	1.22%	0.81%	0.32%
Black	1.40%	0.78%	0.39%	0.33%
Asian/Pacific Islander	2.22%	0.67%	0.52%	0.25%
Hispanic ^a	2.34%	0.84%	0.59%	0.26%

Devcan Version 6.1.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from Hawaii, Seattle, Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.