

Table I-14

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site and Race/Ethnicity

Both Sexes, 17 SEER Areas, 2002-2004

Site	All Races	Whites	Blacks
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	40.93 (40.83, 41.02)	41.39 (41.29, 41.50)	37.22 (36.93, 37.51)
Invasive and In Situ	43.24 (43.14, 43.34)	43.79 (43.68, 43.90)	38.51 (38.22, 38.81)
Oral Cavity and Pharynx	1.02 (1.00, 1.03)	1.03 (1.02, 1.05)	0.89 (0.85, 0.93)
Esophagus	0.50 (0.49, 0.51)	0.51 (0.50, 0.53)	0.52 (0.49, 0.56)
Stomach	0.90 (0.89, 0.91)	0.79 (0.78, 0.81)	1.12 (1.07, 1.17)
Colon and Rectum	5.42 (5.39, 5.46)	5.41 (5.37, 5.45)	5.24 (5.13, 5.35)
Invasive and In Situ	5.73 (5.70, 5.77)	5.71 (5.67, 5.75)	5.58 (5.47, 5.70)
Liver and Intrahepatic Bile Duct	0.66 (0.64, 0.67)	0.55 (0.54, 0.57)	0.68 (0.64, 0.71)
Pancreas	1.31 (1.29, 1.33)	1.30 (1.28, 1.32)	1.40 (1.34, 1.46)
Larynx	0.37 (0.36, 0.38)	0.37 (0.36, 0.38)	0.51 (0.48, 0.54)
Invasive and In Situ	0.40 (0.39, 0.41)	0.40 (0.39, 0.41)	0.53 (0.50, 0.56)
Lung and Bronchus	6.98 (6.94, 7.02)	7.17 (7.12, 7.21)	6.64 (6.52, 6.76)
Melanoma of the Skin	1.72 (1.70, 1.74)	1.99 (1.97, 2.01)	0.09 (0.08, 0.11)
Invasive and In Situ	2.76 (2.73, 2.78)	3.15 (3.12, 3.18)	0.11 (0.09, 0.13)
Breast	6.37 (6.34, 6.41)	6.59 (6.55, 6.63)	5.33 (5.23, 5.44)
Invasive and In Situ	7.62 (7.58, 7.66)	7.86 (7.82, 7.91)	6.34 (6.23, 6.45)
Urinary Bladder(Invasive and In Situ)	2.35 (2.33, 2.37)	2.55 (2.53, 2.58)	1.13 (1.08, 1.19)
Kidney and Renal Pelvis	1.34 (1.33, 1.36)	1.40 (1.38, 1.42)	1.20 (1.15, 1.25)
Brain and Other Nervous System	0.60 (0.59, 0.61)	0.66 (0.65, 0.67)	0.32 (0.30, 0.35)
Thyroid	0.73 (0.72, 0.74)	0.77 (0.76, 0.78)	0.40 (0.37, 0.42)
Hodgkin Lymphoma	0.22 (0.21, 0.23)	0.24 (0.23, 0.25)	0.18 (0.16, 0.19)
Non-Hodgkin Lymphoma	2.02 (2.00, 2.04)	2.12 (2.10, 2.15)	1.21 (1.16, 1.26)
Myeloma	0.61 (0.60, 0.62)	0.56 (0.55, 0.57)	1.02 (0.98, 1.07)
Leukemia	1.26 (1.25, 1.28)	1.31 (1.30, 1.33)	0.86 (0.82, 0.91)
Acute Lymphocytic Leukemia	0.12 (0.12, 0.12)	0.13 (0.12, 0.13)	0.06 (0.05, 0.07)
Chronic Lymphocytic Leukemia	0.45 (0.44, 0.46)	0.48 (0.47, 0.49)	0.27 (0.25, 0.30)
Acute Myeloid Leukemia	0.37 (0.36, 0.38)	0.38 (0.37, 0.39)	0.27 (0.24, 0.29)
Chronic Myeloid Leukemia	0.15 (0.14, 0.16)	0.15 (0.15, 0.16)	0.11 (0.10, 0.13)
Kaposi Sarcoma	0.05 (0.05, 0.06)	0.05 (0.05, 0.05)	0.08 (0.07, 0.09)
Mesothelioma	0.12 (0.12, 0.13)	0.14 (0.13, 0.14)	0.05 (0.04, 0.07)

Devcan Version 6.2.0, April 2007, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Note: Invasive cancer only unless specified otherwise.

Table I-14 - continued

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site and Race/Ethnicity

Both Sexes, 17 SEER Areas, 2002-2004

Site	Asian/Pacific Islanders	American Indian/ Alaska Natives ^a	Hispanics ^b
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	35.73 (35.31, 36.15)	26.34 (25.16, 27.63)	37.25 (36.88, 37.62)
Invasive and In Situ	37.19 (36.77, 37.61)	27.44 (26.24, 28.75)	38.57 (38.19, 38.95)
Oral Cavity and Pharynx	0.88 (0.82, 0.95)	0.61 (0.47, 0.91)	0.68 (0.64, 0.74)
Esophagus	0.36 (0.32, 0.41)	0.41 (0.28, 0.71)	0.41 (0.37, 0.47)
Stomach	2.06 (1.95, 2.17)	1.30 (1.04, 1.70)	1.62 (1.54, 1.72)
Colon and Rectum	5.58 (5.41, 5.76)	4.27 (3.80, 4.86)	4.82 (4.69, 4.97)
Invasive and In Situ	5.83 (5.66, 6.02)	4.45 (3.97, 5.05)	5.08 (4.94, 5.23)
Liver and Intrahepatic Bile Duct	1.78 (1.70, 1.88)	1.08 (0.81, 1.51)	1.11 (1.05, 1.17)
Pancreas	1.34 (1.26, 1.44)	0.96 (0.75, 1.32)	1.46 (1.38, 1.55)
Larynx	0.22 (0.18, 0.27)	0.14 (0.08, 0.41)	0.30 (0.27, 0.34)
Invasive and In Situ	0.22 (0.19, 0.27)	0.14 (0.08, 0.42)	0.31 (0.28, 0.35)
Lung and Bronchus	5.55 (5.39, 5.73)	4.66 (4.18, 5.27)	4.38 (4.25, 4.52)
Melanoma of the Skin	0.18 (0.15, 0.22)	0.31 (0.15, 0.65)	0.50 (0.46, 0.56)
Invasive and In Situ	0.25 (0.21, 0.29)	0.43 (0.26, 0.78)	0.72 (0.67, 0.77)
Breast	4.90 (4.78, 5.03)	3.31 (2.96, 3.79)	4.82 (4.70, 4.95)
Invasive and In Situ	6.11 (5.98, 6.26)	4.03 (3.64, 4.54)	5.66 (5.54, 5.80)
Urinary Bladder(Invasive and In Situ)	1.43 (1.34, 1.53)	0.69 (0.52, 1.02)	1.53 (1.45, 1.62)
Kidney and Renal Pelvis	0.83 (0.77, 0.90)	1.53 (1.28, 1.92)	1.42 (1.36, 1.49)
Brain and Other Nervous System	0.35 (0.32, 0.39)	0.22 (0.14, 0.49)	0.54 (0.50, 0.58)
Thyroid	0.79 (0.75, 0.85)	0.44 (0.34, 0.72)	0.69 (0.65, 0.73)
Hodgkin Lymphoma	0.11 (0.10, 0.14)	0.08 (0.04, 0.35)	0.22 (0.20, 0.25)
Non-Hodgkin Lymphoma	1.73 (1.64, 1.83)	0.91 (0.71, 1.26)	1.94 (1.86, 2.03)
Myeloma	0.43 (0.39, 0.49)	0.55 (0.40, 0.87)	0.70 (0.65, 0.76)
Leukemia	0.88 (0.82, 0.95)	0.69 (0.48, 1.07)	1.12 (1.05, 1.19)
Acute Lymphocytic Leukemia	0.12 (0.11, 0.15)	0.09 (0.06, 0.35)	0.17 (0.16, 0.19)
Chronic Lymphocytic Leukemia	0.13 (0.11, 0.17)	0.29 (0.13, 0.65)	0.28 (0.25, 0.33)
Acute Myeloid Leukemia	0.36 (0.32, 0.41)	0.13 (0.07, 0.41)	0.36 (0.32, 0.40)
Chronic Myeloid Leukemia	0.11 (0.10, 0.15)	0.07 (0.03, 0.35)	0.15 (0.13, 0.19)
Kaposi Sarcoma	0.02 (0.01, 0.04)	0.03 (0.01, 0.30)	0.10 (0.08, 0.12)
Mesothelioma	0.05 (0.03, 0.08)	0.08 (0.03, 0.36)	0.13 (0.11, 0.15)

Devcan Version 6.2.0, April 2007, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Underlying incidence data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.^b Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives.

Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky.

Note: Invasive cancer only unless specified otherwise.

Table I-15

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site and Race/Ethnicity

Males, 17 SEER Areas, 2002-2004

Site	All Races	Whites	Blacks
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	44.94 (44.79, 45.09)	44.89 (44.73, 45.06)	42.33 (41.87, 42.79)
Invasive and In Situ	46.26 (46.10, 46.41)	46.28 (46.12, 46.45)	42.65 (42.19, 43.12)
Oral Cavity and Pharynx	1.40 (1.38, 1.43)	1.43 (1.40, 1.46)	1.30 (1.23, 1.37)
Esophagus	0.78 (0.76, 0.80)	0.80 (0.78, 0.82)	0.76 (0.70, 0.82)
Stomach	1.13 (1.11, 1.15)	1.01 (0.99, 1.04)	1.30 (1.22, 1.38)
Colon and Rectum	5.65 (5.60, 5.70)	5.65 (5.60, 5.71)	5.20 (5.04, 5.36)
Invasive and In Situ	6.00 (5.95, 6.05)	6.00 (5.94, 6.06)	5.53 (5.37, 5.70)
Liver and Intrahepatic Bile Duct	0.91 (0.89, 0.93)	0.77 (0.75, 0.79)	0.98 (0.92, 1.05)
Pancreas	1.30 (1.27, 1.32)	1.30 (1.28, 1.33)	1.25 (1.17, 1.33)
Larynx	0.62 (0.60, 0.64)	0.61 (0.59, 0.63)	0.85 (0.79, 0.91)
Invasive and In Situ	0.66 (0.65, 0.68)	0.66 (0.64, 0.68)	0.89 (0.83, 0.96)
Lung and Bronchus	7.91 (7.85, 7.98)	7.98 (7.92, 8.05)	8.09 (7.90, 8.29)
Melanoma of the Skin	2.10 (2.07, 2.13)	2.42 (2.38, 2.45)	0.07 (0.06, 0.10)
Invasive and In Situ	3.33 (3.29, 3.37)	3.79 (3.75, 3.84)	0.09 (0.07, 0.12)
Breast	0.11 (0.10, 0.12)	0.11 (0.10, 0.12)	0.12 (0.10, 0.15)
Invasive and In Situ	0.12 (0.11, 0.13)	0.12 (0.11, 0.13)	0.14 (0.11, 0.17)
Prostate	16.72 (16.63, 16.81)	16.15 (16.06, 16.25)	19.43 (19.13, 19.74)
Testis	0.36 (0.35, 0.37)	0.43 (0.42, 0.44)	0.09 (0.07, 0.11)
Urinary Bladder (Invasive and In Situ)	3.70 (3.65, 3.74)	4.03 (3.99, 4.08)	1.48 (1.40, 1.57)
Kidney and Renal Pelvis	1.71 (1.68, 1.73)	1.78 (1.75, 1.81)	1.49 (1.42, 1.58)
Brain and Other Nervous System	0.67 (0.65, 0.68)	0.74 (0.72, 0.75)	0.35 (0.31, 0.39)
Thyroid	0.39 (0.37, 0.40)	0.41 (0.40, 0.43)	0.17 (0.15, 0.20)
Hodgkin Lymphoma	0.24 (0.23, 0.25)	0.26 (0.25, 0.27)	0.19 (0.16, 0.21)
Non-Hodgkin Lymphoma	2.19 (2.16, 2.22)	2.30 (2.26, 2.33)	1.31 (1.24, 1.39)
Myeloma	0.69 (0.67, 0.71)	0.65 (0.64, 0.67)	1.09 (1.02, 1.17)
Leukemia	1.50 (1.47, 1.52)	1.56 (1.54, 1.59)	0.98 (0.92, 1.05)
Acute Lymphocytic Leukemia	0.13 (0.13, 0.14)	0.14 (0.14, 0.15)	0.07 (0.06, 0.09)
Chronic Lymphocytic Leukemia	0.56 (0.54, 0.57)	0.59 (0.57, 0.61)	0.34 (0.30, 0.38)
Acute Myeloid Leukemia	0.42 (0.41, 0.44)	0.44 (0.43, 0.46)	0.28 (0.25, 0.32)
Chronic Myeloid Leukemia	0.18 (0.17, 0.19)	0.18 (0.17, 0.19)	0.13 (0.11, 0.16)
Kaposi Sarcoma	0.09 (0.09, 0.10)	0.08 (0.08, 0.09)	0.15 (0.13, 0.18)
Mesothelioma	0.21 (0.20, 0.22)	0.23 (0.22, 0.24)	0.08 (0.06, 0.11)

Devcan Version 6.2.0, April 2007, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Note: Invasive cancer only unless specified otherwise.

Table I-15 - continued

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site and Race/Ethnicity

Males, 17 SEER Areas, 2002-2004

Site	Asian/Pacific Islanders	American Indian/ Alaska Natives ^a	Hispanics ^b
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	39.83 (39.21, 40.48)	26.35 (24.64, 28.34)	41.57 (40.99, 42.18)
Invasive and In Situ	40.27 (39.64, 40.92)	26.83 (25.10, 28.84)	42.14 (41.55, 42.76)
Oral Cavity and Pharynx	1.12 (1.03, 1.23)	0.84 (0.59, 1.53)	0.97 (0.88, 1.08)
Esophagus	0.54 (0.47, 0.63)	0.61 (0.40, 1.29)	0.66 (0.58, 0.76)
Stomach	2.47 (2.31, 2.65)	1.65 (1.22, 2.47)	1.85 (1.73, 1.99)
Colon and Rectum	5.96 (5.72, 6.22)	3.74 (3.17, 4.66)	5.08 (4.88, 5.30)
Invasive and In Situ	6.29 (6.04, 6.55)	3.91 (3.32, 4.83)	5.38 (5.18, 5.61)
Liver and Intrahepatic Bile Duct	2.47 (2.33, 2.62)	1.46 (1.01, 2.32)	1.47 (1.38, 1.58)
Pancreas	1.33 (1.21, 1.46)	0.93 (0.66, 1.64)	1.26 (1.16, 1.38)
Larynx	0.41 (0.34, 0.50)	0.26 (0.14, 0.91)	0.56 (0.50, 0.64)
Invasive and In Situ	0.42 (0.35, 0.51)	0.28 (0.15, 0.93)	0.58 (0.52, 0.66)
Lung and Bronchus	7.06 (6.80, 7.34)	5.01 (4.25, 6.10)	5.32 (5.11, 5.56)
Melanoma of the Skin	0.19 (0.15, 0.26)	0.34 (0.18, 1.01)	0.49 (0.43, 0.58)
Invasive and In Situ	0.25 (0.21, 0.32)	0.47 (0.28, 1.14)	0.69 (0.61, 0.79)
Breast	0.06 (0.04, 0.10)	0.02 (0.00, 0.70)	0.06 (0.04, 0.11)
Invasive and In Situ	0.06 (0.05, 0.11)	0.02 (0.00, 0.70)	0.07 (0.05, 0.12)
Prostate	12.52 (12.18, 12.88)	6.99 (6.14, 8.15)	16.25 (15.90, 16.62)
Testis	0.11 (0.09, 0.15)	0.28 (0.18, 0.91)	0.27 (0.25, 0.31)
Urinary Bladder (Invasive and In Situ)	2.29 (2.13, 2.48)	1.07 (0.77, 1.80)	2.37 (2.22, 2.55)
Kidney and Renal Pelvis	1.11 (1.01, 1.22)	1.78 (1.41, 2.54)	1.77 (1.67, 1.89)
Brain and Other Nervous System	0.37 (0.32, 0.44)	0.30 (0.17, 0.95)	0.60 (0.54, 0.69)
Thyroid	0.41 (0.36, 0.48)	0.26 (0.16, 0.91)	0.31 (0.27, 0.37)
Hodgkin Lymphoma	0.13 (0.10, 0.18)	0.09 (0.05, 0.75)	0.27 (0.23, 0.34)
Non-Hodgkin Lymphoma	1.92 (1.79, 2.08)	0.85 (0.57, 1.57)	2.06 (1.93, 2.20)
Myeloma	0.47 (0.41, 0.56)	0.41 (0.22, 1.09)	0.78 (0.70, 0.87)
Leukemia	1.03 (0.94, 1.14)	0.49 (0.33, 1.14)	1.29 (1.18, 1.42)
Acute Lymphocytic Leukemia	0.14 (0.11, 0.19)	0.11 (0.06, 0.77)	0.17 (0.15, 0.21)
Chronic Lymphocytic Leukemia	0.16 (0.12, 0.22)	0.16 (0.06, 0.83)	0.37 (0.30, 0.46)
Acute Myeloid Leukemia	0.41 (0.36, 0.49)	0.11 (0.05, 0.77)	0.40 (0.34, 0.48)
Chronic Myeloid Leukemia	0.14 (0.11, 0.20)	0.05 (0.01, 0.72)	0.20 (0.14, 0.27)
Kaposi Sarcoma	0.04 (0.03, 0.08)	0.05 (0.02, 0.72)	0.16 (0.13, 0.21)
Mesothelioma	0.08 (0.05, 0.14)	0.06 (0.01, 0.73)	0.20 (0.17, 0.26)

Devcan Version 6.2.0, April 2007, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Underlying incidence data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.^b Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives.

Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky.

Note: Invasive cancer only unless specified otherwise.

Table I-16

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site and Race/Ethnicity

Females, 17 SEER Areas, 2002-2004

Site	All Races	Whites	Blacks
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	37.52 (37.40, 37.65)	38.46 (38.32, 38.60)	32.66 (32.29, 33.03)
Invasive and In Situ	40.82 (40.69, 40.95)	41.86 (41.72, 42.01)	34.86 (34.48, 35.24)
Oral Cavity and Pharynx	0.65 (0.64, 0.67)	0.66 (0.64, 0.68)	0.52 (0.48, 0.57)
Esophagus	0.25 (0.24, 0.26)	0.25 (0.24, 0.26)	0.31 (0.28, 0.35)
Stomach	0.69 (0.68, 0.71)	0.59 (0.58, 0.61)	0.96 (0.90, 1.03)
Colon and Rectum	5.23 (5.18, 5.27)	5.19 (5.14, 5.24)	5.31 (5.16, 5.46)
Invasive and In Situ	5.50 (5.45, 5.55)	5.45 (5.40, 5.51)	5.66 (5.50, 5.81)
Liver and Intrahepatic Bile Duct	0.41 (0.40, 0.43)	0.35 (0.34, 0.36)	0.40 (0.36, 0.44)
Pancreas	1.32 (1.29, 1.34)	1.29 (1.27, 1.32)	1.53 (1.45, 1.62)
Larynx	0.14 (0.14, 0.15)	0.15 (0.14, 0.16)	0.20 (0.17, 0.23)
Invasive and In Situ	0.16 (0.15, 0.16)	0.16 (0.15, 0.17)	0.20 (0.18, 0.23)
Lung and Bronchus	6.18 (6.13, 6.23)	6.49 (6.43, 6.54)	5.35 (5.20, 5.50)
Melanoma of the Skin	1.40 (1.38, 1.42)	1.63 (1.60, 1.66)	0.10 (0.08, 0.13)
Invasive and In Situ	2.26 (2.23, 2.29)	2.60 (2.57, 2.64)	0.13 (0.10, 0.15)
Breast	12.28 (12.22, 12.35)	12.80 (12.72, 12.87)	10.10 (9.91, 10.29)
Invasive and In Situ	14.72 (14.65, 14.80)	15.30 (15.22, 15.39)	12.02 (11.82, 12.24)
Cervix Uteri	0.70 (0.69, 0.72)	0.67 (0.66, 0.69)	0.92 (0.86, 0.97)
Corpus and Uterus, NOS	2.45 (2.42, 2.48)	2.56 (2.53, 2.60)	1.95 (1.86, 2.03)
Invasive and In Situ	2.49 (2.46, 2.52)	2.60 (2.56, 2.63)	1.97 (1.88, 2.06)
Ovary ^a	1.42 (1.39, 1.44)	1.50 (1.47, 1.52)	0.99 (0.92, 1.05)
Urinary Bladder (Invasive and In Situ)	1.17 (1.15, 1.19)	1.25 (1.22, 1.27)	0.82 (0.76, 0.89)
Kidney and Renal Pelvis	1.01 (0.99, 1.03)	1.05 (1.03, 1.08)	0.93 (0.88, 1.00)
Brain and Other Nervous System	0.53 (0.52, 0.55)	0.59 (0.57, 0.60)	0.30 (0.27, 0.34)
Thyroid	1.08 (1.07, 1.10)	1.14 (1.12, 1.17)	0.61 (0.57, 0.66)
Hodgkin Lymphoma	0.20 (0.20, 0.21)	0.22 (0.21, 0.23)	0.17 (0.15, 0.19)
Non-Hodgkin Lymphoma	1.87 (1.85, 1.90)	1.97 (1.94, 2.00)	1.11 (1.05, 1.18)
Myeloma	0.53 (0.52, 0.55)	0.48 (0.47, 0.50)	0.97 (0.90, 1.03)
Leukemia	1.06 (1.04, 1.08)	1.09 (1.07, 1.12)	0.76 (0.70, 0.82)
Acute Lymphocytic Leukemia	0.11 (0.10, 0.11)	0.11 (0.11, 0.12)	0.06 (0.05, 0.07)
Chronic Lymphocytic Leukemia	0.36 (0.35, 0.37)	0.38 (0.36, 0.39)	0.22 (0.19, 0.25)
Acute Myeloid Leukemia	0.32 (0.31, 0.33)	0.33 (0.32, 0.34)	0.26 (0.23, 0.29)
Chronic Myeloid Leukemia	0.12 (0.12, 0.13)	0.13 (0.12, 0.13)	0.10 (0.08, 0.12)
Kaposi Sarcoma	0.01 (0.01, 0.02)	0.01 (0.01, 0.02)	0.01 (0.01, 0.02)
Mesothelioma	0.05 (0.05, 0.05)	0.05 (0.05, 0.06)	0.03 (0.02, 0.04)

Devcan Version 6.2.0, April 2007, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Note: Invasive cancer only unless specified otherwise.

Table I-16 - continued

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site and Race/Ethnicity

Females, 17 SEER Areas, 2002-2004

Site	Asian/Pacific Islanders	American Indian/ Alaska Natives ^a	Hispanics ^b
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	32.32 (31.78, 32.88)	26.45 (24.82, 28.29)	33.95 (33.47, 34.43)
Invasive and In Situ	34.69 (34.14, 35.27)	28.14 (26.47, 30.02)	35.99 (35.50, 36.49)
Oral Cavity and Pharynx	0.66 (0.59, 0.76)	0.39 (0.24, 0.89)	0.43 (0.38, 0.50)
Esophagus	0.20 (0.16, 0.28)	0.22 (0.10, 0.71)	0.21 (0.17, 0.27)
Stomach	1.69 (1.55, 1.85)	0.99 (0.69, 1.59)	1.43 (1.32, 1.56)
Colon and Rectum	5.27 (5.03, 5.54)	4.74 (4.03, 5.68)	4.60 (4.41, 4.80)
Invasive and In Situ	5.46 (5.21, 5.73)	4.93 (4.22, 5.88)	4.82 (4.63, 5.03)
Liver and Intrahepatic Bile Duct	1.19 (1.09, 1.32)	0.73 (0.43, 1.37)	0.77 (0.71, 0.85)
Pancreas	1.35 (1.23, 1.50)	0.97 (0.67, 1.57)	1.63 (1.51, 1.76)
Larynx	0.05 (0.03, 0.10)	0.02 (0.00, 0.52)	0.08 (0.06, 0.12)
Invasive and In Situ	0.05 (0.03, 0.10)	0.02 (0.00, 0.52)	0.09 (0.06, 0.12)
Lung and Bronchus	4.26 (4.05, 4.48)	4.37 (3.75, 5.23)	3.61 (3.45, 3.79)
Melanoma of the Skin	0.17 (0.13, 0.23)	0.27 (0.06, 0.88)	0.53 (0.48, 0.61)
Invasive and In Situ	0.24 (0.19, 0.32)	0.39 (0.15, 0.99)	0.76 (0.70, 0.84)
Breast	9.18 (8.95, 9.43)	6.49 (5.81, 7.40)	9.23 (9.01, 9.46)
Invasive and In Situ	11.45 (11.20, 11.72)	7.89 (7.14, 8.87)	10.86 (10.63, 11.10)
Cervix Uteri	0.76 (0.69, 0.84)	0.57 (0.39, 1.08)	1.19 (1.13, 1.26)
Corpus and Uterus, NOS	1.77 (1.67, 1.88)	1.46 (1.14, 2.05)	1.97 (1.88, 2.08)
Invasive and In Situ	1.79 (1.69, 1.91)	1.47 (1.16, 2.07)	1.99 (1.90, 2.10)
Ovary ^c	1.13 (1.04, 1.23)	0.94 (0.69, 1.49)	1.34 (1.25, 1.43)
Urinary Bladder (Invasive and In Situ)	0.66 (0.58, 0.76)	0.34 (0.19, 0.84)	0.83 (0.75, 0.94)
Kidney and Renal Pelvis	0.58 (0.52, 0.67)	1.28 (0.97, 1.88)	1.12 (1.05, 1.21)
Brain and Other Nervous System	0.34 (0.29, 0.40)	0.14 (0.07, 0.62)	0.49 (0.45, 0.55)
Thyroid	1.14 (1.08, 1.23)	0.62 (0.46, 1.11)	1.07 (1.01, 1.15)
Hodgkin Lymphoma	0.09 (0.08, 0.14)	0.06 (0.01, 0.55)	0.17 (0.15, 0.21)
Non-Hodgkin Lymphoma	1.56 (1.44, 1.70)	0.98 (0.70, 1.55)	1.84 (1.74, 1.97)
Myeloma	0.40 (0.34, 0.48)	0.69 (0.46, 1.24)	0.64 (0.58, 0.72)
Leukemia	0.75 (0.67, 0.85)	0.85 (0.50, 1.52)	0.98 (0.90, 1.08)
Acute Lymphocytic Leukemia	0.11 (0.09, 0.16)	0.07 (0.03, 0.55)	0.17 (0.15, 0.21)
Chronic Lymphocytic Leukemia	0.11 (0.08, 0.17)	0.39 (0.13, 1.03)	0.21 (0.18, 0.27)
Acute Myeloid Leukemia	0.32 (0.26, 0.40)	0.15 (0.05, 0.65)	0.33 (0.28, 0.39)
Chronic Myeloid Leukemia	0.09 (0.07, 0.14)	0.10 (0.04, 0.59)	0.12 (0.09, 0.17)
Kaposi Sarcoma	0.00 (0.00, 0.05)	0.00 (0.00, 0.50)	0.04 (0.02, 0.08)
Mesothelioma	0.02 (0.01, 0.07)	0.10 (0.02, 0.61)	0.06 (0.05, 0.10)

Devcan Version 6.2.0, April 2007, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^a Underlying incidence data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.^b Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives.^c Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky.

Note: Invasive cancer only unless specified otherwise.

Table I-17

Lifetime Risk (Percent) of Dying from Cancer by Site and Race/Ethnicity

Both Sexes, Total U.S., 2002-2004

Site	All Races	Whites	Blacks
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	21.29 (21.26, 21.32)	21.45 (21.42, 21.49)	21.31 (21.22, 21.40)
Oral Cavity and Pharynx	0.28 (0.28, 0.29)	0.28 (0.27, 0.28)	0.33 (0.32, 0.34)
Esophagus	0.48 (0.47, 0.48)	0.48 (0.47, 0.49)	0.51 (0.49, 0.52)
Stomach	0.46 (0.46, 0.47)	0.41 (0.41, 0.42)	0.74 (0.72, 0.76)
Colon and Rectum	2.20 (2.19, 2.21)	2.18 (2.17, 2.19)	2.42 (2.39, 2.46)
Liver and Intrahepatic Bile Duct	0.54 (0.53, 0.54)	0.50 (0.50, 0.51)	0.58 (0.57, 0.60)
Pancreas	1.20 (1.19, 1.21)	1.20 (1.19, 1.20)	1.26 (1.24, 1.29)
Larynx	0.14 (0.13, 0.14)	0.13 (0.13, 0.13)	0.21 (0.20, 0.22)
Lung and Bronchus	5.99 (5.97, 6.00)	6.13 (6.11, 6.15)	5.50 (5.45, 5.55)
Melanoma of the Skin	0.28 (0.27, 0.28)	0.31 (0.31, 0.32)	0.04 (0.03, 0.04)
Breast	1.53 (1.52, 1.54)	1.52 (1.51, 1.53)	1.73 (1.70, 1.76)
Urinary Bladder	0.53 (0.52, 0.53)	0.56 (0.55, 0.56)	0.36 (0.35, 0.38)
Kidney and Renal Pelvis	0.46 (0.46, 0.47)	0.48 (0.47, 0.49)	0.36 (0.35, 0.38)
Brain and Other Nervous System	0.43 (0.43, 0.44)	0.47 (0.46, 0.47)	0.22 (0.21, 0.23)
Thyroid	0.05 (0.05, 0.05)	0.05 (0.05, 0.06)	0.04 (0.04, 0.04)
Hodgkin Lymphoma	0.04 (0.04, 0.05)	0.05 (0.04, 0.05)	0.03 (0.03, 0.03)
Non-Hodgkin Lymphoma	0.83 (0.83, 0.84)	0.88 (0.88, 0.89)	0.45 (0.44, 0.47)
Myeloma	0.42 (0.42, 0.43)	0.40 (0.40, 0.41)	0.64 (0.62, 0.66)
Leukemia	0.83 (0.83, 0.84)	0.87 (0.86, 0.88)	0.60 (0.58, 0.62)
Acute Lymphocytic Leukemia	0.04 (0.04, 0.04)	0.04 (0.04, 0.05)	0.03 (0.02, 0.03)
Chronic Lymphocytic Leukemia	0.19 (0.18, 0.19)	0.20 (0.19, 0.20)	0.14 (0.13, 0.15)
Acute Myeloid Leukemia	0.30 (0.30, 0.30)	0.31 (0.31, 0.32)	0.19 (0.18, 0.20)
Chronic Myeloid Leukemia	0.05 (0.05, 0.05)	0.05 (0.05, 0.05)	0.04 (0.04, 0.05)

Devcan Version 6.2.0, April 2007, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: NCHS public use data file for the total US.

Table I-17 - continued

Lifetime Risk (Percent) of Dying from Cancer by Site and Race/Ethnicity

Both Sexes, Total U.S., 2002-2004

Site	Asian/Pacific Islanders	American Indian/ Alaska Natives ^a	Hispanics ^b
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	18.32 (18.04, 18.61)	16.87 (16.34, 17.43)	17.39 (17.23, 17.55)
Oral Cavity and Pharynx	0.32 (0.28, 0.37)	0.24 (0.18, 0.34)	0.22 (0.20, 0.24)
Esophagus	0.30 (0.26, 0.35)	0.40 (0.32, 0.52)	0.31 (0.29, 0.33)
Stomach	1.32 (1.24, 1.42)	0.73 (0.62, 0.89)	0.91 (0.87, 0.95)
Colon and Rectum	2.17 (2.05, 2.29)	1.86 (1.66, 2.09)	1.92 (1.86, 1.98)
Liver and Intrahepatic Bile Duct	1.54 (1.46, 1.63)	0.88 (0.75, 1.05)	0.99 (0.95, 1.03)
Pancreas	1.21 (1.14, 1.29)	0.80 (0.68, 0.96)	1.13 (1.09, 1.18)
Larynx	0.07 (0.05, 0.10)	0.09 (0.06, 0.17)	0.12 (0.11, 0.14)
Lung and Bronchus	4.12 (3.99, 4.25)	4.13 (3.88, 4.42)	3.17 (3.10, 3.24)
Melanoma of the Skin	0.06 (0.04, 0.09)	0.13 (0.08, 0.24)	0.10 (0.08, 0.11)
Breast	0.94 (0.87, 1.00)	1.01 (0.88, 1.19)	1.11 (1.07, 1.16)
Urinary Bladder	0.36 (0.31, 0.42)	0.23 (0.15, 0.35)	0.39 (0.36, 0.42)
Kidney and Renal Pelvis	0.26 (0.22, 0.30)	0.68 (0.58, 0.82)	0.48 (0.45, 0.51)
Brain and Other Nervous System	0.25 (0.22, 0.29)	0.18 (0.14, 0.27)	0.32 (0.30, 0.34)
Thyroid	0.09 (0.07, 0.12)	0.06 (0.03, 0.14)	0.07 (0.06, 0.08)
Hodgkin Lymphoma	0.04 (0.03, 0.06)	0.04 (0.02, 0.12)	0.05 (0.04, 0.06)
Non-Hodgkin Lymphoma	0.79 (0.72, 0.86)	0.52 (0.41, 0.66)	0.74 (0.71, 0.78)
Myeloma	0.25 (0.22, 0.29)	0.41 (0.32, 0.53)	0.40 (0.38, 0.43)
Leukemia	0.56 (0.51, 0.62)	0.49 (0.40, 0.62)	0.63 (0.60, 0.66)
Acute Lymphocytic Leukemia	0.04 (0.03, 0.06)	0.05 (0.03, 0.13)	0.06 (0.06, 0.07)
Chronic Lymphocytic Leukemia	0.05 (0.03, 0.08)	0.07 (0.04, 0.16)	0.09 (0.07, 0.10)
Acute Myeloid Leukemia	0.24 (0.21, 0.28)	0.18 (0.13, 0.27)	0.21 (0.20, 0.23)
Chronic Myeloid Leukemia	0.04 (0.02, 0.06)	0.04 (0.02, 0.11)	0.04 (0.04, 0.05)

Devcan Version 6.2.0, April 2007, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: NCHS public use data file for the total US.

^a Underlying mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.^b Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table I-18

Lifetime Risk (Percent) of Dying from Cancer by Site and Race/Ethnicity

Males, Total U.S., 2002-2004

Site	All Races	Whites	Blacks
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	23.31 (23.27, 23.36)	23.41 (23.36, 23.46)	23.70 (23.56, 23.84)
Oral Cavity and Pharynx	0.39 (0.38, 0.39)	0.37 (0.36, 0.38)	0.51 (0.49, 0.53)
Esophagus	0.75 (0.74, 0.76)	0.77 (0.76, 0.78)	0.74 (0.72, 0.77)
Stomach	0.56 (0.56, 0.57)	0.51 (0.50, 0.52)	0.87 (0.84, 0.90)
Colon and Rectum	2.29 (2.27, 2.30)	2.28 (2.26, 2.29)	2.41 (2.36, 2.47)
Liver and Intrahepatic Bile Duct	0.69 (0.69, 0.70)	0.65 (0.64, 0.66)	0.78 (0.75, 0.80)
Pancreas	1.20 (1.19, 1.21)	1.21 (1.20, 1.22)	1.18 (1.15, 1.21)
Larynx	0.23 (0.22, 0.23)	0.21 (0.21, 0.22)	0.37 (0.35, 0.39)
Lung and Bronchus	7.12 (7.09, 7.14)	7.20 (7.17, 7.23)	7.09 (7.01, 7.17)
Melanoma of the Skin	0.37 (0.36, 0.37)	0.42 (0.41, 0.42)	0.04 (0.03, 0.04)
Breast	0.03 (0.03, 0.03)	0.03 (0.03, 0.03)	0.04 (0.03, 0.05)
Prostate	2.87 (2.85, 2.89)	2.70 (2.68, 2.72)	4.50 (4.43, 4.58)
Testis	0.02 (0.02, 0.02)	0.02 (0.02, 0.02)	0.01 (0.01, 0.02)
Urinary Bladder	0.78 (0.77, 0.79)	0.83 (0.82, 0.84)	0.40 (0.38, 0.42)
Kidney and Renal Pelvis	0.59 (0.58, 0.60)	0.62 (0.61, 0.63)	0.45 (0.43, 0.47)
Brain and Other Nervous System	0.48 (0.47, 0.49)	0.52 (0.51, 0.53)	0.23 (0.22, 0.25)
Thyroid	0.04 (0.04, 0.05)	0.05 (0.04, 0.05)	0.03 (0.02, 0.03)
Hodgkin Lymphoma	0.05 (0.05, 0.05)	0.05 (0.05, 0.05)	0.04 (0.03, 0.04)
Non-Hodgkin Lymphoma	0.91 (0.90, 0.92)	0.97 (0.96, 0.98)	0.47 (0.45, 0.49)
Myeloma	0.46 (0.45, 0.47)	0.44 (0.44, 0.45)	0.63 (0.60, 0.65)
Leukemia	0.98 (0.97, 0.99)	1.03 (1.01, 1.04)	0.65 (0.62, 0.68)
Acute Lymphocytic Leukemia	0.05 (0.05, 0.05)	0.05 (0.05, 0.05)	0.03 (0.03, 0.04)
Chronic Lymphocytic Leukemia	0.23 (0.22, 0.24)	0.24 (0.24, 0.25)	0.16 (0.15, 0.18)
Acute Myeloid Leukemia	0.35 (0.34, 0.35)	0.37 (0.36, 0.37)	0.20 (0.19, 0.22)
Chronic Myeloid Leukemia	0.05 (0.05, 0.06)	0.05 (0.05, 0.06)	0.05 (0.04, 0.06)

Devcan Version 6.2.0, April 2007, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: NCHS public use data file for the total US.

Table I-18 - continued

Lifetime Risk (Percent) of Dying from Cancer by Site and Race/Ethnicity

Site	Males, Total U.S., 2002-2004		
	Asian/Pacific Islanders Percent (95% C.I.)	American Indian/ Alaska Natives ^a Percent (95% C.I.)	Hispanics ^b Percent (95% C.I.)
All Sites	20.73 (20.32, 21.16)	17.20 (16.42, 18.05)	19.80 (19.55, 20.06)
Oral Cavity and Pharynx	0.43 (0.37, 0.51)	0.32 (0.22, 0.53)	0.32 (0.29, 0.36)
Esophagus	0.47 (0.40, 0.55)	0.62 (0.48, 0.86)	0.48 (0.44, 0.52)
Stomach	1.54 (1.42, 1.69)	0.85 (0.68, 1.10)	1.07 (1.01, 1.14)
Colon and Rectum	2.30 (2.14, 2.48)	1.94 (1.66, 2.31)	2.10 (2.01, 2.20)
Liver and Intrahepatic Bile Duct	1.97 (1.86, 2.10)	0.86 (0.71, 1.12)	1.24 (1.18, 1.31)
Pancreas	1.14 (1.05, 1.25)	0.79 (0.60, 1.07)	1.06 (1.00, 1.12)
Larynx	0.13 (0.09, 0.19)	0.14 (0.08, 0.33)	0.24 (0.21, 0.27)
Lung and Bronchus	5.34 (5.14, 5.55)	4.58 (4.18, 5.06)	4.29 (4.17, 4.41)
Melanoma of the Skin	0.06 (0.04, 0.11)	0.10 (0.05, 0.28)	0.10 (0.08, 0.13)
Breast	0.01 (0.00, 0.05)	0.01 (0.00, 0.20)	0.02 (0.01, 0.04)
Prostate	2.29 (2.10, 2.50)	2.18 (1.82, 2.62)	3.19 (3.05, 3.33)
Testis	0.01 (0.00, 0.05)	0.01 (0.00, 0.19)	0.02 (0.02, 0.04)
Urinary Bladder	0.53 (0.45, 0.64)	0.30 (0.17, 0.54)	0.56 (0.51, 0.62)
Kidney and Renal Pelvis	0.34 (0.28, 0.43)	0.84 (0.69, 1.08)	0.63 (0.59, 0.69)
Brain and Other Nervous System	0.31 (0.26, 0.37)	0.22 (0.16, 0.41)	0.34 (0.31, 0.38)
Thyroid	0.06 (0.04, 0.11)	0.02 (0.00, 0.20)	0.06 (0.05, 0.08)
Hodgkin Lymphoma	0.04 (0.02, 0.08)	0.02 (0.00, 0.20)	0.05 (0.04, 0.07)
Non-Hodgkin Lymphoma	0.86 (0.77, 0.97)	0.44 (0.34, 0.65)	0.76 (0.71, 0.82)
Myeloma	0.26 (0.22, 0.32)	0.45 (0.30, 0.70)	0.42 (0.39, 0.46)
Leukemia	0.66 (0.59, 0.75)	0.49 (0.37, 0.72)	0.70 (0.66, 0.76)
Acute Lymphocytic Leukemia	0.04 (0.03, 0.08)	0.06 (0.03, 0.23)	0.06 (0.06, 0.08)
Chronic Lymphocytic Leukemia	0.08 (0.05, 0.14)	0.07 (0.03, 0.26)	0.11 (0.08, 0.14)
Acute Myeloid Leukemia	0.28 (0.24, 0.33)	0.17 (0.10, 0.36)	0.26 (0.23, 0.29)
Chronic Myeloid Leukemia	0.03 (0.02, 0.07)	0.04 (0.01, 0.22)	0.04 (0.03, 0.06)

Devcan Version 6.2.0, April 2007, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: NCHS public use data file for the total US.

^a Underlying mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.^b Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table I-19

Lifetime Risk (Percent) of Dying from Cancer by Site and Race/Ethnicity

Females, Total U.S., 2002-2004

Site	All Races	Whites	Blacks
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	19.74 (19.70, 19.78)	19.96 (19.91, 20.00)	19.39 (19.27, 19.51)
Oral Cavity and Pharynx	0.19 (0.18, 0.19)	0.19 (0.18, 0.19)	0.16 (0.15, 0.18)
Esophagus	0.22 (0.22, 0.23)	0.21 (0.21, 0.22)	0.30 (0.28, 0.31)
Stomach	0.38 (0.37, 0.38)	0.33 (0.33, 0.34)	0.62 (0.60, 0.65)
Colon and Rectum	2.13 (2.12, 2.14)	2.10 (2.09, 2.12)	2.44 (2.39, 2.49)
Liver and Intrahepatic Bile Duct	0.39 (0.39, 0.40)	0.37 (0.36, 0.38)	0.41 (0.39, 0.43)
Pancreas	1.19 (1.18, 1.20)	1.18 (1.17, 1.19)	1.33 (1.30, 1.37)
Larynx	0.06 (0.05, 0.06)	0.05 (0.05, 0.06)	0.07 (0.07, 0.08)
Lung and Bronchus	5.00 (4.98, 5.02)	5.20 (5.17, 5.22)	4.12 (4.06, 4.18)
Melanoma of the Skin	0.20 (0.19, 0.20)	0.22 (0.22, 0.23)	0.04 (0.03, 0.05)
Breast	2.89 (2.88, 2.91)	2.89 (2.87, 2.90)	3.23 (3.18, 3.28)
Cervix Uteri	0.24 (0.24, 0.24)	0.21 (0.21, 0.22)	0.43 (0.41, 0.45)
Corpus and Uterus, NOS	0.51 (0.51, 0.52)	0.49 (0.48, 0.50)	0.75 (0.73, 0.78)
Ovary	1.05 (1.04, 1.06)	1.10 (1.09, 1.11)	0.77 (0.74, 0.80)
Urinary Bladder	0.32 (0.32, 0.33)	0.33 (0.32, 0.33)	0.33 (0.32, 0.35)
Kidney and Renal Pelvis	0.35 (0.34, 0.35)	0.36 (0.35, 0.36)	0.29 (0.28, 0.31)
Brain and Other Nervous System	0.39 (0.38, 0.39)	0.42 (0.41, 0.43)	0.20 (0.19, 0.21)
Thyroid	0.06 (0.06, 0.06)	0.06 (0.06, 0.06)	0.05 (0.05, 0.06)
Hodgkin Lymphoma	0.04 (0.04, 0.04)	0.04 (0.04, 0.04)	0.03 (0.02, 0.03)
Non-Hodgkin Lymphoma	0.77 (0.76, 0.78)	0.81 (0.80, 0.82)	0.44 (0.42, 0.46)
Myeloma	0.39 (0.39, 0.40)	0.37 (0.36, 0.37)	0.66 (0.63, 0.68)
Leukemia	0.71 (0.70, 0.72)	0.74 (0.73, 0.74)	0.56 (0.54, 0.58)
Acute Lymphocytic Leukemia	0.04 (0.03, 0.04)	0.04 (0.04, 0.04)	0.02 (0.02, 0.03)
Chronic Lymphocytic Leukemia	0.15 (0.15, 0.16)	0.16 (0.15, 0.16)	0.12 (0.11, 0.13)
Acute Myeloid Leukemia	0.26 (0.25, 0.26)	0.27 (0.26, 0.27)	0.19 (0.17, 0.20)
Chronic Myeloid Leukemia	0.04 (0.04, 0.04)	0.04 (0.04, 0.04)	0.04 (0.03, 0.05)

Devcan Version 6.2.0, April 2007, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: NCHS public use data file for the total US.

Table I-19 - continued

Lifetime Risk (Percent) of Dying from Cancer by Site and Race/Ethnicity

Females, Total U.S., 2002-2004

Site	Asian/Pacific Islanders	American Indian/ Alaska Natives ^a	Hispanics ^b
	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Sites	16.26 (15.88, 16.65)	16.71 (15.98, 17.50)	15.62 (15.42, 15.82)
Oral Cavity and Pharynx	0.23 (0.18, 0.30)	0.18 (0.12, 0.33)	0.13 (0.11, 0.15)
Esophagus	0.16 (0.12, 0.22)	0.20 (0.11, 0.37)	0.16 (0.14, 0.19)
Stomach	1.13 (1.02, 1.27)	0.63 (0.48, 0.86)	0.78 (0.73, 0.83)
Colon and Rectum	2.06 (1.89, 2.24)	1.80 (1.53, 2.13)	1.77 (1.70, 1.85)
Liver and Intrahepatic Bile Duct	1.17 (1.06, 1.30)	0.89 (0.70, 1.16)	0.77 (0.72, 0.82)
Pancreas	1.27 (1.16, 1.40)	0.83 (0.66, 1.06)	1.19 (1.13, 1.26)
Larynx	0.02 (0.01, 0.06)	0.04 (0.01, 0.17)	0.03 (0.02, 0.04)
Lung and Bronchus	3.10 (2.94, 3.27)	3.75 (3.42, 4.14)	2.25 (2.17, 2.33)
Melanoma of the Skin	0.06 (0.03, 0.12)	0.16 (0.07, 0.35)	0.09 (0.07, 0.11)
Breast	1.74 (1.63, 1.87)	1.94 (1.69, 2.26)	2.08 (2.01, 2.16)
Cervix Uteri	0.32 (0.28, 0.39)	0.38 (0.29, 0.56)	0.35 (0.33, 0.39)
Corpus and Uterus, NOS	0.34 (0.30, 0.41)	0.29 (0.21, 0.45)	0.45 (0.41, 0.48)
Ovary	0.68 (0.62, 0.76)	0.77 (0.60, 1.00)	0.82 (0.77, 0.86)
Urinary Bladder	0.21 (0.17, 0.28)	0.17 (0.09, 0.35)	0.26 (0.23, 0.30)
Kidney and Renal Pelvis	0.18 (0.15, 0.24)	0.52 (0.39, 0.73)	0.35 (0.32, 0.38)
Brain and Other Nervous System	0.20 (0.17, 0.26)	0.14 (0.09, 0.28)	0.29 (0.27, 0.32)
Thyroid	0.11 (0.08, 0.16)	0.09 (0.04, 0.24)	0.08 (0.07, 0.10)
Hodgkin Lymphoma	0.04 (0.02, 0.08)	0.06 (0.02, 0.20)	0.05 (0.04, 0.06)
Non-Hodgkin Lymphoma	0.73 (0.64, 0.83)	0.57 (0.41, 0.81)	0.72 (0.68, 0.77)
Myeloma	0.24 (0.20, 0.30)	0.39 (0.29, 0.57)	0.39 (0.36, 0.43)
Leukemia	0.47 (0.41, 0.56)	0.49 (0.37, 0.69)	0.57 (0.53, 0.61)
Acute Lymphocytic Leukemia	0.03 (0.02, 0.07)	0.04 (0.02, 0.18)	0.06 (0.05, 0.07)
Chronic Lymphocytic Leukemia	0.03 (0.02, 0.07)	0.07 (0.03, 0.22)	0.07 (0.06, 0.09)
Acute Myeloid Leukemia	0.22 (0.17, 0.29)	0.18 (0.12, 0.33)	0.18 (0.16, 0.21)
Chronic Myeloid Leukemia	0.04 (0.02, 0.09)	0.03 (0.01, 0.17)	0.04 (0.03, 0.06)

Devcan Version 6.2.0, April 2007, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: NCHS public use data file for the total US.

^a Underlying mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.^b Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

Table II-9
ALL CANCER SITES (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.16 (0.16, 0.17)	0.33 (0.33, 0.34)	0.75 (0.74, 0.76)	40.93 (40.83, 41.02)	21.29 (21.26, 21.32)	
	10	0.17 (0.17, 0.18)	0.60 (0.59, 0.60)	1.58 (1.57, 1.60)	41.33 (41.24, 41.43)	21.57 (21.54, 21.60)	
	20	0.43 (0.42, 0.43)	1.42 (1.41, 1.43)	3.93 (3.91, 3.95)	41.39 (41.29, 41.49)	21.63 (21.60, 21.66)	
	30	1.01 (1.00, 1.01)	3.55 (3.53, 3.57)	9.59 (9.55, 9.62)	41.49 (41.39, 41.58)	21.78 (21.75, 21.81)	
	40	2.60 (2.59, 2.62)	8.77 (8.73, 8.80)	20.01 (19.95, 20.07)	41.35 (41.25, 41.45)	21.89 (21.86, 21.93)	
	50	6.47 (6.44, 6.50)	18.27 (18.21, 18.33)	31.33 (31.24, 31.41)	40.67 (40.57, 40.77)	21.85 (21.82, 21.88)	
	60	13.16 (13.11, 13.21)	27.71 (27.63, 27.80)	36.08 (35.98, 36.18)	38.13 (38.02, 38.24)	21.05 (21.02, 21.08)	
	70	18.46 (18.39, 18.54)	29.07 (28.96, 29.17)	- (- , -)	31.67 (31.56, 31.79)	18.54 (18.51, 18.58)	
	80	17.10 (17.00, 17.20)	- (- , -)	- (- , -)	21.30 (21.18, 21.42)	13.89 (13.86, 13.93)	
White	0	0.17 (0.16, 0.17)	0.35 (0.34, 0.36)	0.80 (0.79, 0.81)	41.39 (41.29, 41.50)	21.45 (21.42, 21.49)	
	10	0.18 (0.18, 0.19)	0.64 (0.63, 0.65)	1.66 (1.64, 1.67)	41.73 (41.63, 41.84)	21.68 (21.65, 21.72)	
	20	0.46 (0.45, 0.47)	1.48 (1.47, 1.50)	4.01 (3.99, 4.03)	41.78 (41.67, 41.88)	21.75 (21.71, 21.78)	
	30	1.04 (1.03, 1.05)	3.60 (3.58, 3.62)	9.65 (9.61, 9.68)	41.83 (41.73, 41.94)	21.88 (21.84, 21.91)	
	40	2.61 (2.60, 2.63)	8.79 (8.75, 8.82)	20.22 (20.16, 20.29)	41.66 (41.55, 41.76)	21.98 (21.94, 22.01)	
	50	6.47 (6.44, 6.50)	18.46 (18.40, 18.52)	31.70 (31.61, 31.79)	40.93 (40.82, 41.04)	21.92 (21.88, 21.95)	
	60	13.33 (13.27, 13.39)	28.05 (27.96, 28.14)	36.35 (36.24, 36.47)	38.31 (38.19, 38.43)	21.10 (21.06, 21.13)	
	70	18.66 (18.57, 18.74)	29.18 (29.07, 29.29)	- (- , -)	31.66 (31.54, 31.79)	18.54 (18.50, 18.58)	
	80	17.03 (16.92, 17.13)	- (- , -)	- (- , -)	21.04 (20.91, 21.17)	13.82 (13.78, 13.85)	
Black	0	0.12 (0.11, 0.13)	0.24 (0.23, 0.26)	0.54 (0.52, 0.56)	37.22 (36.93, 37.51)	21.31 (21.22, 21.40)	
	10	0.12 (0.11, 0.13)	0.43 (0.41, 0.45)	1.31 (1.28, 1.35)	38.04 (37.74, 38.33)	21.84 (21.74, 21.93)	
	20	0.31 (0.29, 0.33)	1.20 (1.17, 1.23)	3.86 (3.80, 3.92)	38.17 (37.87, 38.47)	21.93 (21.84, 22.02)	
	30	0.91 (0.88, 0.93)	3.62 (3.56, 3.67)	10.58 (10.48, 10.69)	38.58 (38.27, 38.88)	22.20 (22.11, 22.29)	
	40	2.80 (2.75, 2.84)	9.98 (9.87, 10.09)	21.48 (21.29, 21.66)	38.85 (38.54, 39.16)	22.49 (22.39, 22.58)	
	50	7.71 (7.61, 7.81)	20.05 (19.86, 20.24)	31.40 (31.13, 31.67)	38.70 (38.37, 39.03)	22.63 (22.53, 22.73)	
	60	14.56 (14.38, 14.74)	27.95 (27.67, 28.24)	34.71 (34.37, 35.05)	36.56 (36.20, 36.93)	21.96 (21.85, 22.07)	
	70	18.51 (18.24, 18.78)	27.85 (27.48, 28.22)	- (- , -)	30.41 (30.00, 30.82)	19.62 (19.50, 19.74)	
	80	16.73 (16.36, 17.11)	- (- , -)	- (- , -)	21.31 (20.83, 21.81)	15.32 (15.18, 15.46)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table II-9 - continued
ALL CANCER SITES (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific Islander	0	0.14 (0.12, 0.15)	0.26 (0.24, 0.28)	0.53 (0.51, 0.56)	35.73 (35.31, 36.15)	18.32 (18.04, 18.61)	
	10	0.12 (0.11, 0.14)	0.40 (0.38, 0.42)	1.15 (1.12, 1.18)	35.95 (35.54, 36.38)	18.46 (18.18, 18.75)	
	20	0.28 (0.26, 0.29)	1.03 (1.00, 1.06)	3.00 (2.94, 3.05)	35.96 (35.54, 36.38)	18.48 (18.20, 18.77)	
	30	0.76 (0.73, 0.78)	2.74 (2.69, 2.79)	6.85 (6.76, 6.95)	35.92 (35.51, 36.35)	18.51 (18.23, 18.80)	
	40	2.01 (1.96, 2.05)	6.17 (6.08, 6.26)	13.89 (13.73, 14.05)	35.59 (35.17, 36.02)	18.48 (18.19, 18.77)	
	50	4.29 (4.21, 4.36)	12.24 (12.08, 12.40)	22.90 (22.65, 23.15)	34.58 (34.15, 35.02)	18.24 (17.95, 18.54)	
	60	8.47 (8.33, 8.61)	19.84 (19.59, 20.09)	28.61 (28.26, 28.96)	32.30 (31.84, 32.76)	17.52 (17.22, 17.83)	
	70	13.09 (12.88, 13.31)	23.20 (22.85, 23.55)	- (- , -)	27.44 (26.96, 27.94)	15.90 (15.58, 16.22)	
American Indian/ Alaska Native ^a	80	13.70 (13.37, 14.04)	- (- , -)	- (- , -)	19.46 (18.90, 20.04)	12.82 (12.45, 13.19)	
	0	0.14 (0.10, 0.20)	0.26 (0.21, 0.33)	0.49 (0.42, 0.58)	26.34 (25.16, 27.63)	16.87 (16.34, 17.43)	
	10	0.12 (0.09, 0.16)	0.36 (0.30, 0.42)	1.03 (0.92, 1.15)	26.69 (25.49, 28.00)	17.16 (16.62, 17.73)	
	20	0.23 (0.19, 0.29)	0.91 (0.81, 1.03)	2.45 (2.28, 2.63)	26.80 (25.59, 28.12)	17.29 (16.74, 17.86)	
	30	0.69 (0.61, 0.79)	2.26 (2.09, 2.43)	6.16 (5.84, 6.49)	27.06 (25.84, 28.41)	17.55 (17.00, 18.14)	
	40	1.62 (1.48, 1.76)	5.64 (5.33, 5.96)	12.52 (11.94, 13.11)	27.23 (25.97, 28.62)	17.89 (17.32, 18.50)	
	50	4.24 (3.96, 4.53)	11.48 (10.90, 12.08)	19.72 (18.83, 20.65)	26.97 (25.65, 28.42)	18.20 (17.61, 18.84)	
	60	7.99 (7.47, 8.53)	17.09 (16.18, 18.04)	23.23 (22.02, 24.51)	25.09 (23.69, 26.64)	17.90 (17.25, 18.57)	
Hispanic ^b	70	11.16 (10.33, 12.03)	18.68 (17.40, 20.04)	- (- , -)	20.95 (19.41, 22.69)	16.45 (15.73, 17.22)	
	80	11.13 (9.88, 12.50)	- (- , -)	- (- , -)	14.49 (12.73, 16.58)	13.14 (12.24, 14.11)	
	0	0.16 (0.16, 0.17)	0.32 (0.31, 0.33)	0.64 (0.62, 0.65)	37.25 (36.88, 37.62)	17.39 (17.23, 17.55)	
	10	0.16 (0.15, 0.16)	0.48 (0.46, 0.49)	1.22 (1.20, 1.24)	37.55 (37.18, 37.93)	17.57 (17.41, 17.73)	
	20	0.32 (0.31, 0.33)	1.07 (1.05, 1.09)	2.94 (2.90, 2.98)	37.60 (37.22, 37.98)	17.61 (17.45, 17.77)	
	30	0.75 (0.74, 0.77)	2.65 (2.61, 2.68)	7.14 (7.06, 7.22)	37.67 (37.29, 38.06)	17.70 (17.54, 17.86)	
	40	1.92 (1.89, 1.96)	6.49 (6.42, 6.57)	15.43 (15.28, 15.58)	37.53 (37.14, 37.92)	17.75 (17.59, 17.92)	
	50	4.74 (4.67, 4.81)	14.02 (13.87, 14.17)	25.73 (25.49, 25.98)	36.96 (36.56, 37.36)	17.69 (17.53, 17.86)	
-	60	10.09 (9.95, 10.23)	22.84 (22.59, 23.09)	31.48 (31.15, 31.82)	35.05 (34.63, 35.48)	17.14 (16.97, 17.31)	
	70	15.38 (15.16, 15.61)	25.81 (25.46, 26.17)	- (- , -)	30.12 (29.65, 30.60)	15.57 (15.39, 15.76)	
	80	15.39 (15.04, 15.75)	- (- , -)	- (- , -)	21.75 (21.17, 22.35)	12.47 (12.26, 12.69)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table II-9 - continued
ALL CANCER SITES (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.17 (0.17, 0.18)	0.35 (0.34, 0.36)	0.72 (0.70, 0.73)	44.94 (44.79, 45.09)	23.31 (23.27, 23.36)	
	10	0.18 (0.18, 0.19)	0.55 (0.54, 0.56)	1.26 (1.25, 1.28)	45.45 (45.30, 45.60)	23.65 (23.60, 23.70)	
	20	0.37 (0.36, 0.38)	1.09 (1.08, 1.11)	3.07 (3.05, 3.09)	45.59 (45.44, 45.74)	23.76 (23.71, 23.80)	
	30	0.73 (0.72, 0.74)	2.74 (2.72, 2.76)	9.11 (9.07, 9.16)	45.95 (45.80, 46.10)	24.02 (23.97, 24.06)	
	40	2.06 (2.04, 2.08)	8.58 (8.53, 8.62)	22.17 (22.08, 22.25)	46.26 (46.10, 46.42)	24.29 (24.24, 24.34)	
	50	6.86 (6.82, 6.90)	21.16 (21.07, 21.25)	36.84 (36.70, 36.97)	46.50 (46.34, 46.67)	24.54 (24.49, 24.59)	
	60	16.25 (16.16, 16.33)	34.06 (33.92, 34.20)	43.07 (42.90, 43.23)	45.04 (44.87, 45.22)	24.11 (24.06, 24.17)	
	70	24.10 (23.96, 24.23)	36.28 (36.10, 36.47)	- (- , -)	38.96 (38.76, 39.16)	21.93 (21.87, 21.98)	
	80	22.71 (22.52, 22.91)	- (- , -)	- (- , -)	27.70 (27.45, 27.95)	17.43 (17.36, 17.50)	
White	0	0.18 (0.17, 0.19)	0.37 (0.36, 0.38)	0.77 (0.76, 0.79)	44.89 (44.73, 45.06)	23.41 (23.36, 23.46)	
	10	0.19 (0.19, 0.20)	0.60 (0.58, 0.61)	1.34 (1.32, 1.36)	45.31 (45.15, 45.47)	23.70 (23.65, 23.75)	
	20	0.40 (0.40, 0.41)	1.16 (1.14, 1.17)	3.12 (3.10, 3.15)	45.43 (45.26, 45.60)	23.80 (23.75, 23.85)	
	30	0.76 (0.75, 0.77)	2.76 (2.73, 2.78)	9.02 (8.97, 9.07)	45.72 (45.56, 45.89)	24.03 (23.98, 24.08)	
	40	2.04 (2.02, 2.06)	8.45 (8.40, 8.50)	22.06 (21.96, 22.15)	45.97 (45.80, 46.14)	24.27 (24.22, 24.32)	
	50	6.73 (6.68, 6.77)	21.02 (20.92, 21.12)	36.69 (36.55, 36.84)	46.14 (45.96, 46.32)	24.49 (24.43, 24.54)	
	60	16.16 (16.07, 16.26)	33.88 (33.73, 34.04)	42.70 (42.52, 42.88)	44.57 (44.38, 44.76)	24.03 (23.97, 24.08)	
	70	23.86 (23.71, 24.00)	35.73 (35.53, 35.93)	- (- , -)	38.24 (38.03, 38.46)	21.78 (21.72, 21.84)	
	80	22.07 (21.86, 22.27)	- (- , -)	- (- , -)	26.73 (26.47, 27.00)	17.23 (17.16, 17.30)	
Black	0	0.13 (0.11, 0.14)	0.25 (0.23, 0.27)	0.50 (0.47, 0.53)	42.33 (41.87, 42.79)	23.70 (23.56, 23.84)	
	10	0.13 (0.12, 0.14)	0.38 (0.36, 0.41)	1.01 (0.97, 1.05)	43.38 (42.91, 43.85)	24.35 (24.20, 24.49)	
	20	0.26 (0.24, 0.28)	0.89 (0.85, 0.93)	3.30 (3.23, 3.37)	43.67 (43.19, 44.15)	24.51 (24.37, 24.65)	
	30	0.65 (0.62, 0.68)	3.13 (3.05, 3.20)	11.72 (11.55, 11.88)	44.59 (44.10, 45.08)	25.01 (24.87, 25.16)	
	40	2.57 (2.50, 2.63)	11.48 (11.31, 11.65)	26.49 (26.18, 26.80)	45.57 (45.06, 46.08)	25.61 (25.46, 25.76)	
	50	9.67 (9.51, 9.83)	25.96 (25.63, 26.28)	39.67 (39.22, 40.12)	46.66 (46.12, 47.20)	26.26 (26.10, 26.42)	
	60	20.23 (19.90, 20.55)	37.25 (36.75, 37.76)	44.28 (43.69, 44.87)	45.93 (45.31, 46.57)	26.33 (26.15, 26.50)	
	70	26.64 (26.10, 27.19)	37.64 (36.92, 38.36)	- (- , -)	40.23 (39.44, 41.03)	24.68 (24.47, 24.88)	
	80	24.53 (23.67, 25.41)	- (- , -)	- (- , -)	30.31 (29.19, 31.47)	20.86 (20.59, 21.14)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table II-9 - continued
ALL CANCER SITES (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific Islander	0	0.15 (0.13, 0.17)	0.27 (0.25, 0.30)	0.49 (0.45, 0.52)	39.83 (39.21, 40.48)	20.73 (20.32, 21.16)	
	10	0.13 (0.11, 0.14)	0.34 (0.32, 0.37)	0.80 (0.77, 0.85)	40.13 (39.50, 40.78)	20.90 (20.48, 21.34)	
	20	0.22 (0.20, 0.24)	0.68 (0.65, 0.72)	1.95 (1.89, 2.01)	40.18 (39.54, 40.83)	20.94 (20.52, 21.38)	
	30	0.47 (0.44, 0.49)	1.75 (1.69, 1.81)	5.52 (5.40, 5.64)	40.27 (39.63, 40.93)	21.01 (20.58, 21.44)	
	40	1.29 (1.25, 1.34)	5.11 (4.99, 5.23)	14.39 (14.15, 14.64)	40.24 (39.60, 40.91)	21.02 (20.59, 21.46)	
	50	3.92 (3.81, 4.02)	13.45 (13.20, 13.69)	27.27 (26.85, 27.68)	39.98 (39.32, 40.66)	20.89 (20.46, 21.34)	
	60	10.20 (9.98, 10.43)	25.00 (24.58, 25.42)	35.15 (34.58, 35.72)	38.61 (37.92, 39.33)	20.35 (19.90, 20.81)	
	70	17.68 (17.30, 18.08)	29.81 (29.21, 30.42)	- (- , -)	33.95 (33.18, 34.75)	18.81 (18.33, 19.31)	
American Indian/ Alaska Native ^a	80	18.21 (17.60, 18.84)	- (- , -)	- (- , -)	24.43 (23.49, 25.41)	15.46 (14.89, 16.06)	
	0	0.13 (0.08, 0.21)	0.26 (0.19, 0.36)	0.46 (0.36, 0.58)	26.35 (24.64, 28.34)	17.20 (16.42, 18.05)	
	10	0.14 (0.09, 0.19)	0.33 (0.26, 0.43)	0.88 (0.74, 1.03)	26.78 (25.04, 28.81)	17.54 (16.74, 18.41)	
	20	0.20 (0.14, 0.28)	0.75 (0.62, 0.90)	1.92 (1.70, 2.15)	26.96 (25.20, 29.01)	17.71 (16.91, 18.59)	
	30	0.56 (0.45, 0.69)	1.76 (1.56, 1.98)	5.58 (5.15, 6.04)	27.45 (25.64, 29.55)	18.12 (17.30, 19.02)	
	40	1.25 (1.08, 1.43)	5.22 (4.79, 5.67)	12.91 (12.06, 13.80)	27.98 (26.11, 30.16)	18.71 (17.85, 19.64)	
	50	4.22 (3.82, 4.64)	12.38 (11.51, 13.29)	21.27 (19.92, 22.69)	28.37 (26.40, 30.67)	19.37 (18.46, 20.36)	
	60	9.14 (8.33, 10.00)	19.10 (17.70, 20.59)	25.48 (23.61, 27.48)	27.05 (24.92, 29.57)	19.45 (18.45, 20.53)	
Hispanic ^b	70	12.62 (11.32, 14.02)	20.69 (18.67, 22.89)	- (- , -)	22.68 (20.29, 25.60)	18.33 (17.16, 19.61)	
	80	12.97 (10.84, 15.41)	- (- , -)	- (- , -)	16.17 (13.25, 20.04)	15.20 (13.66, 16.95)	
	0	0.17 (0.16, 0.19)	0.34 (0.33, 0.36)	0.62 (0.60, 0.65)	41.57 (40.99, 42.18)	19.80 (19.55, 20.06)	
	10	0.17 (0.16, 0.18)	0.46 (0.44, 0.47)	0.96 (0.93, 0.98)	41.97 (41.37, 42.58)	20.03 (19.77, 20.29)	
	20	0.29 (0.28, 0.30)	0.79 (0.77, 0.82)	2.12 (2.08, 2.17)	42.10 (41.50, 42.72)	20.11 (19.85, 20.37)	
	30	0.51 (0.49, 0.53)	1.86 (1.81, 1.90)	6.30 (6.19, 6.40)	42.40 (41.79, 43.02)	20.29 (20.03, 20.56)	
	40	1.37 (1.33, 1.41)	5.89 (5.79, 5.99)	16.97 (16.74, 17.20)	42.64 (42.02, 43.28)	20.47 (20.20, 20.74)	
	50	4.70 (4.61, 4.80)	16.23 (15.99, 16.47)	30.99 (30.59, 31.38)	42.94 (42.30, 43.60)	20.65 (20.38, 20.93)	
-	60	12.72 (12.50, 12.95)	29.01 (28.59, 29.43)	38.58 (38.03, 39.14)	42.20 (41.50, 42.92)	20.42 (20.14, 20.72)	
	70	20.76 (20.35, 21.16)	32.96 (32.34, 33.59)	- (- , -)	37.57 (36.75, 38.43)	19.09 (18.77, 19.42)	
	80	20.38 (19.69, 21.08)	- (- , -)	- (- , -)	28.08 (26.95, 29.27)	16.01 (15.61, 16.43)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table II-9 - continued
ALL CANCER SITES (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.15 (0.14, 0.16)	0.31 (0.30, 0.32)	0.79 (0.77, 0.80)	37.52 (37.40, 37.65)	19.74 (19.70, 19.78)	
	10	0.16 (0.16, 0.17)	0.64 (0.63, 0.66)	1.92 (1.90, 1.94)	37.84 (37.72, 37.97)	19.96 (19.92, 20.00)	
	20	0.48 (0.47, 0.49)	1.76 (1.74, 1.78)	4.82 (4.79, 4.85)	37.83 (37.70, 37.95)	19.99 (19.95, 20.03)	
	30	1.29 (1.27, 1.30)	4.38 (4.35, 4.40)	10.08 (10.04, 10.13)	37.69 (37.56, 37.81)	20.04 (20.00, 20.08)	
	40	3.15 (3.12, 3.17)	8.97 (8.93, 9.02)	18.00 (17.93, 18.08)	37.12 (37.00, 37.25)	20.02 (19.98, 20.06)	
	50	6.10 (6.06, 6.13)	15.56 (15.48, 15.63)	26.34 (26.24, 26.45)	35.58 (35.45, 35.71)	19.75 (19.71, 19.79)	
	60	10.36 (10.30, 10.42)	22.18 (22.07, 22.28)	30.10 (29.97, 30.22)	32.29 (32.16, 32.43)	18.70 (18.66, 18.74)	
	70	14.18 (14.09, 14.26)	23.68 (23.56, 23.80)	- (- , -)	26.31 (26.18, 26.45)	16.16 (16.11, 16.20)	
	80	13.84 (13.74, 13.95)	- (- , -)	- (- , -)	17.68 (17.55, 17.82)	11.87 (11.83, 11.91)	
White	0	0.15 (0.15, 0.16)	0.33 (0.32, 0.34)	0.83 (0.82, 0.85)	38.46 (38.32, 38.60)	19.96 (19.91, 20.00)	
	10	0.17 (0.17, 0.18)	0.69 (0.67, 0.70)	2.00 (1.98, 2.02)	38.73 (38.59, 38.87)	20.14 (20.10, 20.19)	
	20	0.52 (0.50, 0.53)	1.83 (1.81, 1.86)	4.94 (4.91, 4.98)	38.71 (38.57, 38.85)	20.17 (20.13, 20.22)	
	30	1.33 (1.31, 1.35)	4.47 (4.44, 4.50)	10.31 (10.25, 10.36)	38.54 (38.40, 38.68)	20.21 (20.17, 20.26)	
	40	3.20 (3.17, 3.23)	9.15 (9.10, 9.20)	18.50 (18.42, 18.59)	37.95 (37.80, 38.09)	20.19 (20.15, 20.23)	
	50	6.23 (6.19, 6.27)	16.01 (15.93, 16.10)	27.13 (27.01, 27.25)	36.35 (36.21, 36.50)	19.90 (19.86, 19.95)	
	60	10.71 (10.63, 10.78)	22.87 (22.75, 22.98)	30.85 (30.71, 30.99)	32.96 (32.81, 33.11)	18.84 (18.80, 18.89)	
	70	14.61 (14.52, 14.71)	24.20 (24.07, 24.34)	- (- , -)	26.74 (26.59, 26.89)	16.24 (16.19, 16.28)	
	80	14.05 (13.93, 14.17)	- (- , -)	- (- , -)	17.77 (17.62, 17.92)	11.85 (11.81, 11.90)	
Black	0	0.12 (0.10, 0.13)	0.23 (0.21, 0.25)	0.58 (0.55, 0.61)	32.66 (32.29, 33.03)	19.39 (19.27, 19.51)	
	10	0.12 (0.11, 0.13)	0.48 (0.45, 0.50)	1.60 (1.55, 1.65)	33.28 (32.91, 33.67)	19.81 (19.69, 19.93)	
	20	0.36 (0.34, 0.38)	1.49 (1.44, 1.54)	4.39 (4.30, 4.47)	33.30 (32.92, 33.68)	19.85 (19.72, 19.97)	
	30	1.14 (1.10, 1.18)	4.07 (3.99, 4.15)	9.59 (9.45, 9.73)	33.30 (32.92, 33.68)	19.94 (19.81, 20.06)	
	40	3.01 (2.94, 3.08)	8.67 (8.53, 8.81)	17.04 (16.81, 17.26)	32.99 (32.60, 33.38)	19.99 (19.87, 20.12)	
	50	6.02 (5.90, 6.14)	14.92 (14.70, 15.14)	24.25 (23.93, 24.57)	31.88 (31.48, 32.29)	19.76 (19.63, 19.89)	
	60	10.06 (9.87, 10.26)	20.62 (20.29, 20.94)	27.19 (26.79, 27.59)	29.25 (28.82, 29.68)	18.72 (18.58, 18.85)	
	70	13.32 (13.03, 13.60)	21.61 (21.21, 22.02)	- (- , -)	24.21 (23.75, 24.67)	16.31 (16.17, 16.45)	
	80	13.07 (12.69, 13.46)	- (- , -)	- (- , -)	17.17 (16.67, 17.68)	12.48 (12.33, 12.65)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table II-9 - continued
ALL CANCER SITES (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific Islander	0	0.13 (0.11, 0.15)	0.25 (0.22, 0.27)	0.58 (0.54, 0.61)	32.32 (31.78, 32.88)	16.26 (15.88, 16.65)	
	10	0.12 (0.11, 0.14)	0.45 (0.42, 0.49)	1.48 (1.42, 1.53)	32.50 (31.95, 33.06)	16.37 (15.99, 16.76)	
	20	0.33 (0.31, 0.36)	1.36 (1.31, 1.41)	3.96 (3.88, 4.05)	32.46 (31.92, 33.03)	16.38 (16.00, 16.77)	
	30	1.03 (0.99, 1.07)	3.65 (3.57, 3.74)	8.07 (7.94, 8.21)	32.31 (31.76, 32.88)	16.38 (16.00, 16.78)	
	40	2.66 (2.59, 2.72)	7.14 (7.00, 7.27)	13.51 (13.30, 13.73)	31.69 (31.14, 32.27)	16.31 (15.93, 16.71)	
	50	4.63 (4.52, 4.73)	11.21 (11.01, 11.42)	19.34 (19.03, 19.65)	29.99 (29.42, 30.57)	15.99 (15.60, 16.39)	
	60	7.00 (6.83, 7.17)	15.63 (15.33, 15.93)	23.16 (22.74, 23.59)	26.94 (26.35, 27.55)	15.15 (14.76, 15.56)	
	70	9.64 (9.40, 9.89)	18.06 (17.65, 18.48)	- (- , -)	22.28 (21.67, 22.93)	13.54 (13.13, 13.97)	
American Indian/ Alaska Native ^a	80	10.63 (10.25, 11.01)	- (- , -)	- (- , -)	15.96 (15.27, 16.69)	10.80 (10.34, 11.28)	
	0	0.15 (0.09, 0.24)	0.26 (0.19, 0.36)	0.53 (0.42, 0.66)	26.45 (24.82, 28.29)	16.71 (15.98, 17.50)	
	10	0.11 (0.07, 0.16)	0.38 (0.29, 0.48)	1.18 (1.02, 1.37)	26.71 (25.06, 28.58)	16.97 (16.22, 17.77)	
	20	0.27 (0.20, 0.36)	1.08 (0.92, 1.26)	2.98 (2.71, 3.26)	26.75 (25.09, 28.63)	17.04 (16.29, 17.85)	
	30	0.82 (0.69, 0.97)	2.74 (2.48, 3.01)	6.72 (6.25, 7.20)	26.79 (25.11, 28.69)	17.17 (16.41, 17.99)	
	40	1.97 (1.76, 2.19)	6.04 (5.60, 6.51)	12.17 (11.39, 12.98)	26.61 (24.90, 28.55)	17.29 (16.52, 18.12)	
	50	4.26 (3.88, 4.67)	10.66 (9.89, 11.47)	18.34 (17.16, 19.58)	25.75 (23.99, 27.76)	17.31 (16.51, 18.17)	
	60	6.97 (6.31, 7.67)	15.33 (14.16, 16.58)	21.30 (19.72, 22.99)	23.40 (21.55, 25.53)	16.71 (15.87, 17.62)	
Hispanic ^b	70	9.97 (8.92, 11.10)	17.08 (15.45, 18.84)	- (- , -)	19.58 (17.59, 21.92)	15.11 (14.18, 16.11)	
	80	9.87 (8.38, 11.56)	- (- , -)	- (- , -)	13.35 (11.20, 16.04)	11.87 (10.79, 13.09)	
	0	0.15 (0.14, 0.17)	0.30 (0.28, 0.31)	0.65 (0.63, 0.68)	33.95 (33.47, 34.43)	15.62 (15.42, 15.82)	
	10	0.14 (0.13, 0.15)	0.50 (0.48, 0.52)	1.52 (1.49, 1.56)	34.18 (33.70, 34.67)	15.76 (15.56, 15.97)	
	20	0.36 (0.35, 0.38)	1.38 (1.35, 1.42)	3.83 (3.77, 3.90)	34.15 (33.67, 34.64)	15.77 (15.56, 15.98)	
	30	1.03 (1.00, 1.06)	3.50 (3.43, 3.56)	8.06 (7.95, 8.18)	34.01 (33.53, 34.51)	15.78 (15.57, 15.99)	
	40	2.50 (2.45, 2.56)	7.14 (7.03, 7.25)	14.18 (13.99, 14.38)	33.47 (32.98, 33.97)	15.72 (15.51, 15.93)	
	50	4.80 (4.70, 4.89)	12.10 (11.91, 12.29)	21.28 (20.99, 21.58)	32.07 (31.57, 32.59)	15.46 (15.25, 15.67)	
-	60	7.84 (7.67, 8.00)	17.70 (17.41, 17.99)	25.64 (25.24, 26.06)	29.28 (28.76, 29.83)	14.70 (14.49, 14.92)	
	70	11.38 (11.12, 11.63)	20.54 (20.13, 20.96)	- (- , -)	24.74 (24.18, 25.33)	13.14 (12.91, 13.36)	
	80	12.42 (12.02, 12.83)	- (- , -)	- (- , -)	18.12 (17.46, 18.80)	10.40 (10.14, 10.65)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table III-9

BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.03 (0.03, 0.04)	0.06 (0.06, 0.06)	0.08 (0.08, 0.09)	0.60 (0.59, 0.61)	0.43 (0.43, 0.44)	
	10	0.02 (0.02, 0.03)	0.05 (0.05, 0.05)	0.08 (0.08, 0.09)	0.57 (0.56, 0.58)	0.43 (0.43, 0.44)	
	20	0.03 (0.02, 0.03)	0.06 (0.06, 0.06)	0.11 (0.11, 0.11)	0.55 (0.54, 0.56)	0.43 (0.42, 0.43)	
	30	0.03 (0.03, 0.04)	0.09 (0.08, 0.09)	0.17 (0.17, 0.17)	0.53 (0.52, 0.54)	0.42 (0.42, 0.43)	
	40	0.05 (0.05, 0.05)	0.14 (0.13, 0.14)	0.27 (0.26, 0.27)	0.50 (0.49, 0.51)	0.41 (0.41, 0.42)	
	50	0.09 (0.09, 0.09)	0.22 (0.21, 0.23)	0.37 (0.36, 0.38)	0.46 (0.45, 0.47)	0.39 (0.39, 0.40)	
	60	0.14 (0.13, 0.15)	0.30 (0.29, 0.31)	0.38 (0.37, 0.39)	0.40 (0.39, 0.41)	0.34 (0.34, 0.35)	
	70	0.18 (0.18, 0.19)	0.28 (0.27, 0.29)	- (- , -)	0.30 (0.29, 0.31)	0.26 (0.26, 0.27)	
	80	0.14 (0.13, 0.14)	- (- , -)	- (- , -)	0.16 (0.16, 0.17)	0.16 (0.15, 0.16)	
White	0	0.04 (0.03, 0.04)	0.06 (0.06, 0.06)	0.09 (0.09, 0.09)	0.66 (0.65, 0.67)	0.47 (0.46, 0.47)	
	10	0.03 (0.02, 0.03)	0.05 (0.05, 0.06)	0.09 (0.09, 0.10)	0.63 (0.62, 0.64)	0.47 (0.46, 0.47)	
	20	0.03 (0.03, 0.03)	0.07 (0.06, 0.07)	0.12 (0.12, 0.13)	0.61 (0.59, 0.62)	0.46 (0.46, 0.47)	
	30	0.04 (0.04, 0.04)	0.10 (0.09, 0.10)	0.19 (0.18, 0.19)	0.58 (0.57, 0.59)	0.46 (0.45, 0.46)	
	40	0.06 (0.05, 0.06)	0.15 (0.15, 0.16)	0.30 (0.29, 0.30)	0.55 (0.54, 0.56)	0.45 (0.44, 0.45)	
	50	0.10 (0.09, 0.10)	0.24 (0.24, 0.25)	0.41 (0.40, 0.42)	0.51 (0.50, 0.52)	0.42 (0.42, 0.43)	
	60	0.16 (0.15, 0.16)	0.33 (0.32, 0.34)	0.42 (0.41, 0.43)	0.43 (0.42, 0.45)	0.37 (0.36, 0.37)	
	70	0.20 (0.19, 0.21)	0.30 (0.29, 0.31)	- (- , -)	0.32 (0.31, 0.33)	0.28 (0.27, 0.28)	
	80	0.14 (0.14, 0.15)	- (- , -)	- (- , -)	0.17 (0.16, 0.18)	0.16 (0.16, 0.17)	
Black	0	0.03 (0.02, 0.03)	0.04 (0.04, 0.05)	0.06 (0.05, 0.06)	0.32 (0.30, 0.35)	0.22 (0.21, 0.23)	
	10	0.02 (0.01, 0.02)	0.03 (0.03, 0.04)	0.05 (0.05, 0.06)	0.30 (0.28, 0.33)	0.22 (0.21, 0.23)	
	20	0.01 (0.01, 0.02)	0.04 (0.03, 0.04)	0.07 (0.06, 0.07)	0.29 (0.27, 0.32)	0.21 (0.20, 0.22)	
	30	0.02 (0.02, 0.03)	0.05 (0.05, 0.06)	0.10 (0.09, 0.11)	0.28 (0.26, 0.31)	0.21 (0.20, 0.22)	
	40	0.03 (0.03, 0.04)	0.08 (0.07, 0.09)	0.15 (0.13, 0.16)	0.27 (0.24, 0.29)	0.20 (0.19, 0.21)	
	50	0.05 (0.05, 0.06)	0.12 (0.11, 0.14)	0.19 (0.17, 0.21)	0.25 (0.22, 0.27)	0.20 (0.19, 0.21)	
	60	0.08 (0.06, 0.09)	0.15 (0.14, 0.17)	0.20 (0.18, 0.23)	0.22 (0.19, 0.24)	0.17 (0.16, 0.18)	
	70	0.10 (0.08, 0.12)	0.16 (0.14, 0.19)	- (- , -)	0.18 (0.15, 0.20)	0.14 (0.13, 0.15)	
	80	0.10 (0.08, 0.12)	- (- , -)	- (- , -)	0.12 (0.10, 0.16)	0.09 (0.08, 0.11)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table III-9 - continued

BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.03 (0.02, 0.03)	0.04 (0.04, 0.05)	0.06 (0.05, 0.07)	0.35 (0.32, 0.39)	0.25 (0.22, 0.29)	
	10	0.02 (0.01, 0.02)	0.03 (0.03, 0.04)	0.05 (0.05, 0.06)	0.33 (0.30, 0.37)	0.25 (0.22, 0.29)	
Islander	20	0.02 (0.01, 0.02)	0.03 (0.03, 0.04)	0.06 (0.06, 0.07)	0.31 (0.28, 0.35)	0.24 (0.22, 0.28)	
	30	0.02 (0.02, 0.02)	0.05 (0.04, 0.06)	0.09 (0.08, 0.10)	0.30 (0.27, 0.34)	0.24 (0.21, 0.28)	
	40	0.03 (0.02, 0.04)	0.07 (0.06, 0.08)	0.14 (0.12, 0.15)	0.28 (0.25, 0.32)	0.24 (0.21, 0.27)	
	50	0.05 (0.04, 0.05)	0.11 (0.09, 0.12)	0.18 (0.16, 0.20)	0.25 (0.22, 0.29)	0.22 (0.19, 0.26)	
	60	0.07 (0.05, 0.08)	0.14 (0.12, 0.16)	0.20 (0.17, 0.23)	0.21 (0.19, 0.26)	0.20 (0.17, 0.24)	
	70	0.08 (0.06, 0.10)	0.15 (0.12, 0.18)	- (-, -)	0.16 (0.13, 0.21)	0.16 (0.13, 0.20)	
	80	0.09 (0.06, 0.11)	- (-, -)	- (-, -)	0.10 (0.08, 0.15)	0.11 (0.08, 0.16)	
American Indian/ Alaska Native ^a	0	0.02 (0.01, 0.04)	0.03 (0.02, 0.06)	0.04 (0.02, 0.08)	0.22 (0.14, 0.49)	0.18 (0.14, 0.27)	
	10	0.01 (0.00, 0.03)	0.03 (0.01, 0.05)	0.04 (0.02, 0.07)	0.20 (0.13, 0.48)	0.17 (0.13, 0.26)	
	20	0.01 (0.00, 0.03)	0.03 (0.01, 0.06)	0.04 (0.02, 0.08)	0.19 (0.12, 0.47)	0.17 (0.13, 0.26)	
	30	0.02 (0.01, 0.04)	0.03 (0.01, 0.06)	0.04 (0.02, 0.08)	0.18 (0.11, 0.47)	0.17 (0.13, 0.26)	
	40	0.01 (0.00, 0.04)	0.03 (0.01, 0.06)	0.11 (0.06, 0.19)	0.17 (0.10, 0.46)	0.17 (0.13, 0.26)	
	50	0.01 (0.00, 0.04)	0.10 (0.05, 0.18)	0.14 (0.08, 0.25)	0.16 (0.09, 0.47)	0.16 (0.11, 0.26)	
	60	0.09 (0.04, 0.17)	0.14 (0.07, 0.25)	0.16 (0.08, 0.30)	0.16 (0.08, 0.49)	0.14 (0.09, 0.24)	
	70	0.06 (0.02, 0.14)	0.08 (0.02, 0.21)	- (-, -)	0.08 (0.02, 0.46)	0.11 (0.06, 0.23)	
	80	0.03 (0.00, 0.20)	- (-, -)	- (-, -)	0.03 (0.00, 0.58)	0.05 (0.02, 0.23)	
Hispanic ^b	0	0.03 (0.03, 0.04)	0.05 (0.05, 0.06)	0.07 (0.06, 0.08)	0.54 (0.50, 0.58)	0.32 (0.30, 0.34)	
	10	0.02 (0.02, 0.02)	0.04 (0.03, 0.04)	0.06 (0.06, 0.07)	0.51 (0.48, 0.56)	0.31 (0.29, 0.33)	
	20	0.02 (0.02, 0.02)	0.04 (0.04, 0.05)	0.08 (0.08, 0.09)	0.50 (0.46, 0.54)	0.31 (0.29, 0.33)	
	30	0.03 (0.02, 0.03)	0.06 (0.06, 0.07)	0.13 (0.12, 0.14)	0.48 (0.44, 0.53)	0.31 (0.29, 0.33)	
	40	0.04 (0.03, 0.04)	0.11 (0.10, 0.12)	0.22 (0.20, 0.24)	0.46 (0.42, 0.51)	0.30 (0.28, 0.32)	
	50	0.07 (0.06, 0.08)	0.19 (0.17, 0.20)	0.32 (0.30, 0.35)	0.43 (0.39, 0.48)	0.29 (0.27, 0.31)	
	60	0.12 (0.11, 0.13)	0.26 (0.24, 0.29)	0.35 (0.32, 0.38)	0.38 (0.34, 0.42)	0.26 (0.24, 0.29)	
	70	0.16 (0.14, 0.18)	0.26 (0.23, 0.29)	- (-, -)	0.29 (0.25, 0.34)	0.21 (0.19, 0.23)	
	80	0.13 (0.10, 0.16)	- (-, -)	- (-, -)	0.17 (0.13, 0.22)	0.14 (0.11, 0.16)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table III-9 - continued

BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.04 (0.03, 0.04)	0.06 (0.06, 0.06)	0.09 (0.08, 0.09)	0.67 (0.65, 0.68)	0.48 (0.47, 0.49)	
	10	0.03 (0.02, 0.03)	0.05 (0.05, 0.06)	0.09 (0.09, 0.10)	0.64 (0.63, 0.66)	0.48 (0.47, 0.49)	
	20	0.03 (0.03, 0.03)	0.06 (0.06, 0.07)	0.12 (0.12, 0.13)	0.62 (0.60, 0.64)	0.48 (0.47, 0.48)	
	30	0.04 (0.04, 0.04)	0.10 (0.09, 0.10)	0.20 (0.19, 0.21)	0.60 (0.58, 0.62)	0.48 (0.47, 0.48)	
	40	0.06 (0.06, 0.07)	0.16 (0.16, 0.17)	0.31 (0.30, 0.32)	0.57 (0.56, 0.59)	0.47 (0.46, 0.47)	
	50	0.11 (0.10, 0.11)	0.26 (0.25, 0.27)	0.43 (0.42, 0.45)	0.53 (0.51, 0.54)	0.44 (0.43, 0.45)	
	60	0.17 (0.16, 0.18)	0.35 (0.34, 0.37)	0.44 (0.42, 0.45)	0.46 (0.44, 0.47)	0.39 (0.38, 0.39)	
	70	0.22 (0.21, 0.23)	0.32 (0.31, 0.34)	- (- , -)	0.34 (0.33, 0.36)	0.30 (0.29, 0.30)	
	80	0.16 (0.15, 0.17)	- (- , -)	- (- , -)	0.19 (0.18, 0.21)	0.18 (0.17, 0.18)	
White	0	0.04 (0.03, 0.04)	0.07 (0.06, 0.07)	0.09 (0.09, 0.10)	0.74 (0.72, 0.75)	0.52 (0.51, 0.53)	
	10	0.03 (0.03, 0.03)	0.06 (0.05, 0.06)	0.10 (0.09, 0.10)	0.71 (0.69, 0.73)	0.52 (0.51, 0.53)	
	20	0.03 (0.03, 0.03)	0.07 (0.07, 0.08)	0.14 (0.13, 0.14)	0.68 (0.66, 0.70)	0.52 (0.51, 0.52)	
	30	0.04 (0.04, 0.05)	0.11 (0.10, 0.11)	0.22 (0.21, 0.23)	0.66 (0.64, 0.68)	0.51 (0.51, 0.52)	
	40	0.07 (0.06, 0.07)	0.18 (0.17, 0.19)	0.35 (0.34, 0.36)	0.63 (0.61, 0.65)	0.50 (0.50, 0.51)	
	50	0.12 (0.11, 0.12)	0.29 (0.28, 0.30)	0.48 (0.46, 0.49)	0.58 (0.56, 0.60)	0.47 (0.47, 0.48)	
	60	0.19 (0.18, 0.20)	0.39 (0.37, 0.40)	0.48 (0.46, 0.50)	0.50 (0.48, 0.52)	0.41 (0.41, 0.42)	
	70	0.24 (0.23, 0.25)	0.35 (0.33, 0.37)	- (- , -)	0.37 (0.35, 0.39)	0.31 (0.31, 0.32)	
	80	0.17 (0.16, 0.18)	- (- , -)	- (- , -)	0.20 (0.19, 0.22)	0.18 (0.18, 0.19)	
Black	0	0.03 (0.02, 0.03)	0.04 (0.04, 0.05)	0.06 (0.05, 0.07)	0.35 (0.31, 0.39)	0.23 (0.22, 0.25)	
	10	0.02 (0.01, 0.02)	0.04 (0.03, 0.04)	0.06 (0.05, 0.07)	0.33 (0.29, 0.37)	0.23 (0.22, 0.25)	
	20	0.02 (0.01, 0.02)	0.04 (0.03, 0.05)	0.07 (0.06, 0.09)	0.31 (0.28, 0.35)	0.23 (0.21, 0.24)	
	30	0.02 (0.02, 0.03)	0.06 (0.05, 0.07)	0.12 (0.10, 0.14)	0.30 (0.27, 0.34)	0.23 (0.21, 0.24)	
	40	0.04 (0.03, 0.05)	0.10 (0.08, 0.12)	0.18 (0.15, 0.20)	0.29 (0.26, 0.33)	0.22 (0.21, 0.24)	
	50	0.06 (0.05, 0.08)	0.15 (0.13, 0.17)	0.23 (0.20, 0.26)	0.27 (0.23, 0.31)	0.21 (0.20, 0.23)	
	60	0.10 (0.08, 0.12)	0.19 (0.16, 0.22)	0.23 (0.20, 0.27)	0.24 (0.20, 0.28)	0.19 (0.17, 0.20)	
	70	0.12 (0.10, 0.16)	0.18 (0.14, 0.23)	- (- , -)	0.19 (0.15, 0.24)	0.15 (0.13, 0.17)	
	80	0.10 (0.07, 0.15)	- (- , -)	- (- , -)	0.11 (0.07, 0.18)	0.09 (0.07, 0.12)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table III-9 - continued

BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific Islander	0	0.02 (0.02, 0.03)	0.04 (0.03, 0.06)	0.06 (0.05, 0.07)	0.37 (0.32, 0.44)	0.31 (0.26, 0.37)	
	10	0.02 (0.01, 0.03)	0.04 (0.03, 0.05)	0.06 (0.05, 0.07)	0.35 (0.30, 0.42)	0.30 (0.26, 0.37)	
	20	0.02 (0.01, 0.02)	0.04 (0.03, 0.05)	0.07 (0.06, 0.09)	0.33 (0.28, 0.40)	0.30 (0.25, 0.36)	
	30	0.02 (0.01, 0.03)	0.06 (0.05, 0.07)	0.11 (0.09, 0.13)	0.31 (0.27, 0.38)	0.29 (0.25, 0.36)	
	40	0.04 (0.03, 0.05)	0.09 (0.08, 0.11)	0.15 (0.13, 0.17)	0.30 (0.25, 0.36)	0.29 (0.24, 0.35)	
	50	0.05 (0.04, 0.07)	0.11 (0.09, 0.14)	0.19 (0.16, 0.22)	0.26 (0.22, 0.33)	0.27 (0.23, 0.34)	
	60	0.06 (0.05, 0.08)	0.14 (0.11, 0.17)	0.21 (0.17, 0.25)	0.22 (0.18, 0.29)	0.24 (0.20, 0.31)	
	70	0.09 (0.06, 0.12)	0.16 (0.12, 0.21)	- (- , -)	0.18 (0.13, 0.25)	0.21 (0.16, 0.28)	
American Indian/ Alaska Native ^a	80	0.10 (0.06, 0.15)	- (- , -)	- (- , -)	0.12 (0.07, 0.21)	0.15 (0.10, 0.24)	
	0	0.01 (0.00, 0.06)	0.02 (0.01, 0.07)	0.03 (0.01, 0.09)	0.30 (0.17, 0.95)	0.22 (0.16, 0.41)	
	10	0.01 (0.00, 0.03)	0.02 (0.01, 0.06)	0.05 (0.02, 0.10)	0.29 (0.16, 0.96)	0.22 (0.15, 0.41)	
	20	0.01 (0.00, 0.05)	0.04 (0.01, 0.08)	0.06 (0.03, 0.11)	0.28 (0.16, 0.96)	0.21 (0.14, 0.41)	
	30	0.02 (0.01, 0.06)	0.05 (0.02, 0.10)	0.07 (0.03, 0.14)	0.28 (0.15, 0.97)	0.21 (0.14, 0.41)	
	40	0.02 (0.01, 0.06)	0.05 (0.02, 0.11)	0.17 (0.08, 0.31)	0.26 (0.13, 0.98)	0.21 (0.14, 0.42)	
	50	0.02 (0.00, 0.08)	0.15 (0.07, 0.29)	0.22 (0.10, 0.42)	0.25 (0.12, 1.01)	0.19 (0.12, 0.41)	
	60	0.14 (0.05, 0.29)	0.21 (0.09, 0.42)	0.25 (0.11, 0.55)	0.25 (0.11, 1.08)	0.16 (0.09, 0.41)	
Hispanic ^b	70	0.09 (0.02, 0.27)	0.13 (0.03, 0.45)	- (- , -)	0.13 (0.03, 1.14)	0.14 (0.06, 0.44)	
	80	0.07 (0.00, 0.51)	- (- , -)	- (- , -)	0.07 (0.00, 1.62)	0.09 (0.02, 0.57)	
	0	0.03 (0.03, 0.04)	0.05 (0.05, 0.06)	0.07 (0.06, 0.08)	0.60 (0.54, 0.69)	0.34 (0.31, 0.38)	
	10	0.02 (0.02, 0.03)	0.04 (0.04, 0.05)	0.07 (0.06, 0.08)	0.58 (0.52, 0.66)	0.34 (0.31, 0.38)	
	20	0.02 (0.02, 0.02)	0.05 (0.04, 0.05)	0.09 (0.08, 0.10)	0.56 (0.50, 0.65)	0.34 (0.31, 0.37)	
	30	0.03 (0.02, 0.03)	0.07 (0.06, 0.08)	0.15 (0.13, 0.16)	0.55 (0.48, 0.63)	0.34 (0.31, 0.37)	
	40	0.04 (0.04, 0.05)	0.12 (0.11, 0.14)	0.25 (0.23, 0.28)	0.53 (0.46, 0.61)	0.33 (0.30, 0.37)	
	50	0.08 (0.07, 0.09)	0.21 (0.19, 0.24)	0.37 (0.33, 0.42)	0.50 (0.43, 0.59)	0.32 (0.29, 0.36)	
-	60	0.14 (0.12, 0.17)	0.31 (0.27, 0.36)	0.40 (0.35, 0.46)	0.45 (0.38, 0.54)	0.29 (0.26, 0.33)	
	70	0.19 (0.16, 0.23)	0.30 (0.25, 0.36)	- (- , -)	0.35 (0.28, 0.45)	0.23 (0.20, 0.27)	
	80	0.15 (0.10, 0.20)	- (- , -)	- (- , -)	0.22 (0.14, 0.35)	0.15 (0.12, 0.21)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table III-9 - continued

BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.03 (0.03, 0.04)	0.05 (0.05, 0.06)	0.08 (0.07, 0.08)	0.53 (0.52, 0.55)	0.39 (0.38, 0.39)	
	10	0.02 (0.02, 0.02)	0.04 (0.04, 0.05)	0.08 (0.07, 0.08)	0.51 (0.49, 0.52)	0.38 (0.38, 0.39)	
	20	0.02 (0.02, 0.03)	0.05 (0.05, 0.06)	0.10 (0.09, 0.10)	0.49 (0.47, 0.50)	0.38 (0.37, 0.39)	
	30	0.03 (0.03, 0.03)	0.07 (0.07, 0.08)	0.14 (0.14, 0.15)	0.47 (0.45, 0.48)	0.38 (0.37, 0.38)	
	40	0.04 (0.04, 0.04)	0.11 (0.11, 0.12)	0.22 (0.21, 0.23)	0.44 (0.42, 0.45)	0.37 (0.36, 0.37)	
	50	0.07 (0.07, 0.07)	0.18 (0.17, 0.19)	0.31 (0.30, 0.32)	0.40 (0.39, 0.42)	0.35 (0.34, 0.36)	
	60	0.11 (0.11, 0.12)	0.25 (0.24, 0.26)	0.33 (0.32, 0.34)	0.35 (0.34, 0.36)	0.31 (0.30, 0.32)	
	70	0.15 (0.14, 0.16)	0.24 (0.23, 0.25)	- (- , -)	0.26 (0.25, 0.28)	0.24 (0.23, 0.24)	
	80	0.12 (0.11, 0.13)	- (- , -)	- (- , -)	0.15 (0.14, 0.16)	0.14 (0.14, 0.15)	
White	0	0.03 (0.03, 0.04)	0.06 (0.05, 0.06)	0.08 (0.08, 0.09)	0.59 (0.57, 0.60)	0.42 (0.41, 0.43)	
	10	0.02 (0.02, 0.03)	0.05 (0.05, 0.05)	0.08 (0.08, 0.09)	0.56 (0.54, 0.57)	0.42 (0.41, 0.42)	
	20	0.03 (0.02, 0.03)	0.06 (0.06, 0.07)	0.11 (0.10, 0.11)	0.54 (0.52, 0.55)	0.41 (0.41, 0.42)	
	30	0.03 (0.03, 0.04)	0.08 (0.08, 0.09)	0.16 (0.15, 0.16)	0.51 (0.50, 0.53)	0.41 (0.40, 0.41)	
	40	0.05 (0.04, 0.05)	0.12 (0.12, 0.13)	0.24 (0.23, 0.25)	0.48 (0.46, 0.50)	0.40 (0.39, 0.40)	
	50	0.08 (0.07, 0.08)	0.20 (0.19, 0.21)	0.34 (0.33, 0.36)	0.44 (0.43, 0.46)	0.38 (0.37, 0.38)	
	60	0.13 (0.12, 0.13)	0.28 (0.27, 0.29)	0.36 (0.35, 0.38)	0.38 (0.36, 0.39)	0.33 (0.33, 0.34)	
	70	0.17 (0.16, 0.18)	0.26 (0.25, 0.28)	- (- , -)	0.28 (0.27, 0.29)	0.25 (0.25, 0.26)	
	80	0.13 (0.12, 0.14)	- (- , -)	- (- , -)	0.15 (0.14, 0.16)	0.15 (0.14, 0.15)	
Black	0	0.03 (0.02, 0.04)	0.04 (0.03, 0.05)	0.05 (0.04, 0.06)	0.30 (0.27, 0.34)	0.20 (0.19, 0.21)	
	10	0.01 (0.01, 0.02)	0.03 (0.02, 0.03)	0.05 (0.04, 0.06)	0.28 (0.25, 0.32)	0.20 (0.19, 0.21)	
	20	0.01 (0.01, 0.02)	0.03 (0.03, 0.04)	0.06 (0.05, 0.07)	0.27 (0.24, 0.31)	0.19 (0.18, 0.21)	
	30	0.02 (0.02, 0.03)	0.05 (0.04, 0.05)	0.09 (0.07, 0.10)	0.26 (0.23, 0.29)	0.19 (0.18, 0.20)	
	40	0.02 (0.02, 0.03)	0.07 (0.06, 0.08)	0.12 (0.10, 0.14)	0.24 (0.21, 0.28)	0.19 (0.17, 0.20)	
	50	0.04 (0.04, 0.06)	0.10 (0.08, 0.12)	0.16 (0.14, 0.19)	0.23 (0.20, 0.26)	0.18 (0.17, 0.19)	
	60	0.06 (0.05, 0.08)	0.13 (0.10, 0.15)	0.18 (0.15, 0.21)	0.20 (0.17, 0.24)	0.16 (0.15, 0.18)	
	70	0.08 (0.06, 0.10)	0.14 (0.12, 0.18)	- (- , -)	0.17 (0.13, 0.21)	0.13 (0.12, 0.15)	
	80	0.09 (0.07, 0.13)	- (- , -)	- (- , -)	0.13 (0.09, 0.17)	0.09 (0.08, 0.11)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table III-9 - continued

BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific Islander	0	0.03 (0.02, 0.04)	0.04 (0.03, 0.06)	0.06 (0.05, 0.07)	0.34 (0.29, 0.40)	0.20 (0.17, 0.26)	
	10	0.02 (0.01, 0.02)	0.03 (0.02, 0.04)	0.05 (0.04, 0.06)	0.31 (0.27, 0.38)	0.20 (0.17, 0.26)	
	20	0.01 (0.01, 0.02)	0.03 (0.03, 0.04)	0.06 (0.05, 0.07)	0.29 (0.25, 0.36)	0.20 (0.16, 0.25)	
	30	0.02 (0.01, 0.03)	0.04 (0.03, 0.05)	0.08 (0.07, 0.09)	0.28 (0.24, 0.35)	0.20 (0.16, 0.25)	
	40	0.02 (0.02, 0.03)	0.06 (0.05, 0.07)	0.13 (0.11, 0.15)	0.26 (0.22, 0.33)	0.19 (0.16, 0.25)	
	50	0.04 (0.03, 0.05)	0.10 (0.08, 0.13)	0.17 (0.14, 0.20)	0.24 (0.20, 0.31)	0.18 (0.15, 0.24)	
	60	0.07 (0.05, 0.09)	0.14 (0.11, 0.17)	0.20 (0.16, 0.24)	0.21 (0.17, 0.28)	0.16 (0.13, 0.22)	
	70	0.07 (0.05, 0.09)	0.14 (0.11, 0.18)	- (- , -)	0.15 (0.12, 0.22)	0.12 (0.09, 0.18)	
American Indian/ Alaska Native ^a	80	0.08 (0.05, 0.12)	- (- , -)	- (- , -)	0.10 (0.06, 0.17)	0.09 (0.05, 0.16)	
	0	0.02 (0.01, 0.07)	0.04 (0.02, 0.09)	0.05 (0.02, 0.11)	0.14 (0.07, 0.62)	0.14 (0.09, 0.28)	
	10	0.02 (0.00, 0.04)	0.03 (0.01, 0.07)	0.04 (0.01, 0.09)	0.12 (0.05, 0.61)	0.14 (0.08, 0.28)	
	20	0.01 (0.00, 0.05)	0.02 (0.01, 0.07)	0.03 (0.01, 0.08)	0.10 (0.04, 0.60)	0.14 (0.08, 0.28)	
	30	0.01 (0.00, 0.04)	0.01 (0.00, 0.05)	0.02 (0.00, 0.06)	0.09 (0.03, 0.59)	0.14 (0.08, 0.28)	
	40	0.01 (0.00, 0.04)	0.01 (0.00, 0.05)	0.06 (0.02, 0.16)	0.08 (0.02, 0.60)	0.13 (0.08, 0.28)	
	50	0.00 (0.00, 0.04)	0.06 (0.01, 0.16)	0.08 (0.02, 0.22)	0.08 (0.02, 0.61)	0.13 (0.08, 0.29)	
	60	0.05 (0.01, 0.16)	0.08 (0.02, 0.22)	0.08 (0.02, 0.27)	0.08 (0.02, 0.64)	0.12 (0.06, 0.28)	
Hispanic ^b	70	0.03 (0.00, 0.16)	0.03 (0.00, 0.23)	- (- , -)	0.03 (0.00, 0.68)	0.09 (0.04, 0.28)	
	80	0.00 (0.00, 0.25)	- (- , -)	- (- , -)	0.00 (0.00, 0.88)	0.03 (0.00, 0.29)	
	0	0.03 (0.03, 0.04)	0.05 (0.04, 0.06)	0.07 (0.06, 0.08)	0.49 (0.45, 0.55)	0.29 (0.27, 0.32)	
	10	0.02 (0.01, 0.02)	0.03 (0.03, 0.04)	0.06 (0.05, 0.07)	0.46 (0.42, 0.52)	0.29 (0.26, 0.32)	
	20	0.02 (0.01, 0.02)	0.04 (0.04, 0.05)	0.07 (0.07, 0.08)	0.44 (0.40, 0.50)	0.28 (0.26, 0.31)	
	30	0.03 (0.02, 0.03)	0.06 (0.05, 0.07)	0.12 (0.11, 0.14)	0.43 (0.38, 0.49)	0.28 (0.26, 0.31)	
	40	0.03 (0.03, 0.04)	0.10 (0.08, 0.11)	0.19 (0.17, 0.21)	0.41 (0.36, 0.47)	0.28 (0.25, 0.31)	
	50	0.07 (0.06, 0.08)	0.16 (0.14, 0.18)	0.28 (0.25, 0.32)	0.38 (0.34, 0.44)	0.27 (0.24, 0.30)	
-	60	0.10 (0.08, 0.12)	0.22 (0.19, 0.26)	0.31 (0.27, 0.35)	0.32 (0.28, 0.38)	0.24 (0.22, 0.27)	
	70	0.14 (0.11, 0.16)	0.23 (0.19, 0.27)	- (- , -)	0.25 (0.21, 0.31)	0.19 (0.17, 0.23)	
	80	0.12 (0.09, 0.15)	- (- , -)	- (- , -)	0.14 (0.10, 0.21)	0.13 (0.10, 0.16)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table IV-14 - continued
FEMALE BREAST CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.06 (0.05, 0.06)	12.28 (12.22, 12.35)	2.89 (2.88, 2.91)	
	10	0.00 (0.00, 0.00)	0.06 (0.05, 0.06)	0.48 (0.47, 0.49)	12.42 (12.36, 12.49)	2.93 (2.91, 2.95)	
	20	0.05 (0.05, 0.06)	0.48 (0.47, 0.49)	1.89 (1.87, 1.91)	12.45 (12.38, 12.52)	2.94 (2.92, 2.95)	
	30	0.43 (0.42, 0.44)	1.84 (1.82, 1.86)	4.24 (4.21, 4.27)	12.46 (12.40, 12.53)	2.95 (2.93, 2.96)	
	40	1.43 (1.42, 1.45)	3.86 (3.83, 3.89)	7.04 (6.99, 7.08)	12.19 (12.12, 12.26)	2.92 (2.90, 2.93)	
	50	2.51 (2.49, 2.54)	5.79 (5.75, 5.84)	8.93 (8.87, 8.98)	11.12 (11.05, 11.19)	2.78 (2.76, 2.79)	
	60	3.51 (3.48, 3.55)	6.87 (6.81, 6.92)	8.76 (8.70, 8.83)	9.21 (9.15, 9.28)	2.47 (2.45, 2.48)	
	70	3.88 (3.83, 3.92)	6.07 (6.01, 6.13)	- (- , -)	6.59 (6.53, 6.65)	2.04 (2.02, 2.06)	
	80	3.04 (3.00, 3.09)	- (- , -)	- (- , -)	3.76 (3.71, 3.81)	1.55 (1.53, 1.57)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.05, 0.06)	12.80 (12.72, 12.87)	2.89 (2.87, 2.90)	
	10	0.00 (0.00, 0.00)	0.05 (0.05, 0.06)	0.48 (0.47, 0.49)	12.92 (12.84, 13.00)	2.92 (2.90, 2.93)	
	20	0.05 (0.05, 0.06)	0.48 (0.47, 0.49)	1.91 (1.88, 1.93)	12.95 (12.87, 13.03)	2.92 (2.91, 2.94)	
	30	0.43 (0.42, 0.44)	1.86 (1.84, 1.88)	4.34 (4.30, 4.37)	12.96 (12.89, 13.04)	2.93 (2.92, 2.95)	
	40	1.45 (1.43, 1.47)	3.96 (3.92, 3.99)	7.31 (7.25, 7.36)	12.69 (12.61, 12.76)	2.91 (2.89, 2.92)	
	50	2.59 (2.56, 2.61)	6.04 (5.99, 6.09)	9.35 (9.28, 9.42)	11.60 (11.52, 11.68)	2.78 (2.76, 2.80)	
	60	3.69 (3.65, 3.73)	7.22 (7.16, 7.29)	9.19 (9.12, 9.26)	9.63 (9.55, 9.70)	2.48 (2.46, 2.50)	
	70	4.08 (4.03, 4.13)	6.35 (6.29, 6.41)	- (- , -)	6.86 (6.79, 6.93)	2.05 (2.03, 2.07)	
	80	3.16 (3.11, 3.21)	- (- , -)	- (- , -)	3.87 (3.81, 3.93)	1.55 (1.53, 1.57)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.07 (0.06, 0.08)	10.10 (9.91, 10.29)	3.23 (3.18, 3.28)	
	10	0.00 (0.00, 0.01)	0.07 (0.06, 0.08)	0.53 (0.50, 0.56)	10.32 (10.13, 10.52)	3.30 (3.25, 3.35)	
	20	0.07 (0.06, 0.08)	0.53 (0.50, 0.56)	1.87 (1.82, 1.93)	10.35 (10.15, 10.55)	3.31 (3.26, 3.37)	
	30	0.46 (0.44, 0.49)	1.82 (1.76, 1.87)	4.04 (3.95, 4.13)	10.37 (10.17, 10.57)	3.33 (3.27, 3.38)	
	40	1.38 (1.34, 1.43)	3.65 (3.56, 3.73)	6.24 (6.11, 6.38)	10.11 (9.91, 10.31)	3.27 (3.22, 3.33)	
	50	2.38 (2.31, 2.46)	5.12 (5.00, 5.25)	7.52 (7.36, 7.70)	9.19 (8.99, 9.40)	3.06 (3.00, 3.11)	
	60	3.04 (2.93, 3.14)	5.70 (5.54, 5.87)	7.14 (6.95, 7.33)	7.55 (7.35, 7.76)	2.64 (2.58, 2.69)	
	70	3.26 (3.13, 3.39)	5.01 (4.84, 5.20)	- (- , -)	5.52 (5.32, 5.72)	2.17 (2.11, 2.23)	
	80	2.66 (2.51, 2.82)	- (- , -)	- (- , -)	3.43 (3.23, 3.63)	1.74 (1.68, 1.81)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table IV-14 - continued
FEMALE BREAST CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/Pacific Islander	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.04 (0.04, 0.05)	9.18 (8.95, 9.43)	1.74 (1.63, 1.87)	
	10	0.00 (0.00, 0.00)	0.05 (0.04, 0.05)	0.44 (0.41, 0.47)	9.26 (9.02, 9.51)	1.76 (1.64, 1.89)	
	20	0.04 (0.04, 0.05)	0.44 (0.41, 0.47)	1.74 (1.68, 1.80)	9.27 (9.04, 9.52)	1.76 (1.64, 1.89)	
	30	0.40 (0.37, 0.42)	1.70 (1.64, 1.76)	3.62 (3.52, 3.71)	9.26 (9.02, 9.51)	1.76 (1.64, 1.89)	
	40	1.31 (1.27, 1.36)	3.24 (3.16, 3.33)	5.47 (5.34, 5.61)	8.93 (8.69, 9.18)	1.73 (1.62, 1.87)	
	50	1.97 (1.90, 2.04)	4.25 (4.12, 4.37)	6.27 (6.11, 6.44)	7.78 (7.55, 8.03)	1.61 (1.50, 1.75)	
	60	2.37 (2.28, 2.47)	4.49 (4.34, 4.64)	5.70 (5.51, 5.90)	6.07 (5.84, 6.31)	1.37 (1.25, 1.50)	
	70	2.30 (2.19, 2.41)	3.62 (3.45, 3.80)	- (-, -)	4.01 (3.80, 4.25)	1.08 (0.96, 1.22)	
	80	1.61 (1.47, 1.75)	- (-, -)	- (-, -)	2.09 (1.88, 2.33)	0.81 (0.68, 0.97)	
American Indian/Alaska Native ^a	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.03 (0.01, 0.09)	6.49 (5.81, 7.40)	1.94 (1.69, 2.26)	
	10	0.00 (0.00, 0.02)	0.03 (0.01, 0.07)	0.25 (0.18, 0.34)	6.58 (5.89, 7.50)	1.97 (1.71, 2.29)	
	20	0.03 (0.01, 0.07)	0.25 (0.18, 0.34)	1.08 (0.92, 1.25)	6.61 (5.92, 7.54)	1.98 (1.72, 2.30)	
	30	0.22 (0.16, 0.31)	1.05 (0.90, 1.23)	2.50 (2.23, 2.81)	6.64 (5.94, 7.58)	1.99 (1.73, 2.32)	
	40	0.85 (0.71, 1.00)	2.32 (2.05, 2.62)	3.97 (3.55, 4.44)	6.55 (5.84, 7.49)	1.98 (1.72, 2.32)	
	50	1.53 (1.31, 1.78)	3.24 (2.84, 3.69)	4.84 (4.27, 5.46)	5.91 (5.20, 6.88)	1.93 (1.66, 2.27)	
	60	1.83 (1.51, 2.20)	3.54 (3.01, 4.14)	4.54 (3.87, 5.31)	4.69 (3.99, 5.68)	1.74 (1.46, 2.10)	
	70	1.99 (1.55, 2.51)	3.15 (2.51, 3.92)	- (-, -)	3.33 (2.65, 4.37)	1.56 (1.25, 1.96)	
	80	1.58 (1.05, 2.30)	- (-, -)	- (-, -)	1.82 (1.20, 2.99)	1.42 (1.06, 1.93)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.04, 0.05)	9.23 (9.01, 9.46)	2.08 (2.01, 2.16)	
	10	0.00 (0.00, 0.00)	0.05 (0.04, 0.05)	0.37 (0.35, 0.39)	9.32 (9.10, 9.55)	2.10 (2.03, 2.18)	
	20	0.04 (0.04, 0.05)	0.37 (0.35, 0.39)	1.45 (1.41, 1.49)	9.34 (9.12, 9.57)	2.11 (2.03, 2.19)	
	30	0.33 (0.31, 0.34)	1.41 (1.37, 1.45)	3.19 (3.11, 3.26)	9.33 (9.11, 9.57)	2.11 (2.04, 2.19)	
	40	1.09 (1.06, 1.13)	2.89 (2.82, 2.96)	5.10 (4.98, 5.21)	9.09 (8.86, 9.32)	2.08 (2.01, 2.16)	
	50	1.84 (1.78, 1.90)	4.10 (3.99, 4.21)	6.30 (6.14, 6.45)	8.19 (7.97, 8.43)	1.97 (1.89, 2.05)	
	60	2.38 (2.29, 2.47)	4.69 (4.55, 4.83)	6.07 (5.89, 6.26)	6.68 (6.46, 6.92)	1.73 (1.66, 1.81)	
	70	2.57 (2.46, 2.69)	4.11 (3.94, 4.28)	- (-, -)	4.79 (4.57, 5.03)	1.42 (1.34, 1.51)	
	80	1.98 (1.84, 2.13)	- (-, -)	- (-, -)	2.85 (2.62, 3.11)	1.12 (1.03, 1.22)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table V-7
CERVIX UTERI CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.04 (0.04, 0.04)	0.70 (0.69, 0.72)	0.24 (0.24, 0.24)	
	10	0.00 (0.00, 0.00)	0.04 (0.04, 0.04)	0.16 (0.15, 0.16)	0.71 (0.70, 0.73)	0.24 (0.24, 0.25)	
	20	0.04 (0.04, 0.04)	0.16 (0.15, 0.16)	0.30 (0.29, 0.31)	0.71 (0.70, 0.73)	0.24 (0.24, 0.25)	
	30	0.12 (0.12, 0.12)	0.26 (0.26, 0.27)	0.40 (0.39, 0.40)	0.68 (0.66, 0.69)	0.24 (0.24, 0.24)	
	40	0.15 (0.14, 0.15)	0.28 (0.27, 0.29)	0.40 (0.39, 0.41)	0.56 (0.55, 0.58)	0.22 (0.22, 0.23)	
	50	0.14 (0.13, 0.14)	0.26 (0.25, 0.27)	0.36 (0.35, 0.37)	0.43 (0.41, 0.44)	0.19 (0.19, 0.20)	
	60	0.13 (0.13, 0.14)	0.24 (0.23, 0.25)	0.29 (0.28, 0.30)	0.30 (0.29, 0.32)	0.15 (0.15, 0.15)	
	70	0.11 (0.11, 0.12)	0.17 (0.17, 0.18)	- (- , -)	0.19 (0.18, 0.20)	0.11 (0.10, 0.11)	
	80	0.08 (0.07, 0.09)	- (- , -)	- (- , -)	0.10 (0.09, 0.11)	0.07 (0.07, 0.08)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.04 (0.04, 0.04)	0.67 (0.66, 0.69)	0.21 (0.21, 0.22)	
	10	0.00 (0.00, 0.00)	0.04 (0.04, 0.04)	0.17 (0.16, 0.18)	0.68 (0.66, 0.70)	0.22 (0.21, 0.22)	
	20	0.04 (0.04, 0.04)	0.17 (0.16, 0.18)	0.31 (0.30, 0.32)	0.68 (0.66, 0.70)	0.22 (0.21, 0.22)	
	30	0.13 (0.12, 0.13)	0.27 (0.26, 0.28)	0.39 (0.38, 0.40)	0.64 (0.63, 0.66)	0.21 (0.21, 0.22)	
	40	0.14 (0.14, 0.15)	0.27 (0.26, 0.28)	0.38 (0.37, 0.39)	0.52 (0.50, 0.53)	0.20 (0.19, 0.20)	
	50	0.13 (0.12, 0.13)	0.24 (0.23, 0.25)	0.33 (0.32, 0.34)	0.39 (0.37, 0.40)	0.17 (0.16, 0.17)	
	60	0.12 (0.11, 0.13)	0.21 (0.20, 0.22)	0.26 (0.25, 0.27)	0.27 (0.26, 0.28)	0.13 (0.13, 0.13)	
	70	0.10 (0.10, 0.11)	0.15 (0.15, 0.16)	- (- , -)	0.17 (0.16, 0.18)	0.09 (0.09, 0.10)	
	80	0.07 (0.06, 0.08)	- (- , -)	- (- , -)	0.09 (0.08, 0.10)	0.06 (0.06, 0.06)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.04 (0.04, 0.05)	0.92 (0.86, 0.97)	0.43 (0.41, 0.45)	
	10	0.00 (0.00, 0.00)	0.05 (0.04, 0.05)	0.16 (0.15, 0.18)	0.94 (0.88, 0.99)	0.44 (0.42, 0.46)	
	20	0.04 (0.04, 0.05)	0.16 (0.14, 0.18)	0.33 (0.31, 0.36)	0.94 (0.88, 0.99)	0.44 (0.42, 0.46)	
	30	0.12 (0.10, 0.13)	0.29 (0.27, 0.31)	0.45 (0.43, 0.49)	0.90 (0.85, 0.96)	0.44 (0.42, 0.46)	
	40	0.18 (0.16, 0.19)	0.34 (0.32, 0.37)	0.53 (0.49, 0.56)	0.80 (0.74, 0.85)	0.42 (0.40, 0.44)	
	50	0.17 (0.16, 0.19)	0.37 (0.33, 0.40)	0.52 (0.48, 0.57)	0.65 (0.60, 0.70)	0.38 (0.36, 0.40)	
	60	0.21 (0.18, 0.24)	0.38 (0.34, 0.42)	0.48 (0.43, 0.53)	0.52 (0.47, 0.57)	0.32 (0.30, 0.34)	
	70	0.20 (0.17, 0.24)	0.32 (0.28, 0.37)	- (- , -)	0.37 (0.32, 0.42)	0.25 (0.24, 0.27)	
	80	0.18 (0.14, 0.22)	- (- , -)	- (- , -)	0.24 (0.19, 0.30)	0.19 (0.17, 0.21)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table V-7 - continued
CERVIX UTERI CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.76 (0.69, 0.84)	0.32 (0.28, 0.39)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.07 (0.06, 0.08)	0.76 (0.70, 0.84)	0.32 (0.28, 0.39)	
Islander	20	0.01 (0.01, 0.02)	0.07 (0.06, 0.08)	0.20 (0.18, 0.23)	0.76 (0.70, 0.85)	0.32 (0.28, 0.39)	
	30	0.06 (0.05, 0.07)	0.19 (0.17, 0.21)	0.34 (0.32, 0.37)	0.75 (0.69, 0.83)	0.32 (0.28, 0.39)	
	40	0.13 (0.12, 0.15)	0.29 (0.26, 0.31)	0.45 (0.42, 0.50)	0.70 (0.64, 0.78)	0.31 (0.27, 0.38)	
	50	0.16 (0.14, 0.18)	0.33 (0.29, 0.36)	0.46 (0.42, 0.51)	0.57 (0.51, 0.65)	0.30 (0.25, 0.36)	
	60	0.17 (0.15, 0.20)	0.32 (0.28, 0.36)	0.40 (0.35, 0.45)	0.42 (0.37, 0.50)	0.26 (0.22, 0.33)	
	70	0.15 (0.12, 0.18)	0.24 (0.20, 0.28)	- (-, -)	0.27 (0.22, 0.34)	0.22 (0.17, 0.29)	
	80	0.10 (0.07, 0.14)	- (-, -)	- (-, -)	0.14 (0.10, 0.22)	0.17 (0.12, 0.25)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.03 (0.01, 0.09)	0.57 (0.39, 1.08)	0.38 (0.29, 0.56)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.03 (0.01, 0.07)	0.11 (0.06, 0.18)	0.58 (0.40, 1.10)	0.39 (0.29, 0.56)	
	20	0.03 (0.01, 0.07)	0.11 (0.06, 0.18)	0.20 (0.14, 0.29)	0.58 (0.40, 1.10)	0.39 (0.29, 0.57)	
	30	0.08 (0.04, 0.13)	0.17 (0.11, 0.25)	0.25 (0.17, 0.36)	0.56 (0.37, 1.08)	0.39 (0.29, 0.57)	
	40	0.10 (0.06, 0.16)	0.18 (0.11, 0.27)	0.25 (0.16, 0.38)	0.49 (0.31, 1.02)	0.37 (0.27, 0.55)	
	50	0.08 (0.04, 0.16)	0.16 (0.08, 0.28)	0.31 (0.18, 0.52)	0.40 (0.23, 0.95)	0.33 (0.23, 0.51)	
	60	0.07 (0.02, 0.18)	0.24 (0.12, 0.45)	0.33 (0.17, 0.62)	0.34 (0.17, 0.92)	0.28 (0.18, 0.48)	
	70	0.19 (0.07, 0.41)	0.30 (0.13, 0.61)	- (-, -)	0.30 (0.13, 0.97)	0.26 (0.16, 0.49)	
	80	0.14 (0.03, 0.47)	- (-, -)	- (-, -)	0.15 (0.03, 1.02)	0.21 (0.09, 0.51)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.04 (0.03, 0.05)	1.19 (1.13, 1.26)	0.35 (0.33, 0.39)	
	10	0.00 (0.00, 0.00)	0.04 (0.03, 0.05)	0.18 (0.17, 0.20)	1.20 (1.14, 1.28)	0.36 (0.33, 0.39)	
	20	0.04 (0.03, 0.04)	0.18 (0.17, 0.19)	0.40 (0.38, 0.42)	1.20 (1.14, 1.28)	0.36 (0.33, 0.39)	
	30	0.14 (0.13, 0.16)	0.37 (0.35, 0.39)	0.61 (0.58, 0.64)	1.17 (1.10, 1.24)	0.35 (0.33, 0.39)	
	40	0.22 (0.21, 0.24)	0.46 (0.44, 0.49)	0.71 (0.67, 0.75)	1.03 (0.97, 1.10)	0.34 (0.31, 0.37)	
	50	0.25 (0.22, 0.27)	0.50 (0.46, 0.54)	0.70 (0.65, 0.75)	0.82 (0.76, 0.89)	0.30 (0.27, 0.33)	
	60	0.26 (0.23, 0.29)	0.47 (0.43, 0.52)	0.57 (0.52, 0.63)	0.60 (0.54, 0.67)	0.25 (0.22, 0.28)	
	70	0.23 (0.20, 0.27)	0.34 (0.29, 0.39)	- (-, -)	0.37 (0.32, 0.43)	0.20 (0.17, 0.23)	
	80	0.13 (0.10, 0.18)	- (-, -)	- (-, -)	0.17 (0.13, 0.24)	0.14 (0.11, 0.18)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VI-15

COLON AND RECTUM CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer						Risk of Dying from Cancer	
		+10 yrs		+20 yrs		+30 yrs		Eventually	Eventually
		Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	5.42 (5.39, 5.46)	2.20 (2.19, 2.21)			
	10	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.07 (0.07, 0.08)	5.49 (5.46, 5.53)	2.23 (2.22, 2.24)			
	20	0.01 (0.01, 0.01)	0.07 (0.07, 0.08)	0.28 (0.27, 0.28)	5.51 (5.48, 5.55)	2.24 (2.23, 2.25)			
	30	0.06 (0.06, 0.06)	0.27 (0.26, 0.27)	0.87 (0.86, 0.88)	5.55 (5.51, 5.58)	2.26 (2.25, 2.27)			
	40	0.21 (0.21, 0.22)	0.82 (0.81, 0.83)	2.04 (2.03, 2.06)	5.56 (5.53, 5.60)	2.27 (2.26, 2.29)			
	50	0.63 (0.62, 0.64)	1.89 (1.87, 1.91)	3.70 (3.67, 3.73)	5.51 (5.48, 5.55)	2.28 (2.27, 2.29)			
	60	1.35 (1.33, 1.36)	3.28 (3.26, 3.31)	4.79 (4.75, 4.82)	5.22 (5.18, 5.25)	2.24 (2.23, 2.25)			
	70	2.26 (2.23, 2.28)	4.00 (3.97, 4.04)	- (- , -)	4.51 (4.47, 4.54)	2.08 (2.07, 2.09)			
	80	2.53 (2.50, 2.56)	- (- , -)	- (- , -)	3.25 (3.22, 3.29)	1.78 (1.77, 1.79)			
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	5.41 (5.37, 5.45)	2.18 (2.17, 2.19)			
	10	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.07 (0.07, 0.08)	5.47 (5.43, 5.51)	2.21 (2.20, 2.22)			
	20	0.01 (0.01, 0.02)	0.07 (0.07, 0.07)	0.27 (0.27, 0.28)	5.49 (5.45, 5.53)	2.22 (2.21, 2.23)			
	30	0.06 (0.06, 0.06)	0.26 (0.25, 0.26)	0.84 (0.83, 0.85)	5.52 (5.48, 5.56)	2.23 (2.22, 2.24)			
	40	0.20 (0.20, 0.21)	0.79 (0.78, 0.80)	2.00 (1.98, 2.02)	5.53 (5.49, 5.57)	2.25 (2.23, 2.26)			
	50	0.60 (0.59, 0.61)	1.85 (1.83, 1.87)	3.68 (3.65, 3.71)	5.48 (5.44, 5.52)	2.25 (2.24, 2.27)			
	60	1.33 (1.31, 1.34)	3.27 (3.24, 3.30)	4.78 (4.74, 4.82)	5.19 (5.15, 5.23)	2.21 (2.20, 2.22)			
	70	2.26 (2.23, 2.28)	4.01 (3.97, 4.04)	- (- , -)	4.49 (4.45, 4.53)	2.05 (2.04, 2.07)			
	80	2.54 (2.50, 2.57)	- (- , -)	- (- , -)	3.23 (3.19, 3.28)	1.75 (1.74, 1.77)			
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	5.24 (5.13, 5.35)	2.42 (2.39, 2.46)			
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.08 (0.07, 0.08)	5.37 (5.26, 5.48)	2.48 (2.45, 2.52)			
	20	0.01 (0.01, 0.01)	0.07 (0.07, 0.08)	0.33 (0.31, 0.34)	5.40 (5.29, 5.51)	2.50 (2.46, 2.53)			
	30	0.07 (0.06, 0.07)	0.32 (0.31, 0.34)	1.12 (1.08, 1.15)	5.48 (5.37, 5.59)	2.53 (2.50, 2.57)			
	40	0.26 (0.25, 0.28)	1.08 (1.04, 1.11)	2.50 (2.43, 2.56)	5.55 (5.43, 5.66)	2.57 (2.54, 2.61)			
	50	0.86 (0.83, 0.89)	2.35 (2.29, 2.42)	4.09 (3.99, 4.18)	5.57 (5.45, 5.70)	2.61 (2.57, 2.65)			
	60	1.69 (1.63, 1.74)	3.64 (3.54, 3.73)	4.92 (4.80, 5.04)	5.31 (5.18, 5.44)	2.59 (2.54, 2.63)			
	70	2.47 (2.39, 2.56)	4.10 (3.98, 4.22)	- (- , -)	4.59 (4.46, 4.73)	2.44 (2.39, 2.48)			
	80	2.64 (2.52, 2.77)	- (- , -)	- (- , -)	3.45 (3.29, 3.62)	2.12 (2.07, 2.18)			

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VI-15 - continued
COLON AND RECTUM CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	5.58 (5.41, 5.76)	2.17 (2.05, 2.29)	
Islander	10	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.08 (0.07, 0.09)	5.63 (5.46, 5.81)	2.19 (2.07, 2.31)	
	20	0.01 (0.01, 0.02)	0.08 (0.07, 0.08)	0.27 (0.25, 0.28)	5.64 (5.47, 5.83)	2.19 (2.08, 2.32)	
	30	0.06 (0.06, 0.07)	0.26 (0.24, 0.27)	0.79 (0.76, 0.82)	5.66 (5.48, 5.84)	2.20 (2.08, 2.33)	
	40	0.19 (0.18, 0.21)	0.73 (0.70, 0.76)	1.77 (1.71, 1.83)	5.63 (5.46, 5.81)	2.19 (2.08, 2.32)	
	50	0.55 (0.52, 0.57)	1.60 (1.55, 1.66)	3.25 (3.16, 3.34)	5.52 (5.34, 5.71)	2.17 (2.05, 2.30)	
	60	1.10 (1.05, 1.14)	2.81 (2.72, 2.90)	4.39 (4.26, 4.52)	5.16 (4.98, 5.36)	2.12 (2.00, 2.25)	
	70	1.87 (1.80, 1.95)	3.60 (3.47, 3.73)	- (- , -)	4.45 (4.27, 4.65)	1.99 (1.86, 2.14)	
	80	2.18 (2.06, 2.30)	- (- , -)	- (- , -)	3.26 (3.05, 3.48)	1.79 (1.64, 1.96)	
American	0	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.02 (0.01, 0.05)	4.27 (3.80, 4.86)	1.86 (1.66, 2.09)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.01)	0.02 (0.01, 0.04)	0.09 (0.06, 0.13)	4.34 (3.86, 4.95)	1.89 (1.69, 2.12)	
	20	0.02 (0.01, 0.04)	0.09 (0.06, 0.13)	0.29 (0.23, 0.36)	4.37 (3.89, 4.98)	1.91 (1.71, 2.14)	
	30	0.08 (0.05, 0.11)	0.28 (0.22, 0.34)	0.79 (0.68, 0.92)	4.43 (3.94, 5.05)	1.94 (1.74, 2.18)	
	40	0.21 (0.16, 0.26)	0.74 (0.63, 0.86)	1.76 (1.55, 1.99)	4.47 (3.97, 5.11)	1.98 (1.77, 2.23)	
	50	0.55 (0.46, 0.66)	1.62 (1.41, 1.85)	3.11 (2.77, 3.49)	4.45 (3.93, 5.11)	2.02 (1.80, 2.28)	
	60	1.15 (0.96, 1.35)	2.76 (2.41, 3.14)	3.93 (3.45, 4.46)	4.19 (3.65, 4.89)	2.00 (1.77, 2.28)	
	70	1.90 (1.58, 2.25)	3.28 (2.78, 3.84)	- (- , -)	3.59 (3.02, 4.36)	1.89 (1.62, 2.21)	
	80	1.96 (1.50, 2.54)	- (- , -)	- (- , -)	2.41 (1.80, 3.34)	1.55 (1.23, 1.96)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	4.82 (4.69, 4.97)	1.92 (1.86, 1.98)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.06 (0.06, 0.07)	4.88 (4.74, 5.03)	1.94 (1.88, 2.00)	
	20	0.01 (0.01, 0.01)	0.06 (0.05, 0.06)	0.22 (0.21, 0.23)	4.90 (4.76, 5.05)	1.95 (1.89, 2.01)	
	30	0.05 (0.04, 0.05)	0.21 (0.20, 0.22)	0.70 (0.67, 0.72)	4.93 (4.78, 5.07)	1.96 (1.90, 2.02)	
	40	0.17 (0.16, 0.17)	0.65 (0.63, 0.68)	1.67 (1.62, 1.72)	4.93 (4.79, 5.08)	1.97 (1.91, 2.03)	
	50	0.50 (0.48, 0.52)	1.54 (1.49, 1.59)	3.05 (2.97, 3.14)	4.88 (4.73, 5.03)	1.97 (1.91, 2.04)	
	60	1.10 (1.06, 1.14)	2.69 (2.61, 2.77)	3.99 (3.88, 4.11)	4.61 (4.46, 4.77)	1.92 (1.86, 1.99)	
	70	1.79 (1.72, 1.86)	3.26 (3.14, 3.38)	- (- , -)	3.96 (3.80, 4.13)	1.78 (1.71, 1.85)	
	80	1.96 (1.85, 2.07)	- (- , -)	- (- , -)	2.89 (2.71, 3.09)	1.52 (1.44, 1.60)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VI-15 - continued

COLON AND RECTUM CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	5.65 (5.60, 5.70)	2.29 (2.27, 2.30)	
	10	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.08 (0.07, 0.08)	5.73 (5.67, 5.78)	2.32 (2.30, 2.34)	
	20	0.01 (0.01, 0.02)	0.07 (0.07, 0.08)	0.29 (0.29, 0.30)	5.76 (5.70, 5.81)	2.33 (2.32, 2.35)	
	30	0.06 (0.06, 0.06)	0.28 (0.28, 0.29)	0.97 (0.95, 0.98)	5.82 (5.77, 5.87)	2.36 (2.35, 2.38)	
	40	0.22 (0.22, 0.23)	0.92 (0.91, 0.93)	2.34 (2.31, 2.37)	5.86 (5.80, 5.91)	2.39 (2.37, 2.41)	
	50	0.72 (0.71, 0.74)	2.19 (2.17, 2.22)	4.20 (4.16, 4.24)	5.85 (5.79, 5.90)	2.41 (2.39, 2.43)	
	60	1.60 (1.57, 1.62)	3.77 (3.73, 3.82)	5.21 (5.16, 5.26)	5.56 (5.51, 5.62)	2.38 (2.36, 2.40)	
	70	2.62 (2.58, 2.66)	4.35 (4.30, 4.41)	- (- , -)	4.78 (4.72, 4.84)	2.20 (2.18, 2.22)	
	80	2.73 (2.68, 2.78)	- (- , -)	- (- , -)	3.39 (3.33, 3.46)	1.86 (1.84, 1.89)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	5.65 (5.60, 5.71)	2.28 (2.26, 2.29)	
	10	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.07 (0.07, 0.08)	5.72 (5.66, 5.78)	2.31 (2.29, 2.32)	
	20	0.01 (0.01, 0.02)	0.07 (0.07, 0.08)	0.29 (0.28, 0.29)	5.75 (5.69, 5.81)	2.32 (2.30, 2.34)	
	30	0.06 (0.06, 0.06)	0.27 (0.27, 0.28)	0.94 (0.92, 0.96)	5.80 (5.75, 5.86)	2.34 (2.33, 2.36)	
	40	0.22 (0.21, 0.23)	0.89 (0.88, 0.91)	2.31 (2.27, 2.34)	5.84 (5.78, 5.90)	2.37 (2.35, 2.39)	
	50	0.70 (0.69, 0.71)	2.16 (2.13, 2.19)	4.18 (4.14, 4.23)	5.83 (5.77, 5.89)	2.39 (2.37, 2.41)	
	60	1.58 (1.55, 1.61)	3.76 (3.72, 3.81)	5.21 (5.15, 5.27)	5.54 (5.48, 5.60)	2.35 (2.33, 2.37)	
	70	2.62 (2.58, 2.66)	4.35 (4.30, 4.41)	- (- , -)	4.75 (4.69, 4.82)	2.17 (2.15, 2.19)	
	80	2.73 (2.68, 2.79)	- (- , -)	- (- , -)	3.36 (3.30, 3.43)	1.84 (1.81, 1.86)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	5.20 (5.04, 5.36)	2.41 (2.36, 2.47)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.08 (0.07, 0.09)	5.34 (5.18, 5.50)	2.48 (2.43, 2.54)	
	20	0.01 (0.01, 0.01)	0.08 (0.07, 0.09)	0.33 (0.31, 0.36)	5.38 (5.22, 5.55)	2.50 (2.45, 2.56)	
	30	0.07 (0.06, 0.08)	0.33 (0.31, 0.36)	1.20 (1.15, 1.25)	5.51 (5.34, 5.68)	2.56 (2.50, 2.61)	
	40	0.27 (0.25, 0.29)	1.17 (1.11, 1.22)	2.70 (2.61, 2.80)	5.62 (5.45, 5.79)	2.61 (2.56, 2.67)	
	50	0.95 (0.91, 1.00)	2.60 (2.50, 2.70)	4.39 (4.25, 4.54)	5.71 (5.53, 5.90)	2.67 (2.62, 2.73)	
	60	1.91 (1.82, 2.01)	4.00 (3.85, 4.15)	5.19 (5.01, 5.38)	5.54 (5.34, 5.74)	2.70 (2.63, 2.76)	
	70	2.82 (2.68, 2.97)	4.43 (4.24, 4.64)	- (- , -)	4.90 (4.67, 5.13)	2.56 (2.49, 2.64)	
	80	2.98 (2.77, 3.21)	- (- , -)	- (- , -)	3.84 (3.56, 4.15)	2.23 (2.13, 2.33)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VI-15 - continued

COLON AND RECTUM CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific Islander	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.03)	5.96 (5.72, 6.22)	2.30 (2.14, 2.48)	
	10	0.00 (0.00, 0.01)	0.02 (0.01, 0.03)	0.08 (0.07, 0.10)	6.02 (5.78, 6.28)	2.32 (2.16, 2.50)	
	20	0.02 (0.01, 0.02)	0.08 (0.07, 0.09)	0.28 (0.25, 0.30)	6.04 (5.79, 6.30)	2.33 (2.16, 2.51)	
	30	0.06 (0.05, 0.08)	0.26 (0.24, 0.28)	0.87 (0.83, 0.92)	6.06 (5.81, 6.32)	2.34 (2.17, 2.52)	
	40	0.20 (0.18, 0.22)	0.82 (0.77, 0.86)	2.06 (1.97, 2.15)	6.04 (5.80, 6.31)	2.34 (2.17, 2.52)	
	50	0.63 (0.59, 0.67)	1.90 (1.81, 1.99)	3.88 (3.73, 4.04)	5.96 (5.71, 6.23)	2.32 (2.15, 2.51)	
	60	1.33 (1.25, 1.41)	3.41 (3.26, 3.56)	4.96 (4.75, 5.17)	5.59 (5.33, 5.86)	2.27 (2.10, 2.47)	
	70	2.33 (2.20, 2.47)	4.07 (3.86, 4.28)	- (-, -)	4.77 (4.51, 5.06)	2.12 (1.94, 2.33)	
American Indian/ Alaska Native ^a	80	2.33 (2.15, 2.52)	- (-, -)	- (-, -)	3.28 (2.99, 3.61)	1.90 (1.67, 2.16)	
	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.01 (0.00, 0.05)	3.74 (3.17, 4.66)	1.94 (1.66, 2.31)	
	10	0.00 (0.00, 0.02)	0.01 (0.00, 0.04)	0.09 (0.05, 0.15)	3.82 (3.23, 4.75)	1.98 (1.69, 2.36)	
	20	0.01 (0.00, 0.04)	0.09 (0.05, 0.15)	0.28 (0.20, 0.37)	3.86 (3.27, 4.80)	2.00 (1.71, 2.39)	
	30	0.09 (0.05, 0.15)	0.27 (0.20, 0.37)	0.79 (0.64, 0.97)	3.95 (3.34, 4.91)	2.05 (1.75, 2.45)	
	40	0.20 (0.14, 0.28)	0.73 (0.58, 0.91)	1.88 (1.57, 2.24)	4.00 (3.38, 5.01)	2.11 (1.80, 2.52)	
	50	0.56 (0.43, 0.72)	1.78 (1.46, 2.14)	3.03 (2.55, 3.59)	4.01 (3.36, 5.06)	2.19 (1.86, 2.62)	
	60	1.33 (1.04, 1.67)	2.70 (2.21, 3.28)	3.74 (3.06, 4.54)	3.78 (3.09, 4.91)	2.18 (1.82, 2.66)	
Hispanic ^b	70	1.66 (1.24, 2.18)	2.91 (2.22, 3.77)	- (-, -)	2.96 (2.25, 4.24)	2.06 (1.64, 2.63)	
	80	1.89 (1.19, 2.90)	- (-, -)	- (-, -)	1.97 (1.23, 3.72)	1.78 (1.24, 2.59)	
	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	5.08 (4.88, 5.30)	2.10 (2.01, 2.20)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.06 (0.05, 0.07)	5.15 (4.94, 5.37)	2.13 (2.04, 2.23)	
	20	0.01 (0.01, 0.01)	0.06 (0.05, 0.07)	0.23 (0.22, 0.25)	5.18 (4.97, 5.40)	2.14 (2.05, 2.24)	
	30	0.05 (0.04, 0.05)	0.22 (0.21, 0.24)	0.78 (0.75, 0.82)	5.22 (5.02, 5.45)	2.16 (2.07, 2.27)	
	40	0.18 (0.16, 0.19)	0.74 (0.71, 0.78)	1.98 (1.90, 2.06)	5.25 (5.04, 5.48)	2.18 (2.09, 2.29)	
	50	0.59 (0.55, 0.62)	1.86 (1.78, 1.94)	3.63 (3.50, 3.76)	5.23 (5.02, 5.47)	2.20 (2.10, 2.30)	
-	60	1.36 (1.29, 1.43)	3.26 (3.13, 3.39)	4.53 (4.35, 4.72)	4.98 (4.75, 5.22)	2.16 (2.05, 2.27)	
	70	2.20 (2.09, 2.32)	3.68 (3.50, 3.87)	- (-, -)	4.20 (3.96, 4.47)	2.00 (1.88, 2.12)	
	80	2.12 (1.94, 2.31)	- (-, -)	- (-, -)	2.86 (2.58, 3.19)	1.70 (1.55, 1.86)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VI-15 - continued
COLON AND RECTUM CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	5.23 (5.18, 5.27)	2.13 (2.12, 2.14)	
	10	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.07 (0.07, 0.08)	5.28 (5.24, 5.33)	2.16 (2.14, 2.17)	
	20	0.01 (0.01, 0.02)	0.07 (0.07, 0.07)	0.27 (0.26, 0.27)	5.30 (5.25, 5.34)	2.16 (2.15, 2.18)	
	30	0.06 (0.05, 0.06)	0.25 (0.25, 0.26)	0.77 (0.76, 0.79)	5.31 (5.26, 5.36)	2.17 (2.16, 2.19)	
	40	0.20 (0.19, 0.20)	0.72 (0.71, 0.74)	1.77 (1.74, 1.79)	5.30 (5.25, 5.35)	2.18 (2.16, 2.19)	
	50	0.53 (0.52, 0.55)	1.60 (1.58, 1.62)	3.26 (3.22, 3.29)	5.21 (5.17, 5.26)	2.17 (2.16, 2.19)	
	60	1.12 (1.10, 1.14)	2.86 (2.83, 2.90)	4.42 (4.38, 4.47)	4.92 (4.87, 4.97)	2.12 (2.11, 2.14)	
	70	1.97 (1.94, 1.99)	3.73 (3.69, 3.78)	- (- , -)	4.30 (4.25, 4.34)	1.99 (1.98, 2.01)	
	80	2.40 (2.36, 2.44)	- (- , -)	- (- , -)	3.17 (3.12, 3.22)	1.73 (1.71, 1.75)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	5.19 (5.14, 5.24)	2.10 (2.09, 2.12)	
	10	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.07 (0.07, 0.08)	5.24 (5.19, 5.29)	2.12 (2.11, 2.14)	
	20	0.01 (0.01, 0.02)	0.07 (0.07, 0.07)	0.26 (0.25, 0.26)	5.25 (5.20, 5.30)	2.13 (2.11, 2.14)	
	30	0.06 (0.05, 0.06)	0.24 (0.24, 0.25)	0.74 (0.72, 0.75)	5.26 (5.21, 5.31)	2.14 (2.12, 2.15)	
	40	0.19 (0.18, 0.20)	0.69 (0.67, 0.70)	1.71 (1.68, 1.73)	5.25 (5.20, 5.30)	2.14 (2.13, 2.16)	
	50	0.51 (0.49, 0.52)	1.55 (1.52, 1.57)	3.21 (3.17, 3.25)	5.17 (5.11, 5.22)	2.14 (2.12, 2.15)	
	60	1.09 (1.07, 1.12)	2.84 (2.80, 2.88)	4.40 (4.36, 4.45)	4.89 (4.83, 4.94)	2.09 (2.07, 2.11)	
	70	1.97 (1.94, 2.00)	3.74 (3.69, 3.78)	- (- , -)	4.28 (4.23, 4.33)	1.96 (1.95, 1.98)	
	80	2.41 (2.37, 2.45)	- (- , -)	- (- , -)	3.15 (3.10, 3.20)	1.71 (1.69, 1.72)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	5.31 (5.16, 5.46)	2.44 (2.39, 2.49)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.07 (0.06, 0.08)	5.42 (5.27, 5.58)	2.49 (2.44, 2.54)	
	20	0.01 (0.01, 0.02)	0.07 (0.06, 0.08)	0.32 (0.30, 0.35)	5.44 (5.28, 5.59)	2.50 (2.45, 2.55)	
	30	0.06 (0.05, 0.07)	0.32 (0.29, 0.34)	1.05 (1.00, 1.09)	5.47 (5.32, 5.63)	2.52 (2.47, 2.57)	
	40	0.26 (0.24, 0.28)	1.00 (0.96, 1.05)	2.31 (2.23, 2.40)	5.50 (5.35, 5.66)	2.54 (2.49, 2.59)	
	50	0.78 (0.74, 0.82)	2.15 (2.06, 2.23)	3.83 (3.71, 3.95)	5.47 (5.31, 5.64)	2.55 (2.50, 2.61)	
	60	1.50 (1.43, 1.58)	3.35 (3.22, 3.47)	4.70 (4.55, 4.86)	5.15 (4.98, 5.32)	2.50 (2.45, 2.56)	
	70	2.22 (2.12, 2.33)	3.86 (3.71, 4.02)	- (- , -)	4.40 (4.22, 4.58)	2.35 (2.29, 2.41)	
	80	2.45 (2.31, 2.60)	- (- , -)	- (- , -)	3.25 (3.06, 3.45)	2.07 (2.00, 2.14)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VI-15 - continued
COLON AND RECTUM CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/Pacific Islander	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	5.27 (5.03, 5.54)	2.06 (1.89, 2.24)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.07 (0.06, 0.08)	5.31 (5.07, 5.58)	2.07 (1.91, 2.26)	
	20	0.01 (0.01, 0.02)	0.07 (0.06, 0.08)	0.26 (0.24, 0.28)	5.32 (5.08, 5.59)	2.08 (1.91, 2.26)	
	30	0.06 (0.05, 0.07)	0.25 (0.23, 0.27)	0.72 (0.68, 0.76)	5.33 (5.08, 5.60)	2.08 (1.91, 2.26)	
	40	0.19 (0.17, 0.21)	0.66 (0.62, 0.70)	1.52 (1.45, 1.60)	5.29 (5.04, 5.56)	2.07 (1.91, 2.26)	
	50	0.47 (0.44, 0.51)	1.35 (1.28, 1.42)	2.73 (2.62, 2.85)	5.16 (4.91, 5.43)	2.05 (1.88, 2.23)	
	60	0.90 (0.84, 0.96)	2.32 (2.21, 2.43)	3.93 (3.76, 4.10)	4.83 (4.57, 5.10)	1.99 (1.82, 2.18)	
	70	1.53 (1.44, 1.62)	3.25 (3.08, 3.42)	- (-, -)	4.21 (3.95, 4.50)	1.89 (1.71, 2.09)	
	80	2.07 (1.92, 2.23)	- (-, -)	- (-, -)	3.24 (2.95, 3.56)	1.71 (1.51, 1.94)	
American Indian/Alaska Native ^a	0	0.00 (0.00, 0.04)	0.01 (0.00, 0.05)	0.03 (0.01, 0.09)	4.74 (4.03, 5.68)	1.80 (1.53, 2.13)	
	10	0.01 (0.00, 0.03)	0.03 (0.01, 0.08)	0.10 (0.06, 0.16)	4.80 (4.09, 5.76)	1.83 (1.56, 2.16)	
	20	0.03 (0.01, 0.07)	0.09 (0.05, 0.15)	0.30 (0.22, 0.40)	4.82 (4.10, 5.79)	1.84 (1.57, 2.18)	
	30	0.07 (0.03, 0.12)	0.28 (0.20, 0.37)	0.80 (0.64, 0.98)	4.84 (4.12, 5.81)	1.85 (1.58, 2.20)	
	40	0.21 (0.15, 0.30)	0.74 (0.59, 0.92)	1.64 (1.37, 1.96)	4.86 (4.13, 5.85)	1.88 (1.60, 2.23)	
	50	0.55 (0.42, 0.70)	1.47 (1.20, 1.78)	3.18 (2.69, 3.73)	4.79 (4.04, 5.81)	1.89 (1.60, 2.25)	
	60	0.98 (0.76, 1.25)	2.79 (2.31, 3.35)	4.09 (3.42, 4.85)	4.51 (3.73, 5.57)	1.87 (1.57, 2.26)	
	70	2.09 (1.65, 2.62)	3.58 (2.90, 4.39)	- (-, -)	4.07 (3.25, 5.24)	1.77 (1.44, 2.20)	
	80	2.01 (1.42, 2.79)	- (-, -)	- (-, -)	2.67 (1.83, 4.04)	1.42 (1.03, 1.96)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	4.60 (4.41, 4.80)	1.77 (1.70, 1.85)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.06 (0.05, 0.07)	4.65 (4.45, 4.85)	1.79 (1.72, 1.87)	
	20	0.01 (0.01, 0.01)	0.06 (0.05, 0.07)	0.21 (0.20, 0.23)	4.65 (4.46, 4.86)	1.80 (1.72, 1.88)	
	30	0.05 (0.04, 0.06)	0.20 (0.19, 0.22)	0.61 (0.58, 0.65)	4.66 (4.47, 4.87)	1.80 (1.72, 1.88)	
	40	0.15 (0.14, 0.17)	0.57 (0.53, 0.60)	1.40 (1.33, 1.46)	4.64 (4.44, 4.85)	1.80 (1.72, 1.88)	
	50	0.42 (0.39, 0.45)	1.26 (1.20, 1.32)	2.56 (2.46, 2.66)	4.56 (4.36, 4.77)	1.80 (1.72, 1.88)	
	60	0.88 (0.82, 0.93)	2.22 (2.12, 2.32)	3.54 (3.40, 3.70)	4.30 (4.09, 4.51)	1.74 (1.66, 1.83)	
	70	1.48 (1.39, 1.56)	2.93 (2.79, 3.08)	- (-, -)	3.76 (3.55, 3.99)	1.63 (1.55, 1.73)	
	80	1.85 (1.71, 1.99)	- (-, -)	- (-, -)	2.90 (2.67, 3.16)	1.41 (1.31, 1.52)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VII-7

CORPUS AND UTERUS, NOS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	2.45 (2.42, 2.48)	0.51 (0.51, 0.52)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.06 (0.06, 0.07)	2.48 (2.45, 2.51)	0.52 (0.51, 0.53)	
	20	0.01 (0.01, 0.01)	0.06 (0.06, 0.07)	0.24 (0.24, 0.25)	2.49 (2.46, 2.52)	0.52 (0.51, 0.53)	
	30	0.05 (0.05, 0.06)	0.24 (0.23, 0.24)	0.75 (0.74, 0.77)	2.49 (2.46, 2.52)	0.52 (0.52, 0.53)	
	40	0.18 (0.18, 0.19)	0.71 (0.69, 0.72)	1.44 (1.42, 1.46)	2.46 (2.43, 2.49)	0.52 (0.52, 0.53)	
	50	0.53 (0.52, 0.54)	1.29 (1.27, 1.31)	1.92 (1.89, 1.94)	2.33 (2.30, 2.36)	0.52 (0.51, 0.53)	
	60	0.79 (0.78, 0.81)	1.46 (1.43, 1.48)	1.81 (1.78, 1.84)	1.89 (1.86, 1.92)	0.49 (0.49, 0.50)	
	70	0.75 (0.73, 0.76)	1.15 (1.12, 1.17)	- (- , -)	1.23 (1.21, 1.26)	0.41 (0.40, 0.42)	
	80	0.54 (0.52, 0.56)	- (- , -)	- (- , -)	0.66 (0.64, 0.68)	0.29 (0.28, 0.30)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	2.56 (2.53, 2.60)	0.49 (0.48, 0.50)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.06 (0.06, 0.07)	2.59 (2.55, 2.62)	0.49 (0.49, 0.50)	
	20	0.01 (0.01, 0.01)	0.06 (0.06, 0.07)	0.25 (0.24, 0.25)	2.59 (2.56, 2.63)	0.49 (0.49, 0.50)	
	30	0.05 (0.05, 0.06)	0.24 (0.23, 0.25)	0.78 (0.77, 0.80)	2.60 (2.56, 2.63)	0.50 (0.49, 0.50)	
	40	0.19 (0.18, 0.19)	0.74 (0.72, 0.75)	1.51 (1.48, 1.53)	2.56 (2.53, 2.60)	0.50 (0.49, 0.51)	
	50	0.56 (0.55, 0.57)	1.35 (1.32, 1.37)	2.01 (1.98, 2.04)	2.43 (2.39, 2.46)	0.49 (0.49, 0.50)	
	60	0.83 (0.81, 0.84)	1.52 (1.49, 1.55)	1.88 (1.85, 1.92)	1.96 (1.93, 1.99)	0.46 (0.46, 0.47)	
	70	0.78 (0.76, 0.80)	1.19 (1.17, 1.22)	- (- , -)	1.28 (1.25, 1.31)	0.39 (0.38, 0.40)	
	80	0.56 (0.54, 0.58)	- (- , -)	- (- , -)	0.67 (0.65, 0.70)	0.28 (0.27, 0.29)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	1.95 (1.86, 2.03)	0.75 (0.73, 0.78)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.05 (0.04, 0.06)	1.99 (1.90, 2.08)	0.77 (0.74, 0.80)	
	20	0.01 (0.01, 0.02)	0.05 (0.04, 0.06)	0.16 (0.15, 0.18)	1.99 (1.91, 2.09)	0.77 (0.74, 0.80)	
	30	0.04 (0.04, 0.05)	0.15 (0.14, 0.17)	0.50 (0.46, 0.53)	2.00 (1.91, 2.09)	0.78 (0.75, 0.80)	
	40	0.11 (0.10, 0.13)	0.46 (0.43, 0.49)	1.16 (1.10, 1.22)	1.99 (1.90, 2.08)	0.79 (0.76, 0.81)	
	50	0.36 (0.33, 0.39)	1.09 (1.03, 1.15)	1.66 (1.58, 1.74)	1.96 (1.87, 2.05)	0.80 (0.77, 0.83)	
	60	0.79 (0.74, 0.85)	1.41 (1.33, 1.50)	1.69 (1.60, 1.78)	1.74 (1.65, 1.84)	0.79 (0.76, 0.82)	
	70	0.74 (0.68, 0.81)	1.07 (0.99, 1.16)	- (- , -)	1.14 (1.06, 1.23)	0.63 (0.60, 0.66)	
	80	0.49 (0.43, 0.56)	- (- , -)	- (- , -)	0.59 (0.51, 0.67)	0.41 (0.38, 0.44)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VII-7 - continued

CORPUS AND UTERUS, NOS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	1.77 (1.67, 1.88)	0.34 (0.30, 0.41)	
Islander	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.08 (0.07, 0.09)	1.78 (1.68, 1.90)	0.35 (0.30, 0.41)	
	20	0.01 (0.01, 0.01)	0.08 (0.07, 0.09)	0.30 (0.28, 0.33)	1.78 (1.69, 1.90)	0.35 (0.30, 0.41)	
	30	0.07 (0.06, 0.08)	0.30 (0.27, 0.32)	0.73 (0.69, 0.77)	1.78 (1.68, 1.90)	0.35 (0.30, 0.41)	
	40	0.23 (0.21, 0.25)	0.66 (0.62, 0.71)	1.13 (1.07, 1.19)	1.72 (1.62, 1.84)	0.35 (0.30, 0.41)	
	50	0.44 (0.41, 0.48)	0.91 (0.85, 0.97)	1.25 (1.17, 1.32)	1.51 (1.41, 1.62)	0.33 (0.29, 0.40)	
	60	0.48 (0.44, 0.53)	0.83 (0.77, 0.89)	1.04 (0.97, 1.13)	1.10 (1.01, 1.21)	0.30 (0.25, 0.36)	
	70	0.37 (0.33, 0.42)	0.60 (0.54, 0.67)	- (- , -)	0.66 (0.58, 0.77)	0.22 (0.17, 0.28)	
	80	0.28 (0.22, 0.34)	- (- , -)	- (- , -)	0.35 (0.27, 0.46)	0.14 (0.09, 0.21)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.02 (0.00, 0.06)	1.46 (1.14, 2.05)	0.29 (0.21, 0.45)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.02 (0.00, 0.05)	0.08 (0.04, 0.14)	1.48 (1.16, 2.08)	0.29 (0.21, 0.46)	
	20	0.02 (0.00, 0.05)	0.08 (0.04, 0.14)	0.23 (0.16, 0.32)	1.49 (1.17, 2.09)	0.29 (0.21, 0.46)	
	30	0.07 (0.03, 0.12)	0.21 (0.15, 0.30)	0.50 (0.38, 0.65)	1.49 (1.16, 2.10)	0.30 (0.21, 0.46)	
	40	0.15 (0.10, 0.22)	0.44 (0.33, 0.59)	0.88 (0.68, 1.13)	1.45 (1.12, 2.07)	0.30 (0.21, 0.47)	
	50	0.30 (0.21, 0.43)	0.75 (0.56, 1.00)	1.11 (0.85, 1.44)	1.33 (1.01, 1.97)	0.28 (0.20, 0.46)	
	60	0.48 (0.32, 0.69)	0.86 (0.61, 1.18)	1.03 (0.74, 1.43)	1.09 (0.77, 1.75)	0.27 (0.18, 0.46)	
	70	0.44 (0.25, 0.71)	0.64 (0.38, 1.03)	- (- , -)	0.71 (0.42, 1.43)	0.23 (0.14, 0.44)	
	80	0.27 (0.10, 0.65)	- (- , -)	- (- , -)	0.36 (0.13, 1.28)	0.13 (0.06, 0.40)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.97 (1.88, 2.08)	0.45 (0.41, 0.48)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.07 (0.06, 0.08)	1.99 (1.89, 2.10)	0.45 (0.42, 0.49)	
	20	0.01 (0.01, 0.01)	0.07 (0.06, 0.07)	0.24 (0.22, 0.25)	1.99 (1.90, 2.10)	0.45 (0.42, 0.49)	
	30	0.06 (0.05, 0.06)	0.23 (0.21, 0.24)	0.63 (0.60, 0.66)	1.99 (1.89, 2.10)	0.45 (0.42, 0.49)	
	40	0.17 (0.16, 0.18)	0.58 (0.54, 0.61)	1.10 (1.05, 1.16)	1.95 (1.85, 2.06)	0.45 (0.42, 0.49)	
	50	0.41 (0.39, 0.44)	0.95 (0.89, 1.00)	1.44 (1.37, 1.52)	1.80 (1.71, 1.92)	0.44 (0.41, 0.48)	
	60	0.55 (0.51, 0.60)	1.07 (1.00, 1.14)	1.36 (1.27, 1.45)	1.45 (1.35, 1.55)	0.42 (0.38, 0.46)	
	70	0.56 (0.51, 0.62)	0.88 (0.81, 0.96)	- (- , -)	0.98 (0.89, 1.08)	0.35 (0.31, 0.39)	
	80	0.40 (0.34, 0.47)	- (- , -)	- (- , -)	0.52 (0.44, 0.63)	0.26 (0.22, 0.30)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VIII-9
ESOPHAGUS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.50 (0.49, 0.51)	0.48 (0.47, 0.48)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.51 (0.50, 0.52)	0.48 (0.48, 0.49)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.51 (0.50, 0.52)	0.48 (0.48, 0.49)	
	30	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.09 (0.09, 0.09)	0.52 (0.51, 0.53)	0.49 (0.48, 0.49)	
	40	0.02 (0.02, 0.02)	0.09 (0.08, 0.09)	0.23 (0.22, 0.23)	0.52 (0.51, 0.53)	0.49 (0.49, 0.50)	
	50	0.07 (0.07, 0.07)	0.22 (0.21, 0.22)	0.39 (0.38, 0.40)	0.52 (0.50, 0.53)	0.49 (0.49, 0.50)	
	60	0.15 (0.15, 0.16)	0.34 (0.33, 0.35)	0.45 (0.44, 0.46)	0.47 (0.46, 0.49)	0.45 (0.45, 0.46)	
	70	0.21 (0.20, 0.22)	0.34 (0.33, 0.35)	- (- , -)	0.37 (0.36, 0.38)	0.36 (0.36, 0.37)	
	80	0.18 (0.17, 0.19)	- (- , -)	- (- , -)	0.22 (0.21, 0.23)	0.23 (0.23, 0.24)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.51 (0.50, 0.53)	0.48 (0.47, 0.49)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.52 (0.51, 0.53)	0.49 (0.48, 0.49)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.52 (0.51, 0.53)	0.49 (0.48, 0.49)	
	30	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.09 (0.08, 0.09)	0.53 (0.51, 0.54)	0.49 (0.49, 0.50)	
	40	0.02 (0.02, 0.02)	0.09 (0.08, 0.09)	0.23 (0.22, 0.24)	0.53 (0.52, 0.54)	0.49 (0.49, 0.50)	
	50	0.07 (0.07, 0.07)	0.22 (0.21, 0.22)	0.39 (0.38, 0.40)	0.52 (0.51, 0.54)	0.49 (0.49, 0.50)	
	60	0.15 (0.15, 0.16)	0.34 (0.33, 0.35)	0.45 (0.44, 0.47)	0.48 (0.47, 0.49)	0.46 (0.45, 0.46)	
	70	0.22 (0.21, 0.23)	0.35 (0.33, 0.36)	- (- , -)	0.38 (0.37, 0.39)	0.37 (0.36, 0.37)	
	80	0.18 (0.17, 0.19)	- (- , -)	- (- , -)	0.23 (0.22, 0.24)	0.23 (0.23, 0.24)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.52 (0.49, 0.56)	0.51 (0.49, 0.52)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.54 (0.50, 0.57)	0.52 (0.50, 0.53)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.03 (0.03, 0.04)	0.54 (0.51, 0.57)	0.52 (0.51, 0.54)	
	30	0.00 (0.00, 0.00)	0.03 (0.03, 0.04)	0.13 (0.12, 0.15)	0.55 (0.51, 0.58)	0.53 (0.51, 0.55)	
	40	0.03 (0.03, 0.03)	0.14 (0.12, 0.15)	0.32 (0.30, 0.34)	0.56 (0.52, 0.60)	0.54 (0.52, 0.56)	
	50	0.11 (0.10, 0.12)	0.31 (0.28, 0.33)	0.48 (0.45, 0.51)	0.56 (0.52, 0.60)	0.54 (0.52, 0.56)	
	60	0.22 (0.20, 0.24)	0.41 (0.38, 0.44)	0.49 (0.45, 0.52)	0.50 (0.46, 0.54)	0.49 (0.47, 0.51)	
	70	0.24 (0.21, 0.27)	0.34 (0.30, 0.37)	- (- , -)	0.35 (0.32, 0.39)	0.36 (0.34, 0.38)	
	80	0.16 (0.13, 0.19)	- (- , -)	- (- , -)	0.18 (0.15, 0.22)	0.21 (0.20, 0.23)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

- Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VIII-9 - continued
ESOPHAGUS CANCER (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer						Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	Percent (95% C.I.)		
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.36 (0.32, 0.41)	0.36 (0.32, 0.41)	0.30 (0.26, 0.35)		
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.36 (0.32, 0.41)	0.36 (0.32, 0.41)	0.30 (0.27, 0.35)		
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	0.36 (0.32, 0.42)	0.36 (0.32, 0.42)	0.30 (0.27, 0.35)		
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.04, 0.05)	0.36 (0.32, 0.42)	0.36 (0.32, 0.42)	0.30 (0.27, 0.35)		
	40	0.01 (0.01, 0.01)	0.04 (0.03, 0.05)	0.11 (0.10, 0.13)	0.36 (0.32, 0.42)	0.36 (0.32, 0.42)	0.31 (0.27, 0.35)		
	50	0.03 (0.03, 0.04)	0.11 (0.09, 0.12)	0.20 (0.17, 0.22)	0.36 (0.32, 0.41)	0.31 (0.27, 0.35)			
	60	0.08 (0.06, 0.09)	0.17 (0.15, 0.19)	0.29 (0.26, 0.33)	0.34 (0.30, 0.39)	0.29 (0.25, 0.34)			
	70	0.10 (0.09, 0.12)	0.24 (0.20, 0.27)	- (-, -)	0.29 (0.24, 0.34)	0.26 (0.22, 0.31)			
	80	0.17 (0.14, 0.20)	- (-, -)	- (-, -)	0.23 (0.18, 0.30)	0.21 (0.16, 0.27)			
American	0	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.41 (0.28, 0.71)	0.40 (0.32, 0.52)			
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.42 (0.29, 0.72)	0.41 (0.32, 0.53)			
	20	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.01 (0.00, 0.03)	0.42 (0.29, 0.73)	0.41 (0.33, 0.54)			
	30	0.00 (0.00, 0.01)	0.01 (0.00, 0.03)	0.04 (0.02, 0.08)	0.43 (0.30, 0.74)	0.42 (0.33, 0.55)			
	40	0.01 (0.00, 0.03)	0.04 (0.02, 0.08)	0.19 (0.12, 0.28)	0.44 (0.30, 0.76)	0.43 (0.34, 0.56)			
	50	0.03 (0.01, 0.06)	0.18 (0.11, 0.28)	0.37 (0.25, 0.52)	0.44 (0.31, 0.78)	0.44 (0.34, 0.57)			
	60	0.16 (0.10, 0.26)	0.36 (0.24, 0.52)	0.44 (0.30, 0.65)	0.44 (0.30, 0.80)	0.43 (0.33, 0.58)			
	70	0.23 (0.14, 0.38)	0.33 (0.19, 0.54)	- (-, -)	0.33 (0.19, 0.73)	0.37 (0.26, 0.54)			
	80	0.13 (0.04, 0.34)	- (-, -)	- (-, -)	0.13 (0.04, 0.68)	0.28 (0.16, 0.52)			
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.41 (0.37, 0.47)	0.31 (0.29, 0.33)			
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.42 (0.38, 0.47)	0.31 (0.29, 0.34)			
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.42 (0.38, 0.47)	0.32 (0.29, 0.34)			
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.04, 0.05)	0.42 (0.38, 0.48)	0.32 (0.30, 0.34)			
	40	0.01 (0.01, 0.01)	0.04 (0.04, 0.05)	0.13 (0.11, 0.14)	0.43 (0.38, 0.48)	0.32 (0.30, 0.35)			
	50	0.03 (0.03, 0.04)	0.12 (0.11, 0.14)	0.25 (0.22, 0.27)	0.43 (0.38, 0.48)	0.32 (0.30, 0.35)			
	60	0.09 (0.08, 0.10)	0.22 (0.20, 0.25)	0.34 (0.31, 0.37)	0.41 (0.36, 0.47)	0.31 (0.28, 0.33)			
	70	0.15 (0.13, 0.17)	0.28 (0.25, 0.31)	- (-, -)	0.36 (0.31, 0.42)	0.27 (0.24, 0.29)			
	80	0.17 (0.14, 0.21)	- (-, -)	- (-, -)	0.28 (0.22, 0.35)	0.20 (0.17, 0.24)			

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VIII-9 - continued
ESOPHAGUS CANCER (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.78 (0.76, 0.80)	0.75 (0.74, 0.76)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.79 (0.77, 0.81)	0.77 (0.76, 0.77)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.04 (0.03, 0.04)	0.79 (0.77, 0.81)	0.77 (0.76, 0.78)	
	30	0.00 (0.00, 0.01)	0.04 (0.03, 0.04)	0.15 (0.14, 0.15)	0.80 (0.78, 0.82)	0.78 (0.77, 0.79)	
	40	0.03 (0.03, 0.03)	0.14 (0.14, 0.15)	0.38 (0.36, 0.39)	0.81 (0.79, 0.83)	0.79 (0.78, 0.80)	
	50	0.12 (0.11, 0.12)	0.36 (0.35, 0.37)	0.63 (0.61, 0.64)	0.81 (0.79, 0.83)	0.79 (0.78, 0.80)	
	60	0.26 (0.25, 0.27)	0.55 (0.53, 0.57)	0.71 (0.69, 0.73)	0.75 (0.73, 0.77)	0.73 (0.72, 0.74)	
	70	0.35 (0.33, 0.36)	0.54 (0.52, 0.56)	- (-, -)	0.58 (0.56, 0.60)	0.59 (0.58, 0.60)	
	80	0.30 (0.28, 0.31)	- (-, -)	- (-, -)	0.36 (0.34, 0.38)	0.38 (0.37, 0.39)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.80 (0.78, 0.82)	0.77 (0.76, 0.78)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.81 (0.79, 0.83)	0.78 (0.77, 0.79)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.04 (0.03, 0.04)	0.81 (0.79, 0.83)	0.78 (0.77, 0.79)	
	30	0.00 (0.00, 0.01)	0.04 (0.03, 0.04)	0.15 (0.14, 0.16)	0.82 (0.80, 0.84)	0.79 (0.78, 0.80)	
	40	0.03 (0.03, 0.03)	0.15 (0.14, 0.15)	0.38 (0.37, 0.39)	0.83 (0.81, 0.85)	0.80 (0.79, 0.81)	
	50	0.12 (0.11, 0.12)	0.36 (0.35, 0.37)	0.64 (0.62, 0.66)	0.83 (0.80, 0.85)	0.80 (0.79, 0.81)	
	60	0.26 (0.25, 0.27)	0.56 (0.54, 0.58)	0.73 (0.71, 0.75)	0.76 (0.74, 0.78)	0.74 (0.73, 0.75)	
	70	0.36 (0.34, 0.37)	0.56 (0.54, 0.58)	- (-, -)	0.60 (0.58, 0.62)	0.60 (0.59, 0.61)	
	80	0.30 (0.29, 0.32)	- (-, -)	- (-, -)	0.37 (0.35, 0.39)	0.39 (0.38, 0.40)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.76 (0.70, 0.82)	0.74 (0.72, 0.77)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.78 (0.72, 0.84)	0.76 (0.74, 0.79)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.04 (0.03, 0.05)	0.79 (0.73, 0.85)	0.77 (0.74, 0.80)	
	30	0.00 (0.00, 0.01)	0.04 (0.03, 0.05)	0.20 (0.18, 0.22)	0.81 (0.75, 0.87)	0.79 (0.76, 0.81)	
	40	0.04 (0.03, 0.05)	0.20 (0.18, 0.23)	0.50 (0.46, 0.54)	0.83 (0.77, 0.90)	0.81 (0.78, 0.84)	
	50	0.17 (0.15, 0.20)	0.49 (0.45, 0.53)	0.73 (0.68, 0.79)	0.84 (0.78, 0.91)	0.82 (0.79, 0.85)	
	60	0.36 (0.33, 0.41)	0.65 (0.59, 0.71)	0.75 (0.69, 0.83)	0.77 (0.70, 0.85)	0.75 (0.72, 0.79)	
	70	0.38 (0.33, 0.43)	0.52 (0.46, 0.59)	- (-, -)	0.54 (0.48, 0.62)	0.55 (0.52, 0.59)	
	80	0.26 (0.20, 0.33)	- (-, -)	- (-, -)	0.30 (0.24, 0.40)	0.33 (0.30, 0.37)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VIII-9 - continued
ESOPHAGUS CANCER (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.54 (0.47, 0.63)	0.47 (0.40, 0.55)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.54 (0.47, 0.63)	0.47 (0.41, 0.55)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.54 (0.47, 0.63)	0.47 (0.41, 0.56)	
	30	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.07 (0.05, 0.08)	0.55 (0.47, 0.64)	0.48 (0.41, 0.56)	
	40	0.01 (0.01, 0.02)	0.07 (0.05, 0.08)	0.19 (0.16, 0.22)	0.55 (0.48, 0.64)	0.48 (0.41, 0.56)	
	50	0.05 (0.04, 0.07)	0.18 (0.15, 0.21)	0.33 (0.29, 0.37)	0.54 (0.47, 0.64)	0.48 (0.41, 0.56)	
	60	0.13 (0.11, 0.16)	0.28 (0.24, 0.33)	0.47 (0.40, 0.53)	0.51 (0.44, 0.61)	0.46 (0.39, 0.54)	
	70	0.17 (0.14, 0.21)	0.37 (0.31, 0.44)	- (-, -)	0.42 (0.35, 0.52)	0.39 (0.32, 0.49)	
	80	0.26 (0.20, 0.34)	- (-, -)	- (-, -)	0.33 (0.25, 0.45)	0.31 (0.23, 0.42)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.61 (0.40, 1.29)	0.62 (0.48, 0.86)	
Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.62 (0.41, 1.31)	0.63 (0.49, 0.88)	
Alaska	20	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.02 (0.00, 0.06)	0.63 (0.41, 1.33)	0.64 (0.50, 0.89)	
	30	0.00 (0.00, 0.03)	0.02 (0.00, 0.06)	0.07 (0.03, 0.14)	0.65 (0.42, 1.36)	0.66 (0.51, 0.91)	
	40	0.02 (0.00, 0.05)	0.07 (0.03, 0.14)	0.31 (0.19, 0.49)	0.67 (0.43, 1.41)	0.68 (0.53, 0.95)	
	50	0.06 (0.02, 0.13)	0.31 (0.18, 0.50)	0.57 (0.36, 0.85)	0.68 (0.44, 1.46)	0.70 (0.54, 0.98)	
	60	0.28 (0.15, 0.47)	0.56 (0.35, 0.86)	0.69 (0.43, 1.09)	0.69 (0.43, 1.54)	0.70 (0.53, 1.01)	
	70	0.33 (0.17, 0.61)	0.49 (0.25, 0.91)	- (-, -)	0.49 (0.25, 1.50)	0.61 (0.42, 0.98)	
	80	0.23 (0.06, 0.73)	- (-, -)	- (-, -)	0.23 (0.06, 1.76)	0.48 (0.26, 1.02)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.66 (0.58, 0.76)	0.48 (0.44, 0.52)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.66 (0.58, 0.77)	0.48 (0.45, 0.53)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.67 (0.59, 0.77)	0.49 (0.45, 0.53)	
	30	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.08 (0.07, 0.09)	0.67 (0.59, 0.78)	0.49 (0.45, 0.54)	
	40	0.02 (0.01, 0.02)	0.08 (0.06, 0.09)	0.23 (0.20, 0.26)	0.68 (0.60, 0.79)	0.50 (0.46, 0.54)	
	50	0.06 (0.05, 0.07)	0.22 (0.19, 0.25)	0.42 (0.37, 0.46)	0.69 (0.60, 0.79)	0.50 (0.46, 0.55)	
	60	0.17 (0.14, 0.19)	0.38 (0.33, 0.43)	0.56 (0.50, 0.63)	0.67 (0.58, 0.78)	0.48 (0.44, 0.53)	
	70	0.24 (0.21, 0.29)	0.45 (0.39, 0.52)	- (-, -)	0.57 (0.48, 0.70)	0.41 (0.37, 0.47)	
	80	0.29 (0.23, 0.37)	- (-, -)	- (-, -)	0.47 (0.35, 0.63)	0.30 (0.25, 0.37)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VIII-9 - continued
ESOPHAGUS CANCER (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.25 (0.24, 0.26)	0.22 (0.22, 0.23)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.26 (0.25, 0.27)	0.22 (0.22, 0.23)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.26 (0.25, 0.27)	0.22 (0.22, 0.23)	
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.26 (0.25, 0.27)	0.22 (0.22, 0.23)	
	40	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.09 (0.08, 0.09)	0.26 (0.25, 0.27)	0.23 (0.22, 0.23)	
	50	0.02 (0.02, 0.03)	0.08 (0.08, 0.09)	0.17 (0.16, 0.18)	0.26 (0.25, 0.27)	0.23 (0.22, 0.23)	
	60	0.06 (0.06, 0.07)	0.15 (0.14, 0.16)	0.22 (0.21, 0.23)	0.24 (0.23, 0.26)	0.22 (0.21, 0.22)	
	70	0.10 (0.10, 0.11)	0.18 (0.17, 0.19)	- (-, -)	0.21 (0.20, 0.22)	0.19 (0.18, 0.19)	
	80	0.10 (0.10, 0.11)	- (-, -)	- (-, -)	0.14 (0.13, 0.15)	0.14 (0.13, 0.14)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.25 (0.24, 0.26)	0.21 (0.21, 0.22)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.25 (0.24, 0.27)	0.22 (0.21, 0.22)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.25 (0.24, 0.27)	0.22 (0.21, 0.22)	
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.02, 0.03)	0.26 (0.24, 0.27)	0.22 (0.21, 0.22)	
	40	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	0.08 (0.08, 0.09)	0.26 (0.24, 0.27)	0.22 (0.21, 0.22)	
	50	0.02 (0.02, 0.02)	0.08 (0.07, 0.08)	0.17 (0.16, 0.18)	0.26 (0.24, 0.27)	0.22 (0.21, 0.22)	
	60	0.06 (0.05, 0.06)	0.15 (0.14, 0.16)	0.22 (0.21, 0.23)	0.24 (0.23, 0.26)	0.21 (0.21, 0.22)	
	70	0.10 (0.10, 0.11)	0.18 (0.17, 0.19)	- (-, -)	0.21 (0.20, 0.22)	0.19 (0.18, 0.19)	
	80	0.10 (0.10, 0.11)	- (-, -)	- (-, -)	0.14 (0.13, 0.15)	0.14 (0.13, 0.14)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.31 (0.28, 0.35)	0.30 (0.28, 0.31)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.32 (0.28, 0.35)	0.30 (0.29, 0.32)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.03)	0.32 (0.28, 0.36)	0.30 (0.29, 0.32)	
	30	0.00 (0.00, 0.00)	0.02 (0.02, 0.03)	0.08 (0.06, 0.09)	0.32 (0.28, 0.36)	0.31 (0.29, 0.32)	
	40	0.02 (0.01, 0.03)	0.08 (0.06, 0.09)	0.16 (0.14, 0.19)	0.32 (0.29, 0.36)	0.31 (0.29, 0.33)	
	50	0.06 (0.05, 0.07)	0.15 (0.13, 0.17)	0.26 (0.23, 0.29)	0.32 (0.28, 0.36)	0.31 (0.29, 0.33)	
	60	0.10 (0.08, 0.12)	0.22 (0.19, 0.25)	0.27 (0.24, 0.31)	0.28 (0.24, 0.32)	0.29 (0.27, 0.31)	
	70	0.14 (0.11, 0.17)	0.20 (0.17, 0.24)	- (-, -)	0.22 (0.18, 0.26)	0.23 (0.21, 0.25)	
	80	0.10 (0.07, 0.13)	- (-, -)	- (-, -)	0.11 (0.08, 0.15)	0.15 (0.13, 0.17)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VIII-9 - continued
ESOPHAGUS CANCER (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.20 (0.16, 0.28)	0.16 (0.12, 0.22)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.20 (0.16, 0.28)	0.16 (0.12, 0.22)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.21 (0.16, 0.28)	0.16 (0.12, 0.22)	
	30	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.01, 0.03)	0.21 (0.16, 0.28)	0.16 (0.12, 0.22)	
	40	0.00 (0.00, 0.01)	0.02 (0.01, 0.03)	0.05 (0.03, 0.06)	0.20 (0.16, 0.28)	0.16 (0.12, 0.22)	
	50	0.01 (0.01, 0.02)	0.04 (0.03, 0.06)	0.09 (0.07, 0.11)	0.20 (0.15, 0.28)	0.16 (0.12, 0.22)	
	60	0.03 (0.02, 0.04)	0.08 (0.06, 0.10)	0.15 (0.12, 0.19)	0.19 (0.15, 0.27)	0.15 (0.11, 0.22)	
	70	0.05 (0.03, 0.07)	0.13 (0.10, 0.17)	- (-, -)	0.17 (0.12, 0.26)	0.15 (0.11, 0.22)	
	80	0.10 (0.07, 0.14)	- (-, -)	- (-, -)	0.15 (0.10, 0.25)	0.13 (0.08, 0.22)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.22 (0.10, 0.71)	0.20 (0.11, 0.37)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.22 (0.10, 0.72)	0.20 (0.11, 0.38)	
	20	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.22 (0.10, 0.73)	0.20 (0.11, 0.38)	
	30	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.01 (0.00, 0.05)	0.22 (0.10, 0.73)	0.20 (0.11, 0.39)	
	40	0.01 (0.00, 0.04)	0.01 (0.00, 0.06)	0.07 (0.02, 0.17)	0.23 (0.10, 0.75)	0.21 (0.11, 0.39)	
	50	0.01 (0.00, 0.04)	0.06 (0.02, 0.16)	0.19 (0.08, 0.37)	0.23 (0.10, 0.76)	0.21 (0.11, 0.40)	
	60	0.06 (0.02, 0.16)	0.19 (0.08, 0.39)	0.23 (0.10, 0.49)	0.23 (0.10, 0.80)	0.20 (0.11, 0.41)	
	70	0.15 (0.05, 0.35)	0.20 (0.07, 0.47)	- (-, -)	0.20 (0.07, 0.85)	0.18 (0.08, 0.42)	
	80	0.06 (0.00, 0.34)	- (-, -)	- (-, -)	0.06 (0.00, 0.93)	0.16 (0.04, 0.49)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.21 (0.17, 0.27)	0.16 (0.14, 0.19)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.21 (0.17, 0.27)	0.17 (0.14, 0.20)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.21 (0.17, 0.28)	0.17 (0.14, 0.20)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.21 (0.17, 0.28)	0.17 (0.14, 0.20)	
	40	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.21 (0.17, 0.28)	0.17 (0.14, 0.20)	
	50	0.01 (0.01, 0.01)	0.03 (0.02, 0.05)	0.10 (0.08, 0.12)	0.21 (0.17, 0.28)	0.17 (0.14, 0.20)	
	60	0.03 (0.02, 0.04)	0.09 (0.07, 0.11)	0.16 (0.13, 0.20)	0.21 (0.16, 0.28)	0.17 (0.14, 0.20)	
	70	0.07 (0.05, 0.09)	0.15 (0.12, 0.19)	- (-, -)	0.20 (0.15, 0.27)	0.16 (0.13, 0.19)	
	80	0.10 (0.07, 0.14)	- (-, -)	- (-, -)	0.16 (0.11, 0.25)	0.14 (0.11, 0.18)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table IX-9
HODGKIN LYMPHOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.06 (0.06, 0.07)	0.22 (0.21, 0.23)	0.04 (0.04, 0.05)	
	10	0.02 (0.02, 0.02)	0.06 (0.06, 0.06)	0.09 (0.09, 0.10)	0.22 (0.22, 0.23)	0.04 (0.04, 0.05)	
	20	0.04 (0.04, 0.04)	0.07 (0.07, 0.08)	0.10 (0.10, 0.10)	0.20 (0.20, 0.21)	0.04 (0.04, 0.05)	
	30	0.03 (0.03, 0.04)	0.06 (0.06, 0.06)	0.08 (0.08, 0.09)	0.16 (0.16, 0.17)	0.04 (0.04, 0.04)	
	40	0.03 (0.03, 0.03)	0.05 (0.05, 0.05)	0.08 (0.08, 0.08)	0.13 (0.12, 0.13)	0.04 (0.04, 0.04)	
	50	0.02 (0.02, 0.03)	0.06 (0.05, 0.06)	0.09 (0.08, 0.09)	0.11 (0.10, 0.11)	0.04 (0.03, 0.04)	
	60	0.03 (0.03, 0.04)	0.07 (0.06, 0.07)	0.08 (0.08, 0.09)	0.09 (0.08, 0.09)	0.03 (0.03, 0.03)	
	70	0.04 (0.03, 0.04)	0.06 (0.06, 0.06)	- (- , -)	0.06 (0.06, 0.07)	0.03 (0.03, 0.03)	
	80	0.03 (0.03, 0.03)	- (- , -)	- (- , -)	0.03 (0.03, 0.04)	0.02 (0.02, 0.02)	
White	0	0.00 (0.00, 0.00)	0.02 (0.02, 0.03)	0.07 (0.07, 0.07)	0.24 (0.23, 0.25)	0.05 (0.04, 0.05)	
	10	0.02 (0.02, 0.02)	0.07 (0.06, 0.07)	0.10 (0.10, 0.11)	0.24 (0.23, 0.25)	0.05 (0.04, 0.05)	
	20	0.05 (0.04, 0.05)	0.08 (0.08, 0.09)	0.11 (0.11, 0.11)	0.22 (0.21, 0.22)	0.05 (0.04, 0.05)	
	30	0.04 (0.04, 0.04)	0.07 (0.06, 0.07)	0.09 (0.09, 0.09)	0.17 (0.17, 0.18)	0.04 (0.04, 0.04)	
	40	0.03 (0.03, 0.03)	0.05 (0.05, 0.06)	0.08 (0.08, 0.09)	0.14 (0.13, 0.14)	0.04 (0.04, 0.04)	
	50	0.02 (0.02, 0.03)	0.06 (0.05, 0.06)	0.09 (0.09, 0.10)	0.11 (0.11, 0.12)	0.04 (0.04, 0.04)	
	60	0.04 (0.03, 0.04)	0.07 (0.07, 0.07)	0.09 (0.09, 0.10)	0.09 (0.09, 0.10)	0.03 (0.03, 0.04)	
	70	0.04 (0.04, 0.04)	0.06 (0.06, 0.07)	- (- , -)	0.07 (0.06, 0.07)	0.03 (0.03, 0.03)	
	80	0.03 (0.03, 0.04)	- (- , -)	- (- , -)	0.04 (0.03, 0.04)	0.02 (0.02, 0.02)	
Black	0	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.05 (0.04, 0.05)	0.18 (0.16, 0.19)	0.03 (0.03, 0.03)	
	10	0.01 (0.01, 0.02)	0.04 (0.04, 0.05)	0.08 (0.07, 0.09)	0.18 (0.16, 0.20)	0.03 (0.03, 0.04)	
	20	0.03 (0.03, 0.04)	0.06 (0.06, 0.07)	0.09 (0.08, 0.10)	0.17 (0.15, 0.18)	0.03 (0.03, 0.03)	
	30	0.03 (0.03, 0.04)	0.06 (0.06, 0.07)	0.09 (0.08, 0.10)	0.14 (0.12, 0.15)	0.03 (0.02, 0.03)	
	40	0.03 (0.03, 0.04)	0.06 (0.05, 0.07)	0.08 (0.07, 0.09)	0.11 (0.09, 0.12)	0.02 (0.02, 0.03)	
	50	0.03 (0.02, 0.03)	0.05 (0.04, 0.06)	0.07 (0.06, 0.09)	0.08 (0.07, 0.09)	0.02 (0.02, 0.02)	
	60	0.03 (0.02, 0.04)	0.05 (0.04, 0.06)	0.06 (0.04, 0.07)	0.06 (0.04, 0.07)	0.02 (0.01, 0.02)	
	70	0.03 (0.02, 0.04)	0.04 (0.03, 0.05)	- (- , -)	0.04 (0.03, 0.05)	0.01 (0.01, 0.02)	
	80	0.01 (0.01, 0.02)	- (- , -)	- (- , -)	0.01 (0.01, 0.03)	0.01 (0.00, 0.01)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table IX-9 - continued

HODGKIN LYMPHOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.03 (0.03, 0.04)	0.11 (0.10, 0.14)	0.04 (0.03, 0.06)	
	10	0.01 (0.01, 0.02)	0.03 (0.03, 0.04)	0.05 (0.04, 0.06)	0.11 (0.10, 0.14)	0.04 (0.03, 0.06)	
Islander	20	0.02 (0.02, 0.03)	0.04 (0.03, 0.04)	0.04 (0.04, 0.05)	0.10 (0.08, 0.13)	0.04 (0.03, 0.06)	
	30	0.01 (0.01, 0.02)	0.02 (0.02, 0.03)	0.03 (0.03, 0.04)	0.08 (0.06, 0.11)	0.03 (0.02, 0.06)	
	40	0.01 (0.01, 0.01)	0.02 (0.02, 0.03)	0.03 (0.03, 0.04)	0.06 (0.05, 0.09)	0.03 (0.02, 0.06)	
	50	0.01 (0.01, 0.01)	0.03 (0.02, 0.03)	0.04 (0.03, 0.05)	0.06 (0.04, 0.08)	0.03 (0.02, 0.06)	
	60	0.02 (0.01, 0.02)	0.03 (0.02, 0.04)	0.05 (0.03, 0.06)	0.05 (0.03, 0.08)	0.03 (0.02, 0.06)	
	70	0.02 (0.01, 0.03)	0.03 (0.02, 0.05)	- (-, -)	0.03 (0.02, 0.06)	0.03 (0.02, 0.05)	
	80	0.02 (0.01, 0.03)	- (-, -)	- (-, -)	0.02 (0.01, 0.06)	0.03 (0.01, 0.06)	
American Indian/ Alaska Native ^a	0	0.00 (0.00, 0.02)	0.01 (0.00, 0.03)	0.02 (0.01, 0.05)	0.08 (0.04, 0.35)	0.04 (0.02, 0.12)	
	10	0.01 (0.00, 0.02)	0.02 (0.01, 0.04)	0.03 (0.01, 0.05)	0.08 (0.04, 0.35)	0.04 (0.02, 0.13)	
	20	0.01 (0.00, 0.03)	0.02 (0.01, 0.04)	0.03 (0.01, 0.06)	0.07 (0.03, 0.35)	0.04 (0.02, 0.13)	
	30	0.01 (0.00, 0.03)	0.02 (0.01, 0.04)	0.04 (0.02, 0.08)	0.06 (0.02, 0.34)	0.04 (0.02, 0.13)	
	40	0.01 (0.00, 0.03)	0.03 (0.01, 0.07)	0.03 (0.01, 0.08)	0.05 (0.02, 0.35)	0.04 (0.02, 0.13)	
	50	0.02 (0.01, 0.06)	0.02 (0.01, 0.07)	0.03 (0.01, 0.09)	0.04 (0.01, 0.35)	0.04 (0.02, 0.14)	
	60	0.00 (0.00, 0.04)	0.01 (0.00, 0.06)	0.02 (0.00, 0.13)	0.02 (0.00, 0.36)	0.04 (0.02, 0.14)	
	70	0.00 (0.00, 0.07)	0.02 (0.00, 0.15)	- (-, -)	0.02 (0.00, 0.42)	0.05 (0.01, 0.17)	
	80	0.03 (0.00, 0.20)	- (-, -)	- (-, -)	0.03 (0.00, 0.58)	0.05 (0.01, 0.23)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.04 (0.04, 0.04)	0.22 (0.20, 0.25)	0.05 (0.04, 0.06)	
	10	0.02 (0.01, 0.02)	0.04 (0.03, 0.04)	0.06 (0.05, 0.06)	0.22 (0.20, 0.25)	0.05 (0.04, 0.06)	
	20	0.02 (0.02, 0.03)	0.04 (0.04, 0.05)	0.06 (0.06, 0.07)	0.21 (0.18, 0.24)	0.05 (0.04, 0.06)	
	30	0.02 (0.02, 0.02)	0.04 (0.03, 0.04)	0.06 (0.06, 0.07)	0.18 (0.16, 0.22)	0.05 (0.04, 0.06)	
	40	0.02 (0.02, 0.02)	0.04 (0.04, 0.05)	0.08 (0.07, 0.09)	0.17 (0.14, 0.20)	0.05 (0.04, 0.06)	
	50	0.02 (0.02, 0.03)	0.06 (0.05, 0.07)	0.11 (0.09, 0.12)	0.15 (0.13, 0.18)	0.05 (0.04, 0.06)	
	60	0.04 (0.03, 0.05)	0.09 (0.07, 0.10)	0.12 (0.10, 0.14)	0.13 (0.11, 0.17)	0.04 (0.04, 0.06)	
	70	0.05 (0.04, 0.07)	0.09 (0.07, 0.11)	- (-, -)	0.10 (0.08, 0.14)	0.04 (0.03, 0.05)	
	80	0.05 (0.03, 0.07)	- (-, -)	- (-, -)	0.07 (0.04, 0.11)	0.03 (0.02, 0.05)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table IX-9 - continued

HODGKIN LYMPHOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.02 (0.02, 0.03)	0.06 (0.06, 0.06)	0.24 (0.23, 0.25)	0.05 (0.05, 0.05)	
	10	0.02 (0.02, 0.02)	0.06 (0.06, 0.06)	0.09 (0.09, 0.10)	0.24 (0.23, 0.25)	0.05 (0.05, 0.05)	
	20	0.04 (0.04, 0.04)	0.07 (0.07, 0.08)	0.10 (0.10, 0.11)	0.22 (0.21, 0.23)	0.05 (0.05, 0.05)	
	30	0.04 (0.03, 0.04)	0.07 (0.06, 0.07)	0.09 (0.09, 0.10)	0.18 (0.18, 0.19)	0.05 (0.04, 0.05)	
	40	0.03 (0.03, 0.03)	0.06 (0.06, 0.06)	0.10 (0.09, 0.10)	0.15 (0.14, 0.16)	0.04 (0.04, 0.05)	
	50	0.03 (0.03, 0.03)	0.07 (0.06, 0.07)	0.10 (0.10, 0.11)	0.12 (0.12, 0.13)	0.04 (0.04, 0.04)	
	60	0.04 (0.04, 0.05)	0.08 (0.07, 0.08)	0.10 (0.09, 0.11)	0.10 (0.09, 0.11)	0.04 (0.04, 0.04)	
	70	0.04 (0.04, 0.05)	0.07 (0.06, 0.07)	- (-, -)	0.07 (0.06, 0.08)	0.03 (0.03, 0.03)	
	80	0.04 (0.03, 0.04)	- (-, -)	- (-, -)	0.04 (0.04, 0.05)	0.02 (0.02, 0.03)	
White	0	0.00 (0.00, 0.00)	0.02 (0.02, 0.03)	0.07 (0.06, 0.07)	0.26 (0.25, 0.27)	0.05 (0.05, 0.05)	
	10	0.02 (0.02, 0.02)	0.06 (0.06, 0.07)	0.10 (0.10, 0.11)	0.26 (0.25, 0.27)	0.05 (0.05, 0.05)	
	20	0.04 (0.04, 0.05)	0.08 (0.08, 0.09)	0.11 (0.11, 0.12)	0.23 (0.23, 0.25)	0.05 (0.05, 0.05)	
	30	0.04 (0.04, 0.04)	0.07 (0.07, 0.08)	0.10 (0.09, 0.10)	0.19 (0.19, 0.20)	0.05 (0.05, 0.05)	
	40	0.03 (0.03, 0.04)	0.06 (0.06, 0.07)	0.10 (0.09, 0.11)	0.16 (0.15, 0.17)	0.05 (0.04, 0.05)	
	50	0.03 (0.03, 0.03)	0.07 (0.07, 0.08)	0.11 (0.10, 0.11)	0.13 (0.12, 0.14)	0.04 (0.04, 0.05)	
	60	0.04 (0.04, 0.05)	0.08 (0.08, 0.09)	0.10 (0.10, 0.11)	0.11 (0.10, 0.12)	0.04 (0.04, 0.04)	
	70	0.05 (0.04, 0.05)	0.07 (0.06, 0.08)	- (-, -)	0.08 (0.07, 0.08)	0.03 (0.03, 0.04)	
	80	0.04 (0.03, 0.05)	- (-, -)	- (-, -)	0.04 (0.04, 0.05)	0.03 (0.02, 0.03)	
Black	0	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.04 (0.03, 0.05)	0.19 (0.16, 0.21)	0.04 (0.03, 0.04)	
	10	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.07 (0.06, 0.08)	0.19 (0.16, 0.22)	0.04 (0.03, 0.04)	
	20	0.03 (0.02, 0.03)	0.06 (0.05, 0.07)	0.09 (0.08, 0.11)	0.18 (0.15, 0.20)	0.04 (0.03, 0.04)	
	30	0.03 (0.03, 0.04)	0.07 (0.06, 0.08)	0.10 (0.09, 0.12)	0.15 (0.13, 0.18)	0.03 (0.03, 0.04)	
	40	0.04 (0.03, 0.04)	0.07 (0.06, 0.08)	0.10 (0.08, 0.11)	0.12 (0.10, 0.15)	0.03 (0.03, 0.04)	
	50	0.04 (0.03, 0.05)	0.06 (0.05, 0.08)	0.08 (0.07, 0.11)	0.09 (0.07, 0.12)	0.03 (0.02, 0.03)	
	60	0.03 (0.02, 0.05)	0.06 (0.04, 0.08)	0.07 (0.05, 0.09)	0.07 (0.05, 0.10)	0.02 (0.02, 0.03)	
	70	0.03 (0.02, 0.05)	0.04 (0.03, 0.07)	- (-, -)	0.04 (0.03, 0.08)	0.01 (0.01, 0.02)	
	80	0.02 (0.01, 0.05)	- (-, -)	- (-, -)	0.02 (0.01, 0.07)	0.01 (0.00, 0.02)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table IX-9 - continued

HODGKIN LYMPHOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.03 (0.02, 0.04)	0.13 (0.10, 0.18)	0.04 (0.02, 0.08)	
	10	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.05 (0.04, 0.06)	0.13 (0.10, 0.18)	0.04 (0.02, 0.08)	
Islander	20	0.02 (0.01, 0.02)	0.03 (0.02, 0.04)	0.04 (0.03, 0.05)	0.11 (0.09, 0.16)	0.04 (0.02, 0.08)	
	30	0.01 (0.01, 0.02)	0.03 (0.02, 0.03)	0.04 (0.03, 0.05)	0.10 (0.08, 0.15)	0.04 (0.02, 0.08)	
	40	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.05 (0.04, 0.07)	0.08 (0.06, 0.13)	0.04 (0.02, 0.08)	
	50	0.02 (0.01, 0.02)	0.04 (0.03, 0.06)	0.06 (0.04, 0.08)	0.07 (0.05, 0.12)	0.04 (0.02, 0.08)	
	60	0.03 (0.02, 0.04)	0.04 (0.03, 0.06)	0.06 (0.04, 0.09)	0.06 (0.04, 0.11)	0.03 (0.02, 0.08)	
	70	0.02 (0.01, 0.04)	0.04 (0.02, 0.07)	- (-, -)	0.04 (0.02, 0.10)	0.03 (0.01, 0.08)	
	80	0.02 (0.01, 0.06)	- (-, -)	- (-, -)	0.03 (0.01, 0.10)	0.02 (0.01, 0.09)	
American Indian/ Alaska Native ^a	0	0.00 (0.00, 0.04)	0.01 (0.00, 0.06)	0.02 (0.01, 0.07)	0.09 (0.05, 0.75)	0.02 (0.00, 0.20)	
	10	0.01 (0.00, 0.03)	0.02 (0.01, 0.06)	0.04 (0.01, 0.09)	0.10 (0.05, 0.77)	0.02 (0.01, 0.21)	
	20	0.01 (0.00, 0.05)	0.03 (0.01, 0.07)	0.05 (0.02, 0.10)	0.09 (0.04, 0.77)	0.02 (0.01, 0.21)	
	30	0.02 (0.00, 0.05)	0.04 (0.01, 0.09)	0.07 (0.03, 0.14)	0.08 (0.03, 0.78)	0.02 (0.01, 0.22)	
	40	0.02 (0.00, 0.06)	0.06 (0.02, 0.13)	0.06 (0.02, 0.15)	0.06 (0.02, 0.79)	0.02 (0.01, 0.22)	
	50	0.04 (0.01, 0.11)	0.04 (0.01, 0.14)	0.04 (0.01, 0.16)	0.04 (0.01, 0.82)	0.02 (0.00, 0.24)	
	60	0.01 (0.00, 0.08)	0.01 (0.00, 0.13)	0.01 (0.00, 0.22)	0.01 (0.00, 0.87)	0.02 (0.00, 0.26)	
	70	0.00 (0.00, 0.15)	0.00 (0.00, 0.27)	- (-, -)	0.00 (0.00, 1.04)	0.02 (0.00, 0.32)	
	80	0.00 (0.00, 0.40)	- (-, -)	- (-, -)	0.00 (0.00, 1.56)	0.00 (0.00, 0.48)	
Hispanic ^b	0	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.04 (0.04, 0.05)	0.27 (0.23, 0.34)	0.05 (0.04, 0.07)	
	10	0.02 (0.01, 0.02)	0.04 (0.03, 0.04)	0.06 (0.05, 0.07)	0.27 (0.23, 0.34)	0.05 (0.04, 0.07)	
	20	0.02 (0.02, 0.03)	0.04 (0.04, 0.05)	0.07 (0.06, 0.07)	0.26 (0.22, 0.32)	0.05 (0.04, 0.07)	
	30	0.02 (0.02, 0.03)	0.04 (0.04, 0.05)	0.08 (0.07, 0.09)	0.24 (0.20, 0.31)	0.05 (0.04, 0.07)	
	40	0.02 (0.02, 0.03)	0.06 (0.05, 0.07)	0.11 (0.09, 0.13)	0.22 (0.18, 0.29)	0.05 (0.04, 0.07)	
	50	0.03 (0.03, 0.04)	0.09 (0.07, 0.11)	0.15 (0.12, 0.18)	0.20 (0.16, 0.27)	0.05 (0.04, 0.07)	
	60	0.06 (0.04, 0.08)	0.12 (0.10, 0.15)	0.16 (0.13, 0.20)	0.18 (0.14, 0.26)	0.05 (0.03, 0.07)	
	70	0.07 (0.05, 0.10)	0.12 (0.09, 0.16)	- (-, -)	0.14 (0.10, 0.22)	0.04 (0.03, 0.07)	
	80	0.06 (0.03, 0.10)	- (-, -)	- (-, -)	0.09 (0.04, 0.21)	0.03 (0.02, 0.06)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table IX-9 - continued

HODGKIN LYMPHOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.06 (0.06, 0.07)	0.20 (0.20, 0.21)	0.04 (0.04, 0.04)	
	10	0.02 (0.02, 0.02)	0.06 (0.06, 0.07)	0.10 (0.09, 0.10)	0.21 (0.20, 0.21)	0.04 (0.04, 0.04)	
	20	0.04 (0.04, 0.05)	0.08 (0.07, 0.08)	0.10 (0.09, 0.10)	0.19 (0.18, 0.19)	0.04 (0.04, 0.04)	
	30	0.03 (0.03, 0.04)	0.06 (0.05, 0.06)	0.07 (0.07, 0.08)	0.14 (0.14, 0.15)	0.04 (0.03, 0.04)	
	40	0.02 (0.02, 0.02)	0.04 (0.04, 0.04)	0.06 (0.06, 0.07)	0.11 (0.10, 0.12)	0.03 (0.03, 0.03)	
	50	0.02 (0.02, 0.02)	0.04 (0.04, 0.05)	0.07 (0.07, 0.08)	0.09 (0.08, 0.10)	0.03 (0.03, 0.03)	
	60	0.03 (0.02, 0.03)	0.05 (0.05, 0.06)	0.07 (0.07, 0.08)	0.07 (0.07, 0.08)	0.03 (0.03, 0.03)	
	70	0.03 (0.03, 0.04)	0.05 (0.05, 0.06)	- (-, -)	0.06 (0.05, 0.06)	0.02 (0.02, 0.03)	
	80	0.03 (0.02, 0.03)	- (-, -)	- (-, -)	0.03 (0.03, 0.04)	0.02 (0.02, 0.02)	
White	0	0.00 (0.00, 0.00)	0.02 (0.02, 0.03)	0.07 (0.07, 0.08)	0.22 (0.21, 0.23)	0.04 (0.04, 0.04)	
	10	0.02 (0.02, 0.02)	0.07 (0.07, 0.07)	0.11 (0.10, 0.11)	0.22 (0.21, 0.23)	0.04 (0.04, 0.04)	
	20	0.05 (0.04, 0.05)	0.08 (0.08, 0.09)	0.11 (0.10, 0.11)	0.20 (0.19, 0.21)	0.04 (0.04, 0.04)	
	30	0.04 (0.03, 0.04)	0.06 (0.06, 0.06)	0.08 (0.07, 0.08)	0.15 (0.15, 0.16)	0.04 (0.04, 0.04)	
	40	0.02 (0.02, 0.03)	0.04 (0.04, 0.05)	0.07 (0.06, 0.07)	0.12 (0.11, 0.13)	0.03 (0.03, 0.04)	
	50	0.02 (0.02, 0.02)	0.05 (0.04, 0.05)	0.08 (0.07, 0.08)	0.10 (0.09, 0.10)	0.03 (0.03, 0.03)	
	60	0.03 (0.02, 0.03)	0.06 (0.05, 0.06)	0.08 (0.07, 0.08)	0.08 (0.07, 0.09)	0.03 (0.03, 0.03)	
	70	0.04 (0.03, 0.04)	0.06 (0.05, 0.06)	- (-, -)	0.06 (0.05, 0.07)	0.03 (0.02, 0.03)	
	80	0.03 (0.02, 0.03)	- (-, -)	- (-, -)	0.03 (0.03, 0.04)	0.02 (0.02, 0.02)	
Black	0	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.05 (0.04, 0.06)	0.17 (0.15, 0.19)	0.03 (0.02, 0.03)	
	10	0.02 (0.01, 0.02)	0.05 (0.04, 0.06)	0.08 (0.07, 0.10)	0.17 (0.15, 0.20)	0.03 (0.02, 0.03)	
	20	0.03 (0.03, 0.04)	0.07 (0.06, 0.08)	0.09 (0.08, 0.11)	0.16 (0.14, 0.18)	0.03 (0.02, 0.03)	
	30	0.03 (0.03, 0.04)	0.06 (0.05, 0.07)	0.08 (0.07, 0.09)	0.12 (0.11, 0.15)	0.02 (0.02, 0.03)	
	40	0.03 (0.02, 0.04)	0.05 (0.04, 0.06)	0.07 (0.06, 0.08)	0.09 (0.08, 0.11)	0.02 (0.01, 0.02)	
	50	0.02 (0.02, 0.03)	0.04 (0.03, 0.06)	0.06 (0.05, 0.08)	0.07 (0.05, 0.09)	0.02 (0.01, 0.02)	
	60	0.02 (0.01, 0.03)	0.04 (0.03, 0.06)	0.05 (0.03, 0.06)	0.05 (0.03, 0.07)	0.01 (0.01, 0.02)	
	70	0.02 (0.01, 0.04)	0.03 (0.02, 0.04)	- (-, -)	0.03 (0.02, 0.05)	0.01 (0.01, 0.02)	
	80	0.01 (0.00, 0.02)	- (-, -)	- (-, -)	0.01 (0.00, 0.03)	0.01 (0.00, 0.01)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table IX-9 - continued

HODGKIN LYMPHOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific Islander	0	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.03 (0.03, 0.04)	0.09 (0.08, 0.14)	0.04 (0.02, 0.08)	
	10	0.01 (0.01, 0.01)	0.03 (0.02, 0.04)	0.05 (0.04, 0.06)	0.09 (0.08, 0.14)	0.04 (0.02, 0.08)	
	20	0.02 (0.02, 0.03)	0.04 (0.03, 0.05)	0.05 (0.04, 0.06)	0.09 (0.07, 0.14)	0.03 (0.02, 0.08)	
	30	0.02 (0.01, 0.02)	0.02 (0.02, 0.03)	0.03 (0.02, 0.04)	0.06 (0.05, 0.11)	0.03 (0.02, 0.08)	
	40	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.02 (0.01, 0.03)	0.05 (0.03, 0.10)	0.03 (0.02, 0.08)	
	50	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.04 (0.03, 0.09)	0.03 (0.02, 0.08)	
	60	0.01 (0.00, 0.02)	0.02 (0.01, 0.04)	0.03 (0.02, 0.06)	0.04 (0.02, 0.09)	0.03 (0.02, 0.07)	
	70	0.02 (0.01, 0.03)	0.03 (0.02, 0.05)	- (-, -)	0.03 (0.02, 0.08)	0.03 (0.01, 0.08)	
American Indian/ Alaska Native ^a	80	0.01 (0.00, 0.04)	- (-, -)	- (-, -)	0.01 (0.00, 0.08)	0.03 (0.01, 0.09)	
	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.05)	0.01 (0.00, 0.06)	0.06 (0.01, 0.55)	0.06 (0.02, 0.20)	
	10	0.00 (0.00, 0.03)	0.01 (0.00, 0.05)	0.01 (0.00, 0.05)	0.06 (0.01, 0.56)	0.06 (0.02, 0.21)	
	20	0.01 (0.00, 0.04)	0.01 (0.00, 0.04)	0.01 (0.00, 0.05)	0.05 (0.01, 0.56)	0.06 (0.02, 0.21)	
	30	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.01 (0.00, 0.06)	0.05 (0.00, 0.56)	0.06 (0.02, 0.21)	
	40	0.00 (0.00, 0.03)	0.01 (0.00, 0.06)	0.01 (0.00, 0.08)	0.04 (0.00, 0.57)	0.06 (0.02, 0.21)	
	50	0.01 (0.00, 0.06)	0.01 (0.00, 0.08)	0.01 (0.00, 0.11)	0.04 (0.00, 0.58)	0.06 (0.02, 0.22)	
	60	0.00 (0.00, 0.07)	0.01 (0.00, 0.11)	0.04 (0.00, 0.23)	0.04 (0.00, 0.61)	0.06 (0.02, 0.23)	
Hispanic ^b	70	0.01 (0.00, 0.12)	0.04 (0.00, 0.27)	- (-, -)	0.04 (0.00, 0.70)	0.07 (0.02, 0.26)	
	80	0.05 (0.00, 0.33)	- (-, -)	- (-, -)	0.05 (0.00, 0.92)	0.08 (0.02, 0.36)	
	0	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.04 (0.03, 0.04)	0.17 (0.15, 0.21)	0.05 (0.04, 0.06)	
	10	0.01 (0.01, 0.02)	0.04 (0.03, 0.04)	0.05 (0.05, 0.06)	0.17 (0.15, 0.21)	0.05 (0.04, 0.06)	
	20	0.02 (0.02, 0.03)	0.04 (0.04, 0.05)	0.06 (0.05, 0.06)	0.16 (0.13, 0.20)	0.05 (0.04, 0.06)	
	30	0.02 (0.01, 0.02)	0.03 (0.03, 0.04)	0.05 (0.04, 0.06)	0.14 (0.11, 0.18)	0.05 (0.03, 0.06)	
	40	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.05 (0.04, 0.07)	0.12 (0.10, 0.16)	0.04 (0.03, 0.06)	
	50	0.02 (0.01, 0.02)	0.04 (0.03, 0.05)	0.07 (0.06, 0.09)	0.11 (0.08, 0.15)	0.04 (0.03, 0.06)	
-	60	0.02 (0.02, 0.03)	0.06 (0.04, 0.08)	0.09 (0.06, 0.11)	0.09 (0.07, 0.14)	0.04 (0.03, 0.06)	
	70	0.04 (0.02, 0.05)	0.07 (0.05, 0.09)	- (-, -)	0.08 (0.05, 0.12)	0.04 (0.03, 0.06)	
	80	0.04 (0.02, 0.07)	- (-, -)	- (-, -)	0.05 (0.03, 0.10)	0.03 (0.02, 0.05)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table X-10
KAPOSI SARCOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.05, 0.06)	0.00 (0.00, 0.00)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.05 (0.05, 0.06)	0.00 (0.00, 0.00)	
	20	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.03 (0.02, 0.03)	0.05 (0.05, 0.06)	0.00 (0.00, 0.00)	
	30	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.03 (0.03, 0.03)	0.05 (0.05, 0.05)	0.00 (0.00, 0.00)	
	40	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.02 (0.02, 0.02)	0.04 (0.04, 0.04)	0.00 (0.00, 0.00)	
	50	0.01 (0.00, 0.01)	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.03 (0.03, 0.03)	0.00 (0.00, 0.00)	
	60	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.02 (0.02, 0.03)	0.00 (0.00, 0.00)	
	70	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	- (- , -)	0.02 (0.02, 0.03)	0.00 (0.00, 0.00)	
	80	0.01 (0.01, 0.02)	- (- , -)	- (- , -)	0.02 (0.02, 0.02)	0.00 (0.00, 0.00)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.05, 0.05)	0.00 (0.00, 0.00)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.05, 0.05)	0.00 (0.00, 0.00)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.05 (0.05, 0.05)	0.00 (0.00, 0.00)	
	30	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.02 (0.02, 0.03)	0.05 (0.04, 0.05)	0.00 (0.00, 0.00)	
	40	0.01 (0.01, 0.01)	0.01 (0.01, 0.02)	0.02 (0.02, 0.02)	0.04 (0.03, 0.04)	0.00 (0.00, 0.00)	
	50	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.03 (0.03, 0.03)	0.00 (0.00, 0.00)	
	60	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.02 (0.02, 0.02)	0.02 (0.02, 0.03)	0.00 (0.00, 0.00)	
	70	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	- (- , -)	0.02 (0.02, 0.03)	0.00 (0.00, 0.00)	
	80	0.01 (0.01, 0.02)	- (- , -)	- (- , -)	0.02 (0.02, 0.02)	0.00 (0.00, 0.00)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.08 (0.07, 0.09)	0.01 (0.01, 0.01)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.04, 0.05)	0.08 (0.07, 0.09)	0.01 (0.01, 0.01)	
	20	0.01 (0.01, 0.01)	0.04 (0.03, 0.05)	0.06 (0.05, 0.07)	0.08 (0.07, 0.09)	0.01 (0.01, 0.01)	
	30	0.03 (0.03, 0.04)	0.05 (0.04, 0.06)	0.06 (0.05, 0.07)	0.07 (0.06, 0.08)	0.01 (0.01, 0.01)	
	40	0.02 (0.02, 0.03)	0.03 (0.03, 0.04)	0.03 (0.03, 0.04)	0.04 (0.04, 0.05)	0.00 (0.00, 0.01)	
	50	0.01 (0.01, 0.01)	0.01 (0.01, 0.02)	0.02 (0.01, 0.02)	0.02 (0.02, 0.03)	0.00 (0.00, 0.00)	
	60	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	70	0.00 (0.00, 0.01)	0.01 (0.00, 0.02)	- (- , -)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	80	0.01 (0.00, 0.02)	- (- , -)	- (- , -)	0.01 (0.00, 0.03)	0.00 (0.00, 0.00)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table X-10 - continued

KAPOSI SARCOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.01, 0.04)	0.00 (0.00, 0.02)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.04)	0.00 (0.00, 0.02)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.02 (0.01, 0.04)	0.00 (0.00, 0.02)	
	30	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.01 (0.01, 0.01)	0.02 (0.01, 0.04)	0.00 (0.00, 0.02)	
	40	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.01 (0.01, 0.01)	0.02 (0.01, 0.04)	0.00 (0.00, 0.02)	
	50	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.01 (0.01, 0.04)	0.00 (0.00, 0.02)	
	60	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.01 (0.01, 0.04)	0.00 (0.00, 0.03)	
	70	0.00 (0.00, 0.01)	0.01 (0.00, 0.02)	- (- , -)	0.01 (0.00, 0.04)	0.00 (0.00, 0.03)	
	80	0.01 (0.00, 0.02)	- (- , -)	- (- , -)	0.01 (0.00, 0.04)	0.00 (0.00, 0.03)	
American Indian/ Alaska Native ^a	0	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.03 (0.01, 0.30)	0.00 (0.00, 0.08)	
	10	0.00 (0.00, 0.01)	0.00 (0.00, 0.02)	0.02 (0.00, 0.04)	0.03 (0.01, 0.30)	0.00 (0.00, 0.08)	
	20	0.00 (0.00, 0.02)	0.02 (0.00, 0.04)	0.03 (0.01, 0.05)	0.03 (0.01, 0.31)	0.00 (0.00, 0.08)	
	30	0.01 (0.00, 0.03)	0.02 (0.01, 0.05)	0.02 (0.01, 0.05)	0.02 (0.01, 0.31)	0.00 (0.00, 0.08)	
	40	0.01 (0.00, 0.03)	0.01 (0.00, 0.04)	0.01 (0.00, 0.05)	0.01 (0.00, 0.31)	0.00 (0.00, 0.08)	
	50	0.00 (0.00, 0.02)	0.00 (0.00, 0.04)	0.00 (0.00, 0.05)	0.00 (0.00, 0.32)	0.00 (0.00, 0.09)	
	60	0.00 (0.00, 0.04)	0.00 (0.00, 0.06)	0.00 (0.00, 0.09)	0.00 (0.00, 0.34)	0.00 (0.00, 0.10)	
	70	0.00 (0.00, 0.07)	0.00 (0.00, 0.11)	- (- , -)	0.00 (0.00, 0.40)	0.00 (0.00, 0.12)	
	80	0.00 (0.00, 0.15)	- (- , -)	- (- , -)	0.00 (0.00, 0.56)	0.00 (0.00, 0.17)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.10 (0.08, 0.12)	0.01 (0.01, 0.02)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.10 (0.08, 0.12)	0.01 (0.01, 0.02)	
	20	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.03 (0.03, 0.04)	0.10 (0.08, 0.13)	0.01 (0.01, 0.02)	
	30	0.01 (0.01, 0.02)	0.03 (0.02, 0.03)	0.04 (0.03, 0.04)	0.10 (0.08, 0.12)	0.01 (0.01, 0.02)	
	40	0.01 (0.01, 0.02)	0.02 (0.02, 0.03)	0.03 (0.03, 0.04)	0.08 (0.07, 0.11)	0.01 (0.01, 0.02)	
	50	0.01 (0.01, 0.01)	0.02 (0.02, 0.03)	0.04 (0.03, 0.05)	0.07 (0.06, 0.10)	0.01 (0.01, 0.02)	
	60	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.06 (0.04, 0.07)	0.06 (0.05, 0.09)	0.01 (0.01, 0.02)	
	70	0.02 (0.02, 0.03)	0.05 (0.04, 0.07)	- (- , -)	0.06 (0.04, 0.09)	0.01 (0.01, 0.02)	
	80	0.03 (0.02, 0.05)	- (- , -)	- (- , -)	0.05 (0.03, 0.08)	0.01 (0.01, 0.03)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table X-10 - continued

KAPOSI SARCOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.09 (0.09, 0.10)	0.01 (0.01, 0.01)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.09 (0.09, 0.10)	0.01 (0.01, 0.01)	
	20	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.05 (0.05, 0.05)	0.09 (0.09, 0.10)	0.01 (0.01, 0.01)	
	30	0.02 (0.02, 0.03)	0.04 (0.04, 0.05)	0.06 (0.05, 0.06)	0.09 (0.08, 0.09)	0.01 (0.01, 0.01)	
	40	0.02 (0.02, 0.02)	0.03 (0.03, 0.04)	0.04 (0.04, 0.04)	0.07 (0.06, 0.07)	0.00 (0.00, 0.01)	
	50	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.03 (0.03, 0.03)	0.05 (0.04, 0.05)	0.00 (0.00, 0.00)	
	60	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.03 (0.03, 0.04)	0.04 (0.03, 0.04)	0.00 (0.00, 0.00)	
	70	0.01 (0.01, 0.02)	0.03 (0.02, 0.03)	- (-, -)	0.03 (0.03, 0.04)	0.00 (0.00, 0.00)	
	80	0.02 (0.02, 0.03)	- (-, -)	- (-, -)	0.03 (0.02, 0.04)	0.00 (0.00, 0.00)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.08 (0.08, 0.09)	0.01 (0.00, 0.01)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.09 (0.08, 0.09)	0.01 (0.00, 0.01)	
	20	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.04 (0.04, 0.05)	0.09 (0.08, 0.09)	0.01 (0.00, 0.01)	
	30	0.02 (0.02, 0.02)	0.04 (0.04, 0.04)	0.05 (0.04, 0.05)	0.08 (0.08, 0.09)	0.01 (0.00, 0.01)	
	40	0.02 (0.02, 0.02)	0.03 (0.03, 0.03)	0.04 (0.03, 0.04)	0.06 (0.06, 0.07)	0.00 (0.00, 0.01)	
	50	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.03 (0.03, 0.03)	0.05 (0.04, 0.05)	0.00 (0.00, 0.00)	
	60	0.01 (0.01, 0.01)	0.02 (0.02, 0.03)	0.03 (0.03, 0.04)	0.04 (0.03, 0.04)	0.00 (0.00, 0.00)	
	70	0.02 (0.01, 0.02)	0.03 (0.03, 0.03)	- (-, -)	0.04 (0.03, 0.04)	0.00 (0.00, 0.00)	
	80	0.02 (0.02, 0.03)	- (-, -)	- (-, -)	0.03 (0.02, 0.04)	0.00 (0.00, 0.01)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.03)	0.15 (0.13, 0.18)	0.02 (0.01, 0.02)	
	10	0.00 (0.00, 0.00)	0.02 (0.02, 0.03)	0.08 (0.07, 0.09)	0.16 (0.14, 0.18)	0.02 (0.01, 0.02)	
	20	0.02 (0.02, 0.03)	0.08 (0.07, 0.09)	0.12 (0.11, 0.14)	0.16 (0.14, 0.18)	0.02 (0.01, 0.02)	
	30	0.06 (0.05, 0.07)	0.10 (0.09, 0.12)	0.12 (0.11, 0.14)	0.14 (0.12, 0.16)	0.01 (0.01, 0.02)	
	40	0.04 (0.03, 0.05)	0.06 (0.05, 0.07)	0.07 (0.06, 0.08)	0.08 (0.06, 0.10)	0.01 (0.01, 0.01)	
	50	0.02 (0.01, 0.03)	0.03 (0.02, 0.04)	0.03 (0.02, 0.05)	0.04 (0.03, 0.06)	0.00 (0.00, 0.01)	
	60	0.01 (0.00, 0.02)	0.02 (0.01, 0.03)	0.02 (0.01, 0.03)	0.02 (0.01, 0.04)	0.00 (0.00, 0.01)	
	70	0.01 (0.00, 0.02)	0.01 (0.00, 0.03)	- (-, -)	0.01 (0.00, 0.04)	0.00 (0.00, 0.01)	
	80	0.01 (0.00, 0.03)	- (-, -)	- (-, -)	0.01 (0.00, 0.07)	0.00 (0.00, 0.01)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table X-10 - continued

KAPOSI SARCOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.04 (0.03, 0.08)	0.00 (0.00, 0.04)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	0.04 (0.03, 0.08)	0.00 (0.00, 0.04)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.01 (0.01, 0.02)	0.04 (0.03, 0.09)	0.00 (0.00, 0.04)	
	30	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.02 (0.01, 0.02)	0.04 (0.02, 0.08)	0.00 (0.00, 0.04)	
	40	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.02 (0.01, 0.03)	0.03 (0.02, 0.08)	0.00 (0.00, 0.04)	
	50	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.02 (0.01, 0.03)	0.03 (0.01, 0.07)	0.00 (0.00, 0.04)	
	60	0.01 (0.00, 0.02)	0.01 (0.01, 0.03)	0.02 (0.01, 0.04)	0.02 (0.01, 0.07)	0.00 (0.00, 0.05)	
	70	0.01 (0.00, 0.02)	0.02 (0.00, 0.04)	- (-, -)	0.02 (0.00, 0.07)	0.00 (0.00, 0.05)	
	80	0.01 (0.00, 0.04)	- (-, -)	- (-, -)	0.01 (0.00, 0.09)	0.00 (0.00, 0.07)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.01 (0.00, 0.05)	0.05 (0.02, 0.72)	0.01 (0.00, 0.19)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.01 (0.00, 0.04)	0.03 (0.01, 0.08)	0.05 (0.02, 0.73)	0.01 (0.00, 0.19)	
	20	0.01 (0.00, 0.04)	0.03 (0.01, 0.08)	0.05 (0.02, 0.11)	0.05 (0.02, 0.74)	0.01 (0.00, 0.20)	
	30	0.02 (0.01, 0.06)	0.04 (0.02, 0.09)	0.04 (0.02, 0.10)	0.04 (0.02, 0.75)	0.01 (0.00, 0.20)	
	40	0.02 (0.00, 0.06)	0.02 (0.00, 0.07)	0.02 (0.00, 0.10)	0.02 (0.00, 0.77)	0.00 (0.00, 0.21)	
	50	0.00 (0.00, 0.04)	0.00 (0.00, 0.07)	0.00 (0.00, 0.11)	0.00 (0.00, 0.79)	0.00 (0.00, 0.22)	
	60	0.00 (0.00, 0.08)	0.00 (0.00, 0.12)	0.00 (0.00, 0.22)	0.00 (0.00, 0.86)	0.00 (0.00, 0.24)	
	70	0.00 (0.00, 0.15)	0.00 (0.00, 0.27)	- (-, -)	0.00 (0.00, 1.03)	0.00 (0.00, 0.30)	
	80	0.00 (0.00, 0.40)	- (-, -)	- (-, -)	0.00 (0.00, 1.56)	0.00 (0.00, 0.47)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.16 (0.13, 0.21)	0.01 (0.01, 0.03)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.03, 0.04)	0.16 (0.13, 0.21)	0.01 (0.01, 0.03)	
	20	0.01 (0.00, 0.01)	0.03 (0.03, 0.04)	0.06 (0.05, 0.07)	0.16 (0.13, 0.21)	0.01 (0.01, 0.03)	
	30	0.03 (0.02, 0.03)	0.05 (0.05, 0.06)	0.07 (0.06, 0.08)	0.15 (0.13, 0.21)	0.01 (0.01, 0.03)	
	40	0.03 (0.02, 0.03)	0.05 (0.04, 0.06)	0.06 (0.05, 0.08)	0.13 (0.10, 0.19)	0.01 (0.01, 0.03)	
	50	0.02 (0.01, 0.03)	0.04 (0.03, 0.05)	0.07 (0.06, 0.09)	0.11 (0.08, 0.16)	0.01 (0.01, 0.03)	
	60	0.02 (0.01, 0.03)	0.06 (0.04, 0.08)	0.08 (0.06, 0.11)	0.09 (0.07, 0.15)	0.01 (0.01, 0.03)	
	70	0.04 (0.03, 0.06)	0.07 (0.05, 0.10)	- (-, -)	0.08 (0.05, 0.15)	0.01 (0.01, 0.03)	
	80	0.04 (0.02, 0.07)	- (-, -)	- (-, -)	0.06 (0.03, 0.15)	0.01 (0.00, 0.04)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table X-10 - continued

KAPOSI SARCOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	40	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	50	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	60	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.01 (0.01, 0.01)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	70	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	- (-, -)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	80	0.01 (0.01, 0.01)	- (-, -)	- (-, -)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	40	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	50	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	60	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.01 (0.01, 0.01)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	70	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	- (-, -)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	80	0.01 (0.01, 0.01)	- (-, -)	- (-, -)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	40	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.00 (0.00, 0.00)	
	50	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.01 (0.00, 0.02)	0.00 (0.00, 0.00)	
	60	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.01 (0.00, 0.02)	0.01 (0.00, 0.02)	0.00 (0.00, 0.00)	
	70	0.00 (0.00, 0.01)	0.01 (0.00, 0.02)	- (-, -)	0.01 (0.00, 0.02)	0.00 (0.00, 0.00)	
	80	0.01 (0.00, 0.03)	- (-, -)	- (-, -)	0.01 (0.00, 0.03)	0.00 (0.00, 0.00)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table X-10 - continued

KAPOSI SARCOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.05)	0.00 (0.00, 0.05)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.05)	0.00 (0.00, 0.05)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.05)	0.00 (0.00, 0.05)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.00 (0.00, 0.05)	0.00 (0.00, 0.05)	
	40	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.00 (0.00, 0.05)	0.00 (0.00, 0.05)	
	50	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.00 (0.00, 0.05)	0.00 (0.00, 0.05)	
	60	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.00 (0.00, 0.02)	0.00 (0.00, 0.06)	0.00 (0.00, 0.05)	
	70	0.00 (0.00, 0.01)	0.00 (0.00, 0.02)	- (- , -)	0.00 (0.00, 0.06)	0.00 (0.00, 0.05)	
	80	0.00 (0.00, 0.02)	- (- , -)	- (- , -)	0.00 (0.00, 0.07)	0.01 (0.00, 0.06)	
American Indian/ Alaska Native ^a	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.50)	0.00 (0.00, 0.14)	
	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.00 (0.00, 0.51)	0.00 (0.00, 0.14)	
	20	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.00 (0.00, 0.51)	0.00 (0.00, 0.14)	
	30	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.00 (0.00, 0.04)	0.00 (0.00, 0.52)	0.00 (0.00, 0.14)	
	40	0.00 (0.00, 0.03)	0.00 (0.00, 0.04)	0.00 (0.00, 0.07)	0.00 (0.00, 0.53)	0.00 (0.00, 0.14)	
	50	0.00 (0.00, 0.04)	0.00 (0.00, 0.07)	0.00 (0.00, 0.10)	0.00 (0.00, 0.54)	0.00 (0.00, 0.15)	
	60	0.00 (0.00, 0.07)	0.00 (0.00, 0.10)	0.00 (0.00, 0.17)	0.00 (0.00, 0.57)	0.00 (0.00, 0.16)	
	70	0.00 (0.00, 0.12)	0.00 (0.00, 0.19)	- (- , -)	0.00 (0.00, 0.66)	0.00 (0.00, 0.19)	
	80	0.00 (0.00, 0.25)	- (- , -)	- (- , -)	0.00 (0.00, 0.88)	0.00 (0.00, 0.26)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.04 (0.02, 0.08)	0.01 (0.00, 0.03)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.04 (0.02, 0.08)	0.01 (0.00, 0.03)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.04 (0.02, 0.08)	0.01 (0.00, 0.03)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.04 (0.02, 0.08)	0.01 (0.00, 0.03)	
	40	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.04 (0.02, 0.08)	0.01 (0.00, 0.03)	
	50	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.04 (0.02, 0.08)	0.01 (0.00, 0.03)	
	60	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.03 (0.02, 0.06)	0.04 (0.02, 0.08)	0.01 (0.00, 0.03)	
	70	0.01 (0.00, 0.02)	0.03 (0.02, 0.06)	- (- , -)	0.04 (0.02, 0.09)	0.01 (0.00, 0.03)	
	80	0.03 (0.01, 0.05)	- (- , -)	- (- , -)	0.04 (0.02, 0.09)	0.01 (0.01, 0.04)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XI-9

KIDNEY AND RENAL PELVIS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.01 (0.01, 0.01)	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	1.34 (1.33, 1.36)	0.46 (0.46, 0.47)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	1.35 (1.33, 1.37)	0.47 (0.46, 0.47)	
	20	0.01 (0.00, 0.01)	0.03 (0.03, 0.03)	0.12 (0.12, 0.12)	1.35 (1.34, 1.37)	0.47 (0.46, 0.47)	
	30	0.03 (0.02, 0.03)	0.12 (0.11, 0.12)	0.33 (0.33, 0.34)	1.36 (1.34, 1.38)	0.47 (0.47, 0.48)	
	40	0.09 (0.09, 0.09)	0.31 (0.30, 0.31)	0.69 (0.68, 0.70)	1.35 (1.33, 1.37)	0.48 (0.47, 0.48)	
	50	0.22 (0.22, 0.23)	0.61 (0.60, 0.62)	1.03 (1.02, 1.04)	1.30 (1.28, 1.31)	0.47 (0.47, 0.48)	
	60	0.41 (0.41, 0.42)	0.86 (0.84, 0.87)	1.10 (1.08, 1.11)	1.14 (1.13, 1.16)	0.45 (0.44, 0.45)	
	70	0.51 (0.50, 0.52)	0.79 (0.77, 0.80)	- (- , -)	0.84 (0.83, 0.86)	0.38 (0.37, 0.38)	
	80	0.39 (0.38, 0.41)	- (- , -)	- (- , -)	0.47 (0.46, 0.48)	0.28 (0.27, 0.28)	
White	0	0.01 (0.01, 0.01)	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	1.40 (1.38, 1.42)	0.48 (0.47, 0.49)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	1.40 (1.39, 1.42)	0.48 (0.48, 0.49)	
	20	0.01 (0.00, 0.01)	0.03 (0.03, 0.03)	0.12 (0.12, 0.13)	1.41 (1.39, 1.43)	0.49 (0.48, 0.49)	
	30	0.03 (0.03, 0.03)	0.12 (0.11, 0.12)	0.34 (0.33, 0.35)	1.41 (1.40, 1.43)	0.49 (0.48, 0.50)	
	40	0.09 (0.09, 0.10)	0.32 (0.31, 0.33)	0.71 (0.70, 0.73)	1.40 (1.39, 1.42)	0.49 (0.49, 0.50)	
	50	0.23 (0.23, 0.24)	0.64 (0.63, 0.65)	1.07 (1.06, 1.09)	1.35 (1.33, 1.37)	0.49 (0.48, 0.50)	
	60	0.43 (0.42, 0.44)	0.89 (0.88, 0.91)	1.14 (1.12, 1.16)	1.19 (1.17, 1.20)	0.46 (0.46, 0.47)	
	70	0.53 (0.52, 0.55)	0.82 (0.80, 0.83)	- (- , -)	0.87 (0.85, 0.89)	0.39 (0.38, 0.39)	
	80	0.41 (0.39, 0.42)	- (- , -)	- (- , -)	0.48 (0.47, 0.50)	0.28 (0.27, 0.29)	
Black	0	0.01 (0.01, 0.01)	0.01 (0.01, 0.02)	0.02 (0.01, 0.02)	1.20 (1.15, 1.25)	0.36 (0.35, 0.38)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.03, 0.05)	1.22 (1.17, 1.27)	0.37 (0.36, 0.39)	
	20	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	0.14 (0.13, 0.15)	1.22 (1.18, 1.28)	0.37 (0.36, 0.39)	
	30	0.03 (0.03, 0.04)	0.14 (0.12, 0.15)	0.37 (0.35, 0.39)	1.24 (1.19, 1.29)	0.38 (0.37, 0.39)	
	40	0.11 (0.10, 0.12)	0.35 (0.33, 0.37)	0.72 (0.69, 0.76)	1.24 (1.19, 1.29)	0.38 (0.37, 0.40)	
	50	0.26 (0.24, 0.27)	0.65 (0.62, 0.68)	1.00 (0.95, 1.04)	1.19 (1.14, 1.25)	0.38 (0.37, 0.40)	
	60	0.44 (0.41, 0.47)	0.83 (0.79, 0.88)	1.01 (0.96, 1.06)	1.05 (1.00, 1.11)	0.37 (0.35, 0.38)	
	70	0.49 (0.45, 0.53)	0.71 (0.67, 0.77)	- (- , -)	0.76 (0.71, 0.82)	0.32 (0.31, 0.34)	
	80	0.36 (0.32, 0.41)	- (- , -)	- (- , -)	0.44 (0.39, 0.50)	0.25 (0.23, 0.27)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XI-9 - continued

KIDNEY AND RENAL PELVIS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/Pacific Islander	0	0.01 (0.00, 0.01)	0.01 (0.00, 0.01)	0.01 (0.00, 0.01)	0.83 (0.77, 0.90)	0.26 (0.22, 0.30)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.83 (0.77, 0.90)	0.26 (0.22, 0.31)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.06 (0.05, 0.06)	0.83 (0.77, 0.90)	0.26 (0.22, 0.31)	
	30	0.01 (0.01, 0.02)	0.05 (0.05, 0.06)	0.16 (0.15, 0.18)	0.83 (0.78, 0.90)	0.26 (0.23, 0.31)	
	40	0.04 (0.04, 0.05)	0.15 (0.14, 0.17)	0.34 (0.32, 0.37)	0.83 (0.77, 0.90)	0.26 (0.22, 0.31)	
	50	0.11 (0.10, 0.12)	0.31 (0.28, 0.33)	0.55 (0.52, 0.59)	0.80 (0.74, 0.87)	0.26 (0.22, 0.31)	
	60	0.20 (0.18, 0.22)	0.46 (0.42, 0.50)	0.64 (0.60, 0.70)	0.71 (0.65, 0.78)	0.25 (0.21, 0.30)	
	70	0.28 (0.25, 0.31)	0.48 (0.44, 0.53)	- (- , -)	0.55 (0.49, 0.62)	0.22 (0.18, 0.27)	
	80	0.25 (0.21, 0.29)	- (- , -)	- (- , -)	0.34 (0.28, 0.42)	0.17 (0.12, 0.23)	
American Indian/Alaska Native ^a	0	0.02 (0.01, 0.05)	0.02 (0.01, 0.05)	0.02 (0.01, 0.06)	1.53 (1.28, 1.92)	0.68 (0.58, 0.82)	
	10	0.00 (0.00, 0.01)	0.00 (0.00, 0.02)	0.04 (0.02, 0.07)	1.53 (1.28, 1.93)	0.69 (0.59, 0.84)	
	20	0.00 (0.00, 0.02)	0.04 (0.02, 0.07)	0.15 (0.11, 0.20)	1.55 (1.29, 1.94)	0.70 (0.59, 0.84)	
	30	0.04 (0.02, 0.07)	0.15 (0.11, 0.20)	0.41 (0.33, 0.50)	1.57 (1.31, 1.97)	0.71 (0.60, 0.86)	
	40	0.11 (0.08, 0.16)	0.38 (0.30, 0.47)	0.75 (0.62, 0.91)	1.57 (1.31, 1.99)	0.73 (0.62, 0.88)	
	50	0.28 (0.21, 0.36)	0.67 (0.54, 0.82)	1.20 (0.99, 1.45)	1.52 (1.25, 1.95)	0.74 (0.62, 0.89)	
	60	0.42 (0.31, 0.55)	1.00 (0.79, 1.24)	1.30 (1.04, 1.62)	1.34 (1.06, 1.79)	0.70 (0.58, 0.87)	
	70	0.68 (0.50, 0.91)	1.03 (0.77, 1.36)	- (- , -)	1.08 (0.80, 1.58)	0.60 (0.47, 0.80)	
	80	0.49 (0.29, 0.81)	- (- , -)	- (- , -)	0.56 (0.32, 1.17)	0.39 (0.25, 0.63)	
Hispanic ^b	0	0.01 (0.01, 0.01)	0.01 (0.01, 0.01)	0.01 (0.01, 0.02)	1.42 (1.36, 1.49)	0.48 (0.45, 0.51)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	1.43 (1.37, 1.50)	0.48 (0.46, 0.51)	
	20	0.00 (0.00, 0.01)	0.03 (0.02, 0.03)	0.11 (0.10, 0.12)	1.44 (1.37, 1.51)	0.48 (0.46, 0.51)	
	30	0.02 (0.02, 0.02)	0.11 (0.10, 0.11)	0.32 (0.31, 0.34)	1.44 (1.38, 1.51)	0.49 (0.46, 0.52)	
	40	0.09 (0.08, 0.09)	0.30 (0.29, 0.32)	0.68 (0.65, 0.71)	1.44 (1.37, 1.51)	0.49 (0.46, 0.52)	
	50	0.22 (0.21, 0.24)	0.61 (0.58, 0.64)	1.06 (1.01, 1.10)	1.38 (1.32, 1.45)	0.49 (0.46, 0.52)	
	60	0.40 (0.38, 0.43)	0.88 (0.83, 0.92)	1.16 (1.10, 1.22)	1.22 (1.15, 1.29)	0.46 (0.43, 0.49)	
	70	0.53 (0.49, 0.57)	0.85 (0.79, 0.91)	- (- , -)	0.91 (0.85, 0.99)	0.39 (0.36, 0.42)	
	80	0.42 (0.37, 0.47)	- (- , -)	- (- , -)	0.51 (0.44, 0.59)	0.29 (0.26, 0.33)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XI-9 - continued

KIDNEY AND RENAL PELVIS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.01 (0.01, 0.01)	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	1.71 (1.68, 1.73)	0.59 (0.58, 0.60)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	1.72 (1.69, 1.75)	0.60 (0.59, 0.61)	
	20	0.01 (0.00, 0.01)	0.04 (0.03, 0.04)	0.15 (0.14, 0.16)	1.73 (1.70, 1.75)	0.60 (0.60, 0.61)	
	30	0.03 (0.03, 0.03)	0.15 (0.14, 0.15)	0.43 (0.42, 0.44)	1.74 (1.72, 1.77)	0.61 (0.60, 0.62)	
	40	0.12 (0.11, 0.12)	0.41 (0.40, 0.42)	0.91 (0.89, 0.93)	1.74 (1.71, 1.77)	0.62 (0.61, 0.63)	
	50	0.30 (0.29, 0.31)	0.82 (0.81, 0.84)	1.36 (1.34, 1.39)	1.69 (1.66, 1.71)	0.62 (0.61, 0.63)	
	60	0.57 (0.55, 0.58)	1.15 (1.13, 1.17)	1.45 (1.42, 1.47)	1.50 (1.47, 1.53)	0.58 (0.57, 0.59)	
	70	0.70 (0.68, 0.72)	1.05 (1.03, 1.08)	- (- , -)	1.12 (1.09, 1.14)	0.49 (0.48, 0.50)	
	80	0.55 (0.52, 0.57)	- (- , -)	- (- , -)	0.65 (0.62, 0.67)	0.37 (0.36, 0.38)	
White	0	0.01 (0.01, 0.01)	0.01 (0.01, 0.02)	0.02 (0.02, 0.02)	1.78 (1.75, 1.81)	0.62 (0.61, 0.63)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.04, 0.04)	1.78 (1.75, 1.82)	0.62 (0.62, 0.63)	
	20	0.01 (0.00, 0.01)	0.04 (0.03, 0.04)	0.15 (0.15, 0.16)	1.79 (1.76, 1.82)	0.63 (0.62, 0.64)	
	30	0.03 (0.03, 0.04)	0.15 (0.14, 0.16)	0.44 (0.43, 0.45)	1.81 (1.78, 1.84)	0.63 (0.63, 0.64)	
	40	0.12 (0.11, 0.12)	0.41 (0.40, 0.42)	0.94 (0.92, 0.96)	1.81 (1.77, 1.84)	0.64 (0.63, 0.65)	
	50	0.30 (0.29, 0.31)	0.85 (0.83, 0.87)	1.42 (1.39, 1.44)	1.75 (1.72, 1.78)	0.64 (0.63, 0.65)	
	60	0.59 (0.57, 0.60)	1.20 (1.17, 1.23)	1.50 (1.47, 1.53)	1.56 (1.52, 1.59)	0.60 (0.59, 0.61)	
	70	0.73 (0.71, 0.75)	1.09 (1.06, 1.12)	- (- , -)	1.15 (1.12, 1.18)	0.51 (0.50, 0.52)	
	80	0.56 (0.54, 0.58)	- (- , -)	- (- , -)	0.66 (0.63, 0.69)	0.38 (0.36, 0.39)	
Black	0	0.01 (0.01, 0.01)	0.01 (0.01, 0.02)	0.02 (0.01, 0.03)	1.49 (1.42, 1.58)	0.45 (0.43, 0.47)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.04, 0.05)	1.52 (1.45, 1.61)	0.46 (0.44, 0.48)	
	20	0.01 (0.00, 0.01)	0.04 (0.04, 0.05)	0.18 (0.16, 0.20)	1.54 (1.46, 1.62)	0.46 (0.44, 0.48)	
	30	0.04 (0.03, 0.04)	0.18 (0.16, 0.19)	0.51 (0.48, 0.55)	1.57 (1.49, 1.65)	0.47 (0.45, 0.49)	
	40	0.14 (0.13, 0.16)	0.49 (0.46, 0.53)	0.99 (0.94, 1.05)	1.58 (1.50, 1.67)	0.48 (0.45, 0.50)	
	50	0.37 (0.34, 0.40)	0.90 (0.85, 0.96)	1.31 (1.24, 1.39)	1.53 (1.45, 1.63)	0.48 (0.46, 0.51)	
	60	0.62 (0.57, 0.67)	1.09 (1.02, 1.17)	1.31 (1.22, 1.40)	1.35 (1.26, 1.45)	0.46 (0.44, 0.49)	
	70	0.64 (0.57, 0.71)	0.92 (0.83, 1.01)	- (- , -)	0.98 (0.88, 1.08)	0.41 (0.38, 0.44)	
	80	0.52 (0.43, 0.61)	- (- , -)	- (- , -)	0.62 (0.52, 0.75)	0.31 (0.27, 0.35)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XI-9 - continued

KIDNEY AND RENAL PELVIS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.01 (0.00, 0.01)	0.01 (0.00, 0.01)	0.01 (0.00, 0.01)	1.11 (1.01, 1.22)	0.34 (0.28, 0.43)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	1.11 (1.02, 1.23)	0.35 (0.29, 0.43)	
	20	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.07 (0.06, 0.08)	1.12 (1.02, 1.23)	0.35 (0.29, 0.43)	
	30	0.02 (0.01, 0.02)	0.07 (0.06, 0.08)	0.22 (0.20, 0.25)	1.12 (1.03, 1.24)	0.35 (0.29, 0.43)	
	40	0.05 (0.04, 0.06)	0.21 (0.19, 0.24)	0.49 (0.44, 0.53)	1.12 (1.02, 1.24)	0.35 (0.29, 0.43)	
	50	0.16 (0.14, 0.18)	0.44 (0.40, 0.49)	0.77 (0.70, 0.83)	1.08 (0.98, 1.20)	0.35 (0.28, 0.43)	
	60	0.29 (0.26, 0.33)	0.63 (0.57, 0.70)	0.88 (0.80, 0.98)	0.97 (0.87, 1.09)	0.34 (0.27, 0.43)	
	70	0.38 (0.33, 0.43)	0.66 (0.58, 0.75)	- (- , -)	0.75 (0.65, 0.88)	0.29 (0.22, 0.39)	
	80	0.37 (0.30, 0.45)	- (- , -)	- (- , -)	0.49 (0.38, 0.63)	0.23 (0.15, 0.35)	
American	0	0.01 (0.00, 0.06)	0.01 (0.00, 0.06)	0.02 (0.00, 0.07)	1.78 (1.41, 2.54)	0.84 (0.69, 1.08)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.01 (0.00, 0.04)	0.06 (0.03, 0.11)	1.81 (1.43, 2.58)	0.86 (0.70, 1.11)	
	20	0.01 (0.00, 0.04)	0.06 (0.03, 0.11)	0.18 (0.12, 0.27)	1.83 (1.45, 2.61)	0.87 (0.71, 1.12)	
	30	0.05 (0.02, 0.10)	0.18 (0.12, 0.26)	0.49 (0.37, 0.65)	1.87 (1.48, 2.66)	0.89 (0.73, 1.15)	
	40	0.13 (0.08, 0.20)	0.46 (0.34, 0.61)	0.92 (0.71, 1.17)	1.88 (1.48, 2.70)	0.92 (0.75, 1.19)	
	50	0.34 (0.24, 0.48)	0.83 (0.62, 1.09)	1.49 (1.16, 1.90)	1.84 (1.43, 2.71)	0.96 (0.78, 1.25)	
	60	0.53 (0.36, 0.76)	1.26 (0.93, 1.68)	1.63 (1.21, 2.20)	1.64 (1.21, 2.58)	0.93 (0.74, 1.25)	
	70	0.87 (0.57, 1.28)	1.32 (0.89, 1.93)	- (- , -)	1.33 (0.89, 2.42)	0.78 (0.58, 1.15)	
	80	0.68 (0.32, 1.34)	- (- , -)	- (- , -)	0.69 (0.32, 2.25)	0.42 (0.23, 0.93)	
Hispanic ^b	0	0.01 (0.01, 0.01)	0.01 (0.01, 0.01)	0.02 (0.01, 0.02)	1.77 (1.67, 1.89)	0.63 (0.59, 0.69)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.03, 0.04)	1.78 (1.68, 1.90)	0.64 (0.60, 0.70)	
	20	0.01 (0.00, 0.01)	0.03 (0.03, 0.04)	0.14 (0.12, 0.15)	1.79 (1.69, 1.91)	0.64 (0.60, 0.70)	
	30	0.03 (0.02, 0.03)	0.13 (0.12, 0.14)	0.40 (0.37, 0.43)	1.80 (1.70, 1.93)	0.65 (0.60, 0.71)	
	40	0.11 (0.10, 0.12)	0.38 (0.35, 0.40)	0.87 (0.82, 0.92)	1.80 (1.70, 1.93)	0.66 (0.61, 0.71)	
	50	0.28 (0.26, 0.30)	0.79 (0.74, 0.84)	1.37 (1.29, 1.45)	1.75 (1.64, 1.88)	0.66 (0.61, 0.71)	
	60	0.54 (0.50, 0.59)	1.16 (1.09, 1.24)	1.49 (1.39, 1.60)	1.57 (1.46, 1.70)	0.63 (0.58, 0.69)	
	70	0.72 (0.65, 0.79)	1.09 (1.00, 1.20)	- (- , -)	1.19 (1.07, 1.33)	0.54 (0.48, 0.61)	
	80	0.54 (0.45, 0.63)	- (- , -)	- (- , -)	0.67 (0.55, 0.83)	0.41 (0.35, 0.50)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XI-9 - continued

KIDNEY AND RENAL PELVIS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.01 (0.01, 0.01)	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	1.01 (0.99, 1.03)	0.35 (0.34, 0.35)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	1.01 (0.99, 1.03)	0.35 (0.34, 0.35)	
	20	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	0.09 (0.09, 0.09)	1.01 (0.99, 1.03)	0.35 (0.34, 0.35)	
	30	0.02 (0.02, 0.02)	0.08 (0.08, 0.09)	0.23 (0.23, 0.24)	1.01 (0.99, 1.03)	0.35 (0.34, 0.36)	
	40	0.06 (0.06, 0.07)	0.21 (0.21, 0.22)	0.47 (0.46, 0.48)	1.00 (0.98, 1.02)	0.35 (0.35, 0.36)	
	50	0.15 (0.15, 0.16)	0.42 (0.40, 0.43)	0.73 (0.71, 0.74)	0.95 (0.93, 0.97)	0.35 (0.34, 0.36)	
	60	0.28 (0.27, 0.29)	0.60 (0.59, 0.62)	0.80 (0.78, 0.82)	0.84 (0.82, 0.86)	0.34 (0.33, 0.34)	
	70	0.37 (0.35, 0.38)	0.59 (0.57, 0.60)	- (-, -)	0.63 (0.62, 0.65)	0.29 (0.29, 0.30)	
	80	0.30 (0.28, 0.31)	- (-, -)	- (-, -)	0.36 (0.35, 0.38)	0.22 (0.21, 0.23)	
White	0	0.01 (0.01, 0.01)	0.01 (0.01, 0.02)	0.02 (0.02, 0.02)	1.05 (1.03, 1.08)	0.36 (0.35, 0.36)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	1.05 (1.03, 1.08)	0.36 (0.35, 0.37)	
	20	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	0.09 (0.09, 0.10)	1.05 (1.03, 1.08)	0.36 (0.35, 0.37)	
	30	0.02 (0.02, 0.02)	0.09 (0.08, 0.09)	0.24 (0.24, 0.25)	1.05 (1.03, 1.08)	0.36 (0.35, 0.37)	
	40	0.07 (0.06, 0.07)	0.22 (0.22, 0.23)	0.49 (0.48, 0.51)	1.04 (1.02, 1.06)	0.36 (0.36, 0.37)	
	50	0.16 (0.15, 0.17)	0.44 (0.42, 0.45)	0.76 (0.74, 0.78)	0.99 (0.97, 1.02)	0.36 (0.35, 0.37)	
	60	0.29 (0.28, 0.30)	0.62 (0.61, 0.64)	0.83 (0.81, 0.85)	0.87 (0.85, 0.89)	0.34 (0.34, 0.35)	
	70	0.38 (0.36, 0.39)	0.60 (0.59, 0.62)	- (-, -)	0.65 (0.63, 0.67)	0.30 (0.29, 0.31)	
	80	0.31 (0.29, 0.32)	- (-, -)	- (-, -)	0.37 (0.35, 0.39)	0.22 (0.22, 0.23)	
Black	0	0.01 (0.01, 0.02)	0.01 (0.01, 0.02)	0.02 (0.01, 0.03)	0.93 (0.88, 1.00)	0.29 (0.28, 0.31)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	0.94 (0.88, 1.01)	0.30 (0.28, 0.32)	
	20	0.01 (0.00, 0.01)	0.03 (0.03, 0.04)	0.10 (0.09, 0.12)	0.95 (0.89, 1.01)	0.30 (0.28, 0.32)	
	30	0.03 (0.02, 0.03)	0.10 (0.09, 0.11)	0.24 (0.22, 0.27)	0.95 (0.89, 1.01)	0.30 (0.28, 0.32)	
	40	0.07 (0.06, 0.08)	0.22 (0.20, 0.25)	0.49 (0.45, 0.52)	0.94 (0.88, 1.00)	0.30 (0.29, 0.32)	
	50	0.16 (0.14, 0.18)	0.43 (0.39, 0.47)	0.72 (0.67, 0.78)	0.90 (0.84, 0.97)	0.30 (0.29, 0.32)	
	60	0.30 (0.27, 0.33)	0.62 (0.57, 0.67)	0.77 (0.71, 0.84)	0.81 (0.75, 0.88)	0.30 (0.28, 0.31)	
	70	0.38 (0.34, 0.43)	0.57 (0.51, 0.63)	- (-, -)	0.61 (0.55, 0.68)	0.27 (0.25, 0.29)	
	80	0.27 (0.23, 0.32)	- (-, -)	- (-, -)	0.34 (0.28, 0.41)	0.22 (0.19, 0.24)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XI-9 - continued

KIDNEY AND RENAL PELVIS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.01 (0.00, 0.01)	0.01 (0.00, 0.01)	0.01 (0.00, 0.02)	0.58 (0.52, 0.67)	0.18 (0.15, 0.24)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.58 (0.52, 0.67)	0.18 (0.15, 0.24)	
Islander	20	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.04 (0.04, 0.06)	0.58 (0.52, 0.67)	0.19 (0.15, 0.25)	
	30	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.11 (0.10, 0.13)	0.58 (0.51, 0.67)	0.19 (0.15, 0.25)	
	40	0.03 (0.02, 0.04)	0.10 (0.09, 0.12)	0.22 (0.19, 0.25)	0.57 (0.51, 0.66)	0.19 (0.15, 0.25)	
	50	0.07 (0.06, 0.08)	0.19 (0.16, 0.22)	0.38 (0.34, 0.42)	0.55 (0.48, 0.64)	0.18 (0.14, 0.24)	
	60	0.12 (0.10, 0.15)	0.32 (0.28, 0.36)	0.44 (0.39, 0.50)	0.49 (0.42, 0.58)	0.17 (0.13, 0.24)	
	70	0.21 (0.17, 0.24)	0.34 (0.29, 0.40)	- (-, -)	0.39 (0.33, 0.48)	0.16 (0.12, 0.22)	
	80	0.16 (0.13, 0.21)	- (-, -)	- (-, -)	0.22 (0.16, 0.32)	0.12 (0.08, 0.20)	
American Indian/ Alaska Native ^a	0	0.03 (0.01, 0.09)	0.03 (0.01, 0.09)	0.03 (0.01, 0.09)	1.28 (0.97, 1.88)	0.52 (0.39, 0.73)	
	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.03 (0.01, 0.07)	1.27 (0.95, 1.88)	0.53 (0.40, 0.74)	
	20	0.00 (0.00, 0.03)	0.03 (0.01, 0.07)	0.12 (0.07, 0.19)	1.27 (0.96, 1.88)	0.53 (0.40, 0.75)	
	30	0.02 (0.01, 0.06)	0.12 (0.07, 0.19)	0.32 (0.23, 0.45)	1.28 (0.96, 1.90)	0.54 (0.40, 0.75)	
	40	0.09 (0.05, 0.16)	0.30 (0.21, 0.43)	0.60 (0.44, 0.80)	1.28 (0.96, 1.91)	0.55 (0.41, 0.77)	
	50	0.22 (0.14, 0.33)	0.52 (0.36, 0.72)	0.94 (0.69, 1.27)	1.22 (0.89, 1.87)	0.53 (0.39, 0.76)	
	60	0.32 (0.19, 0.50)	0.77 (0.53, 1.09)	1.00 (0.70, 1.42)	1.07 (0.74, 1.74)	0.50 (0.35, 0.74)	
	70	0.52 (0.31, 0.82)	0.79 (0.50, 1.22)	- (-, -)	0.86 (0.54, 1.61)	0.46 (0.30, 0.73)	
	80	0.36 (0.14, 0.76)	- (-, -)	- (-, -)	0.45 (0.18, 1.38)	0.36 (0.18, 0.72)	
Hispanic ^b	0	0.01 (0.01, 0.01)	0.01 (0.01, 0.01)	0.01 (0.01, 0.02)	1.12 (1.05, 1.21)	0.35 (0.32, 0.38)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	1.13 (1.05, 1.22)	0.35 (0.32, 0.39)	
	20	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.08 (0.07, 0.09)	1.13 (1.05, 1.22)	0.35 (0.32, 0.39)	
	30	0.02 (0.01, 0.02)	0.08 (0.07, 0.09)	0.25 (0.23, 0.27)	1.13 (1.05, 1.22)	0.35 (0.32, 0.39)	
	40	0.06 (0.06, 0.07)	0.23 (0.21, 0.25)	0.50 (0.47, 0.54)	1.12 (1.04, 1.21)	0.35 (0.32, 0.39)	
	50	0.17 (0.15, 0.19)	0.44 (0.41, 0.48)	0.79 (0.73, 0.84)	1.07 (0.99, 1.16)	0.35 (0.32, 0.38)	
	60	0.29 (0.26, 0.32)	0.64 (0.59, 0.69)	0.89 (0.82, 0.96)	0.93 (0.86, 1.03)	0.33 (0.30, 0.36)	
	70	0.39 (0.34, 0.43)	0.66 (0.59, 0.73)	- (-, -)	0.71 (0.63, 0.80)	0.28 (0.25, 0.32)	
	80	0.34 (0.29, 0.41)	- (-, -)	- (-, -)	0.41 (0.34, 0.50)	0.22 (0.18, 0.26)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XII-9
LARYNX CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.37 (0.36, 0.38)	0.14 (0.13, 0.14)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.38 (0.37, 0.39)	0.14 (0.14, 0.14)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.38 (0.37, 0.39)	0.14 (0.14, 0.14)	
	30	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.09 (0.08, 0.09)	0.38 (0.37, 0.39)	0.14 (0.14, 0.14)	
	40	0.02 (0.02, 0.02)	0.09 (0.08, 0.09)	0.21 (0.20, 0.22)	0.38 (0.37, 0.39)	0.14 (0.14, 0.14)	
	50	0.07 (0.06, 0.07)	0.20 (0.19, 0.20)	0.31 (0.31, 0.32)	0.37 (0.36, 0.38)	0.14 (0.14, 0.14)	
	60	0.14 (0.13, 0.14)	0.26 (0.25, 0.27)	0.32 (0.31, 0.33)	0.33 (0.32, 0.33)	0.13 (0.13, 0.13)	
	70	0.14 (0.14, 0.15)	0.21 (0.20, 0.21)	- (- , -)	0.22 (0.21, 0.23)	0.10 (0.09, 0.10)	
	80	0.09 (0.08, 0.09)	- (- , -)	- (- , -)	0.10 (0.10, 0.11)	0.06 (0.05, 0.06)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.37 (0.36, 0.38)	0.13 (0.13, 0.13)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.38 (0.37, 0.39)	0.13 (0.13, 0.13)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.38 (0.37, 0.39)	0.13 (0.13, 0.13)	
	30	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.08 (0.08, 0.09)	0.38 (0.37, 0.39)	0.13 (0.13, 0.14)	
	40	0.02 (0.02, 0.02)	0.08 (0.08, 0.09)	0.21 (0.20, 0.21)	0.38 (0.37, 0.39)	0.13 (0.13, 0.14)	
	50	0.06 (0.06, 0.07)	0.19 (0.19, 0.20)	0.31 (0.30, 0.32)	0.37 (0.36, 0.38)	0.13 (0.13, 0.14)	
	60	0.14 (0.13, 0.14)	0.26 (0.25, 0.27)	0.32 (0.31, 0.33)	0.33 (0.32, 0.33)	0.12 (0.12, 0.13)	
	70	0.14 (0.14, 0.15)	0.21 (0.20, 0.22)	- (- , -)	0.22 (0.21, 0.23)	0.09 (0.09, 0.10)	
	80	0.09 (0.08, 0.10)	- (- , -)	- (- , -)	0.10 (0.10, 0.11)	0.06 (0.05, 0.06)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.51 (0.48, 0.54)	0.21 (0.20, 0.22)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.52 (0.49, 0.55)	0.22 (0.21, 0.23)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.04 (0.04, 0.05)	0.52 (0.49, 0.56)	0.22 (0.21, 0.23)	
	30	0.00 (0.00, 0.01)	0.04 (0.04, 0.05)	0.16 (0.15, 0.18)	0.53 (0.50, 0.56)	0.22 (0.21, 0.23)	
	40	0.04 (0.03, 0.05)	0.16 (0.15, 0.18)	0.35 (0.32, 0.37)	0.54 (0.51, 0.57)	0.23 (0.22, 0.24)	
	50	0.13 (0.12, 0.14)	0.32 (0.30, 0.35)	0.48 (0.45, 0.51)	0.53 (0.49, 0.56)	0.23 (0.22, 0.24)	
	60	0.22 (0.20, 0.24)	0.39 (0.36, 0.42)	0.44 (0.40, 0.47)	0.44 (0.41, 0.48)	0.21 (0.19, 0.22)	
	70	0.21 (0.19, 0.24)	0.27 (0.24, 0.31)	- (- , -)	0.28 (0.25, 0.32)	0.15 (0.14, 0.16)	
	80	0.10 (0.08, 0.12)	- (- , -)	- (- , -)	0.11 (0.09, 0.14)	0.09 (0.08, 0.10)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

- Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XII-9 - continued
LARYNX CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer						Risk of Dying from Cancer	
		+10 yrs		+20 yrs		+30 yrs		Eventually	Eventually
		Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
Asian/ Pacific Islander	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.22 (0.18, 0.27)	0.07 (0.05, 0.10)		
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.22 (0.19, 0.27)	0.07 (0.05, 0.10)		
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.02, 0.03)	0.22 (0.19, 0.27)	0.07 (0.05, 0.11)		
	30	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.02, 0.03)	0.02 (0.06, 0.09)	0.22 (0.19, 0.27)	0.07 (0.05, 0.11)		
	40	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.07 (0.06, 0.09)	0.22 (0.19, 0.27)	0.22 (0.19, 0.27)	0.07 (0.05, 0.11)		
	50	0.02 (0.01, 0.02)	0.07 (0.06, 0.08)	0.13 (0.12, 0.16)	0.22 (0.19, 0.27)	0.22 (0.19, 0.27)	0.07 (0.05, 0.11)		
	60	0.05 (0.04, 0.06)	0.12 (0.10, 0.14)	0.17 (0.15, 0.20)	0.21 (0.17, 0.26)	0.21 (0.17, 0.26)	0.07 (0.05, 0.11)		
	70	0.07 (0.06, 0.09)	0.13 (0.11, 0.15)	- (-, -)	0.17 (0.13, 0.22)	0.17 (0.13, 0.22)	0.07 (0.04, 0.11)		
	80	0.07 (0.05, 0.09)	- (-, -)	- (-, -)	0.12 (0.08, 0.18)	0.12 (0.08, 0.18)	0.07 (0.04, 0.12)		
American	0	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.14 (0.08, 0.41)	0.14 (0.08, 0.41)	0.09 (0.06, 0.17)		
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.14 (0.08, 0.41)	0.14 (0.08, 0.41)	0.09 (0.06, 0.17)		
	20	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.01 (0.00, 0.03)	0.14 (0.08, 0.42)	0.14 (0.08, 0.42)	0.09 (0.06, 0.17)		
	30	0.00 (0.00, 0.01)	0.01 (0.00, 0.03)	0.04 (0.02, 0.08)	0.14 (0.08, 0.43)	0.14 (0.08, 0.43)	0.09 (0.06, 0.18)		
	40	0.01 (0.00, 0.03)	0.04 (0.02, 0.08)	0.08 (0.05, 0.15)	0.15 (0.08, 0.44)	0.15 (0.08, 0.44)	0.09 (0.06, 0.18)		
	50	0.03 (0.01, 0.07)	0.08 (0.04, 0.14)	0.14 (0.07, 0.24)	0.14 (0.07, 0.44)	0.14 (0.07, 0.44)	0.10 (0.06, 0.19)		
	60	0.05 (0.02, 0.11)	0.11 (0.05, 0.22)	0.11 (0.05, 0.24)	0.11 (0.05, 0.44)	0.11 (0.05, 0.44)	0.09 (0.05, 0.19)		
	70	0.08 (0.03, 0.18)	0.08 (0.03, 0.22)	- (-, -)	0.08 (0.03, 0.46)	0.08 (0.03, 0.46)	0.07 (0.03, 0.18)		
	80	0.00 (0.00, 0.16)	- (-, -)	- (-, -)	0.00 (0.00, 0.56)	0.00 (0.00, 0.56)	0.04 (0.01, 0.21)		
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.30 (0.27, 0.34)	0.30 (0.27, 0.34)	0.12 (0.11, 0.14)		
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.31 (0.28, 0.34)	0.31 (0.28, 0.34)	0.12 (0.11, 0.14)		
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.31 (0.28, 0.35)	0.31 (0.28, 0.35)	0.13 (0.11, 0.14)		
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.05, 0.06)	0.31 (0.28, 0.35)	0.31 (0.28, 0.35)	0.13 (0.11, 0.14)		
	40	0.01 (0.01, 0.01)	0.05 (0.04, 0.06)	0.13 (0.12, 0.14)	0.31 (0.28, 0.35)	0.31 (0.28, 0.35)	0.13 (0.11, 0.14)		
	50	0.04 (0.04, 0.05)	0.12 (0.11, 0.14)	0.23 (0.21, 0.25)	0.31 (0.28, 0.35)	0.31 (0.28, 0.35)	0.13 (0.11, 0.14)		
	60	0.08 (0.07, 0.10)	0.20 (0.17, 0.22)	0.26 (0.23, 0.29)	0.28 (0.25, 0.32)	0.28 (0.25, 0.32)	0.12 (0.11, 0.14)		
	70	0.12 (0.11, 0.14)	0.20 (0.17, 0.23)	- (-, -)	0.22 (0.19, 0.26)	0.22 (0.19, 0.26)	0.10 (0.09, 0.12)		
	80	0.10 (0.07, 0.13)	- (-, -)	- (-, -)	0.12 (0.09, 0.17)	0.12 (0.09, 0.17)	0.07 (0.06, 0.09)		

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XII-9 - continued
LARYNX CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.62 (0.60, 0.64)	0.23 (0.22, 0.23)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.63 (0.61, 0.64)	0.23 (0.22, 0.23)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.04 (0.03, 0.04)	0.63 (0.62, 0.65)	0.23 (0.23, 0.24)	
	30	0.00 (0.00, 0.00)	0.04 (0.03, 0.04)	0.14 (0.14, 0.15)	0.64 (0.62, 0.66)	0.23 (0.23, 0.24)	
	40	0.03 (0.03, 0.03)	0.14 (0.13, 0.15)	0.35 (0.33, 0.36)	0.65 (0.63, 0.66)	0.24 (0.23, 0.24)	
	50	0.11 (0.11, 0.12)	0.33 (0.31, 0.34)	0.53 (0.51, 0.54)	0.64 (0.62, 0.65)	0.24 (0.23, 0.24)	
	60	0.23 (0.22, 0.24)	0.45 (0.44, 0.46)	0.55 (0.53, 0.57)	0.57 (0.55, 0.58)	0.22 (0.22, 0.23)	
	70	0.26 (0.25, 0.27)	0.38 (0.36, 0.40)	- (-, -)	0.40 (0.39, 0.42)	0.18 (0.17, 0.18)	
	80	0.18 (0.17, 0.19)	- (-, -)	- (-, -)	0.22 (0.20, 0.23)	0.12 (0.11, 0.12)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.61 (0.59, 0.63)	0.21 (0.21, 0.22)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.62 (0.60, 0.64)	0.22 (0.21, 0.22)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.03 (0.03, 0.04)	0.62 (0.60, 0.64)	0.22 (0.21, 0.22)	
	30	0.00 (0.00, 0.01)	0.03 (0.03, 0.04)	0.14 (0.13, 0.14)	0.63 (0.61, 0.65)	0.22 (0.21, 0.23)	
	40	0.03 (0.03, 0.03)	0.13 (0.13, 0.14)	0.34 (0.32, 0.35)	0.63 (0.62, 0.65)	0.22 (0.22, 0.23)	
	50	0.11 (0.10, 0.11)	0.32 (0.31, 0.33)	0.52 (0.50, 0.54)	0.63 (0.61, 0.65)	0.22 (0.22, 0.23)	
	60	0.23 (0.22, 0.24)	0.44 (0.43, 0.46)	0.54 (0.52, 0.56)	0.56 (0.54, 0.58)	0.21 (0.20, 0.21)	
	70	0.26 (0.25, 0.27)	0.38 (0.36, 0.39)	- (-, -)	0.40 (0.38, 0.41)	0.17 (0.16, 0.17)	
	80	0.18 (0.17, 0.20)	- (-, -)	- (-, -)	0.21 (0.20, 0.23)	0.11 (0.10, 0.12)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.85 (0.79, 0.91)	0.37 (0.35, 0.39)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.87 (0.81, 0.94)	0.38 (0.36, 0.40)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.06 (0.05, 0.07)	0.88 (0.82, 0.95)	0.39 (0.37, 0.41)	
	30	0.00 (0.00, 0.01)	0.06 (0.05, 0.08)	0.27 (0.24, 0.30)	0.90 (0.84, 0.97)	0.39 (0.37, 0.42)	
	40	0.06 (0.05, 0.07)	0.27 (0.25, 0.30)	0.60 (0.55, 0.64)	0.93 (0.86, 1.00)	0.41 (0.38, 0.43)	
	50	0.23 (0.20, 0.25)	0.57 (0.53, 0.62)	0.83 (0.77, 0.89)	0.92 (0.86, 1.00)	0.41 (0.39, 0.44)	
	60	0.40 (0.36, 0.44)	0.70 (0.64, 0.76)	0.79 (0.72, 0.86)	0.81 (0.74, 0.88)	0.38 (0.36, 0.41)	
	70	0.40 (0.35, 0.46)	0.52 (0.45, 0.59)	- (-, -)	0.54 (0.48, 0.62)	0.30 (0.28, 0.33)	
	80	0.21 (0.16, 0.28)	- (-, -)	- (-, -)	0.26 (0.20, 0.35)	0.19 (0.17, 0.22)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XII-9 - continued
LARYNX CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.41 (0.34, 0.50)	0.13 (0.09, 0.19)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.41 (0.34, 0.50)	0.13 (0.09, 0.19)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.41 (0.34, 0.50)	0.13 (0.09, 0.20)	
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.05 (0.04, 0.06)	0.42 (0.35, 0.51)	0.13 (0.09, 0.20)	
	40	0.01 (0.01, 0.02)	0.05 (0.03, 0.06)	0.14 (0.11, 0.16)	0.42 (0.35, 0.51)	0.13 (0.09, 0.20)	
	50	0.04 (0.03, 0.05)	0.13 (0.11, 0.16)	0.26 (0.22, 0.30)	0.41 (0.34, 0.51)	0.13 (0.09, 0.20)	
	60	0.10 (0.08, 0.12)	0.23 (0.19, 0.27)	0.33 (0.28, 0.38)	0.40 (0.32, 0.49)	0.13 (0.09, 0.20)	
	70	0.15 (0.12, 0.18)	0.25 (0.21, 0.31)	- (-, -)	0.33 (0.26, 0.44)	0.13 (0.08, 0.21)	
	80	0.14 (0.10, 0.19)	- (-, -)	- (-, -)	0.24 (0.16, 0.37)	0.13 (0.08, 0.23)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.26 (0.14, 0.91)	0.14 (0.08, 0.33)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.27 (0.14, 0.93)	0.14 (0.09, 0.34)	
	20	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.02 (0.00, 0.06)	0.27 (0.15, 0.94)	0.15 (0.09, 0.34)	
	30	0.00 (0.00, 0.03)	0.02 (0.00, 0.06)	0.07 (0.03, 0.14)	0.28 (0.15, 0.96)	0.15 (0.09, 0.35)	
	40	0.02 (0.00, 0.05)	0.07 (0.03, 0.14)	0.16 (0.08, 0.28)	0.28 (0.15, 1.00)	0.16 (0.09, 0.36)	
	50	0.06 (0.02, 0.13)	0.15 (0.07, 0.28)	0.28 (0.14, 0.50)	0.28 (0.15, 1.03)	0.16 (0.09, 0.38)	
	60	0.10 (0.04, 0.22)	0.24 (0.11, 0.47)	0.25 (0.11, 0.54)	0.25 (0.11, 1.07)	0.15 (0.08, 0.39)	
	70	0.17 (0.06, 0.40)	0.18 (0.06, 0.49)	- (-, -)	0.18 (0.06, 1.18)	0.12 (0.05, 0.42)	
	80	0.01 (0.00, 0.41)	- (-, -)	- (-, -)	0.01 (0.00, 1.57)	0.09 (0.02, 0.57)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.56 (0.50, 0.64)	0.24 (0.21, 0.27)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.57 (0.50, 0.65)	0.24 (0.21, 0.28)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.02, 0.02)	0.57 (0.51, 0.66)	0.24 (0.21, 0.28)	
	30	0.00 (0.00, 0.01)	0.02 (0.02, 0.02)	0.09 (0.08, 0.10)	0.58 (0.51, 0.66)	0.24 (0.22, 0.28)	
	40	0.02 (0.01, 0.02)	0.09 (0.08, 0.10)	0.24 (0.21, 0.27)	0.58 (0.52, 0.67)	0.25 (0.22, 0.29)	
	50	0.07 (0.06, 0.09)	0.23 (0.20, 0.26)	0.43 (0.39, 0.48)	0.58 (0.52, 0.67)	0.25 (0.22, 0.29)	
	60	0.16 (0.14, 0.19)	0.38 (0.34, 0.43)	0.51 (0.45, 0.58)	0.54 (0.47, 0.64)	0.24 (0.21, 0.28)	
	70	0.25 (0.21, 0.29)	0.40 (0.34, 0.46)	- (-, -)	0.43 (0.36, 0.54)	0.21 (0.17, 0.25)	
	80	0.21 (0.15, 0.28)	- (-, -)	- (-, -)	0.26 (0.18, 0.39)	0.16 (0.12, 0.22)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XII-9 - continued
LARYNX CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.14 (0.14, 0.15)	0.06 (0.05, 0.06)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.15 (0.14, 0.15)	0.06 (0.05, 0.06)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.15 (0.14, 0.15)	0.06 (0.05, 0.06)	
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	0.15 (0.14, 0.15)	0.06 (0.05, 0.06)	
	40	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	0.08 (0.08, 0.09)	0.15 (0.14, 0.15)	0.06 (0.05, 0.06)	
	50	0.02 (0.02, 0.03)	0.07 (0.07, 0.08)	0.12 (0.11, 0.13)	0.14 (0.13, 0.15)	0.06 (0.05, 0.06)	
	60	0.05 (0.05, 0.06)	0.10 (0.09, 0.11)	0.12 (0.11, 0.13)	0.12 (0.11, 0.13)	0.05 (0.05, 0.05)	
	70	0.05 (0.05, 0.06)	0.07 (0.07, 0.08)	- (-, -)	0.08 (0.07, 0.08)	0.04 (0.04, 0.04)	
	80	0.03 (0.03, 0.03)	- (-, -)	- (-, -)	0.03 (0.03, 0.04)	0.02 (0.02, 0.02)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.15 (0.14, 0.16)	0.05 (0.05, 0.06)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.15 (0.14, 0.16)	0.05 (0.05, 0.06)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.15 (0.14, 0.16)	0.06 (0.05, 0.06)	
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	0.15 (0.14, 0.16)	0.06 (0.05, 0.06)	
	40	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.08 (0.08, 0.09)	0.15 (0.14, 0.16)	0.06 (0.05, 0.06)	
	50	0.02 (0.02, 0.03)	0.08 (0.07, 0.08)	0.12 (0.11, 0.13)	0.14 (0.13, 0.15)	0.06 (0.05, 0.06)	
	60	0.06 (0.05, 0.06)	0.10 (0.09, 0.11)	0.12 (0.11, 0.13)	0.13 (0.12, 0.13)	0.05 (0.05, 0.05)	
	70	0.05 (0.05, 0.06)	0.08 (0.07, 0.08)	- (-, -)	0.08 (0.07, 0.09)	0.04 (0.04, 0.04)	
	80	0.03 (0.03, 0.04)	- (-, -)	- (-, -)	0.04 (0.03, 0.04)	0.02 (0.02, 0.02)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.20 (0.17, 0.23)	0.07 (0.07, 0.08)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.20 (0.17, 0.23)	0.08 (0.07, 0.08)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.20 (0.17, 0.23)	0.08 (0.07, 0.08)	
	30	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.07 (0.06, 0.08)	0.20 (0.18, 0.23)	0.08 (0.07, 0.09)	
	40	0.02 (0.02, 0.03)	0.07 (0.05, 0.08)	0.13 (0.11, 0.15)	0.20 (0.18, 0.23)	0.08 (0.07, 0.09)	
	50	0.05 (0.04, 0.06)	0.11 (0.09, 0.13)	0.17 (0.15, 0.20)	0.19 (0.16, 0.22)	0.08 (0.07, 0.08)	
	60	0.07 (0.06, 0.09)	0.14 (0.11, 0.16)	0.15 (0.13, 0.18)	0.15 (0.13, 0.19)	0.07 (0.06, 0.08)	
	70	0.08 (0.06, 0.10)	0.10 (0.08, 0.13)	- (-, -)	0.10 (0.08, 0.13)	0.05 (0.04, 0.06)	
	80	0.03 (0.02, 0.05)	- (-, -)	- (-, -)	0.03 (0.02, 0.05)	0.03 (0.02, 0.04)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XII-9 - continued
LARYNX CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.03, 0.10)	0.02 (0.01, 0.06)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.03, 0.10)	0.02 (0.01, 0.06)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.03, 0.10)	0.02 (0.01, 0.06)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.05 (0.03, 0.10)	0.02 (0.01, 0.06)	
	40	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.03)	0.05 (0.03, 0.10)	0.02 (0.01, 0.06)	
	50	0.00 (0.00, 0.01)	0.02 (0.01, 0.03)	0.03 (0.02, 0.05)	0.05 (0.03, 0.10)	0.02 (0.01, 0.06)	
	60	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.04 (0.03, 0.07)	0.05 (0.03, 0.10)	0.02 (0.01, 0.06)	
	70	0.02 (0.01, 0.03)	0.03 (0.02, 0.05)	- (-, -)	0.04 (0.02, 0.09)	0.02 (0.01, 0.06)	
	80	0.02 (0.01, 0.04)	- (-, -)	- (-, -)	0.02 (0.01, 0.09)	0.02 (0.00, 0.07)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.02 (0.00, 0.52)	0.04 (0.01, 0.17)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.02 (0.00, 0.52)	0.04 (0.01, 0.18)	
	20	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.02 (0.00, 0.52)	0.04 (0.01, 0.18)	
	30	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.02 (0.00, 0.07)	0.02 (0.00, 0.53)	0.04 (0.01, 0.18)	
	40	0.01 (0.00, 0.04)	0.02 (0.00, 0.07)	0.02 (0.00, 0.09)	0.02 (0.00, 0.54)	0.04 (0.01, 0.18)	
	50	0.01 (0.00, 0.06)	0.01 (0.00, 0.08)	0.01 (0.00, 0.11)	0.01 (0.00, 0.55)	0.04 (0.01, 0.19)	
	60	0.00 (0.00, 0.07)	0.00 (0.00, 0.10)	0.00 (0.00, 0.17)	0.00 (0.00, 0.57)	0.04 (0.01, 0.19)	
	70	0.00 (0.00, 0.12)	0.00 (0.00, 0.19)	- (-, -)	0.00 (0.00, 0.66)	0.03 (0.00, 0.21)	
	80	0.00 (0.00, 0.25)	- (-, -)	- (-, -)	0.00 (0.00, 0.88)	0.00 (0.00, 0.27)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.08 (0.06, 0.12)	0.03 (0.02, 0.04)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.08 (0.06, 0.12)	0.03 (0.02, 0.04)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.08 (0.06, 0.12)	0.03 (0.02, 0.04)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.08 (0.06, 0.12)	0.03 (0.02, 0.04)	
	40	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.08 (0.06, 0.12)	0.03 (0.02, 0.04)	
	50	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.05 (0.04, 0.06)	0.08 (0.06, 0.12)	0.03 (0.02, 0.04)	
	60	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.06 (0.04, 0.08)	0.07 (0.05, 0.11)	0.03 (0.02, 0.04)	
	70	0.03 (0.02, 0.04)	0.05 (0.03, 0.07)	- (-, -)	0.06 (0.04, 0.10)	0.03 (0.02, 0.04)	
	80	0.03 (0.01, 0.05)	- (-, -)	- (-, -)	0.04 (0.02, 0.09)	0.02 (0.01, 0.04)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XIII-17

LEUKEMIA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.06 (0.05, 0.06)	0.09 (0.08, 0.09)	0.11 (0.11, 0.11)	1.26 (1.25, 1.28)	0.83 (0.83, 0.84)	
	10	0.03 (0.03, 0.03)	0.05 (0.05, 0.06)	0.09 (0.08, 0.09)	1.22 (1.21, 1.24)	0.84 (0.83, 0.84)	
	20	0.02 (0.02, 0.02)	0.06 (0.05, 0.06)	0.11 (0.11, 0.12)	1.20 (1.18, 1.21)	0.83 (0.82, 0.84)	
	30	0.03 (0.03, 0.03)	0.09 (0.09, 0.09)	0.21 (0.20, 0.21)	1.19 (1.17, 1.20)	0.83 (0.82, 0.84)	
	40	0.06 (0.06, 0.06)	0.18 (0.17, 0.18)	0.42 (0.42, 0.43)	1.17 (1.15, 1.18)	0.83 (0.82, 0.83)	
	50	0.12 (0.12, 0.13)	0.38 (0.37, 0.39)	0.75 (0.74, 0.77)	1.14 (1.13, 1.16)	0.83 (0.82, 0.83)	
	60	0.27 (0.26, 0.28)	0.67 (0.66, 0.68)	0.98 (0.97, 1.00)	1.08 (1.07, 1.10)	0.82 (0.81, 0.83)	
	70	0.46 (0.45, 0.47)	0.82 (0.81, 0.84)	- (- , -)	0.94 (0.92, 0.96)	0.77 (0.77, 0.78)	
	80	0.52 (0.50, 0.53)	- (- , -)	- (- , -)	0.68 (0.66, 0.70)	0.63 (0.63, 0.64)	
White	0	0.06 (0.06, 0.06)	0.09 (0.09, 0.09)	0.11 (0.11, 0.12)	1.31 (1.30, 1.33)	0.87 (0.86, 0.88)	
	10	0.03 (0.03, 0.03)	0.06 (0.05, 0.06)	0.09 (0.08, 0.09)	1.27 (1.25, 1.29)	0.87 (0.86, 0.88)	
	20	0.02 (0.02, 0.03)	0.06 (0.05, 0.06)	0.11 (0.11, 0.12)	1.24 (1.23, 1.26)	0.87 (0.86, 0.87)	
	30	0.03 (0.03, 0.03)	0.09 (0.09, 0.09)	0.21 (0.21, 0.22)	1.23 (1.21, 1.25)	0.86 (0.86, 0.87)	
	40	0.06 (0.06, 0.06)	0.18 (0.18, 0.19)	0.44 (0.43, 0.45)	1.21 (1.19, 1.23)	0.86 (0.85, 0.87)	
	50	0.13 (0.12, 0.13)	0.39 (0.38, 0.40)	0.78 (0.77, 0.80)	1.18 (1.17, 1.20)	0.86 (0.85, 0.87)	
	60	0.28 (0.27, 0.29)	0.70 (0.68, 0.71)	1.02 (1.00, 1.04)	1.12 (1.10, 1.14)	0.85 (0.84, 0.86)	
	70	0.48 (0.47, 0.49)	0.85 (0.84, 0.87)	- (- , -)	0.97 (0.95, 0.99)	0.80 (0.79, 0.81)	
	80	0.53 (0.52, 0.55)	- (- , -)	- (- , -)	0.70 (0.68, 0.72)	0.65 (0.64, 0.66)	
Black	0	0.03 (0.03, 0.04)	0.05 (0.05, 0.06)	0.07 (0.06, 0.08)	0.86 (0.82, 0.91)	0.60 (0.58, 0.62)	
	10	0.02 (0.02, 0.02)	0.04 (0.04, 0.05)	0.07 (0.07, 0.08)	0.85 (0.81, 0.90)	0.61 (0.59, 0.63)	
	20	0.02 (0.02, 0.03)	0.05 (0.05, 0.06)	0.10 (0.10, 0.11)	0.83 (0.79, 0.88)	0.61 (0.59, 0.62)	
	30	0.03 (0.03, 0.04)	0.09 (0.08, 0.09)	0.19 (0.18, 0.20)	0.83 (0.78, 0.87)	0.61 (0.59, 0.62)	
	40	0.05 (0.05, 0.06)	0.16 (0.15, 0.18)	0.35 (0.33, 0.38)	0.81 (0.77, 0.86)	0.60 (0.59, 0.62)	
	50	0.11 (0.10, 0.12)	0.32 (0.29, 0.34)	0.56 (0.53, 0.60)	0.80 (0.76, 0.85)	0.61 (0.59, 0.63)	
	60	0.23 (0.21, 0.25)	0.50 (0.47, 0.54)	0.70 (0.66, 0.74)	0.77 (0.72, 0.82)	0.61 (0.59, 0.63)	
	70	0.35 (0.32, 0.38)	0.59 (0.55, 0.64)	- (- , -)	0.68 (0.63, 0.74)	0.58 (0.56, 0.60)	
	80	0.39 (0.35, 0.44)	- (- , -)	- (- , -)	0.53 (0.47, 0.60)	0.48 (0.46, 0.51)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table X1II-17 - continued

LEUKEMIA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.06 (0.05, 0.07)	0.08 (0.07, 0.09)	0.10 (0.09, 0.11)	0.88 (0.82, 0.95)	0.56 (0.51, 0.62)	
	10	0.02 (0.02, 0.03)	0.04 (0.04, 0.05)	0.07 (0.06, 0.08)	0.83 (0.77, 0.90)	0.56 (0.51, 0.61)	
Islander	20	0.02 (0.02, 0.02)	0.05 (0.04, 0.06)	0.10 (0.09, 0.11)	0.81 (0.75, 0.88)	0.55 (0.50, 0.61)	
	30	0.03 (0.03, 0.04)	0.08 (0.07, 0.09)	0.14 (0.13, 0.16)	0.79 (0.73, 0.86)	0.54 (0.50, 0.60)	
	40	0.05 (0.04, 0.05)	0.11 (0.10, 0.13)	0.25 (0.23, 0.28)	0.77 (0.70, 0.84)	0.54 (0.49, 0.59)	
	50	0.07 (0.06, 0.08)	0.21 (0.19, 0.23)	0.45 (0.41, 0.48)	0.73 (0.67, 0.80)	0.53 (0.48, 0.58)	
	60	0.15 (0.13, 0.17)	0.39 (0.36, 0.42)	0.60 (0.55, 0.65)	0.68 (0.62, 0.76)	0.51 (0.46, 0.57)	
	70	0.26 (0.24, 0.29)	0.49 (0.44, 0.54)	- (- , -)	0.58 (0.52, 0.66)	0.46 (0.41, 0.53)	
	80	0.28 (0.24, 0.33)	- (- , -)	- (- , -)	0.40 (0.33, 0.48)	0.36 (0.30, 0.43)	
American Indian/ Alaska Native ^a	0	0.04 (0.02, 0.07)	0.08 (0.05, 0.11)	0.10 (0.07, 0.14)	0.69 (0.48, 1.07)	0.49 (0.40, 0.62)	
	10	0.03 (0.02, 0.05)	0.06 (0.03, 0.09)	0.08 (0.05, 0.12)	0.66 (0.45, 1.05)	0.50 (0.40, 0.63)	
	20	0.02 (0.01, 0.04)	0.05 (0.03, 0.08)	0.07 (0.05, 0.11)	0.63 (0.42, 1.02)	0.49 (0.40, 0.62)	
	30	0.02 (0.01, 0.05)	0.05 (0.03, 0.09)	0.12 (0.08, 0.17)	0.62 (0.40, 1.02)	0.48 (0.39, 0.62)	
	40	0.03 (0.01, 0.06)	0.10 (0.06, 0.15)	0.21 (0.14, 0.30)	0.61 (0.39, 1.02)	0.48 (0.39, 0.62)	
	50	0.07 (0.04, 0.12)	0.18 (0.12, 0.28)	0.27 (0.18, 0.40)	0.61 (0.38, 1.03)	0.49 (0.39, 0.64)	
	60	0.12 (0.07, 0.21)	0.21 (0.13, 0.34)	0.41 (0.26, 0.62)	0.58 (0.34, 1.03)	0.49 (0.38, 0.64)	
	70	0.11 (0.05, 0.21)	0.34 (0.19, 0.57)	- (- , -)	0.53 (0.27, 1.06)	0.50 (0.38, 0.68)	
	80	0.32 (0.15, 0.62)	- (- , -)	- (- , -)	0.60 (0.26, 1.34)	0.51 (0.36, 0.75)	
Hispanic ^b	0	0.07 (0.06, 0.07)	0.11 (0.10, 0.11)	0.13 (0.13, 0.14)	1.12 (1.05, 1.19)	0.63 (0.60, 0.66)	
	10	0.04 (0.04, 0.04)	0.07 (0.06, 0.07)	0.10 (0.09, 0.11)	1.06 (1.00, 1.14)	0.62 (0.59, 0.66)	
	20	0.03 (0.03, 0.03)	0.06 (0.06, 0.07)	0.11 (0.10, 0.12)	1.03 (0.96, 1.10)	0.61 (0.58, 0.65)	
	30	0.03 (0.03, 0.04)	0.08 (0.08, 0.09)	0.17 (0.16, 0.18)	1.01 (0.94, 1.08)	0.61 (0.58, 0.64)	
	40	0.05 (0.04, 0.05)	0.14 (0.13, 0.15)	0.33 (0.31, 0.35)	0.98 (0.92, 1.06)	0.60 (0.57, 0.63)	
	50	0.09 (0.08, 0.10)	0.29 (0.27, 0.31)	0.55 (0.52, 0.58)	0.96 (0.89, 1.03)	0.59 (0.56, 0.62)	
	60	0.21 (0.19, 0.23)	0.48 (0.45, 0.52)	0.74 (0.69, 0.79)	0.91 (0.84, 0.99)	0.57 (0.54, 0.61)	
	70	0.31 (0.28, 0.34)	0.59 (0.55, 0.65)	- (- , -)	0.78 (0.71, 0.87)	0.52 (0.49, 0.56)	
	80	0.38 (0.33, 0.43)	- (- , -)	- (- , -)	0.63 (0.54, 0.74)	0.44 (0.40, 0.49)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table X1II-17 - continued

LEUKEMIA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.06 (0.06, 0.07)	0.10 (0.09, 0.10)	0.12 (0.12, 0.13)	1.50 (1.47, 1.52)	0.98 (0.97, 0.99)	
	10	0.04 (0.03, 0.04)	0.06 (0.06, 0.07)	0.10 (0.10, 0.10)	1.46 (1.43, 1.48)	0.98 (0.97, 1.00)	
	20	0.03 (0.03, 0.03)	0.06 (0.06, 0.07)	0.13 (0.12, 0.14)	1.43 (1.40, 1.46)	0.98 (0.97, 0.99)	
	30	0.04 (0.03, 0.04)	0.10 (0.10, 0.11)	0.25 (0.24, 0.25)	1.42 (1.39, 1.45)	0.98 (0.97, 0.99)	
	40	0.07 (0.06, 0.07)	0.21 (0.21, 0.22)	0.52 (0.51, 0.54)	1.41 (1.38, 1.43)	0.98 (0.97, 0.99)	
	50	0.15 (0.15, 0.16)	0.47 (0.46, 0.49)	0.95 (0.93, 0.97)	1.39 (1.36, 1.42)	0.99 (0.98, 1.00)	
	60	0.35 (0.34, 0.36)	0.86 (0.84, 0.89)	1.23 (1.21, 1.26)	1.34 (1.31, 1.36)	1.00 (0.99, 1.01)	
	70	0.62 (0.60, 0.64)	1.06 (1.03, 1.08)	- (- , -)	1.18 (1.15, 1.21)	0.96 (0.95, 0.98)	
	80	0.68 (0.65, 0.70)	- (- , -)	- (- , -)	0.87 (0.83, 0.90)	0.80 (0.79, 0.82)	
White	0	0.06 (0.06, 0.07)	0.10 (0.10, 0.11)	0.13 (0.13, 0.14)	1.56 (1.54, 1.59)	1.03 (1.01, 1.04)	
	10	0.04 (0.04, 0.04)	0.07 (0.06, 0.07)	0.10 (0.10, 0.11)	1.52 (1.49, 1.55)	1.03 (1.02, 1.04)	
	20	0.03 (0.03, 0.03)	0.07 (0.06, 0.07)	0.13 (0.13, 0.14)	1.49 (1.46, 1.52)	1.03 (1.01, 1.04)	
	30	0.04 (0.03, 0.04)	0.10 (0.10, 0.11)	0.25 (0.24, 0.26)	1.48 (1.45, 1.51)	1.03 (1.01, 1.04)	
	40	0.07 (0.06, 0.07)	0.22 (0.21, 0.22)	0.54 (0.52, 0.55)	1.46 (1.43, 1.49)	1.03 (1.02, 1.04)	
	50	0.15 (0.15, 0.16)	0.49 (0.47, 0.50)	0.99 (0.97, 1.01)	1.45 (1.42, 1.48)	1.04 (1.02, 1.05)	
	60	0.36 (0.35, 0.37)	0.90 (0.88, 0.92)	1.28 (1.26, 1.31)	1.39 (1.36, 1.42)	1.04 (1.03, 1.06)	
	70	0.64 (0.62, 0.66)	1.10 (1.07, 1.13)	- (- , -)	1.23 (1.19, 1.26)	1.00 (0.98, 1.01)	
	80	0.71 (0.68, 0.74)	- (- , -)	- (- , -)	0.90 (0.87, 0.94)	0.83 (0.81, 0.85)	
Black	0	0.04 (0.03, 0.05)	0.06 (0.05, 0.07)	0.08 (0.07, 0.10)	0.98 (0.92, 1.05)	0.65 (0.62, 0.68)	
	10	0.02 (0.02, 0.03)	0.05 (0.04, 0.06)	0.09 (0.08, 0.10)	0.97 (0.90, 1.04)	0.66 (0.64, 0.69)	
	20	0.03 (0.02, 0.04)	0.07 (0.06, 0.08)	0.13 (0.11, 0.14)	0.96 (0.89, 1.03)	0.66 (0.63, 0.69)	
	30	0.04 (0.03, 0.05)	0.10 (0.09, 0.12)	0.24 (0.22, 0.26)	0.95 (0.89, 1.03)	0.66 (0.63, 0.69)	
	40	0.07 (0.06, 0.08)	0.21 (0.18, 0.23)	0.45 (0.41, 0.49)	0.94 (0.87, 1.02)	0.66 (0.64, 0.69)	
	50	0.15 (0.13, 0.17)	0.41 (0.37, 0.45)	0.71 (0.65, 0.77)	0.94 (0.86, 1.02)	0.68 (0.65, 0.71)	
	60	0.30 (0.26, 0.34)	0.65 (0.59, 0.71)	0.84 (0.77, 0.92)	0.91 (0.83, 1.00)	0.69 (0.66, 0.72)	
	70	0.46 (0.41, 0.52)	0.72 (0.64, 0.80)	- (- , -)	0.81 (0.72, 0.92)	0.67 (0.63, 0.71)	
	80	0.46 (0.39, 0.56)	- (- , -)	- (- , -)	0.63 (0.52, 0.77)	0.56 (0.51, 0.62)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table X1II-17 - continued

LEUKEMIA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.06 (0.05, 0.07)	0.08 (0.07, 0.10)	0.11 (0.09, 0.12)	1.03 (0.94, 1.14)	0.66 (0.59, 0.75)	
	10	0.03 (0.02, 0.04)	0.05 (0.04, 0.06)	0.09 (0.07, 0.10)	0.98 (0.89, 1.09)	0.66 (0.59, 0.75)	
Islander	20	0.02 (0.02, 0.03)	0.06 (0.05, 0.07)	0.11 (0.10, 0.13)	0.96 (0.87, 1.07)	0.65 (0.58, 0.74)	
	30	0.03 (0.03, 0.04)	0.09 (0.07, 0.10)	0.16 (0.15, 0.19)	0.94 (0.85, 1.05)	0.65 (0.58, 0.74)	
	40	0.05 (0.04, 0.06)	0.13 (0.11, 0.15)	0.31 (0.28, 0.35)	0.91 (0.82, 1.02)	0.64 (0.57, 0.73)	
	50	0.08 (0.07, 0.10)	0.27 (0.23, 0.30)	0.57 (0.51, 0.63)	0.88 (0.79, 0.99)	0.63 (0.56, 0.72)	
	60	0.19 (0.16, 0.23)	0.51 (0.46, 0.57)	0.75 (0.68, 0.84)	0.83 (0.74, 0.95)	0.62 (0.55, 0.71)	
	70	0.35 (0.30, 0.41)	0.62 (0.55, 0.71)	- (- , -)	0.71 (0.61, 0.83)	0.56 (0.49, 0.67)	
	80	0.36 (0.29, 0.44)	- (- , -)	- (- , -)	0.47 (0.37, 0.61)	0.42 (0.33, 0.54)	
American	0	0.06 (0.03, 0.11)	0.10 (0.06, 0.17)	0.11 (0.07, 0.19)	0.49 (0.33, 1.14)	0.49 (0.37, 0.72)	
Indian/ Alaska Native ^a	10	0.04 (0.02, 0.08)	0.06 (0.03, 0.11)	0.07 (0.04, 0.12)	0.44 (0.28, 1.11)	0.49 (0.37, 0.73)	
	20	0.01 (0.00, 0.05)	0.02 (0.01, 0.07)	0.05 (0.02, 0.10)	0.40 (0.24, 1.08)	0.49 (0.36, 0.72)	
	30	0.01 (0.00, 0.04)	0.03 (0.01, 0.08)	0.11 (0.05, 0.19)	0.40 (0.24, 1.09)	0.48 (0.35, 0.72)	
	40	0.03 (0.01, 0.07)	0.10 (0.05, 0.18)	0.24 (0.14, 0.40)	0.40 (0.24, 1.12)	0.48 (0.35, 0.73)	
	50	0.08 (0.03, 0.16)	0.23 (0.12, 0.39)	0.32 (0.19, 0.54)	0.40 (0.23, 1.15)	0.50 (0.36, 0.76)	
	60	0.16 (0.07, 0.32)	0.27 (0.13, 0.49)	0.35 (0.18, 0.67)	0.35 (0.18, 1.18)	0.50 (0.35, 0.79)	
	70	0.13 (0.04, 0.32)	0.22 (0.08, 0.57)	- (- , -)	0.22 (0.08, 1.23)	0.51 (0.34, 0.88)	
	80	0.14 (0.02, 0.61)	- (- , -)	- (- , -)	0.14 (0.02, 1.68)	0.54 (0.31, 1.07)	
Hispanic ^b	0	0.07 (0.06, 0.08)	0.12 (0.11, 0.13)	0.15 (0.14, 0.16)	1.29 (1.18, 1.42)	0.70 (0.66, 0.76)	
	10	0.05 (0.04, 0.05)	0.08 (0.07, 0.09)	0.12 (0.11, 0.13)	1.23 (1.12, 1.37)	0.70 (0.65, 0.76)	
	20	0.04 (0.03, 0.04)	0.07 (0.06, 0.08)	0.12 (0.11, 0.13)	1.19 (1.08, 1.33)	0.69 (0.64, 0.75)	
	30	0.04 (0.03, 0.04)	0.09 (0.08, 0.10)	0.18 (0.16, 0.19)	1.17 (1.06, 1.31)	0.68 (0.63, 0.74)	
	40	0.05 (0.05, 0.06)	0.14 (0.13, 0.16)	0.35 (0.32, 0.39)	1.15 (1.04, 1.29)	0.68 (0.63, 0.73)	
	50	0.09 (0.08, 0.11)	0.31 (0.28, 0.34)	0.64 (0.59, 0.70)	1.13 (1.02, 1.28)	0.67 (0.62, 0.73)	
	60	0.23 (0.20, 0.26)	0.58 (0.53, 0.64)	0.91 (0.82, 0.99)	1.11 (0.99, 1.26)	0.66 (0.61, 0.72)	
	70	0.41 (0.36, 0.46)	0.78 (0.69, 0.87)	- (- , -)	1.01 (0.88, 1.18)	0.62 (0.56, 0.69)	
	80	0.53 (0.44, 0.63)	- (- , -)	- (- , -)	0.86 (0.68, 1.08)	0.53 (0.46, 0.62)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table X1II-17 - continued

LEUKEMIA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.05 (0.05, 0.05)	0.07 (0.07, 0.08)	0.09 (0.09, 0.10)	1.06 (1.04, 1.08)	0.71 (0.70, 0.72)	
	10	0.02 (0.02, 0.03)	0.04 (0.04, 0.04)	0.07 (0.07, 0.07)	1.02 (1.00, 1.04)	0.71 (0.70, 0.72)	
	20	0.02 (0.02, 0.02)	0.05 (0.04, 0.05)	0.09 (0.09, 0.10)	1.00 (0.98, 1.02)	0.71 (0.70, 0.71)	
	30	0.03 (0.03, 0.03)	0.08 (0.07, 0.08)	0.17 (0.16, 0.17)	0.98 (0.96, 1.00)	0.70 (0.69, 0.71)	
	40	0.05 (0.05, 0.05)	0.14 (0.14, 0.15)	0.33 (0.32, 0.34)	0.96 (0.94, 0.98)	0.70 (0.69, 0.71)	
	50	0.10 (0.09, 0.10)	0.29 (0.28, 0.30)	0.58 (0.56, 0.59)	0.94 (0.92, 0.96)	0.69 (0.68, 0.70)	
	60	0.20 (0.19, 0.21)	0.50 (0.49, 0.52)	0.78 (0.76, 0.80)	0.88 (0.86, 0.90)	0.68 (0.67, 0.69)	
	70	0.34 (0.33, 0.35)	0.65 (0.63, 0.66)	- (-, -)	0.76 (0.74, 0.78)	0.64 (0.63, 0.65)	
	80	0.41 (0.40, 0.43)	- (-, -)	- (-, -)	0.57 (0.55, 0.59)	0.53 (0.52, 0.54)	
White	0	0.05 (0.05, 0.06)	0.08 (0.07, 0.08)	0.10 (0.09, 0.10)	1.09 (1.07, 1.12)	0.74 (0.73, 0.74)	
	10	0.02 (0.02, 0.03)	0.04 (0.04, 0.05)	0.07 (0.07, 0.08)	1.05 (1.03, 1.07)	0.74 (0.73, 0.75)	
	20	0.02 (0.02, 0.02)	0.05 (0.04, 0.05)	0.10 (0.09, 0.10)	1.03 (1.01, 1.05)	0.73 (0.72, 0.74)	
	30	0.03 (0.03, 0.03)	0.08 (0.07, 0.08)	0.17 (0.17, 0.18)	1.01 (0.99, 1.04)	0.72 (0.72, 0.73)	
	40	0.05 (0.05, 0.05)	0.15 (0.14, 0.15)	0.34 (0.33, 0.35)	0.99 (0.97, 1.02)	0.72 (0.71, 0.73)	
	50	0.10 (0.09, 0.10)	0.30 (0.29, 0.31)	0.60 (0.58, 0.62)	0.96 (0.94, 0.99)	0.71 (0.70, 0.72)	
	60	0.21 (0.20, 0.22)	0.52 (0.51, 0.54)	0.80 (0.78, 0.82)	0.90 (0.88, 0.93)	0.70 (0.69, 0.71)	
	70	0.35 (0.34, 0.36)	0.66 (0.64, 0.68)	- (-, -)	0.78 (0.76, 0.80)	0.65 (0.64, 0.66)	
	80	0.42 (0.41, 0.44)	- (-, -)	- (-, -)	0.58 (0.55, 0.60)	0.55 (0.54, 0.56)	
Black	0	0.03 (0.02, 0.03)	0.05 (0.04, 0.06)	0.06 (0.05, 0.07)	0.76 (0.70, 0.82)	0.56 (0.54, 0.58)	
	10	0.02 (0.01, 0.03)	0.03 (0.03, 0.04)	0.06 (0.05, 0.07)	0.75 (0.69, 0.81)	0.57 (0.55, 0.59)	
	20	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.08 (0.07, 0.10)	0.73 (0.67, 0.79)	0.56 (0.54, 0.59)	
	30	0.03 (0.02, 0.03)	0.07 (0.06, 0.08)	0.15 (0.13, 0.16)	0.72 (0.67, 0.78)	0.56 (0.54, 0.58)	
	40	0.04 (0.04, 0.05)	0.12 (0.11, 0.14)	0.27 (0.24, 0.30)	0.71 (0.65, 0.77)	0.56 (0.53, 0.58)	
	50	0.08 (0.07, 0.09)	0.23 (0.21, 0.26)	0.44 (0.40, 0.48)	0.69 (0.63, 0.75)	0.56 (0.53, 0.58)	
	60	0.17 (0.14, 0.19)	0.39 (0.35, 0.43)	0.59 (0.53, 0.64)	0.67 (0.61, 0.73)	0.55 (0.52, 0.58)	
	70	0.27 (0.23, 0.30)	0.50 (0.45, 0.56)	- (-, -)	0.59 (0.53, 0.66)	0.52 (0.49, 0.55)	
	80	0.35 (0.29, 0.41)	- (-, -)	- (-, -)	0.49 (0.41, 0.57)	0.44 (0.41, 0.48)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

- Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table X1II-17 - continued

LEUKEMIA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.06 (0.05, 0.08)	0.08 (0.07, 0.10)	0.10 (0.08, 0.11)	0.75 (0.67, 0.85)	0.47 (0.41, 0.56)	
Islander	10	0.02 (0.01, 0.03)	0.03 (0.03, 0.04)	0.06 (0.05, 0.07)	0.69 (0.61, 0.80)	0.47 (0.40, 0.55)	
	20	0.02 (0.01, 0.02)	0.04 (0.03, 0.05)	0.08 (0.07, 0.10)	0.68 (0.60, 0.78)	0.46 (0.40, 0.55)	
	30	0.03 (0.02, 0.03)	0.07 (0.06, 0.08)	0.13 (0.11, 0.14)	0.66 (0.58, 0.77)	0.46 (0.39, 0.54)	
	40	0.04 (0.03, 0.05)	0.10 (0.09, 0.12)	0.20 (0.18, 0.23)	0.64 (0.56, 0.74)	0.45 (0.38, 0.54)	
	50	0.06 (0.05, 0.07)	0.16 (0.14, 0.19)	0.34 (0.30, 0.39)	0.61 (0.52, 0.71)	0.44 (0.37, 0.53)	
	60	0.11 (0.09, 0.13)	0.29 (0.25, 0.33)	0.47 (0.41, 0.53)	0.56 (0.48, 0.67)	0.42 (0.35, 0.51)	
	70	0.20 (0.17, 0.23)	0.38 (0.33, 0.45)	- (-, -)	0.48 (0.40, 0.59)	0.39 (0.32, 0.48)	
	80	0.22 (0.18, 0.28)	- (-, -)	- (-, -)	0.34 (0.25, 0.47)	0.31 (0.23, 0.42)	
American	0	0.03 (0.01, 0.08)	0.05 (0.02, 0.11)	0.08 (0.04, 0.14)	0.85 (0.50, 1.52)	0.49 (0.37, 0.69)	
Indian/ Alaska Native ^a	10	0.02 (0.01, 0.05)	0.05 (0.02, 0.10)	0.09 (0.05, 0.15)	0.83 (0.48, 1.52)	0.49 (0.37, 0.69)	
	20	0.03 (0.01, 0.07)	0.07 (0.03, 0.13)	0.10 (0.06, 0.17)	0.82 (0.47, 1.50)	0.49 (0.36, 0.69)	
	30	0.04 (0.02, 0.08)	0.07 (0.04, 0.13)	0.13 (0.07, 0.22)	0.79 (0.44, 1.49)	0.48 (0.35, 0.68)	
	40	0.03 (0.01, 0.08)	0.09 (0.05, 0.17)	0.17 (0.10, 0.30)	0.77 (0.41, 1.47)	0.47 (0.35, 0.68)	
	50	0.06 (0.02, 0.13)	0.14 (0.07, 0.27)	0.22 (0.11, 0.40)	0.76 (0.40, 1.48)	0.49 (0.35, 0.70)	
	60	0.09 (0.03, 0.20)	0.17 (0.07, 0.34)	0.45 (0.24, 0.78)	0.73 (0.36, 1.50)	0.47 (0.34, 0.70)	
	70	0.09 (0.02, 0.25)	0.41 (0.19, 0.78)	- (-, -)	0.74 (0.33, 1.62)	0.49 (0.33, 0.75)	
	80	0.43 (0.17, 0.89)	- (-, -)	- (-, -)	0.87 (0.35, 2.04)	0.48 (0.30, 0.82)	
Hispanic ^b	0	0.06 (0.05, 0.07)	0.09 (0.09, 0.10)	0.11 (0.10, 0.12)	0.98 (0.90, 1.08)	0.57 (0.53, 0.61)	
	10	0.03 (0.03, 0.04)	0.05 (0.05, 0.06)	0.08 (0.08, 0.09)	0.93 (0.85, 1.03)	0.56 (0.52, 0.61)	
	20	0.02 (0.02, 0.02)	0.05 (0.04, 0.06)	0.10 (0.09, 0.11)	0.90 (0.81, 1.00)	0.55 (0.51, 0.60)	
	30	0.03 (0.02, 0.03)	0.08 (0.07, 0.09)	0.16 (0.15, 0.18)	0.88 (0.80, 0.98)	0.55 (0.51, 0.59)	
	40	0.05 (0.04, 0.05)	0.13 (0.12, 0.15)	0.31 (0.28, 0.34)	0.86 (0.77, 0.96)	0.54 (0.50, 0.58)	
	50	0.09 (0.08, 0.10)	0.27 (0.24, 0.30)	0.47 (0.43, 0.52)	0.82 (0.74, 0.92)	0.52 (0.48, 0.57)	
	60	0.19 (0.17, 0.22)	0.40 (0.36, 0.44)	0.61 (0.55, 0.67)	0.76 (0.68, 0.86)	0.50 (0.46, 0.55)	
	70	0.23 (0.20, 0.26)	0.46 (0.40, 0.52)	- (-, -)	0.62 (0.54, 0.73)	0.46 (0.41, 0.51)	
	80	0.29 (0.24, 0.35)	- (-, -)	- (-, -)	0.50 (0.40, 0.62)	0.38 (0.34, 0.44)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XIV-9
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.01 (0.01, 0.01)	0.66 (0.64, 0.67)	0.54 (0.53, 0.54)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.66 (0.65, 0.67)	0.54 (0.54, 0.55)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.05, 0.05)	0.66 (0.65, 0.67)	0.55 (0.54, 0.55)	
	30	0.01 (0.01, 0.01)	0.05 (0.05, 0.05)	0.17 (0.16, 0.17)	0.66 (0.65, 0.68)	0.55 (0.54, 0.56)	
	40	0.04 (0.04, 0.05)	0.16 (0.16, 0.17)	0.33 (0.33, 0.34)	0.67 (0.65, 0.68)	0.55 (0.55, 0.56)	
	50	0.12 (0.12, 0.13)	0.30 (0.29, 0.30)	0.50 (0.49, 0.51)	0.64 (0.63, 0.65)	0.54 (0.54, 0.55)	
	60	0.19 (0.18, 0.19)	0.40 (0.40, 0.41)	0.52 (0.51, 0.53)	0.55 (0.54, 0.56)	0.50 (0.49, 0.50)	
	70	0.25 (0.25, 0.26)	0.39 (0.38, 0.40)	- (- , -)	0.42 (0.41, 0.43)	0.42 (0.42, 0.43)	
	80	0.19 (0.19, 0.20)	- (- , -)	- (- , -)	0.24 (0.23, 0.25)	0.29 (0.29, 0.30)	
White	0	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.01 (0.01, 0.01)	0.55 (0.54, 0.57)	0.50 (0.50, 0.51)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.56 (0.54, 0.57)	0.51 (0.50, 0.51)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.04, 0.04)	0.56 (0.54, 0.57)	0.51 (0.51, 0.52)	
	30	0.00 (0.00, 0.01)	0.04 (0.04, 0.04)	0.14 (0.13, 0.14)	0.56 (0.55, 0.57)	0.51 (0.51, 0.52)	
	40	0.04 (0.03, 0.04)	0.13 (0.13, 0.14)	0.27 (0.26, 0.28)	0.56 (0.55, 0.57)	0.52 (0.51, 0.52)	
	50	0.10 (0.10, 0.11)	0.24 (0.24, 0.25)	0.42 (0.41, 0.43)	0.54 (0.53, 0.55)	0.51 (0.50, 0.52)	
	60	0.15 (0.14, 0.16)	0.34 (0.33, 0.35)	0.44 (0.43, 0.45)	0.46 (0.45, 0.48)	0.47 (0.47, 0.48)	
	70	0.22 (0.21, 0.22)	0.34 (0.33, 0.35)	- (- , -)	0.36 (0.35, 0.37)	0.40 (0.40, 0.41)	
	80	0.17 (0.16, 0.18)	- (- , -)	- (- , -)	0.21 (0.20, 0.22)	0.28 (0.27, 0.29)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.68 (0.64, 0.71)	0.58 (0.57, 0.60)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.01, 0.02)	0.69 (0.65, 0.73)	0.60 (0.58, 0.61)	
	20	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.07 (0.06, 0.08)	0.69 (0.66, 0.73)	0.60 (0.58, 0.62)	
	30	0.01 (0.01, 0.01)	0.07 (0.06, 0.08)	0.25 (0.23, 0.27)	0.70 (0.66, 0.74)	0.61 (0.59, 0.62)	
	40	0.06 (0.05, 0.07)	0.25 (0.23, 0.26)	0.44 (0.42, 0.47)	0.71 (0.67, 0.75)	0.61 (0.60, 0.63)	
	50	0.20 (0.18, 0.21)	0.40 (0.38, 0.43)	0.57 (0.54, 0.61)	0.68 (0.64, 0.72)	0.60 (0.58, 0.62)	
	60	0.23 (0.21, 0.25)	0.42 (0.39, 0.45)	0.52 (0.49, 0.56)	0.54 (0.51, 0.59)	0.52 (0.50, 0.54)	
	70	0.24 (0.21, 0.26)	0.37 (0.33, 0.40)	- (- , -)	0.39 (0.35, 0.43)	0.44 (0.42, 0.46)	
	80	0.21 (0.18, 0.25)	- (- , -)	- (- , -)	0.25 (0.21, 0.30)	0.32 (0.30, 0.34)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XIV-9 - continued

LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.01 (0.00, 0.01)	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	1.78 (1.70, 1.88)	1.54 (1.46, 1.63)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	1.80 (1.71, 1.89)	1.55 (1.47, 1.64)	
Islander	20	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	0.11 (0.10, 0.12)	1.80 (1.71, 1.89)	1.55 (1.48, 1.64)	
	30	0.02 (0.02, 0.03)	0.11 (0.10, 0.12)	0.32 (0.30, 0.34)	1.80 (1.72, 1.90)	1.56 (1.48, 1.64)	
	40	0.09 (0.08, 0.10)	0.30 (0.28, 0.32)	0.74 (0.71, 0.78)	1.79 (1.70, 1.89)	1.55 (1.47, 1.64)	
	50	0.22 (0.20, 0.23)	0.67 (0.63, 0.70)	1.25 (1.19, 1.30)	1.73 (1.64, 1.82)	1.52 (1.44, 1.60)	
	60	0.47 (0.44, 0.50)	1.07 (1.01, 1.12)	1.44 (1.36, 1.51)	1.57 (1.48, 1.66)	1.42 (1.34, 1.51)	
	70	0.65 (0.61, 0.70)	1.05 (0.99, 1.13)	- (-, -)	1.20 (1.11, 1.29)	1.19 (1.10, 1.28)	
	80	0.50 (0.45, 0.56)	- (-, -)	- (-, -)	0.68 (0.59, 0.78)	0.82 (0.73, 0.93)	
American	0	0.01 (0.00, 0.04)	0.01 (0.00, 0.04)	0.02 (0.01, 0.05)	1.08 (0.81, 1.51)	0.88 (0.75, 1.05)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.01)	0.01 (0.00, 0.03)	0.01 (0.00, 0.04)	1.09 (0.81, 1.52)	0.90 (0.77, 1.07)	
	20	0.01 (0.00, 0.02)	0.01 (0.00, 0.03)	0.08 (0.05, 0.12)	1.09 (0.82, 1.53)	0.91 (0.77, 1.08)	
	30	0.01 (0.00, 0.02)	0.07 (0.05, 0.11)	0.28 (0.21, 0.36)	1.11 (0.82, 1.55)	0.92 (0.79, 1.10)	
	40	0.07 (0.04, 0.10)	0.28 (0.21, 0.36)	0.51 (0.40, 0.64)	1.13 (0.84, 1.59)	0.95 (0.80, 1.13)	
	50	0.22 (0.16, 0.30)	0.45 (0.35, 0.58)	0.71 (0.56, 0.89)	1.10 (0.81, 1.58)	0.96 (0.81, 1.15)	
	60	0.25 (0.17, 0.36)	0.52 (0.38, 0.70)	0.76 (0.56, 1.02)	0.95 (0.64, 1.46)	0.92 (0.77, 1.13)	
	70	0.32 (0.20, 0.48)	0.59 (0.39, 0.87)	- (-, -)	0.82 (0.49, 1.40)	0.88 (0.71, 1.12)	
	80	0.39 (0.20, 0.69)	- (-, -)	- (-, -)	0.70 (0.31, 1.50)	0.77 (0.55, 1.08)	
Hispanic ^b	0	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.01 (0.01, 0.01)	1.11 (1.05, 1.17)	0.99 (0.95, 1.03)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.12 (1.06, 1.18)	1.00 (0.96, 1.04)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.07 (0.06, 0.07)	1.12 (1.06, 1.18)	1.00 (0.97, 1.04)	
	30	0.01 (0.01, 0.01)	0.07 (0.06, 0.07)	0.24 (0.23, 0.26)	1.13 (1.07, 1.19)	1.01 (0.98, 1.05)	
	40	0.06 (0.05, 0.07)	0.24 (0.22, 0.25)	0.52 (0.49, 0.55)	1.13 (1.08, 1.20)	1.02 (0.98, 1.06)	
	50	0.18 (0.17, 0.20)	0.47 (0.44, 0.50)	0.83 (0.79, 0.87)	1.10 (1.04, 1.16)	1.01 (0.97, 1.05)	
	60	0.30 (0.28, 0.32)	0.68 (0.64, 0.72)	0.90 (0.85, 0.96)	0.96 (0.90, 1.03)	0.94 (0.90, 0.98)	
	70	0.43 (0.39, 0.46)	0.68 (0.63, 0.73)	- (-, -)	0.74 (0.68, 0.80)	0.80 (0.75, 0.84)	
	80	0.33 (0.29, 0.38)	- (-, -)	- (-, -)	0.41 (0.36, 0.48)	0.57 (0.52, 0.62)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XIV-9 - continued

LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.01 (0.00, 0.01)	0.01 (0.01, 0.01)	0.01 (0.01, 0.01)	0.91 (0.89, 0.93)	0.69 (0.69, 0.70)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.92 (0.90, 0.94)	0.70 (0.70, 0.71)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.08 (0.08, 0.09)	0.92 (0.90, 0.94)	0.71 (0.70, 0.72)	
	30	0.01 (0.01, 0.01)	0.08 (0.08, 0.08)	0.27 (0.27, 0.28)	0.93 (0.91, 0.95)	0.72 (0.71, 0.72)	
	40	0.07 (0.07, 0.08)	0.27 (0.26, 0.28)	0.52 (0.50, 0.53)	0.94 (0.92, 0.96)	0.72 (0.71, 0.73)	
	50	0.20 (0.20, 0.21)	0.46 (0.45, 0.47)	0.74 (0.72, 0.76)	0.90 (0.88, 0.92)	0.71 (0.70, 0.72)	
	60	0.28 (0.27, 0.29)	0.58 (0.56, 0.59)	0.72 (0.70, 0.74)	0.75 (0.73, 0.77)	0.64 (0.63, 0.64)	
	70	0.36 (0.35, 0.37)	0.53 (0.51, 0.55)	- (- , -)	0.56 (0.54, 0.58)	0.53 (0.52, 0.54)	
	80	0.26 (0.25, 0.28)	- (- , -)	- (- , -)	0.32 (0.30, 0.34)	0.36 (0.35, 0.37)	
White	0	0.01 (0.00, 0.01)	0.01 (0.01, 0.01)	0.01 (0.01, 0.01)	0.77 (0.75, 0.79)	0.65 (0.64, 0.66)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.77 (0.75, 0.79)	0.65 (0.65, 0.66)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.06 (0.06, 0.07)	0.77 (0.75, 0.79)	0.66 (0.65, 0.67)	
	30	0.01 (0.01, 0.01)	0.06 (0.06, 0.07)	0.22 (0.21, 0.23)	0.78 (0.76, 0.80)	0.66 (0.66, 0.67)	
	40	0.06 (0.05, 0.06)	0.22 (0.21, 0.23)	0.42 (0.41, 0.43)	0.79 (0.77, 0.81)	0.67 (0.66, 0.68)	
	50	0.17 (0.16, 0.18)	0.37 (0.36, 0.39)	0.62 (0.60, 0.63)	0.76 (0.73, 0.78)	0.66 (0.65, 0.67)	
	60	0.22 (0.21, 0.23)	0.48 (0.47, 0.50)	0.61 (0.59, 0.63)	0.63 (0.61, 0.65)	0.60 (0.59, 0.61)	
	70	0.31 (0.29, 0.32)	0.46 (0.44, 0.48)	- (- , -)	0.49 (0.47, 0.51)	0.51 (0.50, 0.52)	
	80	0.23 (0.21, 0.25)	- (- , -)	- (- , -)	0.27 (0.26, 0.29)	0.35 (0.34, 0.36)	
Black	0	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	0.98 (0.92, 1.05)	0.78 (0.75, 0.80)	
	10	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.02 (0.02, 0.03)	1.00 (0.94, 1.07)	0.80 (0.77, 0.83)	
	20	0.00 (0.00, 0.01)	0.02 (0.01, 0.03)	0.12 (0.10, 0.13)	1.01 (0.95, 1.08)	0.80 (0.78, 0.83)	
	30	0.02 (0.01, 0.02)	0.11 (0.10, 0.13)	0.43 (0.39, 0.46)	1.03 (0.97, 1.10)	0.82 (0.79, 0.85)	
	40	0.10 (0.09, 0.11)	0.42 (0.39, 0.46)	0.72 (0.67, 0.77)	1.05 (0.98, 1.12)	0.83 (0.80, 0.86)	
	50	0.34 (0.31, 0.37)	0.66 (0.61, 0.71)	0.89 (0.83, 0.95)	1.01 (0.94, 1.09)	0.81 (0.78, 0.84)	
	60	0.37 (0.33, 0.41)	0.63 (0.57, 0.69)	0.74 (0.68, 0.81)	0.77 (0.71, 0.85)	0.68 (0.64, 0.71)	
	70	0.35 (0.30, 0.40)	0.50 (0.44, 0.57)	- (- , -)	0.54 (0.47, 0.62)	0.56 (0.52, 0.59)	
	80	0.28 (0.22, 0.35)	- (- , -)	- (- , -)	0.35 (0.27, 0.46)	0.40 (0.36, 0.45)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XIV-9 - continued

LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.02 (0.01, 0.03)	2.47 (2.33, 2.62)	1.97 (1.86, 2.10)	
	10	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.05 (0.04, 0.06)	2.48 (2.35, 2.64)	1.99 (1.87, 2.12)	
Islander	20	0.01 (0.00, 0.01)	0.05 (0.04, 0.06)	0.20 (0.18, 0.22)	2.49 (2.35, 2.64)	1.99 (1.88, 2.12)	
	30	0.04 (0.03, 0.05)	0.19 (0.17, 0.21)	0.55 (0.51, 0.59)	2.50 (2.36, 2.65)	2.00 (1.88, 2.13)	
	40	0.16 (0.14, 0.17)	0.52 (0.48, 0.56)	1.17 (1.10, 1.24)	2.48 (2.34, 2.63)	1.99 (1.87, 2.12)	
	50	0.37 (0.34, 0.40)	1.03 (0.97, 1.10)	1.83 (1.72, 1.93)	2.37 (2.23, 2.52)	1.92 (1.81, 2.06)	
	60	0.69 (0.64, 0.75)	1.52 (1.42, 1.62)	1.97 (1.85, 2.10)	2.09 (1.95, 2.25)	1.75 (1.63, 1.89)	
	70	0.92 (0.84, 1.01)	1.43 (1.31, 1.55)	- (- , -)	1.55 (1.42, 1.71)	1.43 (1.31, 1.57)	
	80	0.67 (0.57, 0.77)	- (- , -)	- (- , -)	0.83 (0.70, 1.00)	0.93 (0.80, 1.09)	
American	0	0.02 (0.00, 0.08)	0.03 (0.01, 0.09)	0.03 (0.01, 0.09)	1.46 (1.01, 2.32)	0.86 (0.71, 1.12)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.02 (0.01, 0.06)	1.47 (1.01, 2.34)	0.88 (0.72, 1.14)	
	20	0.01 (0.00, 0.04)	0.02 (0.00, 0.05)	0.12 (0.07, 0.19)	1.48 (1.01, 2.36)	0.89 (0.72, 1.15)	
	30	0.01 (0.00, 0.04)	0.11 (0.07, 0.18)	0.45 (0.33, 0.60)	1.51 (1.03, 2.42)	0.91 (0.74, 1.18)	
	40	0.11 (0.06, 0.17)	0.46 (0.34, 0.61)	0.74 (0.56, 0.97)	1.55 (1.06, 2.49)	0.94 (0.76, 1.22)	
	50	0.37 (0.26, 0.51)	0.67 (0.49, 0.90)	0.95 (0.71, 1.27)	1.52 (1.01, 2.51)	0.95 (0.77, 1.25)	
	60	0.33 (0.19, 0.52)	0.64 (0.42, 0.94)	1.01 (0.67, 1.50)	1.26 (0.73, 2.34)	0.87 (0.67, 1.19)	
	70	0.37 (0.20, 0.66)	0.82 (0.47, 1.37)	- (- , -)	1.12 (0.54, 2.42)	0.82 (0.60, 1.22)	
	80	0.68 (0.30, 1.37)	- (- , -)	- (- , -)	1.12 (0.38, 3.06)	0.62 (0.37, 1.17)	
Hispanic ^b	0	0.01 (0.00, 0.01)	0.01 (0.00, 0.01)	0.01 (0.01, 0.01)	1.47 (1.38, 1.58)	1.24 (1.18, 1.31)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	1.48 (1.40, 1.59)	1.26 (1.20, 1.32)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.11 (0.10, 0.12)	1.49 (1.40, 1.60)	1.26 (1.20, 1.33)	
	30	0.01 (0.01, 0.01)	0.11 (0.10, 0.12)	0.40 (0.37, 0.43)	1.51 (1.42, 1.62)	1.28 (1.22, 1.35)	
	40	0.10 (0.09, 0.11)	0.40 (0.37, 0.42)	0.80 (0.75, 0.85)	1.52 (1.43, 1.63)	1.29 (1.23, 1.36)	
	50	0.30 (0.28, 0.33)	0.72 (0.68, 0.77)	1.19 (1.12, 1.26)	1.46 (1.37, 1.58)	1.28 (1.21, 1.35)	
	60	0.45 (0.41, 0.49)	0.94 (0.87, 1.02)	1.17 (1.09, 1.27)	1.24 (1.14, 1.35)	1.17 (1.10, 1.24)	
	70	0.57 (0.51, 0.63)	0.84 (0.76, 0.93)	- (- , -)	0.91 (0.81, 1.03)	0.97 (0.90, 1.05)	
	80	0.38 (0.31, 0.46)	- (- , -)	- (- , -)	0.48 (0.38, 0.62)	0.67 (0.59, 0.77)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XIV-9 - continued

LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.41 (0.40, 0.43)	0.39 (0.39, 0.40)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.41 (0.40, 0.43)	0.40 (0.39, 0.40)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.02, 0.02)	0.41 (0.40, 0.43)	0.40 (0.39, 0.40)	
	30	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.06 (0.06, 0.07)	0.41 (0.40, 0.43)	0.40 (0.39, 0.40)	
	40	0.02 (0.01, 0.02)	0.06 (0.06, 0.06)	0.16 (0.15, 0.16)	0.41 (0.40, 0.43)	0.40 (0.39, 0.40)	
	50	0.04 (0.04, 0.05)	0.14 (0.14, 0.15)	0.29 (0.28, 0.30)	0.41 (0.39, 0.42)	0.40 (0.39, 0.40)	
	60	0.10 (0.10, 0.11)	0.25 (0.24, 0.26)	0.35 (0.34, 0.37)	0.38 (0.37, 0.39)	0.38 (0.37, 0.39)	
	70	0.17 (0.16, 0.18)	0.28 (0.27, 0.29)	- (- , -)	0.31 (0.30, 0.32)	0.34 (0.33, 0.34)	
	80	0.15 (0.14, 0.16)	- (- , -)	- (- , -)	0.19 (0.18, 0.20)	0.25 (0.24, 0.25)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.35 (0.34, 0.36)	0.37 (0.36, 0.38)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.35 (0.34, 0.36)	0.37 (0.37, 0.38)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.02, 0.02)	0.35 (0.34, 0.36)	0.37 (0.37, 0.38)	
	30	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.05 (0.05, 0.06)	0.35 (0.34, 0.36)	0.37 (0.37, 0.38)	
	40	0.01 (0.01, 0.01)	0.05 (0.05, 0.05)	0.13 (0.12, 0.13)	0.35 (0.34, 0.36)	0.38 (0.37, 0.38)	
	50	0.04 (0.03, 0.04)	0.12 (0.11, 0.12)	0.24 (0.23, 0.25)	0.34 (0.33, 0.36)	0.37 (0.37, 0.38)	
	60	0.08 (0.08, 0.09)	0.21 (0.20, 0.22)	0.30 (0.28, 0.31)	0.32 (0.31, 0.33)	0.36 (0.35, 0.37)	
	70	0.14 (0.13, 0.15)	0.24 (0.23, 0.25)	- (- , -)	0.27 (0.25, 0.28)	0.32 (0.31, 0.33)	
	80	0.13 (0.12, 0.14)	- (- , -)	- (- , -)	0.17 (0.16, 0.18)	0.24 (0.23, 0.24)	
Black	0	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.40 (0.36, 0.44)	0.41 (0.39, 0.43)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	0.41 (0.37, 0.45)	0.42 (0.40, 0.44)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.02, 0.04)	0.41 (0.37, 0.45)	0.42 (0.40, 0.44)	
	30	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	0.09 (0.08, 0.11)	0.41 (0.37, 0.45)	0.42 (0.40, 0.44)	
	40	0.02 (0.02, 0.03)	0.09 (0.08, 0.11)	0.19 (0.17, 0.22)	0.41 (0.37, 0.45)	0.42 (0.40, 0.45)	
	50	0.07 (0.06, 0.09)	0.18 (0.16, 0.21)	0.30 (0.27, 0.34)	0.40 (0.36, 0.45)	0.42 (0.40, 0.45)	
	60	0.12 (0.10, 0.14)	0.25 (0.22, 0.29)	0.35 (0.30, 0.39)	0.36 (0.32, 0.41)	0.40 (0.38, 0.42)	
	70	0.16 (0.13, 0.19)	0.27 (0.23, 0.31)	- (- , -)	0.29 (0.25, 0.34)	0.36 (0.34, 0.39)	
	80	0.17 (0.13, 0.21)	- (- , -)	- (- , -)	0.20 (0.15, 0.25)	0.28 (0.26, 0.31)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XIV-9 - continued

LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	1.19 (1.09, 1.32)	1.17 (1.06, 1.30)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	1.20 (1.09, 1.33)	1.18 (1.07, 1.31)	
	20	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	1.20 (1.09, 1.33)	1.18 (1.07, 1.31)	
	30	0.01 (0.00, 0.01)	0.03 (0.02, 0.04)	0.11 (0.09, 0.13)	1.20 (1.09, 1.33)	1.18 (1.07, 1.31)	
	40	0.03 (0.02, 0.03)	0.10 (0.09, 0.12)	0.37 (0.33, 0.41)	1.20 (1.09, 1.33)	1.18 (1.07, 1.31)	
	50	0.08 (0.07, 0.09)	0.35 (0.31, 0.38)	0.76 (0.70, 0.82)	1.18 (1.08, 1.31)	1.18 (1.07, 1.30)	
	60	0.27 (0.24, 0.31)	0.69 (0.64, 0.76)	0.99 (0.91, 1.08)	1.13 (1.03, 1.27)	1.15 (1.03, 1.28)	
	70	0.45 (0.40, 0.50)	0.77 (0.69, 0.85)	- (- , -)	0.92 (0.81, 1.05)	1.00 (0.89, 1.14)	
	80	0.38 (0.32, 0.45)	- (- , -)	- (- , -)	0.56 (0.45, 0.70)	0.75 (0.62, 0.90)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.01 (0.00, 0.05)	0.73 (0.43, 1.37)	0.89 (0.70, 1.16)	
Native ^a	10	0.00 (0.00, 0.02)	0.01 (0.00, 0.04)	0.01 (0.00, 0.04)	0.75 (0.43, 1.39)	0.90 (0.71, 1.17)	
Alaska	20	0.01 (0.00, 0.04)	0.01 (0.00, 0.04)	0.04 (0.01, 0.09)	0.75 (0.43, 1.39)	0.91 (0.71, 1.18)	
	30	0.00 (0.00, 0.03)	0.03 (0.01, 0.07)	0.11 (0.06, 0.20)	0.75 (0.43, 1.40)	0.92 (0.72, 1.19)	
	40	0.03 (0.01, 0.07)	0.11 (0.06, 0.20)	0.28 (0.17, 0.44)	0.76 (0.44, 1.42)	0.93 (0.73, 1.22)	
	50	0.09 (0.04, 0.16)	0.26 (0.15, 0.41)	0.49 (0.31, 0.73)	0.75 (0.42, 1.43)	0.95 (0.74, 1.24)	
	60	0.18 (0.09, 0.32)	0.42 (0.25, 0.67)	0.54 (0.33, 0.87)	0.70 (0.37, 1.43)	0.96 (0.73, 1.27)	
	70	0.28 (0.14, 0.51)	0.41 (0.21, 0.76)	- (- , -)	0.60 (0.25, 1.42)	0.91 (0.66, 1.27)	
	80	0.18 (0.04, 0.54)	- (- , -)	- (- , -)	0.43 (0.08, 1.52)	0.85 (0.55, 1.31)	
Hispanic ^b	0	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.77 (0.71, 0.85)	0.77 (0.72, 0.82)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.78 (0.71, 0.86)	0.77 (0.73, 0.83)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.02, 0.03)	0.78 (0.71, 0.86)	0.78 (0.73, 0.83)	
	30	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.09 (0.08, 0.10)	0.78 (0.71, 0.86)	0.78 (0.73, 0.83)	
	40	0.02 (0.01, 0.02)	0.09 (0.07, 0.10)	0.25 (0.22, 0.28)	0.78 (0.71, 0.86)	0.78 (0.73, 0.83)	
	50	0.07 (0.06, 0.08)	0.24 (0.21, 0.26)	0.52 (0.47, 0.56)	0.78 (0.71, 0.86)	0.78 (0.73, 0.83)	
	60	0.17 (0.15, 0.20)	0.46 (0.42, 0.51)	0.68 (0.61, 0.74)	0.73 (0.66, 0.82)	0.76 (0.71, 0.81)	
	70	0.32 (0.28, 0.36)	0.55 (0.49, 0.62)	- (- , -)	0.61 (0.54, 0.70)	0.67 (0.62, 0.73)	
	80	0.30 (0.24, 0.36)	- (- , -)	- (- , -)	0.37 (0.30, 0.46)	0.51 (0.45, 0.57)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XV-15

LUNG AND BRONCHUS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	6.98 (6.94, 7.02)	5.99 (5.97, 6.00)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.03, 0.03)	7.07 (7.03, 7.11)	6.07 (6.05, 6.09)	
	20	0.00 (0.00, 0.01)	0.03 (0.03, 0.03)	0.22 (0.21, 0.22)	7.10 (7.06, 7.13)	6.10 (6.08, 6.12)	
	30	0.03 (0.03, 0.03)	0.21 (0.21, 0.22)	0.94 (0.93, 0.95)	7.16 (7.12, 7.20)	6.15 (6.14, 6.17)	
	40	0.19 (0.19, 0.19)	0.93 (0.92, 0.94)	2.88 (2.86, 2.90)	7.22 (7.18, 7.26)	6.22 (6.20, 6.24)	
	50	0.76 (0.75, 0.77)	2.77 (2.75, 2.79)	5.51 (5.48, 5.54)	7.24 (7.20, 7.28)	6.25 (6.24, 6.27)	
	60	2.14 (2.12, 2.16)	5.06 (5.03, 5.10)	6.63 (6.59, 6.67)	6.90 (6.86, 6.95)	6.01 (5.99, 6.03)	
	70	3.39 (3.36, 3.42)	5.21 (5.17, 5.25)	- (- , -)	5.52 (5.48, 5.56)	4.94 (4.92, 4.95)	
	80	2.61 (2.58, 2.64)	- (- , -)	- (- , -)	3.06 (3.02, 3.09)	2.94 (2.92, 2.96)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	7.17 (7.12, 7.21)	6.13 (6.11, 6.15)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.03, 0.03)	7.25 (7.20, 7.29)	6.20 (6.18, 6.22)	
	20	0.00 (0.00, 0.01)	0.03 (0.03, 0.03)	0.21 (0.21, 0.21)	7.27 (7.23, 7.32)	6.23 (6.21, 6.25)	
	30	0.03 (0.02, 0.03)	0.21 (0.20, 0.21)	0.93 (0.91, 0.94)	7.33 (7.29, 7.38)	6.28 (6.26, 6.30)	
	40	0.18 (0.18, 0.19)	0.91 (0.90, 0.92)	2.94 (2.91, 2.96)	7.39 (7.35, 7.44)	6.34 (6.32, 6.36)	
	50	0.75 (0.74, 0.76)	2.83 (2.80, 2.85)	5.68 (5.64, 5.71)	7.41 (7.36, 7.46)	6.37 (6.35, 6.39)	
	60	2.21 (2.19, 2.23)	5.24 (5.20, 5.27)	6.82 (6.77, 6.86)	7.08 (7.03, 7.12)	6.12 (6.10, 6.14)	
	70	3.50 (3.47, 3.53)	5.33 (5.29, 5.38)	- (- , -)	5.63 (5.59, 5.68)	5.01 (4.99, 5.03)	
	80	2.63 (2.60, 2.67)	- (- , -)	- (- , -)	3.06 (3.03, 3.10)	2.96 (2.94, 2.98)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	6.64 (6.52, 6.76)	5.50 (5.45, 5.55)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.04 (0.03, 0.04)	6.80 (6.68, 6.93)	5.64 (5.59, 5.69)	
	20	0.01 (0.00, 0.01)	0.04 (0.03, 0.04)	0.33 (0.32, 0.35)	6.84 (6.72, 6.97)	5.67 (5.62, 5.73)	
	30	0.03 (0.03, 0.04)	0.34 (0.32, 0.35)	1.39 (1.35, 1.43)	6.95 (6.82, 7.08)	5.76 (5.71, 5.81)	
	40	0.31 (0.29, 0.32)	1.39 (1.35, 1.43)	3.51 (3.44, 3.59)	7.08 (6.96, 7.21)	5.88 (5.83, 5.93)	
	50	1.14 (1.11, 1.18)	3.38 (3.30, 3.46)	5.79 (5.68, 5.91)	7.14 (7.01, 7.28)	5.95 (5.89, 6.00)	
	60	2.51 (2.44, 2.58)	5.22 (5.11, 5.34)	6.51 (6.38, 6.65)	6.74 (6.60, 6.89)	5.68 (5.63, 5.74)	
	70	3.41 (3.31, 3.52)	5.04 (4.91, 5.18)	- (- , -)	5.33 (5.19, 5.47)	4.63 (4.57, 4.69)	
	80	2.61 (2.50, 2.74)	- (- , -)	- (- , -)	3.08 (2.94, 3.22)	2.84 (2.78, 2.90)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XV-15 - continued

LUNG AND BRONCHUS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	5.55 (5.39, 5.73)	4.12 (3.99, 4.25)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.03 (0.03, 0.04)	5.60 (5.44, 5.78)	4.16 (4.03, 4.29)	
	20	0.00 (0.00, 0.00)	0.03 (0.03, 0.04)	0.15 (0.14, 0.16)	5.62 (5.45, 5.79)	4.17 (4.04, 4.30)	
	30	0.03 (0.02, 0.03)	0.14 (0.13, 0.16)	0.58 (0.55, 0.60)	5.64 (5.47, 5.82)	4.18 (4.05, 4.32)	
	40	0.12 (0.11, 0.13)	0.55 (0.52, 0.58)	1.66 (1.60, 1.72)	5.65 (5.48, 5.82)	4.19 (4.06, 4.32)	
	50	0.44 (0.42, 0.46)	1.56 (1.51, 1.62)	3.46 (3.36, 3.56)	5.61 (5.44, 5.79)	4.17 (4.04, 4.31)	
	60	1.16 (1.12, 1.22)	3.13 (3.03, 3.23)	4.79 (4.65, 4.93)	5.36 (5.18, 5.54)	4.04 (3.91, 4.18)	
	70	2.14 (2.06, 2.22)	3.95 (3.82, 4.09)	- (- , -)	4.57 (4.39, 4.76)	3.60 (3.47, 3.75)	
	80	2.26 (2.14, 2.39)	- (- , -)	- (- , -)	3.04 (2.85, 3.24)	2.56 (2.41, 2.73)	
American	0	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.01 (0.00, 0.03)	4.66 (4.18, 5.27)	4.13 (3.88, 4.42)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.01)	0.01 (0.00, 0.02)	0.02 (0.01, 0.04)	4.74 (4.25, 5.36)	4.21 (3.95, 4.50)	
	20	0.01 (0.00, 0.02)	0.02 (0.01, 0.04)	0.10 (0.07, 0.14)	4.78 (4.28, 5.40)	4.24 (3.98, 4.54)	
	30	0.01 (0.00, 0.03)	0.09 (0.06, 0.13)	0.61 (0.51, 0.72)	4.85 (4.35, 5.49)	4.32 (4.06, 4.62)	
	40	0.08 (0.06, 0.12)	0.62 (0.52, 0.73)	1.96 (1.73, 2.21)	4.97 (4.46, 5.62)	4.44 (4.17, 4.75)	
	50	0.55 (0.46, 0.66)	1.95 (1.72, 2.21)	3.88 (3.49, 4.31)	5.09 (4.55, 5.76)	4.57 (4.28, 4.89)	
	60	1.50 (1.29, 1.74)	3.58 (3.18, 4.02)	4.71 (4.20, 5.29)	4.88 (4.32, 5.59)	4.52 (4.21, 4.86)	
	70	2.44 (2.08, 2.85)	3.77 (3.25, 4.37)	- (- , -)	3.96 (3.39, 4.73)	3.72 (3.40, 4.09)	
	80	1.88 (1.44, 2.44)	- (- , -)	- (- , -)	2.15 (1.60, 3.02)	2.29 (1.95, 2.72)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	4.38 (4.25, 4.52)	3.17 (3.10, 3.24)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	4.43 (4.30, 4.57)	3.21 (3.14, 3.28)	
	20	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.09 (0.08, 0.10)	4.45 (4.32, 4.59)	3.22 (3.15, 3.29)	
	30	0.01 (0.01, 0.01)	0.09 (0.08, 0.09)	0.40 (0.39, 0.42)	4.48 (4.35, 4.62)	3.25 (3.18, 3.32)	
	40	0.08 (0.07, 0.08)	0.40 (0.38, 0.41)	1.35 (1.31, 1.40)	4.51 (4.38, 4.65)	3.27 (3.20, 3.35)	
	50	0.33 (0.31, 0.35)	1.31 (1.26, 1.35)	2.94 (2.86, 3.02)	4.54 (4.40, 4.68)	3.30 (3.23, 3.37)	
	60	1.03 (0.99, 1.07)	2.75 (2.66, 2.83)	3.99 (3.88, 4.11)	4.43 (4.29, 4.58)	3.24 (3.17, 3.32)	
	70	1.93 (1.86, 2.00)	3.33 (3.21, 3.45)	- (- , -)	3.82 (3.67, 3.98)	2.89 (2.81, 2.97)	
	80	1.86 (1.75, 1.97)	- (- , -)	- (- , -)	2.51 (2.35, 2.69)	2.05 (1.96, 2.14)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XV-15 - continued

LUNG AND BRONCHUS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	7.91 (7.85, 7.98)	7.12 (7.09, 7.14)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.03 (0.03, 0.03)	8.02 (7.96, 8.08)	7.23 (7.20, 7.26)	
	20	0.00 (0.00, 0.01)	0.03 (0.03, 0.03)	0.22 (0.22, 0.23)	8.07 (8.01, 8.13)	7.27 (7.24, 7.30)	
	30	0.03 (0.02, 0.03)	0.22 (0.22, 0.23)	1.04 (1.02, 1.05)	8.17 (8.11, 8.23)	7.37 (7.34, 7.40)	
	40	0.20 (0.19, 0.21)	1.03 (1.02, 1.05)	3.28 (3.24, 3.31)	8.28 (8.22, 8.35)	7.48 (7.45, 7.51)	
	50	0.86 (0.85, 0.88)	3.19 (3.16, 3.22)	6.40 (6.35, 6.45)	8.38 (8.32, 8.45)	7.59 (7.56, 7.62)	
	60	2.52 (2.49, 2.55)	5.99 (5.94, 6.05)	7.83 (7.77, 7.90)	8.14 (8.07, 8.21)	7.42 (7.39, 7.45)	
	70	4.16 (4.12, 4.21)	6.37 (6.31, 6.43)	- (- , -)	6.74 (6.67, 6.81)	6.29 (6.26, 6.33)	
	80	3.43 (3.38, 3.49)	- (- , -)	- (- , -)	4.00 (3.94, 4.07)	4.01 (3.98, 4.05)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	7.98 (7.92, 8.05)	7.20 (7.17, 7.23)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.03 (0.03, 0.03)	8.08 (8.01, 8.14)	7.29 (7.27, 7.32)	
	20	0.00 (0.00, 0.01)	0.03 (0.03, 0.03)	0.21 (0.20, 0.22)	8.12 (8.05, 8.19)	7.34 (7.31, 7.37)	
	30	0.02 (0.02, 0.03)	0.21 (0.20, 0.21)	0.99 (0.97, 1.01)	8.21 (8.14, 8.28)	7.43 (7.40, 7.45)	
	40	0.19 (0.18, 0.19)	0.98 (0.96, 1.00)	3.25 (3.22, 3.29)	8.32 (8.25, 8.39)	7.53 (7.50, 7.56)	
	50	0.82 (0.81, 0.84)	3.17 (3.14, 3.21)	6.45 (6.39, 6.51)	8.42 (8.35, 8.49)	7.63 (7.60, 7.67)	
	60	2.53 (2.50, 2.57)	6.06 (6.00, 6.12)	7.90 (7.83, 7.97)	8.19 (8.12, 8.27)	7.46 (7.42, 7.49)	
	70	4.22 (4.16, 4.27)	6.41 (6.34, 6.47)	- (- , -)	6.76 (6.68, 6.83)	6.32 (6.29, 6.35)	
	80	3.41 (3.36, 3.47)	- (- , -)	- (- , -)	3.96 (3.89, 4.04)	4.01 (3.97, 4.04)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	8.09 (7.90, 8.29)	7.09 (7.01, 7.17)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.04 (0.03, 0.05)	8.31 (8.11, 8.51)	7.29 (7.20, 7.37)	
	20	0.01 (0.00, 0.01)	0.04 (0.03, 0.05)	0.38 (0.36, 0.41)	8.38 (8.18, 8.59)	7.35 (7.26, 7.43)	
	30	0.03 (0.03, 0.04)	0.39 (0.36, 0.41)	1.76 (1.70, 1.83)	8.59 (8.38, 8.80)	7.51 (7.43, 7.60)	
	40	0.36 (0.34, 0.39)	1.78 (1.72, 1.85)	4.49 (4.36, 4.61)	8.83 (8.62, 9.05)	7.73 (7.64, 7.81)	
	50	1.52 (1.46, 1.58)	4.40 (4.28, 4.54)	7.41 (7.22, 7.60)	9.04 (8.82, 9.27)	7.93 (7.84, 8.02)	
	60	3.35 (3.23, 3.47)	6.84 (6.64, 7.04)	8.46 (8.23, 8.69)	8.73 (8.49, 8.98)	7.77 (7.67, 7.87)	
	70	4.68 (4.50, 4.87)	6.86 (6.61, 7.11)	- (- , -)	7.23 (6.97, 7.50)	6.58 (6.47, 6.70)	
	80	3.97 (3.72, 4.22)	- (- , -)	- (- , -)	4.64 (4.35, 4.96)	4.34 (4.22, 4.47)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XV-15 - continued

LUNG AND BRONCHUS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	7.06 (6.80, 7.34)	5.34 (5.14, 5.55)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.03 (0.02, 0.04)	7.13 (6.87, 7.41)	5.39 (5.19, 5.60)	
	20	0.00 (0.00, 0.00)	0.03 (0.02, 0.04)	0.17 (0.15, 0.19)	7.16 (6.89, 7.44)	5.41 (5.21, 5.62)	
	30	0.03 (0.02, 0.04)	0.17 (0.15, 0.18)	0.70 (0.66, 0.75)	7.20 (6.94, 7.48)	5.43 (5.23, 5.65)	
	40	0.14 (0.12, 0.15)	0.68 (0.64, 0.72)	2.14 (2.05, 2.24)	7.23 (6.96, 7.51)	5.45 (5.25, 5.67)	
	50	0.55 (0.51, 0.59)	2.04 (1.95, 2.14)	4.69 (4.52, 4.86)	7.22 (6.95, 7.51)	5.45 (5.25, 5.67)	
	60	1.56 (1.47, 1.64)	4.32 (4.16, 4.50)	6.42 (6.18, 6.66)	6.97 (6.70, 7.27)	5.33 (5.12, 5.56)	
	70	3.09 (2.94, 3.24)	5.43 (5.19, 5.67)	- (- , -)	6.04 (5.76, 6.35)	4.82 (4.60, 5.07)	
	80	3.11 (2.90, 3.34)	- (- , -)	- (- , -)	3.93 (3.63, 4.27)	3.40 (3.15, 3.67)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.01 (0.00, 0.06)	5.01 (4.25, 6.10)	4.58 (4.18, 5.06)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.01 (0.00, 0.05)	0.03 (0.01, 0.07)	5.11 (4.34, 6.23)	4.68 (4.27, 5.17)	
	20	0.01 (0.00, 0.04)	0.03 (0.01, 0.07)	0.11 (0.07, 0.17)	5.17 (4.39, 6.29)	4.73 (4.32, 5.23)	
	30	0.02 (0.00, 0.05)	0.10 (0.06, 0.16)	0.71 (0.56, 0.89)	5.28 (4.48, 6.43)	4.86 (4.43, 5.36)	
	40	0.09 (0.05, 0.14)	0.72 (0.56, 0.90)	2.07 (1.74, 2.45)	5.46 (4.63, 6.65)	5.04 (4.60, 5.56)	
	50	0.66 (0.51, 0.84)	2.09 (1.74, 2.48)	4.19 (3.60, 4.85)	5.65 (4.78, 6.91)	5.25 (4.79, 5.81)	
	60	1.56 (1.25, 1.92)	3.85 (3.25, 4.54)	5.18 (4.38, 6.11)	5.45 (4.53, 6.81)	5.25 (4.74, 5.87)	
	70	2.76 (2.20, 3.42)	4.36 (3.52, 5.37)	- (- , -)	4.69 (3.69, 6.23)	4.57 (4.00, 5.28)	
	80	2.43 (1.65, 3.50)	- (- , -)	- (- , -)	2.93 (1.83, 5.01)	3.14 (2.46, 4.07)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	5.32 (5.11, 5.56)	4.29 (4.17, 4.41)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	5.39 (5.17, 5.62)	4.34 (4.22, 4.47)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.09 (0.08, 0.10)	5.42 (5.20, 5.66)	4.37 (4.25, 4.49)	
	30	0.01 (0.01, 0.01)	0.09 (0.08, 0.10)	0.44 (0.42, 0.47)	5.48 (5.26, 5.72)	4.42 (4.30, 4.55)	
	40	0.08 (0.07, 0.08)	0.44 (0.41, 0.47)	1.63 (1.56, 1.70)	5.54 (5.32, 5.79)	4.47 (4.35, 4.60)	
	50	0.37 (0.35, 0.40)	1.60 (1.53, 1.68)	3.70 (3.57, 3.84)	5.64 (5.41, 5.89)	4.55 (4.43, 4.69)	
	60	1.31 (1.24, 1.38)	3.55 (3.41, 3.69)	5.10 (4.91, 5.30)	5.62 (5.38, 5.88)	4.53 (4.40, 4.67)	
	70	2.59 (2.46, 2.72)	4.38 (4.18, 4.59)	- (- , -)	4.98 (4.72, 5.27)	4.11 (3.97, 4.27)	
	80	2.54 (2.35, 2.75)	- (- , -)	- (- , -)	3.39 (3.09, 3.74)	2.99 (2.81, 3.18)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XV-15 - continued

LUNG AND BRONCHUS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	6.18 (6.13, 6.23)	5.00 (4.98, 5.02)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.03, 0.04)	6.25 (6.20, 6.30)	5.06 (5.04, 5.09)	
	20	0.00 (0.00, 0.01)	0.03 (0.03, 0.04)	0.21 (0.21, 0.22)	6.27 (6.22, 6.32)	5.08 (5.06, 5.10)	
	30	0.03 (0.03, 0.03)	0.21 (0.20, 0.21)	0.85 (0.83, 0.86)	6.29 (6.24, 6.35)	5.10 (5.08, 5.13)	
	40	0.18 (0.18, 0.19)	0.82 (0.81, 0.84)	2.51 (2.48, 2.54)	6.32 (6.27, 6.37)	5.13 (5.11, 5.16)	
	50	0.66 (0.65, 0.67)	2.38 (2.35, 2.41)	4.73 (4.68, 4.77)	6.27 (6.22, 6.33)	5.12 (5.09, 5.14)	
	60	1.81 (1.78, 1.83)	4.27 (4.23, 4.31)	5.64 (5.59, 5.69)	5.90 (5.84, 5.95)	4.86 (4.84, 4.88)	
	70	2.78 (2.74, 2.81)	4.32 (4.28, 4.37)	- (- , -)	4.61 (4.56, 4.66)	3.92 (3.90, 3.94)	
	80	2.08 (2.05, 2.12)	- (- , -)	- (- , -)	2.47 (2.43, 2.51)	2.29 (2.27, 2.31)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	6.49 (6.43, 6.54)	5.20 (5.17, 5.22)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.03, 0.04)	6.55 (6.49, 6.61)	5.25 (5.23, 5.28)	
	20	0.00 (0.00, 0.01)	0.03 (0.03, 0.04)	0.21 (0.20, 0.22)	6.56 (6.51, 6.62)	5.27 (5.24, 5.29)	
	30	0.03 (0.03, 0.03)	0.21 (0.20, 0.21)	0.86 (0.85, 0.88)	6.59 (6.53, 6.65)	5.29 (5.27, 5.31)	
	40	0.18 (0.17, 0.19)	0.84 (0.83, 0.86)	2.63 (2.60, 2.67)	6.61 (6.56, 6.67)	5.32 (5.29, 5.34)	
	50	0.67 (0.66, 0.69)	2.50 (2.47, 2.54)	4.99 (4.94, 5.04)	6.56 (6.50, 6.62)	5.29 (5.27, 5.32)	
	60	1.92 (1.89, 1.94)	4.52 (4.47, 4.57)	5.92 (5.86, 5.98)	6.17 (6.11, 6.23)	5.02 (5.00, 5.05)	
	70	2.93 (2.89, 2.97)	4.51 (4.46, 4.56)	- (- , -)	4.79 (4.73, 4.84)	4.03 (4.01, 4.05)	
	80	2.13 (2.10, 2.17)	- (- , -)	- (- , -)	2.51 (2.47, 2.55)	2.33 (2.31, 2.35)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	5.35 (5.20, 5.50)	4.12 (4.06, 4.18)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.04 (0.03, 0.05)	5.47 (5.32, 5.62)	4.21 (4.15, 4.27)	
	20	0.00 (0.00, 0.01)	0.04 (0.03, 0.04)	0.29 (0.27, 0.31)	5.48 (5.33, 5.64)	4.23 (4.16, 4.29)	
	30	0.03 (0.03, 0.04)	0.29 (0.27, 0.31)	1.06 (1.01, 1.11)	5.52 (5.37, 5.68)	4.26 (4.20, 4.32)	
	40	0.26 (0.24, 0.28)	1.05 (1.00, 1.09)	2.65 (2.56, 2.74)	5.58 (5.43, 5.74)	4.31 (4.25, 4.38)	
	50	0.82 (0.78, 0.86)	2.49 (2.40, 2.59)	4.41 (4.28, 4.55)	5.55 (5.39, 5.71)	4.31 (4.25, 4.37)	
	60	1.83 (1.75, 1.91)	3.93 (3.79, 4.06)	4.96 (4.81, 5.12)	5.17 (5.01, 5.34)	4.07 (4.01, 4.14)	
	70	2.51 (2.40, 2.63)	3.76 (3.61, 3.91)	- (- , -)	4.00 (3.84, 4.17)	3.30 (3.23, 3.37)	
	80	1.85 (1.72, 1.98)	- (- , -)	- (- , -)	2.22 (2.07, 2.37)	2.03 (1.96, 2.09)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XV-15 - continued

LUNG AND BRONCHUS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	4.26 (4.05, 4.48)	3.10 (2.94, 3.27)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.03 (0.02, 0.04)	4.29 (4.09, 4.52)	3.12 (2.96, 3.30)	
	20	0.00 (0.00, 0.01)	0.03 (0.02, 0.04)	0.13 (0.11, 0.15)	4.30 (4.09, 4.53)	3.13 (2.97, 3.30)	
	30	0.03 (0.02, 0.03)	0.13 (0.11, 0.14)	0.46 (0.43, 0.50)	4.31 (4.10, 4.54)	3.13 (2.97, 3.31)	
	40	0.10 (0.09, 0.11)	0.44 (0.40, 0.47)	1.24 (1.17, 1.31)	4.30 (4.09, 4.53)	3.13 (2.97, 3.31)	
	50	0.34 (0.31, 0.37)	1.15 (1.09, 1.22)	2.46 (2.35, 2.57)	4.25 (4.04, 4.48)	3.10 (2.94, 3.28)	
	60	0.83 (0.78, 0.89)	2.17 (2.07, 2.28)	3.45 (3.30, 3.62)	4.01 (3.80, 4.25)	2.99 (2.82, 3.17)	
	70	1.43 (1.35, 1.52)	2.80 (2.65, 2.96)	- (-, -)	3.40 (3.18, 3.64)	2.65 (2.48, 2.84)	
	80	1.64 (1.51, 1.78)	- (-, -)	- (-, -)	2.35 (2.13, 2.61)	1.94 (1.76, 2.15)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	4.37 (3.75, 5.23)	3.75 (3.42, 4.14)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	4.44 (3.81, 5.31)	3.81 (3.48, 4.20)	
	20	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.09 (0.05, 0.15)	4.46 (3.83, 5.33)	3.83 (3.50, 4.23)	
	30	0.01 (0.00, 0.04)	0.09 (0.05, 0.15)	0.52 (0.40, 0.67)	4.50 (3.86, 5.38)	3.87 (3.54, 4.27)	
	40	0.08 (0.04, 0.14)	0.52 (0.40, 0.67)	1.86 (1.55, 2.21)	4.57 (3.92, 5.47)	3.94 (3.60, 4.34)	
	50	0.45 (0.34, 0.60)	1.83 (1.52, 2.19)	3.61 (3.09, 4.20)	4.62 (3.96, 5.55)	4.01 (3.65, 4.43)	
	60	1.46 (1.17, 1.79)	3.35 (2.83, 3.95)	4.33 (3.67, 5.09)	4.42 (3.74, 5.38)	3.94 (3.57, 4.38)	
	70	2.18 (1.73, 2.72)	3.31 (2.66, 4.08)	- (-, -)	3.41 (2.74, 4.42)	3.10 (2.73, 3.56)	
	80	1.51 (1.01, 2.19)	- (-, -)	- (-, -)	1.65 (1.09, 2.74)	1.76 (1.40, 2.26)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	3.61 (3.45, 3.79)	2.25 (2.17, 2.33)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.01, 0.02)	3.65 (3.49, 3.83)	2.27 (2.19, 2.35)	
	20	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.09 (0.08, 0.10)	3.66 (3.49, 3.83)	2.27 (2.19, 2.36)	
	30	0.01 (0.01, 0.02)	0.09 (0.08, 0.10)	0.37 (0.34, 0.39)	3.67 (3.50, 3.84)	2.28 (2.20, 2.36)	
	40	0.08 (0.07, 0.08)	0.36 (0.33, 0.38)	1.11 (1.05, 1.16)	3.67 (3.51, 3.85)	2.29 (2.21, 2.37)	
	50	0.28 (0.26, 0.31)	1.05 (0.99, 1.11)	2.30 (2.20, 2.40)	3.65 (3.49, 3.83)	2.28 (2.20, 2.37)	
	60	0.79 (0.74, 0.84)	2.09 (1.99, 2.19)	3.10 (2.96, 3.24)	3.49 (3.32, 3.68)	2.23 (2.14, 2.31)	
	70	1.42 (1.34, 1.51)	2.53 (2.40, 2.67)	- (-, -)	2.96 (2.79, 3.15)	2.00 (1.92, 2.09)	
	80	1.40 (1.29, 1.53)	- (-, -)	- (-, -)	1.95 (1.77, 2.16)	1.47 (1.38, 1.57)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVI-9

MELANOMA OF THE SKIN (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.07 (0.06, 0.07)	1.72 (1.70, 1.74)	0.28 (0.27, 0.28)	
	10	0.01 (0.01, 0.01)	0.07 (0.06, 0.07)	0.17 (0.17, 0.17)	1.74 (1.72, 1.76)	0.28 (0.28, 0.28)	
	20	0.05 (0.05, 0.06)	0.16 (0.15, 0.16)	0.34 (0.34, 0.35)	1.74 (1.72, 1.76)	0.28 (0.28, 0.28)	
	30	0.11 (0.10, 0.11)	0.29 (0.29, 0.30)	0.57 (0.56, 0.58)	1.70 (1.68, 1.72)	0.28 (0.28, 0.28)	
	40	0.19 (0.18, 0.19)	0.47 (0.47, 0.48)	0.86 (0.85, 0.87)	1.62 (1.60, 1.64)	0.28 (0.27, 0.28)	
	50	0.29 (0.29, 0.30)	0.69 (0.68, 0.70)	1.13 (1.12, 1.15)	1.47 (1.45, 1.49)	0.26 (0.26, 0.27)	
	60	0.42 (0.41, 0.43)	0.89 (0.88, 0.91)	1.18 (1.17, 1.20)	1.26 (1.24, 1.27)	0.24 (0.24, 0.24)	
	70	0.54 (0.53, 0.56)	0.88 (0.87, 0.90)	- (- , -)	0.97 (0.95, 0.98)	0.20 (0.20, 0.21)	
	80	0.48 (0.47, 0.50)	- (- , -)	- (- , -)	0.60 (0.59, 0.62)	0.15 (0.14, 0.15)	
White	0	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.08 (0.08, 0.08)	1.99 (1.97, 2.01)	0.31 (0.31, 0.32)	
	10	0.01 (0.01, 0.02)	0.08 (0.08, 0.08)	0.20 (0.20, 0.21)	2.01 (1.99, 2.04)	0.32 (0.31, 0.32)	
	20	0.07 (0.06, 0.07)	0.19 (0.18, 0.19)	0.41 (0.40, 0.41)	2.01 (1.99, 2.03)	0.32 (0.31, 0.32)	
	30	0.13 (0.12, 0.13)	0.35 (0.34, 0.35)	0.67 (0.66, 0.68)	1.96 (1.94, 1.98)	0.32 (0.31, 0.32)	
	40	0.22 (0.22, 0.23)	0.56 (0.55, 0.56)	1.01 (0.99, 1.02)	1.86 (1.84, 1.88)	0.31 (0.31, 0.32)	
	50	0.34 (0.34, 0.35)	0.80 (0.79, 0.82)	1.31 (1.29, 1.33)	1.68 (1.66, 1.70)	0.30 (0.29, 0.30)	
	60	0.49 (0.48, 0.50)	1.03 (1.01, 1.05)	1.35 (1.33, 1.37)	1.42 (1.40, 1.45)	0.27 (0.26, 0.27)	
	70	0.62 (0.61, 0.64)	0.99 (0.97, 1.01)	- (- , -)	1.08 (1.06, 1.10)	0.22 (0.22, 0.23)	
	80	0.53 (0.52, 0.54)	- (- , -)	- (- , -)	0.65 (0.64, 0.67)	0.16 (0.16, 0.16)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.09 (0.08, 0.11)	0.04 (0.03, 0.04)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.09 (0.08, 0.11)	0.04 (0.04, 0.04)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.09 (0.08, 0.11)	0.04 (0.04, 0.04)	
	30	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	0.02 (0.02, 0.03)	0.09 (0.08, 0.11)	0.04 (0.04, 0.05)	
	40	0.01 (0.00, 0.01)	0.02 (0.01, 0.02)	0.03 (0.03, 0.04)	0.09 (0.07, 0.10)	0.04 (0.04, 0.05)	
	50	0.01 (0.01, 0.01)	0.03 (0.02, 0.04)	0.06 (0.05, 0.07)	0.09 (0.07, 0.10)	0.04 (0.04, 0.05)	
	60	0.02 (0.02, 0.03)	0.05 (0.04, 0.07)	0.07 (0.06, 0.09)	0.09 (0.07, 0.10)	0.04 (0.03, 0.05)	
	70	0.04 (0.03, 0.05)	0.06 (0.05, 0.08)	- (- , -)	0.08 (0.06, 0.10)	0.04 (0.03, 0.04)	
	80	0.04 (0.02, 0.05)	- (- , -)	- (- , -)	0.06 (0.04, 0.09)	0.03 (0.03, 0.04)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVI-9 - continued
MELANOMA OF THE SKIN (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.18 (0.15, 0.22)	0.06 (0.04, 0.09)	
Islander	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.01, 0.02)	0.18 (0.15, 0.22)	0.06 (0.04, 0.09)	
	20	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.03 (0.02, 0.04)	0.18 (0.15, 0.22)	0.06 (0.04, 0.09)	
	30	0.01 (0.01, 0.01)	0.03 (0.02, 0.03)	0.04 (0.04, 0.05)	0.18 (0.15, 0.22)	0.06 (0.04, 0.09)	
	40	0.01 (0.01, 0.02)	0.03 (0.03, 0.04)	0.06 (0.05, 0.07)	0.17 (0.14, 0.21)	0.06 (0.04, 0.09)	
	50	0.02 (0.02, 0.03)	0.05 (0.04, 0.06)	0.09 (0.07, 0.10)	0.15 (0.13, 0.19)	0.06 (0.04, 0.09)	
	60	0.03 (0.02, 0.04)	0.07 (0.06, 0.09)	0.11 (0.09, 0.14)	0.14 (0.11, 0.18)	0.06 (0.04, 0.09)	
	70	0.05 (0.04, 0.06)	0.09 (0.07, 0.12)	- (-, -)	0.12 (0.09, 0.17)	0.05 (0.03, 0.09)	
	80	0.06 (0.04, 0.08)	- (-, -)	- (-, -)	0.09 (0.06, 0.14)	0.05 (0.03, 0.09)	
American	0	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.31 (0.15, 0.65)	0.13 (0.08, 0.24)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.01)	0.00 (0.00, 0.02)	0.01 (0.00, 0.04)	0.31 (0.15, 0.66)	0.14 (0.08, 0.25)	
	20	0.00 (0.00, 0.02)	0.01 (0.00, 0.04)	0.04 (0.02, 0.07)	0.31 (0.15, 0.67)	0.14 (0.08, 0.25)	
	30	0.01 (0.00, 0.03)	0.03 (0.02, 0.06)	0.07 (0.04, 0.11)	0.31 (0.15, 0.68)	0.14 (0.08, 0.25)	
	40	0.02 (0.01, 0.05)	0.06 (0.03, 0.10)	0.09 (0.05, 0.15)	0.31 (0.15, 0.68)	0.14 (0.08, 0.26)	
	50	0.04 (0.02, 0.08)	0.07 (0.03, 0.13)	0.13 (0.07, 0.23)	0.30 (0.13, 0.69)	0.15 (0.08, 0.27)	
	60	0.03 (0.01, 0.09)	0.10 (0.05, 0.20)	0.19 (0.09, 0.35)	0.28 (0.11, 0.70)	0.14 (0.07, 0.28)	
	70	0.08 (0.03, 0.19)	0.18 (0.08, 0.37)	- (-, -)	0.29 (0.10, 0.78)	0.14 (0.06, 0.30)	
	80	0.14 (0.04, 0.36)	- (-, -)	- (-, -)	0.29 (0.06, 0.98)	0.12 (0.03, 0.36)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.50 (0.46, 0.56)	0.10 (0.08, 0.11)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.04 (0.04, 0.04)	0.51 (0.47, 0.56)	0.10 (0.08, 0.11)	
	20	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	0.08 (0.07, 0.09)	0.51 (0.47, 0.56)	0.10 (0.08, 0.11)	
	30	0.03 (0.02, 0.03)	0.07 (0.06, 0.08)	0.13 (0.12, 0.14)	0.50 (0.46, 0.55)	0.10 (0.08, 0.11)	
	40	0.04 (0.04, 0.05)	0.10 (0.09, 0.11)	0.19 (0.18, 0.21)	0.48 (0.44, 0.53)	0.10 (0.08, 0.11)	
	50	0.06 (0.05, 0.07)	0.15 (0.14, 0.17)	0.29 (0.26, 0.31)	0.45 (0.40, 0.50)	0.09 (0.08, 0.11)	
	60	0.10 (0.08, 0.11)	0.24 (0.22, 0.26)	0.35 (0.32, 0.39)	0.41 (0.36, 0.46)	0.09 (0.07, 0.11)	
	70	0.16 (0.14, 0.18)	0.29 (0.25, 0.32)	- (-, -)	0.35 (0.30, 0.41)	0.08 (0.07, 0.10)	
	80	0.17 (0.14, 0.20)	- (-, -)	- (-, -)	0.25 (0.20, 0.32)	0.07 (0.05, 0.09)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVI-9 - continued
MELANOMA OF THE SKIN (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.04, 0.05)	2.10 (2.07, 2.13)	0.37 (0.36, 0.37)	
	10	0.01 (0.01, 0.01)	0.04 (0.04, 0.05)	0.13 (0.12, 0.13)	2.12 (2.09, 2.15)	0.37 (0.37, 0.38)	
	20	0.04 (0.03, 0.04)	0.12 (0.11, 0.12)	0.30 (0.29, 0.30)	2.13 (2.10, 2.16)	0.37 (0.37, 0.38)	
	30	0.08 (0.08, 0.09)	0.27 (0.26, 0.27)	0.60 (0.59, 0.61)	2.12 (2.09, 2.15)	0.38 (0.37, 0.38)	
	40	0.18 (0.18, 0.19)	0.53 (0.51, 0.54)	1.03 (1.02, 1.05)	2.07 (2.04, 2.10)	0.37 (0.37, 0.38)	
	50	0.35 (0.35, 0.36)	0.88 (0.86, 0.90)	1.50 (1.47, 1.52)	1.96 (1.93, 1.99)	0.36 (0.35, 0.37)	
	60	0.57 (0.56, 0.59)	1.24 (1.21, 1.26)	1.64 (1.61, 1.67)	1.74 (1.71, 1.77)	0.33 (0.33, 0.34)	
	70	0.80 (0.78, 0.82)	1.28 (1.25, 1.31)	- (- , -)	1.40 (1.37, 1.43)	0.29 (0.28, 0.30)	
	80	0.75 (0.73, 0.78)	- (- , -)	- (- , -)	0.93 (0.90, 0.96)	0.22 (0.22, 0.23)	
White	0	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.05 (0.05, 0.06)	2.42 (2.38, 2.45)	0.42 (0.41, 0.42)	
	10	0.01 (0.01, 0.01)	0.05 (0.05, 0.06)	0.15 (0.15, 0.16)	2.44 (2.41, 2.48)	0.42 (0.42, 0.43)	
	20	0.04 (0.04, 0.05)	0.14 (0.14, 0.15)	0.35 (0.34, 0.36)	2.44 (2.41, 2.48)	0.43 (0.42, 0.43)	
	30	0.10 (0.10, 0.11)	0.31 (0.30, 0.32)	0.70 (0.69, 0.72)	2.43 (2.40, 2.47)	0.43 (0.42, 0.43)	
	40	0.22 (0.21, 0.22)	0.61 (0.60, 0.62)	1.20 (1.18, 1.22)	2.37 (2.33, 2.41)	0.42 (0.42, 0.43)	
	50	0.41 (0.40, 0.42)	1.02 (1.00, 1.04)	1.73 (1.70, 1.76)	2.23 (2.20, 2.27)	0.41 (0.40, 0.41)	
	60	0.66 (0.64, 0.68)	1.42 (1.39, 1.45)	1.87 (1.83, 1.90)	1.97 (1.93, 2.01)	0.37 (0.37, 0.38)	
	70	0.91 (0.89, 0.93)	1.44 (1.41, 1.47)	- (- , -)	1.56 (1.53, 1.60)	0.32 (0.31, 0.33)	
	80	0.83 (0.80, 0.86)	- (- , -)	- (- , -)	1.02 (0.98, 1.06)	0.24 (0.23, 0.25)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.07 (0.06, 0.10)	0.04 (0.03, 0.04)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.07 (0.06, 0.10)	0.04 (0.03, 0.05)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.08 (0.06, 0.10)	0.04 (0.03, 0.05)	
	30	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.02 (0.01, 0.03)	0.08 (0.06, 0.10)	0.04 (0.03, 0.05)	
	40	0.01 (0.00, 0.01)	0.02 (0.01, 0.02)	0.03 (0.02, 0.05)	0.07 (0.06, 0.10)	0.04 (0.03, 0.05)	
	50	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.05 (0.04, 0.07)	0.07 (0.05, 0.10)	0.04 (0.03, 0.05)	
	60	0.02 (0.01, 0.04)	0.05 (0.04, 0.07)	0.07 (0.05, 0.09)	0.07 (0.05, 0.10)	0.04 (0.03, 0.05)	
	70	0.04 (0.02, 0.06)	0.06 (0.04, 0.09)	- (- , -)	0.06 (0.04, 0.10)	0.04 (0.03, 0.05)	
	80	0.04 (0.02, 0.07)	- (- , -)	- (- , -)	0.05 (0.02, 0.11)	0.04 (0.02, 0.05)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

- Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVI-9 - continued
MELANOMA OF THE SKIN (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.19 (0.15, 0.26)	0.06 (0.04, 0.11)	
Islander	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.19 (0.15, 0.26)	0.06 (0.04, 0.11)	
	20	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	0.02 (0.02, 0.03)	0.19 (0.15, 0.26)	0.06 (0.04, 0.11)	
	30	0.01 (0.00, 0.01)	0.02 (0.01, 0.03)	0.04 (0.03, 0.05)	0.19 (0.15, 0.25)	0.06 (0.04, 0.11)	
	40	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.06 (0.05, 0.08)	0.19 (0.14, 0.25)	0.06 (0.04, 0.11)	
	50	0.02 (0.01, 0.03)	0.05 (0.04, 0.07)	0.11 (0.08, 0.13)	0.18 (0.13, 0.24)	0.06 (0.04, 0.11)	
	60	0.03 (0.02, 0.05)	0.09 (0.07, 0.12)	0.13 (0.10, 0.17)	0.16 (0.12, 0.23)	0.06 (0.04, 0.11)	
	70	0.06 (0.04, 0.09)	0.11 (0.08, 0.15)	- (-, -)	0.14 (0.10, 0.22)	0.05 (0.03, 0.11)	
	80	0.06 (0.04, 0.10)	- (-, -)	- (-, -)	0.11 (0.06, 0.20)	0.04 (0.02, 0.11)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.34 (0.18, 1.01)	0.10 (0.05, 0.28)	
Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.01 (0.00, 0.05)	0.34 (0.18, 1.03)	0.10 (0.06, 0.29)	
Alaska	20	0.00 (0.00, 0.02)	0.02 (0.00, 0.05)	0.05 (0.02, 0.10)	0.35 (0.18, 1.04)	0.10 (0.06, 0.29)	
	30	0.01 (0.00, 0.05)	0.05 (0.02, 0.10)	0.11 (0.05, 0.19)	0.36 (0.18, 1.06)	0.10 (0.06, 0.30)	
	40	0.03 (0.01, 0.08)	0.10 (0.05, 0.18)	0.12 (0.06, 0.23)	0.35 (0.18, 1.09)	0.11 (0.06, 0.31)	
	50	0.06 (0.02, 0.14)	0.09 (0.03, 0.20)	0.19 (0.09, 0.38)	0.34 (0.16, 1.11)	0.11 (0.06, 0.32)	
	60	0.03 (0.00, 0.13)	0.14 (0.05, 0.34)	0.29 (0.11, 0.64)	0.30 (0.12, 1.15)	0.10 (0.05, 0.34)	
	70	0.14 (0.04, 0.36)	0.32 (0.12, 0.73)	- (-, -)	0.33 (0.12, 1.35)	0.07 (0.02, 0.37)	
	80	0.27 (0.06, 0.82)	- (-, -)	- (-, -)	0.28 (0.06, 1.83)	0.05 (0.01, 0.52)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.49 (0.43, 0.58)	0.10 (0.08, 0.13)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.01, 0.02)	0.49 (0.43, 0.58)	0.10 (0.08, 0.13)	
	20	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.04 (0.04, 0.05)	0.50 (0.43, 0.58)	0.10 (0.09, 0.13)	
	30	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.09 (0.08, 0.10)	0.50 (0.43, 0.59)	0.10 (0.09, 0.13)	
	40	0.03 (0.02, 0.03)	0.08 (0.07, 0.09)	0.16 (0.14, 0.19)	0.49 (0.42, 0.58)	0.10 (0.09, 0.13)	
	50	0.05 (0.04, 0.06)	0.14 (0.12, 0.16)	0.30 (0.26, 0.34)	0.48 (0.41, 0.57)	0.10 (0.08, 0.13)	
	60	0.09 (0.08, 0.12)	0.26 (0.23, 0.30)	0.39 (0.34, 0.45)	0.46 (0.38, 0.55)	0.09 (0.07, 0.12)	
	70	0.19 (0.16, 0.23)	0.34 (0.29, 0.41)	- (-, -)	0.42 (0.34, 0.53)	0.08 (0.06, 0.12)	
	80	0.21 (0.16, 0.28)	- (-, -)	- (-, -)	0.31 (0.22, 0.46)	0.07 (0.05, 0.12)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVI-9 - continued
MELANOMA OF THE SKIN (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.09 (0.08, 0.09)	1.40 (1.38, 1.42)	0.20 (0.19, 0.20)	
	10	0.02 (0.01, 0.02)	0.09 (0.08, 0.09)	0.21 (0.21, 0.22)	1.42 (1.39, 1.44)	0.20 (0.19, 0.20)	
	20	0.07 (0.07, 0.08)	0.20 (0.19, 0.21)	0.39 (0.38, 0.40)	1.40 (1.38, 1.43)	0.20 (0.19, 0.20)	
	30	0.13 (0.12, 0.13)	0.32 (0.31, 0.33)	0.55 (0.53, 0.56)	1.34 (1.32, 1.36)	0.20 (0.19, 0.20)	
	40	0.19 (0.19, 0.20)	0.42 (0.41, 0.43)	0.69 (0.68, 0.71)	1.22 (1.20, 1.24)	0.19 (0.19, 0.20)	
	50	0.23 (0.23, 0.24)	0.51 (0.50, 0.52)	0.80 (0.78, 0.82)	1.05 (1.03, 1.07)	0.18 (0.18, 0.19)	
	60	0.29 (0.28, 0.30)	0.59 (0.58, 0.61)	0.80 (0.78, 0.82)	0.86 (0.84, 0.88)	0.16 (0.16, 0.17)	
	70	0.34 (0.33, 0.36)	0.57 (0.56, 0.59)	- (- , -)	0.64 (0.62, 0.66)	0.14 (0.14, 0.14)	
	80	0.31 (0.30, 0.32)	- (- , -)	- (- , -)	0.40 (0.38, 0.42)	0.10 (0.10, 0.11)	
White	0	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.11 (0.10, 0.11)	1.63 (1.60, 1.66)	0.22 (0.22, 0.23)	
	10	0.02 (0.02, 0.02)	0.11 (0.10, 0.11)	0.26 (0.25, 0.27)	1.64 (1.62, 1.67)	0.22 (0.22, 0.23)	
	20	0.09 (0.08, 0.09)	0.24 (0.23, 0.25)	0.47 (0.46, 0.48)	1.63 (1.60, 1.66)	0.22 (0.22, 0.23)	
	30	0.15 (0.15, 0.16)	0.38 (0.37, 0.39)	0.65 (0.63, 0.66)	1.55 (1.52, 1.58)	0.22 (0.22, 0.23)	
	40	0.23 (0.22, 0.24)	0.50 (0.49, 0.51)	0.82 (0.80, 0.83)	1.41 (1.39, 1.44)	0.22 (0.21, 0.22)	
	50	0.28 (0.27, 0.28)	0.60 (0.58, 0.61)	0.93 (0.91, 0.95)	1.21 (1.18, 1.23)	0.20 (0.20, 0.21)	
	60	0.34 (0.32, 0.35)	0.69 (0.67, 0.71)	0.91 (0.89, 0.93)	0.97 (0.95, 1.00)	0.18 (0.18, 0.19)	
	70	0.39 (0.38, 0.41)	0.64 (0.63, 0.66)	- (- , -)	0.71 (0.69, 0.74)	0.15 (0.15, 0.16)	
	80	0.34 (0.32, 0.35)	- (- , -)	- (- , -)	0.43 (0.42, 0.45)	0.11 (0.11, 0.12)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.10 (0.08, 0.13)	0.04 (0.03, 0.05)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	0.10 (0.08, 0.13)	0.04 (0.04, 0.05)	
	20	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.10 (0.08, 0.13)	0.04 (0.04, 0.05)	
	30	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.02 (0.01, 0.03)	0.10 (0.08, 0.13)	0.04 (0.04, 0.05)	
	40	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.10 (0.08, 0.12)	0.04 (0.04, 0.05)	
	50	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.06 (0.05, 0.08)	0.10 (0.08, 0.12)	0.04 (0.04, 0.05)	
	60	0.02 (0.01, 0.03)	0.06 (0.04, 0.08)	0.08 (0.06, 0.10)	0.09 (0.07, 0.12)	0.04 (0.03, 0.05)	
	70	0.04 (0.03, 0.06)	0.07 (0.05, 0.09)	- (- , -)	0.09 (0.07, 0.12)	0.04 (0.03, 0.05)	
	80	0.04 (0.02, 0.06)	- (- , -)	- (- , -)	0.07 (0.04, 0.11)	0.03 (0.02, 0.04)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

- Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVI-9 - continued
MELANOMA OF THE SKIN (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.17 (0.13, 0.23)	0.06 (0.03, 0.12)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.02, 0.03)	0.17 (0.13, 0.24)	0.06 (0.03, 0.12)	
Islander	20	0.01 (0.00, 0.01)	0.02 (0.02, 0.03)	0.04 (0.03, 0.05)	0.17 (0.13, 0.24)	0.06 (0.03, 0.12)	
	30	0.02 (0.01, 0.02)	0.03 (0.02, 0.04)	0.05 (0.04, 0.06)	0.16 (0.13, 0.23)	0.06 (0.03, 0.12)	
	40	0.02 (0.01, 0.02)	0.03 (0.03, 0.04)	0.06 (0.04, 0.07)	0.15 (0.11, 0.22)	0.06 (0.03, 0.12)	
	50	0.02 (0.01, 0.03)	0.04 (0.03, 0.06)	0.07 (0.06, 0.10)	0.13 (0.10, 0.20)	0.06 (0.03, 0.12)	
	60	0.02 (0.01, 0.03)	0.06 (0.04, 0.08)	0.10 (0.07, 0.13)	0.12 (0.08, 0.19)	0.06 (0.03, 0.12)	
	70	0.04 (0.02, 0.05)	0.08 (0.06, 0.11)	- (-, -)	0.10 (0.06, 0.18)	0.05 (0.02, 0.12)	
	80	0.05 (0.03, 0.08)	- (-, -)	- (-, -)	0.08 (0.04, 0.16)	0.06 (0.02, 0.13)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.01 (0.00, 0.05)	0.27 (0.06, 0.88)	0.16 (0.07, 0.35)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.01 (0.00, 0.04)	0.01 (0.00, 0.05)	0.27 (0.06, 0.89)	0.16 (0.07, 0.36)	
	20	0.01 (0.00, 0.04)	0.01 (0.00, 0.05)	0.03 (0.01, 0.07)	0.27 (0.06, 0.90)	0.16 (0.07, 0.36)	
	30	0.01 (0.00, 0.04)	0.02 (0.00, 0.06)	0.03 (0.01, 0.09)	0.27 (0.06, 0.90)	0.16 (0.07, 0.36)	
	40	0.01 (0.00, 0.05)	0.03 (0.01, 0.08)	0.06 (0.02, 0.16)	0.26 (0.05, 0.91)	0.17 (0.07, 0.37)	
	50	0.01 (0.00, 0.06)	0.05 (0.01, 0.15)	0.08 (0.02, 0.22)	0.26 (0.05, 0.93)	0.17 (0.07, 0.38)	
	60	0.04 (0.01, 0.14)	0.07 (0.02, 0.21)	0.10 (0.03, 0.31)	0.26 (0.04, 0.97)	0.17 (0.07, 0.40)	
	70	0.04 (0.00, 0.17)	0.07 (0.01, 0.30)	- (-, -)	0.25 (0.03, 1.07)	0.18 (0.06, 0.44)	
	80	0.05 (0.00, 0.33)	- (-, -)	- (-, -)	0.29 (0.02, 1.38)	0.17 (0.03, 0.53)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.53 (0.48, 0.61)	0.09 (0.07, 0.11)	
	10	0.00 (0.00, 0.00)	0.02 (0.02, 0.03)	0.06 (0.06, 0.07)	0.54 (0.48, 0.61)	0.09 (0.07, 0.12)	
	20	0.02 (0.02, 0.03)	0.06 (0.05, 0.07)	0.12 (0.11, 0.13)	0.54 (0.48, 0.61)	0.09 (0.07, 0.12)	
	30	0.04 (0.03, 0.05)	0.10 (0.09, 0.11)	0.17 (0.15, 0.18)	0.52 (0.46, 0.59)	0.09 (0.07, 0.12)	
	40	0.06 (0.05, 0.07)	0.13 (0.11, 0.14)	0.22 (0.20, 0.24)	0.48 (0.42, 0.55)	0.09 (0.07, 0.11)	
	50	0.07 (0.06, 0.08)	0.16 (0.14, 0.18)	0.28 (0.25, 0.31)	0.43 (0.37, 0.50)	0.09 (0.07, 0.11)	
	60	0.10 (0.08, 0.12)	0.22 (0.19, 0.25)	0.32 (0.28, 0.37)	0.37 (0.31, 0.45)	0.08 (0.06, 0.11)	
	70	0.13 (0.11, 0.16)	0.24 (0.20, 0.29)	- (-, -)	0.30 (0.24, 0.38)	0.08 (0.06, 0.10)	
	80	0.14 (0.10, 0.18)	- (-, -)	- (-, -)	0.21 (0.15, 0.30)	0.07 (0.04, 0.10)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVII-8
MESOTHELIOMA (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.12 (0.12, 0.13)	0.10 (0.09, 0.10)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.12 (0.12, 0.13)	0.10 (0.10, 0.10)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.13 (0.12, 0.13)	0.10 (0.10, 0.10)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.13 (0.12, 0.13)	0.10 (0.10, 0.10)	
	40	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.04, 0.04)	0.13 (0.12, 0.13)	0.10 (0.10, 0.10)	
	50	0.01 (0.01, 0.01)	0.04 (0.04, 0.04)	0.09 (0.09, 0.09)	0.13 (0.12, 0.13)	0.10 (0.10, 0.10)	
	60	0.03 (0.03, 0.03)	0.08 (0.08, 0.09)	0.12 (0.11, 0.12)	0.13 (0.12, 0.13)	0.10 (0.10, 0.10)	
	70	0.06 (0.06, 0.07)	0.10 (0.10, 0.11)	- (- , -)	0.11 (0.10, 0.12)	0.09 (0.09, 0.09)	
	80	0.06 (0.05, 0.06)	- (- , -)	- (- , -)	0.07 (0.06, 0.07)	0.06 (0.06, 0.06)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.14 (0.13, 0.14)	0.11 (0.10, 0.11)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.14 (0.13, 0.15)	0.11 (0.10, 0.11)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.14 (0.13, 0.15)	0.11 (0.10, 0.11)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.14 (0.13, 0.15)	0.11 (0.11, 0.11)	
	40	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.05 (0.04, 0.05)	0.14 (0.14, 0.15)	0.11 (0.11, 0.11)	
	50	0.01 (0.01, 0.01)	0.04 (0.04, 0.05)	0.10 (0.10, 0.11)	0.14 (0.14, 0.15)	0.11 (0.11, 0.11)	
	60	0.03 (0.03, 0.04)	0.09 (0.09, 0.10)	0.13 (0.13, 0.14)	0.14 (0.13, 0.15)	0.11 (0.11, 0.11)	
	70	0.07 (0.06, 0.07)	0.11 (0.11, 0.12)	- (- , -)	0.12 (0.11, 0.13)	0.10 (0.09, 0.10)	
	80	0.06 (0.06, 0.07)	- (- , -)	- (- , -)	0.07 (0.07, 0.08)	0.06 (0.06, 0.07)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.04, 0.07)	0.03 (0.03, 0.04)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.04, 0.07)	0.03 (0.03, 0.04)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.04, 0.07)	0.03 (0.03, 0.04)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.04, 0.07)	0.03 (0.03, 0.04)	
	40	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.02 (0.01, 0.02)	0.06 (0.04, 0.07)	0.03 (0.03, 0.04)	
	50	0.01 (0.00, 0.01)	0.02 (0.01, 0.02)	0.04 (0.03, 0.05)	0.06 (0.05, 0.07)	0.04 (0.03, 0.04)	
	60	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.05 (0.04, 0.07)	0.06 (0.04, 0.07)	0.04 (0.03, 0.04)	
	70	0.03 (0.02, 0.04)	0.05 (0.04, 0.07)	- (- , -)	0.06 (0.04, 0.07)	0.03 (0.03, 0.04)	
	80	0.04 (0.02, 0.05)	- (- , -)	- (- , -)	0.04 (0.03, 0.07)	0.02 (0.02, 0.03)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

- Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVII-8 - continued
MESOTHELIOMA (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/Pacific Islander	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.03, 0.08)	0.03 (0.02, 0.06)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.03, 0.08)	0.03 (0.02, 0.06)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.05 (0.03, 0.08)	0.03 (0.02, 0.06)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.05 (0.03, 0.08)	0.03 (0.02, 0.06)	
	40	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.01, 0.03)	0.05 (0.03, 0.08)	0.03 (0.02, 0.06)	
	50	0.00 (0.00, 0.01)	0.02 (0.01, 0.03)	0.03 (0.03, 0.05)	0.05 (0.03, 0.08)	0.03 (0.02, 0.06)	
	60	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.04 (0.03, 0.05)	0.04 (0.03, 0.08)	0.03 (0.02, 0.06)	
	70	0.02 (0.01, 0.03)	0.02 (0.02, 0.04)	- (-, -)	0.03 (0.02, 0.07)	0.03 (0.01, 0.06)	
	80	0.01 (0.00, 0.02)	- (-, -)	- (-, -)	0.02 (0.00, 0.06)	0.02 (0.00, 0.05)	
American Indian/Alaska Native ^a	0	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.08 (0.03, 0.36)	0.04 (0.02, 0.12)	
	10	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.08 (0.03, 0.37)	0.05 (0.02, 0.13)	
	20	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.00 (0.00, 0.02)	0.09 (0.03, 0.37)	0.05 (0.02, 0.13)	
	30	0.00 (0.00, 0.01)	0.00 (0.00, 0.02)	0.01 (0.00, 0.03)	0.09 (0.03, 0.38)	0.05 (0.02, 0.13)	
	40	0.00 (0.00, 0.02)	0.01 (0.00, 0.04)	0.02 (0.01, 0.07)	0.09 (0.03, 0.39)	0.05 (0.02, 0.13)	
	50	0.01 (0.00, 0.03)	0.02 (0.00, 0.07)	0.05 (0.01, 0.12)	0.09 (0.03, 0.40)	0.05 (0.02, 0.14)	
	60	0.01 (0.00, 0.06)	0.04 (0.01, 0.12)	0.09 (0.03, 0.22)	0.09 (0.03, 0.43)	0.05 (0.02, 0.15)	
	70	0.03 (0.01, 0.12)	0.08 (0.02, 0.24)	- (-, -)	0.09 (0.02, 0.48)	0.05 (0.02, 0.17)	
	80	0.07 (0.01, 0.27)	- (-, -)	- (-, -)	0.08 (0.01, 0.63)	0.03 (0.00, 0.21)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.13 (0.11, 0.15)	0.06 (0.06, 0.08)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.13 (0.11, 0.16)	0.07 (0.06, 0.08)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.13 (0.11, 0.16)	0.07 (0.06, 0.08)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.13 (0.11, 0.16)	0.07 (0.06, 0.08)	
	40	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.13 (0.11, 0.16)	0.07 (0.06, 0.08)	
	50	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.09 (0.08, 0.11)	0.13 (0.11, 0.16)	0.07 (0.06, 0.08)	
	60	0.03 (0.02, 0.04)	0.08 (0.07, 0.10)	0.12 (0.10, 0.14)	0.12 (0.10, 0.15)	0.07 (0.06, 0.08)	
	70	0.06 (0.05, 0.08)	0.10 (0.08, 0.12)	- (-, -)	0.11 (0.09, 0.14)	0.06 (0.05, 0.07)	
	80	0.05 (0.03, 0.07)	- (-, -)	- (-, -)	0.06 (0.04, 0.10)	0.04 (0.03, 0.05)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVII-8 - continued
MESOTHELIOMA (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.21 (0.20, 0.22)	0.17 (0.16, 0.17)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.21 (0.20, 0.22)	0.17 (0.16, 0.17)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.21 (0.20, 0.22)	0.17 (0.17, 0.17)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.21 (0.20, 0.23)	0.17 (0.17, 0.18)	
	40	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.07 (0.06, 0.07)	0.22 (0.21, 0.23)	0.17 (0.17, 0.18)	
	50	0.02 (0.01, 0.02)	0.06 (0.06, 0.07)	0.15 (0.15, 0.16)	0.22 (0.21, 0.23)	0.18 (0.17, 0.18)	
	60	0.05 (0.05, 0.06)	0.15 (0.14, 0.16)	0.21 (0.20, 0.22)	0.22 (0.21, 0.24)	0.18 (0.18, 0.19)	
	70	0.11 (0.11, 0.12)	0.19 (0.18, 0.20)	- (-, -)	0.20 (0.19, 0.22)	0.17 (0.17, 0.18)	
	80	0.11 (0.10, 0.12)	- (-, -)	- (-, -)	0.14 (0.12, 0.15)	0.12 (0.12, 0.13)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.23 (0.22, 0.24)	0.18 (0.18, 0.19)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.23 (0.22, 0.25)	0.19 (0.18, 0.19)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.24 (0.22, 0.25)	0.19 (0.18, 0.19)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.02, 0.02)	0.24 (0.23, 0.25)	0.19 (0.18, 0.19)	
	40	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.07 (0.07, 0.08)	0.24 (0.23, 0.25)	0.19 (0.19, 0.20)	
	50	0.02 (0.01, 0.02)	0.07 (0.07, 0.08)	0.17 (0.16, 0.18)	0.25 (0.23, 0.26)	0.20 (0.19, 0.20)	
	60	0.06 (0.05, 0.06)	0.17 (0.16, 0.18)	0.23 (0.22, 0.25)	0.25 (0.23, 0.26)	0.20 (0.19, 0.20)	
	70	0.13 (0.12, 0.14)	0.21 (0.19, 0.22)	- (-, -)	0.22 (0.21, 0.24)	0.18 (0.18, 0.19)	
	80	0.12 (0.11, 0.13)	- (-, -)	- (-, -)	0.15 (0.13, 0.16)	0.13 (0.12, 0.14)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.08 (0.06, 0.11)	0.06 (0.05, 0.07)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.09 (0.07, 0.11)	0.06 (0.05, 0.07)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.09 (0.07, 0.11)	0.06 (0.05, 0.07)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.09 (0.07, 0.12)	0.06 (0.05, 0.07)	
	40	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.09 (0.07, 0.12)	0.06 (0.05, 0.07)	
	50	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.06 (0.05, 0.08)	0.09 (0.07, 0.12)	0.07 (0.06, 0.08)	
	60	0.02 (0.01, 0.03)	0.06 (0.04, 0.08)	0.09 (0.07, 0.12)	0.10 (0.07, 0.13)	0.07 (0.06, 0.08)	
	70	0.05 (0.03, 0.08)	0.09 (0.07, 0.13)	- (-, -)	0.10 (0.07, 0.14)	0.07 (0.06, 0.08)	
	80	0.08 (0.05, 0.12)	- (-, -)	- (-, -)	0.09 (0.05, 0.15)	0.05 (0.04, 0.07)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

- Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVII-8 - continued
MESOTHELIOMA (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.08 (0.05, 0.14)	0.06 (0.04, 0.11)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.08 (0.05, 0.14)	0.06 (0.04, 0.11)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.08 (0.05, 0.14)	0.06 (0.04, 0.11)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.08 (0.05, 0.14)	0.06 (0.04, 0.11)	
	40	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.03 (0.02, 0.04)	0.08 (0.05, 0.14)	0.06 (0.04, 0.11)	
	50	0.01 (0.00, 0.01)	0.03 (0.02, 0.04)	0.06 (0.04, 0.08)	0.08 (0.05, 0.14)	0.06 (0.04, 0.11)	
	60	0.02 (0.01, 0.04)	0.06 (0.04, 0.08)	0.07 (0.05, 0.09)	0.08 (0.05, 0.14)	0.06 (0.04, 0.12)	
	70	0.04 (0.02, 0.06)	0.05 (0.03, 0.08)	- (-, -)	0.06 (0.03, 0.13)	0.06 (0.03, 0.12)	
	80	0.01 (0.00, 0.04)	- (-, -)	- (-, -)	0.03 (0.01, 0.12)	0.03 (0.01, 0.11)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.06 (0.01, 0.73)	0.03 (0.01, 0.21)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.06 (0.01, 0.75)	0.03 (0.01, 0.22)	
	20	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.06 (0.01, 0.76)	0.04 (0.01, 0.22)	
	30	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.02 (0.00, 0.07)	0.07 (0.02, 0.77)	0.04 (0.01, 0.23)	
	40	0.01 (0.00, 0.04)	0.02 (0.00, 0.07)	0.04 (0.01, 0.13)	0.07 (0.02, 0.80)	0.04 (0.01, 0.24)	
	50	0.01 (0.00, 0.07)	0.04 (0.01, 0.13)	0.06 (0.01, 0.21)	0.06 (0.01, 0.84)	0.04 (0.01, 0.25)	
	60	0.03 (0.00, 0.13)	0.06 (0.01, 0.21)	0.06 (0.01, 0.29)	0.06 (0.01, 0.91)	0.04 (0.01, 0.28)	
	70	0.03 (0.00, 0.20)	0.03 (0.00, 0.31)	- (-, -)	0.03 (0.00, 1.06)	0.02 (0.00, 0.32)	
	80	0.00 (0.00, 0.40)	- (-, -)	- (-, -)	0.00 (0.00, 1.56)	0.00 (0.00, 0.47)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.20 (0.17, 0.26)	0.11 (0.09, 0.13)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.20 (0.17, 0.26)	0.11 (0.09, 0.13)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.20 (0.17, 0.26)	0.11 (0.09, 0.14)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.21 (0.17, 0.27)	0.11 (0.10, 0.14)	
	40	0.00 (0.00, 0.01)	0.02 (0.01, 0.03)	0.06 (0.05, 0.08)	0.21 (0.17, 0.27)	0.11 (0.10, 0.14)	
	50	0.02 (0.01, 0.02)	0.06 (0.05, 0.08)	0.15 (0.13, 0.18)	0.21 (0.17, 0.28)	0.11 (0.10, 0.14)	
	60	0.04 (0.03, 0.06)	0.14 (0.12, 0.17)	0.20 (0.16, 0.24)	0.21 (0.17, 0.27)	0.12 (0.10, 0.14)	
	70	0.11 (0.09, 0.14)	0.17 (0.14, 0.22)	- (-, -)	0.19 (0.14, 0.26)	0.11 (0.09, 0.14)	
	80	0.08 (0.06, 0.13)	- (-, -)	- (-, -)	0.10 (0.06, 0.20)	0.06 (0.05, 0.10)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVII-8 - continued
MESOTHELIOMA (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.05, 0.05)	0.03 (0.03, 0.04)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.05, 0.05)	0.03 (0.03, 0.04)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.05, 0.05)	0.03 (0.03, 0.04)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.05, 0.06)	0.03 (0.03, 0.04)	
	40	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.05 (0.05, 0.05)	0.03 (0.03, 0.04)	
	50	0.01 (0.00, 0.01)	0.02 (0.01, 0.02)	0.03 (0.03, 0.04)	0.05 (0.04, 0.05)	0.03 (0.03, 0.04)	
	60	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.04 (0.04, 0.05)	0.05 (0.04, 0.05)	0.03 (0.03, 0.03)	
	70	0.02 (0.02, 0.02)	0.04 (0.03, 0.04)	- (-, -)	0.04 (0.03, 0.04)	0.03 (0.03, 0.03)	
	80	0.02 (0.02, 0.03)	- (-, -)	- (-, -)	0.02 (0.02, 0.03)	0.02 (0.02, 0.02)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.05, 0.06)	0.04 (0.04, 0.04)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.05, 0.06)	0.04 (0.04, 0.04)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.05 (0.05, 0.06)	0.04 (0.04, 0.04)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.06 (0.05, 0.06)	0.04 (0.04, 0.04)	
	40	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.05 (0.05, 0.06)	0.04 (0.04, 0.04)	
	50	0.01 (0.00, 0.01)	0.02 (0.01, 0.02)	0.04 (0.03, 0.04)	0.05 (0.05, 0.06)	0.04 (0.04, 0.04)	
	60	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	0.05 (0.04, 0.05)	0.05 (0.04, 0.06)	0.04 (0.03, 0.04)	
	70	0.02 (0.02, 0.03)	0.04 (0.04, 0.05)	- (-, -)	0.04 (0.04, 0.05)	0.03 (0.03, 0.03)	
	80	0.02 (0.02, 0.03)	- (-, -)	- (-, -)	0.03 (0.02, 0.03)	0.02 (0.02, 0.02)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.03 (0.02, 0.04)	0.01 (0.01, 0.01)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.03 (0.02, 0.04)	0.01 (0.01, 0.01)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.03 (0.02, 0.04)	0.01 (0.01, 0.01)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.03 (0.02, 0.04)	0.01 (0.01, 0.01)	
	40	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.03 (0.02, 0.04)	0.01 (0.01, 0.01)	
	50	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.02 (0.01, 0.03)	0.03 (0.02, 0.04)	0.01 (0.01, 0.01)	
	60	0.00 (0.00, 0.01)	0.02 (0.01, 0.03)	0.02 (0.01, 0.04)	0.03 (0.02, 0.04)	0.01 (0.01, 0.01)	
	70	0.01 (0.01, 0.03)	0.02 (0.01, 0.04)	- (-, -)	0.03 (0.01, 0.05)	0.01 (0.01, 0.01)	
	80	0.01 (0.00, 0.03)	- (-, -)	- (-, -)	0.02 (0.01, 0.04)	0.01 (0.00, 0.01)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

- Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVII-8 - continued
MESOTHELIOMA (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.07)	0.01 (0.00, 0.05)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.07)	0.01 (0.00, 0.05)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.07)	0.01 (0.00, 0.05)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.02 (0.01, 0.07)	0.01 (0.00, 0.05)	
	40	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.02 (0.01, 0.07)	0.01 (0.00, 0.05)	
	50	0.00 (0.00, 0.01)	0.01 (0.00, 0.02)	0.01 (0.01, 0.02)	0.02 (0.01, 0.06)	0.01 (0.00, 0.05)	
	60	0.01 (0.00, 0.02)	0.01 (0.01, 0.02)	0.01 (0.01, 0.03)	0.01 (0.01, 0.06)	0.01 (0.00, 0.05)	
	70	0.00 (0.00, 0.01)	0.01 (0.00, 0.02)	- (-, -)	0.01 (0.00, 0.06)	0.00 (0.00, 0.05)	
	80	0.00 (0.00, 0.02)	- (-, -)	- (-, -)	0.00 (0.00, 0.07)	0.00 (0.00, 0.06)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.10 (0.02, 0.61)	0.05 (0.02, 0.19)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.10 (0.02, 0.61)	0.05 (0.02, 0.20)	
	20	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.10 (0.02, 0.62)	0.05 (0.02, 0.20)	
	30	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.00 (0.00, 0.04)	0.10 (0.02, 0.62)	0.05 (0.02, 0.20)	
	40	0.00 (0.00, 0.03)	0.00 (0.00, 0.04)	0.00 (0.00, 0.07)	0.10 (0.02, 0.63)	0.05 (0.02, 0.21)	
	50	0.00 (0.00, 0.04)	0.00 (0.00, 0.07)	0.03 (0.00, 0.15)	0.11 (0.02, 0.65)	0.06 (0.02, 0.21)	
	60	0.00 (0.00, 0.07)	0.03 (0.00, 0.15)	0.11 (0.02, 0.34)	0.11 (0.02, 0.69)	0.06 (0.02, 0.23)	
	70	0.03 (0.00, 0.17)	0.12 (0.02, 0.39)	- (-, -)	0.13 (0.02, 0.79)	0.07 (0.02, 0.27)	
	80	0.12 (0.01, 0.45)	- (-, -)	- (-, -)	0.13 (0.01, 1.00)	0.05 (0.01, 0.32)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.06 (0.05, 0.10)	0.03 (0.02, 0.04)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.06 (0.05, 0.10)	0.03 (0.02, 0.04)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.06 (0.05, 0.10)	0.03 (0.02, 0.04)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.06 (0.05, 0.10)	0.03 (0.02, 0.04)	
	40	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	0.02 (0.02, 0.03)	0.06 (0.05, 0.10)	0.03 (0.02, 0.04)	
	50	0.01 (0.00, 0.01)	0.02 (0.01, 0.03)	0.04 (0.03, 0.05)	0.06 (0.04, 0.10)	0.03 (0.02, 0.04)	
	60	0.01 (0.01, 0.02)	0.03 (0.02, 0.05)	0.05 (0.04, 0.08)	0.06 (0.04, 0.10)	0.03 (0.02, 0.04)	
	70	0.02 (0.01, 0.03)	0.04 (0.03, 0.06)	- (-, -)	0.05 (0.03, 0.09)	0.02 (0.01, 0.04)	
	80	0.03 (0.01, 0.05)	- (-, -)	- (-, -)	0.03 (0.01, 0.08)	0.02 (0.01, 0.03)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVIII-9

MYELOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.61 (0.60, 0.62)	0.42 (0.42, 0.43)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.61 (0.60, 0.63)	0.43 (0.42, 0.43)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.62 (0.60, 0.63)	0.43 (0.42, 0.43)	
	30	0.01 (0.00, 0.01)	0.03 (0.03, 0.03)	0.10 (0.10, 0.10)	0.62 (0.61, 0.63)	0.43 (0.43, 0.44)	
	40	0.02 (0.02, 0.03)	0.10 (0.09, 0.10)	0.25 (0.24, 0.25)	0.62 (0.61, 0.64)	0.44 (0.43, 0.44)	
	50	0.07 (0.07, 0.08)	0.23 (0.22, 0.24)	0.45 (0.44, 0.46)	0.62 (0.61, 0.63)	0.44 (0.44, 0.45)	
	60	0.16 (0.16, 0.17)	0.39 (0.39, 0.40)	0.55 (0.54, 0.56)	0.58 (0.56, 0.59)	0.44 (0.43, 0.44)	
	70	0.27 (0.26, 0.27)	0.44 (0.43, 0.45)	- (- , -)	0.48 (0.46, 0.49)	0.40 (0.39, 0.40)	
	80	0.25 (0.24, 0.26)	- (- , -)	- (- , -)	0.30 (0.29, 0.31)	0.29 (0.28, 0.29)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.56 (0.55, 0.57)	0.40 (0.40, 0.41)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.57 (0.56, 0.58)	0.41 (0.40, 0.41)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	0.57 (0.56, 0.58)	0.41 (0.40, 0.41)	
	30	0.00 (0.00, 0.01)	0.03 (0.02, 0.03)	0.09 (0.09, 0.09)	0.57 (0.56, 0.59)	0.41 (0.41, 0.42)	
	40	0.02 (0.02, 0.02)	0.09 (0.08, 0.09)	0.22 (0.22, 0.23)	0.58 (0.56, 0.59)	0.41 (0.41, 0.42)	
	50	0.07 (0.06, 0.07)	0.21 (0.20, 0.21)	0.41 (0.40, 0.42)	0.57 (0.56, 0.58)	0.42 (0.41, 0.42)	
	60	0.15 (0.14, 0.16)	0.36 (0.35, 0.37)	0.51 (0.50, 0.52)	0.53 (0.52, 0.55)	0.41 (0.41, 0.42)	
	70	0.25 (0.24, 0.26)	0.41 (0.40, 0.42)	- (- , -)	0.44 (0.43, 0.46)	0.38 (0.37, 0.38)	
	80	0.24 (0.23, 0.25)	- (- , -)	- (- , -)	0.28 (0.27, 0.29)	0.27 (0.27, 0.28)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.02 (0.98, 1.07)	0.64 (0.62, 0.66)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	1.05 (1.00, 1.10)	0.66 (0.64, 0.68)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.06 (0.06, 0.07)	1.05 (1.01, 1.10)	0.66 (0.64, 0.68)	
	30	0.01 (0.01, 0.02)	0.06 (0.06, 0.07)	0.21 (0.20, 0.23)	1.07 (1.02, 1.12)	0.67 (0.65, 0.69)	
	40	0.05 (0.05, 0.06)	0.20 (0.19, 0.22)	0.50 (0.47, 0.52)	1.08 (1.03, 1.13)	0.68 (0.67, 0.70)	
	50	0.16 (0.15, 0.17)	0.47 (0.44, 0.50)	0.83 (0.78, 0.87)	1.09 (1.03, 1.14)	0.70 (0.68, 0.72)	
	60	0.34 (0.32, 0.37)	0.75 (0.71, 0.79)	0.99 (0.94, 1.04)	1.04 (0.98, 1.10)	0.70 (0.68, 0.73)	
	70	0.51 (0.47, 0.55)	0.81 (0.76, 0.86)	- (- , -)	0.87 (0.81, 0.93)	0.65 (0.63, 0.68)	
	80	0.48 (0.44, 0.54)	- (- , -)	- (- , -)	0.58 (0.52, 0.65)	0.49 (0.46, 0.52)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVIII-9 - continued

MYELOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.43 (0.39, 0.49)	0.25 (0.22, 0.29)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.44 (0.40, 0.49)	0.25 (0.22, 0.29)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.44 (0.40, 0.49)	0.25 (0.22, 0.29)	
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.06 (0.05, 0.07)	0.44 (0.40, 0.49)	0.25 (0.22, 0.29)	
	40	0.01 (0.01, 0.02)	0.06 (0.05, 0.07)	0.16 (0.14, 0.17)	0.44 (0.40, 0.50)	0.26 (0.23, 0.29)	
	50	0.05 (0.04, 0.05)	0.15 (0.13, 0.16)	0.29 (0.27, 0.32)	0.44 (0.39, 0.49)	0.26 (0.23, 0.29)	
	60	0.10 (0.09, 0.12)	0.26 (0.23, 0.29)	0.37 (0.33, 0.41)	0.40 (0.36, 0.46)	0.25 (0.21, 0.29)	
	70	0.17 (0.15, 0.19)	0.29 (0.26, 0.33)	- (-, -)	0.33 (0.28, 0.38)	0.22 (0.19, 0.26)	
	80	0.15 (0.12, 0.19)	- (-, -)	- (-, -)	0.20 (0.15, 0.26)	0.15 (0.12, 0.20)	
American	0	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.55 (0.40, 0.87)	0.41 (0.32, 0.53)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.01)	0.00 (0.00, 0.02)	0.01 (0.00, 0.03)	0.56 (0.40, 0.89)	0.41 (0.33, 0.54)	
	20	0.00 (0.00, 0.02)	0.01 (0.00, 0.03)	0.03 (0.01, 0.05)	0.57 (0.41, 0.90)	0.42 (0.33, 0.55)	
	30	0.01 (0.00, 0.03)	0.02 (0.01, 0.05)	0.09 (0.06, 0.14)	0.57 (0.41, 0.91)	0.43 (0.34, 0.56)	
	40	0.02 (0.01, 0.04)	0.09 (0.05, 0.14)	0.20 (0.13, 0.29)	0.58 (0.41, 0.92)	0.44 (0.35, 0.57)	
	50	0.07 (0.04, 0.12)	0.19 (0.12, 0.29)	0.42 (0.30, 0.58)	0.59 (0.41, 0.94)	0.45 (0.35, 0.59)	
	60	0.13 (0.07, 0.21)	0.38 (0.25, 0.54)	0.52 (0.36, 0.74)	0.55 (0.38, 0.93)	0.43 (0.33, 0.58)	
	70	0.29 (0.17, 0.46)	0.45 (0.29, 0.70)	- (-, -)	0.50 (0.31, 0.93)	0.42 (0.30, 0.59)	
	80	0.23 (0.10, 0.47)	- (-, -)	- (-, -)	0.29 (0.12, 0.87)	0.29 (0.16, 0.52)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.70 (0.65, 0.76)	0.40 (0.38, 0.43)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.71 (0.66, 0.77)	0.41 (0.39, 0.44)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.03, 0.03)	0.71 (0.66, 0.77)	0.41 (0.39, 0.44)	
	30	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	0.10 (0.09, 0.11)	0.72 (0.67, 0.78)	0.41 (0.39, 0.44)	
	40	0.02 (0.02, 0.03)	0.09 (0.09, 0.10)	0.26 (0.24, 0.28)	0.72 (0.67, 0.78)	0.42 (0.39, 0.44)	
	50	0.07 (0.06, 0.08)	0.24 (0.22, 0.26)	0.48 (0.45, 0.51)	0.71 (0.66, 0.77)	0.42 (0.40, 0.45)	
	60	0.18 (0.16, 0.20)	0.43 (0.39, 0.46)	0.62 (0.57, 0.66)	0.67 (0.62, 0.73)	0.41 (0.38, 0.44)	
	70	0.28 (0.25, 0.31)	0.49 (0.45, 0.54)	- (-, -)	0.55 (0.50, 0.62)	0.36 (0.34, 0.39)	
	80	0.28 (0.24, 0.33)	- (-, -)	- (-, -)	0.36 (0.31, 0.43)	0.26 (0.23, 0.29)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVIII-9 - continued

MYELOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.69 (0.67, 0.71)	0.46 (0.45, 0.47)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.70 (0.68, 0.72)	0.46 (0.46, 0.47)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	0.71 (0.69, 0.72)	0.47 (0.46, 0.48)	
	30	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	0.12 (0.11, 0.12)	0.71 (0.70, 0.73)	0.47 (0.47, 0.48)	
	40	0.03 (0.03, 0.03)	0.11 (0.11, 0.12)	0.29 (0.28, 0.30)	0.72 (0.70, 0.74)	0.48 (0.47, 0.49)	
	50	0.09 (0.08, 0.09)	0.27 (0.26, 0.28)	0.52 (0.51, 0.54)	0.72 (0.70, 0.74)	0.49 (0.48, 0.50)	
	60	0.20 (0.19, 0.21)	0.47 (0.46, 0.49)	0.65 (0.63, 0.66)	0.68 (0.66, 0.70)	0.49 (0.48, 0.50)	
	70	0.33 (0.31, 0.34)	0.53 (0.52, 0.55)	- (-, -)	0.58 (0.56, 0.60)	0.45 (0.44, 0.46)	
	80	0.32 (0.30, 0.34)	- (-, -)	- (-, -)	0.38 (0.36, 0.41)	0.34 (0.33, 0.35)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.65 (0.64, 0.67)	0.44 (0.44, 0.45)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.66 (0.64, 0.68)	0.45 (0.44, 0.46)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.67 (0.65, 0.69)	0.45 (0.44, 0.46)	
	30	0.01 (0.00, 0.01)	0.03 (0.03, 0.03)	0.11 (0.10, 0.11)	0.67 (0.65, 0.69)	0.46 (0.45, 0.47)	
	40	0.03 (0.02, 0.03)	0.10 (0.10, 0.11)	0.26 (0.25, 0.28)	0.68 (0.66, 0.70)	0.46 (0.46, 0.47)	
	50	0.08 (0.07, 0.08)	0.25 (0.24, 0.26)	0.49 (0.47, 0.51)	0.68 (0.66, 0.70)	0.47 (0.46, 0.48)	
	60	0.18 (0.17, 0.19)	0.44 (0.43, 0.46)	0.61 (0.59, 0.63)	0.64 (0.62, 0.66)	0.47 (0.46, 0.48)	
	70	0.31 (0.29, 0.32)	0.51 (0.49, 0.53)	- (-, -)	0.55 (0.52, 0.57)	0.43 (0.42, 0.44)	
	80	0.31 (0.29, 0.32)	- (-, -)	- (-, -)	0.37 (0.35, 0.39)	0.33 (0.32, 0.34)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.09 (1.02, 1.17)	0.63 (0.60, 0.65)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	1.12 (1.05, 1.20)	0.64 (0.62, 0.67)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.07 (0.06, 0.08)	1.13 (1.06, 1.21)	0.65 (0.62, 0.68)	
	30	0.01 (0.01, 0.02)	0.07 (0.06, 0.08)	0.24 (0.22, 0.27)	1.16 (1.08, 1.24)	0.66 (0.64, 0.69)	
	40	0.06 (0.05, 0.07)	0.23 (0.21, 0.26)	0.56 (0.52, 0.61)	1.18 (1.10, 1.26)	0.68 (0.65, 0.71)	
	50	0.19 (0.17, 0.21)	0.54 (0.50, 0.59)	0.94 (0.87, 1.00)	1.20 (1.12, 1.28)	0.71 (0.68, 0.74)	
	60	0.41 (0.36, 0.45)	0.86 (0.80, 0.94)	1.12 (1.04, 1.21)	1.17 (1.08, 1.26)	0.72 (0.68, 0.75)	
	70	0.61 (0.55, 0.68)	0.96 (0.87, 1.05)	- (-, -)	1.02 (0.92, 1.13)	0.68 (0.64, 0.71)	
	80	0.63 (0.53, 0.73)	- (-, -)	- (-, -)	0.74 (0.62, 0.87)	0.55 (0.50, 0.60)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

- Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVIII-9 - continued

MYELOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.47 (0.41, 0.56)	0.26 (0.22, 0.32)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.48 (0.41, 0.56)	0.27 (0.22, 0.33)	
Islander	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.48 (0.41, 0.56)	0.27 (0.23, 0.33)	
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.06 (0.05, 0.08)	0.48 (0.42, 0.57)	0.27 (0.23, 0.33)	
	40	0.01 (0.01, 0.02)	0.06 (0.05, 0.08)	0.18 (0.15, 0.21)	0.48 (0.42, 0.57)	0.27 (0.23, 0.33)	
	50	0.05 (0.04, 0.07)	0.17 (0.14, 0.20)	0.34 (0.29, 0.38)	0.48 (0.41, 0.57)	0.27 (0.23, 0.33)	
	60	0.12 (0.10, 0.15)	0.30 (0.25, 0.34)	0.41 (0.36, 0.48)	0.45 (0.38, 0.54)	0.26 (0.22, 0.33)	
	70	0.19 (0.16, 0.23)	0.32 (0.27, 0.39)	- (-, -)	0.36 (0.29, 0.45)	0.23 (0.19, 0.30)	
	80	0.17 (0.12, 0.23)	- (-, -)	- (-, -)	0.22 (0.15, 0.33)	0.16 (0.11, 0.25)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.41 (0.22, 1.09)	0.45 (0.30, 0.70)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.01 (0.00, 0.05)	0.42 (0.23, 1.11)	0.45 (0.31, 0.71)	
	20	0.00 (0.00, 0.02)	0.01 (0.00, 0.05)	0.02 (0.01, 0.07)	0.42 (0.23, 1.12)	0.46 (0.31, 0.72)	
	30	0.01 (0.00, 0.05)	0.02 (0.01, 0.07)	0.07 (0.03, 0.14)	0.43 (0.24, 1.15)	0.47 (0.32, 0.74)	
	40	0.01 (0.00, 0.04)	0.06 (0.02, 0.12)	0.13 (0.06, 0.26)	0.43 (0.23, 1.17)	0.49 (0.33, 0.77)	
	50	0.05 (0.01, 0.12)	0.13 (0.05, 0.26)	0.27 (0.14, 0.48)	0.44 (0.23, 1.23)	0.50 (0.34, 0.80)	
	60	0.09 (0.03, 0.21)	0.24 (0.11, 0.46)	0.43 (0.21, 0.81)	0.43 (0.21, 1.29)	0.50 (0.32, 0.83)	
	70	0.18 (0.06, 0.42)	0.41 (0.17, 0.85)	- (-, -)	0.42 (0.18, 1.44)	0.53 (0.31, 0.94)	
	80	0.34 (0.10, 0.91)	- (-, -)	- (-, -)	0.35 (0.10, 1.89)	0.50 (0.22, 1.11)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.78 (0.70, 0.87)	0.42 (0.39, 0.46)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.79 (0.71, 0.89)	0.43 (0.39, 0.47)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	0.79 (0.72, 0.89)	0.43 (0.39, 0.47)	
	30	0.01 (0.00, 0.01)	0.03 (0.03, 0.04)	0.11 (0.09, 0.12)	0.80 (0.72, 0.90)	0.43 (0.40, 0.48)	
	40	0.03 (0.02, 0.03)	0.10 (0.09, 0.12)	0.30 (0.27, 0.33)	0.81 (0.73, 0.91)	0.44 (0.40, 0.48)	
	50	0.08 (0.06, 0.09)	0.28 (0.25, 0.31)	0.54 (0.49, 0.59)	0.80 (0.72, 0.91)	0.44 (0.41, 0.49)	
	60	0.22 (0.19, 0.25)	0.49 (0.44, 0.55)	0.71 (0.64, 0.79)	0.77 (0.69, 0.88)	0.44 (0.40, 0.48)	
	70	0.32 (0.28, 0.37)	0.57 (0.50, 0.65)	- (-, -)	0.64 (0.56, 0.76)	0.39 (0.35, 0.44)	
	80	0.36 (0.29, 0.44)	- (-, -)	- (-, -)	0.46 (0.36, 0.61)	0.29 (0.24, 0.35)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVIII-9 - continued

MYELOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.53 (0.52, 0.55)	0.39 (0.39, 0.40)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.54 (0.52, 0.55)	0.40 (0.39, 0.40)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.03 (0.02, 0.03)	0.54 (0.53, 0.56)	0.40 (0.39, 0.41)	
	30	0.00 (0.00, 0.01)	0.03 (0.02, 0.03)	0.08 (0.08, 0.09)	0.54 (0.53, 0.56)	0.40 (0.39, 0.41)	
	40	0.02 (0.02, 0.02)	0.08 (0.08, 0.09)	0.21 (0.20, 0.22)	0.54 (0.53, 0.56)	0.40 (0.40, 0.41)	
	50	0.06 (0.06, 0.06)	0.19 (0.18, 0.20)	0.37 (0.36, 0.39)	0.53 (0.52, 0.55)	0.41 (0.40, 0.41)	
	60	0.14 (0.13, 0.14)	0.33 (0.32, 0.34)	0.47 (0.45, 0.48)	0.49 (0.48, 0.51)	0.40 (0.39, 0.40)	
	70	0.22 (0.21, 0.23)	0.37 (0.36, 0.38)	- (- , -)	0.40 (0.39, 0.42)	0.36 (0.35, 0.37)	
	80	0.21 (0.20, 0.22)	- (- , -)	- (- , -)	0.25 (0.24, 0.26)	0.26 (0.25, 0.26)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.48 (0.47, 0.50)	0.37 (0.36, 0.37)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.49 (0.47, 0.50)	0.37 (0.36, 0.38)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.49 (0.47, 0.50)	0.37 (0.36, 0.38)	
	30	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.07 (0.07, 0.08)	0.49 (0.47, 0.51)	0.37 (0.37, 0.38)	
	40	0.02 (0.02, 0.02)	0.07 (0.07, 0.07)	0.18 (0.17, 0.19)	0.49 (0.48, 0.51)	0.37 (0.37, 0.38)	
	50	0.05 (0.05, 0.06)	0.17 (0.16, 0.18)	0.34 (0.32, 0.35)	0.48 (0.47, 0.50)	0.38 (0.37, 0.38)	
	60	0.12 (0.11, 0.13)	0.30 (0.28, 0.31)	0.42 (0.41, 0.44)	0.45 (0.43, 0.46)	0.37 (0.36, 0.38)	
	70	0.20 (0.19, 0.21)	0.34 (0.33, 0.35)	- (- , -)	0.37 (0.35, 0.38)	0.33 (0.33, 0.34)	
	80	0.19 (0.18, 0.20)	- (- , -)	- (- , -)	0.23 (0.22, 0.24)	0.24 (0.24, 0.25)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.97 (0.90, 1.03)	0.66 (0.63, 0.68)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.99 (0.92, 1.05)	0.67 (0.65, 0.70)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.06 (0.05, 0.07)	0.99 (0.93, 1.06)	0.67 (0.65, 0.70)	
	30	0.01 (0.01, 0.02)	0.06 (0.05, 0.07)	0.18 (0.17, 0.20)	1.00 (0.93, 1.06)	0.68 (0.65, 0.71)	
	40	0.05 (0.04, 0.06)	0.18 (0.16, 0.20)	0.44 (0.40, 0.47)	1.00 (0.94, 1.07)	0.69 (0.66, 0.72)	
	50	0.13 (0.12, 0.15)	0.40 (0.37, 0.44)	0.73 (0.68, 0.79)	0.99 (0.93, 1.06)	0.70 (0.67, 0.73)	
	60	0.29 (0.26, 0.33)	0.65 (0.60, 0.71)	0.88 (0.82, 0.95)	0.94 (0.87, 1.01)	0.69 (0.67, 0.72)	
	70	0.43 (0.39, 0.48)	0.70 (0.64, 0.77)	- (- , -)	0.77 (0.70, 0.85)	0.64 (0.61, 0.67)	
	80	0.40 (0.35, 0.47)	- (- , -)	- (- , -)	0.50 (0.43, 0.58)	0.46 (0.43, 0.49)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

- Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVIII-9 - continued

MYELOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.40 (0.34, 0.48)	0.24 (0.20, 0.30)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.40 (0.35, 0.49)	0.24 (0.20, 0.30)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.40 (0.35, 0.49)	0.24 (0.20, 0.30)	
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.06 (0.04, 0.07)	0.41 (0.35, 0.49)	0.24 (0.20, 0.31)	
	40	0.01 (0.01, 0.02)	0.05 (0.04, 0.07)	0.14 (0.12, 0.16)	0.41 (0.35, 0.49)	0.24 (0.20, 0.31)	
	50	0.04 (0.03, 0.05)	0.13 (0.10, 0.15)	0.26 (0.23, 0.30)	0.40 (0.34, 0.48)	0.24 (0.20, 0.30)	
	60	0.09 (0.07, 0.11)	0.23 (0.19, 0.26)	0.34 (0.29, 0.39)	0.36 (0.31, 0.45)	0.23 (0.19, 0.30)	
	70	0.15 (0.12, 0.18)	0.26 (0.22, 0.32)	- (-, -)	0.30 (0.24, 0.38)	0.21 (0.16, 0.27)	
	80	0.14 (0.10, 0.19)	- (-, -)	- (-, -)	0.18 (0.12, 0.27)	0.14 (0.10, 0.22)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.01 (0.00, 0.05)	0.69 (0.46, 1.24)	0.39 (0.29, 0.57)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.01 (0.00, 0.04)	0.01 (0.00, 0.04)	0.70 (0.46, 1.26)	0.39 (0.29, 0.58)	
	20	0.01 (0.00, 0.04)	0.01 (0.00, 0.04)	0.03 (0.01, 0.08)	0.71 (0.47, 1.26)	0.40 (0.29, 0.58)	
	30	0.00 (0.00, 0.03)	0.02 (0.01, 0.06)	0.11 (0.06, 0.20)	0.70 (0.47, 1.27)	0.40 (0.30, 0.58)	
	40	0.02 (0.01, 0.06)	0.11 (0.06, 0.20)	0.27 (0.16, 0.42)	0.72 (0.47, 1.29)	0.41 (0.30, 0.60)	
	50	0.09 (0.05, 0.18)	0.25 (0.15, 0.41)	0.57 (0.37, 0.83)	0.71 (0.46, 1.30)	0.42 (0.31, 0.61)	
	60	0.17 (0.08, 0.30)	0.50 (0.31, 0.77)	0.60 (0.38, 0.94)	0.65 (0.40, 1.28)	0.38 (0.27, 0.59)	
	70	0.38 (0.20, 0.65)	0.50 (0.28, 0.85)	- (-, -)	0.56 (0.31, 1.26)	0.35 (0.23, 0.58)	
	80	0.16 (0.05, 0.46)	- (-, -)	- (-, -)	0.24 (0.06, 1.13)	0.16 (0.07, 0.45)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.64 (0.58, 0.72)	0.39 (0.36, 0.43)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.65 (0.59, 0.73)	0.40 (0.37, 0.43)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.65 (0.59, 0.73)	0.40 (0.37, 0.43)	
	30	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.09 (0.08, 0.10)	0.65 (0.59, 0.73)	0.40 (0.37, 0.44)	
	40	0.02 (0.02, 0.02)	0.09 (0.08, 0.10)	0.23 (0.20, 0.25)	0.65 (0.59, 0.73)	0.40 (0.37, 0.44)	
	50	0.07 (0.06, 0.08)	0.21 (0.18, 0.24)	0.43 (0.39, 0.47)	0.64 (0.58, 0.72)	0.40 (0.37, 0.44)	
	60	0.15 (0.12, 0.17)	0.37 (0.33, 0.41)	0.54 (0.49, 0.60)	0.59 (0.53, 0.67)	0.39 (0.35, 0.43)	
	70	0.25 (0.21, 0.28)	0.43 (0.38, 0.49)	- (-, -)	0.49 (0.42, 0.57)	0.34 (0.31, 0.38)	
	80	0.23 (0.19, 0.29)	- (-, -)	- (-, -)	0.30 (0.24, 0.39)	0.24 (0.21, 0.29)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XIX-13
NON-HODGKIN LYMPHOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.05 (0.05, 0.05)	2.02 (2.00, 2.04)	0.83 (0.83, 0.84)	
	10	0.01 (0.01, 0.02)	0.04 (0.04, 0.05)	0.10 (0.10, 0.11)	2.04 (2.02, 2.06)	0.84 (0.84, 0.85)	
	20	0.03 (0.03, 0.03)	0.09 (0.09, 0.09)	0.21 (0.21, 0.22)	2.03 (2.01, 2.05)	0.85 (0.84, 0.85)	
	30	0.06 (0.06, 0.06)	0.19 (0.18, 0.19)	0.44 (0.43, 0.45)	2.02 (2.00, 2.04)	0.85 (0.84, 0.86)	
	40	0.13 (0.12, 0.13)	0.38 (0.38, 0.39)	0.85 (0.84, 0.86)	1.99 (1.97, 2.01)	0.85 (0.84, 0.86)	
	50	0.27 (0.26, 0.27)	0.74 (0.73, 0.76)	1.38 (1.36, 1.39)	1.92 (1.90, 1.94)	0.85 (0.85, 0.86)	
	60	0.51 (0.50, 0.52)	1.18 (1.17, 1.20)	1.66 (1.64, 1.68)	1.76 (1.74, 1.78)	0.84 (0.83, 0.85)	
	70	0.78 (0.77, 0.79)	1.33 (1.31, 1.35)	- (- , -)	1.45 (1.43, 1.47)	0.78 (0.77, 0.79)	
	80	0.78 (0.77, 0.80)	- (- , -)	- (- , -)	0.95 (0.93, 0.98)	0.62 (0.62, 0.63)	
White	0	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.05 (0.05, 0.05)	2.12 (2.10, 2.15)	0.88 (0.88, 0.89)	
	10	0.02 (0.01, 0.02)	0.04 (0.04, 0.05)	0.10 (0.10, 0.11)	2.14 (2.12, 2.16)	0.89 (0.89, 0.90)	
	20	0.03 (0.03, 0.03)	0.09 (0.08, 0.09)	0.21 (0.21, 0.22)	2.13 (2.11, 2.16)	0.90 (0.89, 0.90)	
	30	0.06 (0.06, 0.06)	0.19 (0.18, 0.19)	0.45 (0.44, 0.46)	2.12 (2.10, 2.15)	0.90 (0.89, 0.91)	
	40	0.13 (0.13, 0.13)	0.40 (0.39, 0.40)	0.89 (0.88, 0.91)	2.09 (2.07, 2.11)	0.90 (0.89, 0.91)	
	50	0.27 (0.27, 0.28)	0.78 (0.77, 0.80)	1.45 (1.44, 1.47)	2.02 (1.99, 2.04)	0.90 (0.89, 0.91)	
	60	0.54 (0.53, 0.55)	1.25 (1.24, 1.27)	1.75 (1.72, 1.77)	1.85 (1.83, 1.87)	0.89 (0.88, 0.90)	
	70	0.82 (0.81, 0.84)	1.39 (1.37, 1.41)	- (- , -)	1.51 (1.49, 1.53)	0.82 (0.81, 0.83)	
	80	0.82 (0.80, 0.83)	- (- , -)	- (- , -)	0.99 (0.97, 1.01)	0.65 (0.64, 0.66)	
Black	0	0.01 (0.00, 0.01)	0.02 (0.02, 0.02)	0.06 (0.05, 0.06)	1.21 (1.16, 1.26)	0.45 (0.44, 0.47)	
	10	0.01 (0.01, 0.02)	0.05 (0.04, 0.06)	0.13 (0.12, 0.14)	1.23 (1.18, 1.28)	0.46 (0.45, 0.48)	
	20	0.04 (0.03, 0.04)	0.11 (0.10, 0.12)	0.25 (0.24, 0.27)	1.22 (1.18, 1.27)	0.46 (0.45, 0.48)	
	30	0.08 (0.07, 0.09)	0.22 (0.20, 0.23)	0.44 (0.42, 0.46)	1.21 (1.16, 1.26)	0.46 (0.45, 0.48)	
	40	0.14 (0.13, 0.15)	0.37 (0.35, 0.39)	0.66 (0.63, 0.69)	1.16 (1.11, 1.21)	0.46 (0.44, 0.47)	
	50	0.24 (0.22, 0.26)	0.55 (0.52, 0.58)	0.87 (0.83, 0.92)	1.07 (1.02, 1.12)	0.45 (0.44, 0.47)	
	60	0.35 (0.32, 0.37)	0.71 (0.67, 0.75)	0.89 (0.84, 0.94)	0.93 (0.88, 0.99)	0.43 (0.42, 0.45)	
	70	0.46 (0.42, 0.50)	0.68 (0.63, 0.73)	- (- , -)	0.73 (0.68, 0.79)	0.38 (0.36, 0.40)	
	80	0.36 (0.31, 0.40)	- (- , -)	- (- , -)	0.45 (0.39, 0.51)	0.28 (0.26, 0.30)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XIX-13 - continued

NON-HODGKIN LYMPHOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer						Risk of Dying from Cancer	
		+10 yrs		+20 yrs		+30 yrs		Eventually	Eventually
		Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
Asian/ Pacific Islander	0	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.04 (0.04, 0.05)	1.73 (1.64, 1.83)	0.79 (0.72, 0.86)			
	10	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.08 (0.07, 0.09)	1.74 (1.65, 1.85)	0.79 (0.73, 0.87)			
	20	0.02 (0.02, 0.03)	0.06 (0.06, 0.07)	0.14 (0.13, 0.15)	1.73 (1.64, 1.84)	0.79 (0.73, 0.87)			
	30	0.04 (0.03, 0.04)	0.11 (0.10, 0.12)	0.28 (0.27, 0.30)	1.72 (1.63, 1.82)	0.79 (0.73, 0.87)			
	40	0.08 (0.07, 0.08)	0.25 (0.23, 0.27)	0.55 (0.52, 0.58)	1.69 (1.60, 1.79)	0.79 (0.72, 0.86)			
	50	0.17 (0.16, 0.19)	0.48 (0.45, 0.51)	0.97 (0.92, 1.03)	1.64 (1.54, 1.74)	0.79 (0.72, 0.86)			
	60	0.32 (0.29, 0.35)	0.83 (0.78, 0.88)	1.31 (1.24, 1.38)	1.52 (1.42, 1.62)	0.77 (0.70, 0.85)			
	70	0.55 (0.51, 0.60)	1.07 (1.01, 1.15)	- (-, -)	1.30 (1.20, 1.41)	0.73 (0.66, 0.81)			
American Indian/ Alaska Native ^a	80	0.65 (0.59, 0.72)	- (-, -)	- (-, -)	0.93 (0.83, 1.06)	0.59 (0.51, 0.69)			
	0	0.00 (0.00, 0.02)	0.01 (0.00, 0.03)	0.01 (0.00, 0.04)	0.91 (0.71, 1.26)	0.52 (0.41, 0.66)			
	10	0.01 (0.00, 0.02)	0.01 (0.00, 0.03)	0.04 (0.02, 0.07)	0.92 (0.72, 1.28)	0.53 (0.42, 0.67)			
	20	0.00 (0.00, 0.02)	0.03 (0.02, 0.06)	0.10 (0.07, 0.14)	0.92 (0.72, 1.29)	0.53 (0.42, 0.68)			
	30	0.03 (0.01, 0.06)	0.10 (0.07, 0.14)	0.26 (0.19, 0.33)	0.94 (0.73, 1.30)	0.54 (0.43, 0.69)			
	40	0.07 (0.05, 0.11)	0.23 (0.17, 0.31)	0.45 (0.35, 0.57)	0.93 (0.72, 1.31)	0.54 (0.43, 0.70)			
	50	0.17 (0.12, 0.23)	0.39 (0.29, 0.51)	0.61 (0.47, 0.78)	0.89 (0.68, 1.29)	0.55 (0.43, 0.71)			
	60	0.24 (0.16, 0.35)	0.47 (0.34, 0.65)	0.71 (0.52, 0.96)	0.78 (0.56, 1.20)	0.54 (0.41, 0.72)			
Hispanic ^b	70	0.27 (0.17, 0.43)	0.55 (0.36, 0.81)	- (-, -)	0.63 (0.41, 1.10)	0.50 (0.36, 0.71)			
	80	0.38 (0.20, 0.69)	- (-, -)	- (-, -)	0.51 (0.26, 1.14)	0.39 (0.22, 0.68)			
	0	0.01 (0.01, 0.01)	0.02 (0.02, 0.02)	0.04 (0.04, 0.05)	1.94 (1.86, 2.03)	0.74 (0.71, 0.78)			
	10	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	0.09 (0.08, 0.09)	1.95 (1.87, 2.04)	0.75 (0.71, 0.78)			
	20	0.02 (0.02, 0.03)	0.07 (0.07, 0.08)	0.18 (0.17, 0.19)	1.95 (1.87, 2.04)	0.75 (0.72, 0.79)			
	30	0.05 (0.05, 0.06)	0.16 (0.15, 0.17)	0.38 (0.36, 0.40)	1.94 (1.86, 2.03)	0.75 (0.72, 0.79)			
	40	0.11 (0.10, 0.12)	0.33 (0.32, 0.35)	0.70 (0.67, 0.73)	1.91 (1.83, 2.00)	0.75 (0.72, 0.79)			
	50	0.23 (0.21, 0.24)	0.60 (0.57, 0.63)	1.20 (1.15, 1.26)	1.84 (1.76, 1.94)	0.74 (0.71, 0.78)			
-	60	0.39 (0.37, 0.42)	1.03 (0.98, 1.08)	1.52 (1.45, 1.59)	1.70 (1.61, 1.79)	0.73 (0.69, 0.76)			
	70	0.71 (0.67, 0.76)	1.26 (1.19, 1.33)	- (-, -)	1.46 (1.37, 1.56)	0.67 (0.63, 0.71)			
	80	0.73 (0.66, 0.80)	- (-, -)	- (-, -)	1.00 (0.90, 1.11)	0.52 (0.48, 0.57)			

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XIX-13 - continued

NON-HODGKIN LYMPHOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.06 (0.06, 0.06)	2.19 (2.16, 2.22)	0.91 (0.90, 0.92)	
	10	0.02 (0.02, 0.02)	0.05 (0.05, 0.06)	0.12 (0.12, 0.13)	2.21 (2.18, 2.24)	0.92 (0.91, 0.93)	
	20	0.03 (0.03, 0.04)	0.11 (0.10, 0.11)	0.26 (0.25, 0.26)	2.20 (2.17, 2.23)	0.93 (0.92, 0.94)	
	30	0.07 (0.07, 0.08)	0.22 (0.22, 0.23)	0.51 (0.50, 0.53)	2.20 (2.17, 2.23)	0.93 (0.92, 0.94)	
	40	0.15 (0.15, 0.16)	0.45 (0.44, 0.46)	0.96 (0.94, 0.98)	2.16 (2.13, 2.19)	0.94 (0.93, 0.95)	
	50	0.31 (0.30, 0.32)	0.84 (0.82, 0.85)	1.53 (1.50, 1.55)	2.08 (2.05, 2.12)	0.94 (0.93, 0.95)	
	60	0.57 (0.56, 0.59)	1.32 (1.30, 1.35)	1.82 (1.79, 1.85)	1.92 (1.89, 1.96)	0.93 (0.92, 0.95)	
	70	0.89 (0.87, 0.91)	1.49 (1.46, 1.52)	- (- , -)	1.61 (1.58, 1.65)	0.87 (0.86, 0.88)	
	80	0.93 (0.90, 0.96)	- (- , -)	- (- , -)	1.11 (1.08, 1.15)	0.72 (0.70, 0.73)	
White	0	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.06 (0.06, 0.07)	2.30 (2.26, 2.33)	0.97 (0.96, 0.98)	
	10	0.02 (0.02, 0.02)	0.05 (0.05, 0.06)	0.12 (0.12, 0.13)	2.31 (2.28, 2.35)	0.98 (0.97, 0.99)	
	20	0.03 (0.03, 0.04)	0.10 (0.10, 0.11)	0.26 (0.25, 0.27)	2.31 (2.27, 2.34)	0.98 (0.97, 0.99)	
	30	0.07 (0.07, 0.08)	0.23 (0.22, 0.23)	0.53 (0.51, 0.54)	2.30 (2.27, 2.34)	0.99 (0.98, 1.00)	
	40	0.16 (0.15, 0.16)	0.46 (0.45, 0.47)	1.01 (0.99, 1.03)	2.27 (2.23, 2.30)	0.99 (0.98, 1.01)	
	50	0.32 (0.31, 0.33)	0.88 (0.86, 0.90)	1.61 (1.58, 1.64)	2.19 (2.15, 2.22)	1.00 (0.99, 1.01)	
	60	0.61 (0.59, 0.63)	1.39 (1.37, 1.42)	1.92 (1.88, 1.95)	2.02 (1.98, 2.05)	0.99 (0.98, 1.00)	
	70	0.94 (0.91, 0.96)	1.56 (1.52, 1.59)	- (- , -)	1.68 (1.64, 1.71)	0.92 (0.90, 0.93)	
	80	0.96 (0.93, 1.00)	- (- , -)	- (- , -)	1.15 (1.11, 1.19)	0.75 (0.74, 0.77)	
Black	0	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	0.07 (0.06, 0.08)	1.31 (1.24, 1.39)	0.47 (0.45, 0.49)	
	10	0.02 (0.01, 0.02)	0.06 (0.05, 0.08)	0.16 (0.14, 0.17)	1.34 (1.27, 1.42)	0.48 (0.46, 0.50)	
	20	0.05 (0.04, 0.06)	0.14 (0.12, 0.15)	0.31 (0.29, 0.33)	1.33 (1.26, 1.41)	0.48 (0.46, 0.51)	
	30	0.09 (0.08, 0.11)	0.27 (0.25, 0.29)	0.54 (0.51, 0.58)	1.32 (1.25, 1.40)	0.49 (0.46, 0.51)	
	40	0.18 (0.17, 0.20)	0.46 (0.43, 0.50)	0.79 (0.74, 0.84)	1.27 (1.19, 1.35)	0.48 (0.46, 0.51)	
	50	0.30 (0.27, 0.33)	0.64 (0.60, 0.69)	0.99 (0.92, 1.05)	1.16 (1.08, 1.24)	0.47 (0.45, 0.50)	
	60	0.40 (0.36, 0.44)	0.80 (0.73, 0.86)	0.97 (0.89, 1.05)	1.00 (0.92, 1.08)	0.45 (0.43, 0.48)	
	70	0.53 (0.47, 0.59)	0.76 (0.68, 0.84)	- (- , -)	0.79 (0.71, 0.89)	0.40 (0.37, 0.43)	
	80	0.42 (0.34, 0.51)	- (- , -)	- (- , -)	0.49 (0.40, 0.60)	0.30 (0.27, 0.34)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XIX-13 - continued

NON-HODGKIN LYMPHOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.01 (0.00, 0.01)	0.02 (0.02, 0.03)	0.05 (0.04, 0.06)	1.92 (1.79, 2.08)	0.86 (0.77, 0.97)	
	10	0.02 (0.01, 0.02)	0.05 (0.04, 0.06)	0.09 (0.08, 0.11)	1.94 (1.80, 2.09)	0.86 (0.77, 0.97)	
Islander	20	0.03 (0.02, 0.04)	0.07 (0.06, 0.09)	0.16 (0.14, 0.18)	1.93 (1.79, 2.08)	0.86 (0.77, 0.97)	
	30	0.05 (0.04, 0.05)	0.13 (0.11, 0.15)	0.31 (0.28, 0.34)	1.91 (1.77, 2.07)	0.86 (0.77, 0.97)	
	40	0.09 (0.07, 0.10)	0.27 (0.24, 0.30)	0.60 (0.55, 0.65)	1.88 (1.74, 2.04)	0.86 (0.77, 0.97)	
	50	0.18 (0.16, 0.21)	0.52 (0.48, 0.57)	1.10 (1.02, 1.18)	1.83 (1.69, 1.99)	0.86 (0.76, 0.97)	
	60	0.35 (0.31, 0.40)	0.95 (0.88, 1.04)	1.49 (1.38, 1.61)	1.72 (1.57, 1.88)	0.84 (0.74, 0.96)	
	70	0.67 (0.60, 0.74)	1.27 (1.16, 1.39)	- (-, -)	1.52 (1.37, 1.69)	0.80 (0.69, 0.92)	
	80	0.79 (0.69, 0.91)	- (-, -)	- (-, -)	1.12 (0.96, 1.33)	0.66 (0.54, 0.81)	
American	0	0.00 (0.00, 0.04)	0.01 (0.00, 0.05)	0.01 (0.00, 0.06)	0.85 (0.57, 1.57)	0.44 (0.34, 0.65)	
Indian/ Alaska Native ^a	10	0.01 (0.00, 0.03)	0.01 (0.00, 0.04)	0.05 (0.03, 0.11)	0.87 (0.58, 1.60)	0.45 (0.34, 0.66)	
	20	0.01 (0.00, 0.03)	0.05 (0.02, 0.10)	0.13 (0.08, 0.20)	0.87 (0.58, 1.61)	0.45 (0.35, 0.67)	
	30	0.04 (0.02, 0.09)	0.12 (0.07, 0.20)	0.30 (0.20, 0.42)	0.89 (0.59, 1.64)	0.46 (0.35, 0.68)	
	40	0.08 (0.05, 0.14)	0.26 (0.17, 0.38)	0.40 (0.27, 0.57)	0.87 (0.57, 1.66)	0.46 (0.35, 0.69)	
	50	0.19 (0.11, 0.30)	0.33 (0.21, 0.51)	0.50 (0.33, 0.75)	0.83 (0.52, 1.66)	0.47 (0.35, 0.71)	
	60	0.16 (0.07, 0.31)	0.34 (0.19, 0.59)	0.61 (0.35, 1.02)	0.71 (0.39, 1.61)	0.45 (0.32, 0.72)	
	70	0.22 (0.09, 0.46)	0.54 (0.26, 1.01)	- (-, -)	0.65 (0.31, 1.74)	0.34 (0.22, 0.66)	
	80	0.48 (0.18, 1.11)	- (-, -)	- (-, -)	0.66 (0.23, 2.29)	0.13 (0.04, 0.60)	
Hispanic ^b	0	0.01 (0.01, 0.01)	0.03 (0.02, 0.03)	0.05 (0.05, 0.06)	2.06 (1.93, 2.20)	0.76 (0.71, 0.82)	
	10	0.02 (0.01, 0.02)	0.04 (0.04, 0.05)	0.10 (0.10, 0.11)	2.07 (1.94, 2.22)	0.77 (0.72, 0.83)	
	20	0.03 (0.02, 0.03)	0.09 (0.08, 0.10)	0.22 (0.20, 0.23)	2.07 (1.94, 2.21)	0.77 (0.72, 0.83)	
	30	0.06 (0.05, 0.07)	0.19 (0.18, 0.21)	0.44 (0.41, 0.46)	2.06 (1.93, 2.21)	0.78 (0.73, 0.83)	
	40	0.13 (0.12, 0.15)	0.38 (0.36, 0.41)	0.76 (0.72, 0.81)	2.03 (1.90, 2.18)	0.78 (0.73, 0.83)	
	50	0.25 (0.23, 0.28)	0.65 (0.60, 0.70)	1.28 (1.20, 1.36)	1.96 (1.82, 2.11)	0.77 (0.72, 0.83)	
	60	0.42 (0.38, 0.46)	1.09 (1.02, 1.17)	1.61 (1.51, 1.73)	1.82 (1.68, 1.98)	0.75 (0.70, 0.81)	
	70	0.78 (0.71, 0.85)	1.38 (1.26, 1.50)	- (-, -)	1.61 (1.46, 1.79)	0.69 (0.63, 0.76)	
	80	0.85 (0.74, 0.97)	- (-, -)	- (-, -)	1.18 (1.00, 1.41)	0.53 (0.46, 0.62)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XIX-13 - continued

NON-HODGKIN LYMPHOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer						Risk of Dying from Cancer	
		+10 yrs		+20 yrs		+30 yrs		Eventually	Eventually
		Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)	Percent (95% C.I.)
All Races	0	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.04 (0.03, 0.04)	1.87 (1.85, 1.90)	0.77 (0.76, 0.78)			
	10	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	0.08 (0.08, 0.08)	1.89 (1.86, 1.92)	0.78 (0.77, 0.79)			
	20	0.02 (0.02, 0.02)	0.07 (0.07, 0.07)	0.17 (0.16, 0.17)	1.88 (1.86, 1.91)	0.78 (0.77, 0.79)			
	30	0.05 (0.04, 0.05)	0.15 (0.14, 0.15)	0.36 (0.36, 0.37)	1.87 (1.84, 1.90)	0.78 (0.77, 0.79)			
	40	0.10 (0.10, 0.10)	0.32 (0.31, 0.33)	0.74 (0.73, 0.76)	1.84 (1.81, 1.87)	0.78 (0.77, 0.79)			
	50	0.23 (0.22, 0.23)	0.66 (0.64, 0.67)	1.24 (1.22, 1.26)	1.78 (1.75, 1.81)	0.78 (0.77, 0.79)			
	60	0.45 (0.44, 0.47)	1.06 (1.04, 1.09)	1.52 (1.49, 1.55)	1.63 (1.60, 1.66)	0.77 (0.76, 0.78)			
	70	0.69 (0.67, 0.71)	1.20 (1.18, 1.23)	- (-, -)	1.33 (1.30, 1.35)	0.72 (0.71, 0.73)			
	80	0.69 (0.67, 0.71)	- (-, -)	- (-, -)	0.86 (0.83, 0.88)	0.57 (0.56, 0.58)			
White	0	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.04 (0.03, 0.04)	1.97 (1.94, 2.00)	0.81 (0.80, 0.82)			
	10	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.08 (0.07, 0.08)	1.98 (1.95, 2.02)	0.82 (0.81, 0.83)			
	20	0.02 (0.02, 0.02)	0.07 (0.06, 0.07)	0.17 (0.16, 0.17)	1.98 (1.95, 2.01)	0.82 (0.81, 0.83)			
	30	0.05 (0.04, 0.05)	0.14 (0.14, 0.15)	0.37 (0.36, 0.38)	1.97 (1.94, 2.00)	0.82 (0.81, 0.83)			
	40	0.10 (0.10, 0.11)	0.33 (0.32, 0.34)	0.78 (0.76, 0.80)	1.94 (1.91, 1.97)	0.82 (0.81, 0.83)			
	50	0.23 (0.22, 0.24)	0.69 (0.68, 0.71)	1.31 (1.29, 1.34)	1.87 (1.84, 1.91)	0.82 (0.81, 0.83)			
	60	0.48 (0.47, 0.50)	1.13 (1.11, 1.16)	1.61 (1.58, 1.64)	1.72 (1.69, 1.75)	0.81 (0.80, 0.82)			
	70	0.73 (0.71, 0.75)	1.26 (1.24, 1.29)	- (-, -)	1.39 (1.36, 1.42)	0.75 (0.74, 0.76)			
	80	0.72 (0.70, 0.74)	- (-, -)	- (-, -)	0.89 (0.86, 0.92)	0.60 (0.59, 0.61)			
Black	0	0.01 (0.00, 0.01)	0.02 (0.01, 0.02)	0.04 (0.03, 0.05)	1.11 (1.05, 1.18)	0.44 (0.42, 0.46)			
	10	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	0.10 (0.09, 0.11)	1.13 (1.06, 1.20)	0.45 (0.43, 0.47)			
	20	0.03 (0.02, 0.03)	0.09 (0.08, 0.10)	0.19 (0.18, 0.21)	1.12 (1.06, 1.19)	0.44 (0.42, 0.47)			
	30	0.06 (0.05, 0.07)	0.17 (0.15, 0.19)	0.35 (0.32, 0.37)	1.11 (1.04, 1.17)	0.44 (0.42, 0.46)			
	40	0.11 (0.10, 0.12)	0.29 (0.26, 0.31)	0.55 (0.51, 0.59)	1.06 (1.00, 1.13)	0.44 (0.42, 0.46)			
	50	0.19 (0.17, 0.21)	0.46 (0.43, 0.50)	0.78 (0.72, 0.83)	0.99 (0.93, 1.06)	0.44 (0.41, 0.46)			
	60	0.30 (0.27, 0.34)	0.64 (0.59, 0.70)	0.82 (0.76, 0.89)	0.88 (0.81, 0.95)	0.42 (0.40, 0.44)			
	70	0.41 (0.36, 0.46)	0.62 (0.56, 0.69)	- (-, -)	0.69 (0.63, 0.76)	0.37 (0.35, 0.40)			
	80	0.32 (0.27, 0.38)	- (-, -)	- (-, -)	0.42 (0.36, 0.49)	0.27 (0.25, 0.30)			

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XIX-13 - continued

NON-HODGKIN LYMPHOMA

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.03 (0.03, 0.04)	1.56 (1.44, 1.70)	0.73 (0.64, 0.83)	
	10	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.06 (0.05, 0.08)	1.57 (1.45, 1.71)	0.73 (0.64, 0.84)	
Islander	20	0.02 (0.01, 0.03)	0.05 (0.04, 0.06)	0.12 (0.10, 0.13)	1.56 (1.44, 1.70)	0.73 (0.64, 0.84)	
	30	0.03 (0.03, 0.04)	0.10 (0.09, 0.11)	0.26 (0.24, 0.29)	1.55 (1.42, 1.69)	0.73 (0.64, 0.84)	
	40	0.07 (0.06, 0.08)	0.23 (0.21, 0.25)	0.51 (0.47, 0.55)	1.52 (1.40, 1.66)	0.73 (0.64, 0.84)	
	50	0.17 (0.15, 0.19)	0.45 (0.41, 0.49)	0.87 (0.81, 0.94)	1.47 (1.35, 1.61)	0.72 (0.63, 0.83)	
	60	0.29 (0.26, 0.32)	0.73 (0.67, 0.79)	1.15 (1.06, 1.25)	1.34 (1.22, 1.49)	0.71 (0.62, 0.82)	
	70	0.47 (0.42, 0.52)	0.92 (0.84, 1.01)	- (-, -)	1.12 (1.00, 1.27)	0.67 (0.58, 0.79)	
	80	0.54 (0.47, 0.63)	- (-, -)	- (-, -)	0.78 (0.65, 0.95)	0.54 (0.43, 0.67)	
American Indian/ Alaska Native ^a	0	0.00 (0.00, 0.04)	0.01 (0.00, 0.06)	0.01 (0.00, 0.06)	0.98 (0.70, 1.55)	0.57 (0.41, 0.81)	
	10	0.01 (0.00, 0.03)	0.01 (0.00, 0.04)	0.03 (0.01, 0.07)	0.99 (0.71, 1.57)	0.58 (0.42, 0.83)	
	20	0.00 (0.00, 0.03)	0.02 (0.00, 0.05)	0.08 (0.04, 0.13)	0.99 (0.71, 1.57)	0.58 (0.42, 0.83)	
	30	0.02 (0.00, 0.05)	0.08 (0.04, 0.13)	0.22 (0.14, 0.32)	0.99 (0.71, 1.58)	0.59 (0.42, 0.84)	
	40	0.06 (0.03, 0.11)	0.20 (0.13, 0.31)	0.49 (0.34, 0.68)	1.00 (0.71, 1.60)	0.59 (0.42, 0.84)	
	50	0.15 (0.08, 0.24)	0.44 (0.30, 0.64)	0.71 (0.50, 0.99)	0.96 (0.67, 1.58)	0.59 (0.41, 0.86)	
	60	0.31 (0.19, 0.49)	0.59 (0.39, 0.87)	0.80 (0.54, 1.17)	0.86 (0.57, 1.51)	0.59 (0.41, 0.88)	
	70	0.32 (0.17, 0.56)	0.56 (0.32, 0.93)	- (-, -)	0.63 (0.36, 1.35)	0.60 (0.39, 0.93)	
	80	0.32 (0.11, 0.73)	- (-, -)	- (-, -)	0.42 (0.15, 1.34)	0.54 (0.29, 0.98)	
Hispanic ^b	0	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	0.03 (0.02, 0.03)	1.84 (1.74, 1.97)	0.72 (0.68, 0.77)	
	10	0.01 (0.01, 0.01)	0.02 (0.02, 0.03)	0.06 (0.06, 0.07)	1.86 (1.75, 1.98)	0.73 (0.68, 0.78)	
	20	0.02 (0.01, 0.02)	0.06 (0.05, 0.07)	0.14 (0.13, 0.15)	1.86 (1.75, 1.98)	0.73 (0.68, 0.78)	
	30	0.04 (0.03, 0.05)	0.12 (0.11, 0.14)	0.32 (0.30, 0.35)	1.85 (1.74, 1.97)	0.73 (0.68, 0.78)	
	40	0.08 (0.08, 0.09)	0.28 (0.26, 0.31)	0.63 (0.59, 0.68)	1.82 (1.71, 1.94)	0.73 (0.68, 0.78)	
	50	0.20 (0.18, 0.22)	0.56 (0.52, 0.60)	1.14 (1.07, 1.21)	1.76 (1.65, 1.88)	0.72 (0.67, 0.77)	
	60	0.37 (0.34, 0.40)	0.97 (0.91, 1.04)	1.44 (1.35, 1.54)	1.61 (1.50, 1.74)	0.70 (0.66, 0.76)	
	70	0.66 (0.61, 0.72)	1.17 (1.08, 1.27)	- (-, -)	1.36 (1.25, 1.49)	0.65 (0.60, 0.71)	
	80	0.65 (0.57, 0.73)	- (-, -)	- (-, -)	0.89 (0.77, 1.03)	0.51 (0.46, 0.58)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XX-9

CANCER OF THE ORAL CAVITY AND PHARYNX (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.02 (1.00, 1.03)	0.28 (0.28, 0.29)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	1.03 (1.01, 1.04)	0.29 (0.28, 0.29)	
	20	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.12 (0.12, 0.12)	1.03 (1.01, 1.04)	0.29 (0.28, 0.29)	
	30	0.02 (0.02, 0.03)	0.11 (0.11, 0.11)	0.31 (0.30, 0.31)	1.03 (1.02, 1.04)	0.29 (0.29, 0.29)	
	40	0.09 (0.09, 0.09)	0.29 (0.28, 0.29)	0.56 (0.56, 0.57)	1.02 (1.00, 1.03)	0.29 (0.29, 0.29)	
	50	0.21 (0.20, 0.21)	0.49 (0.48, 0.50)	0.76 (0.75, 0.77)	0.96 (0.94, 0.97)	0.28 (0.28, 0.29)	
	60	0.30 (0.30, 0.31)	0.59 (0.58, 0.60)	0.76 (0.74, 0.77)	0.80 (0.79, 0.82)	0.26 (0.25, 0.26)	
	70	0.33 (0.33, 0.34)	0.52 (0.51, 0.54)	- (- , -)	0.58 (0.56, 0.59)	0.21 (0.20, 0.21)	
	80	0.27 (0.26, 0.28)	- (- , -)	- (- , -)	0.35 (0.33, 0.36)	0.15 (0.14, 0.15)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.03 (1.02, 1.05)	0.28 (0.27, 0.28)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	1.04 (1.03, 1.06)	0.28 (0.28, 0.28)	
	20	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.11 (0.11, 0.12)	1.05 (1.03, 1.06)	0.28 (0.28, 0.28)	
	30	0.02 (0.02, 0.02)	0.11 (0.10, 0.11)	0.30 (0.30, 0.31)	1.05 (1.03, 1.06)	0.28 (0.28, 0.29)	
	40	0.09 (0.08, 0.09)	0.29 (0.28, 0.29)	0.57 (0.56, 0.58)	1.04 (1.02, 1.05)	0.28 (0.28, 0.29)	
	50	0.21 (0.20, 0.21)	0.50 (0.49, 0.51)	0.78 (0.77, 0.79)	0.98 (0.96, 0.99)	0.28 (0.27, 0.28)	
	60	0.31 (0.30, 0.32)	0.61 (0.60, 0.62)	0.78 (0.76, 0.79)	0.82 (0.81, 0.84)	0.25 (0.25, 0.26)	
	70	0.35 (0.34, 0.36)	0.54 (0.52, 0.55)	- (- , -)	0.59 (0.58, 0.60)	0.21 (0.20, 0.21)	
	80	0.27 (0.26, 0.29)	- (- , -)	- (- , -)	0.35 (0.33, 0.36)	0.15 (0.14, 0.15)	
Black	0	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.89 (0.85, 0.93)	0.33 (0.32, 0.34)	
	10	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.04 (0.03, 0.04)	0.91 (0.87, 0.96)	0.34 (0.32, 0.35)	
	20	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	0.13 (0.12, 0.14)	0.91 (0.87, 0.96)	0.34 (0.33, 0.35)	
	30	0.02 (0.02, 0.03)	0.13 (0.12, 0.14)	0.36 (0.34, 0.38)	0.92 (0.88, 0.96)	0.34 (0.33, 0.35)	
	40	0.10 (0.10, 0.11)	0.34 (0.32, 0.36)	0.60 (0.57, 0.63)	0.92 (0.87, 0.96)	0.34 (0.33, 0.36)	
	50	0.25 (0.23, 0.27)	0.52 (0.49, 0.55)	0.74 (0.70, 0.78)	0.86 (0.81, 0.90)	0.33 (0.32, 0.34)	
	60	0.31 (0.28, 0.33)	0.55 (0.51, 0.59)	0.65 (0.61, 0.70)	0.68 (0.64, 0.73)	0.28 (0.27, 0.30)	
	70	0.30 (0.28, 0.34)	0.44 (0.40, 0.48)	- (- , -)	0.47 (0.43, 0.51)	0.20 (0.18, 0.21)	
	80	0.21 (0.18, 0.25)	- (- , -)	- (- , -)	0.26 (0.22, 0.31)	0.13 (0.12, 0.15)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XX-9 - continued

CANCER OF THE ORAL CAVITY AND PHARYNX (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.88 (0.82, 0.95)	0.32 (0.28, 0.37)	
	10	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.05 (0.05, 0.06)	0.88 (0.83, 0.95)	0.33 (0.29, 0.37)	
Islander	20	0.01 (0.01, 0.02)	0.05 (0.04, 0.06)	0.13 (0.12, 0.14)	0.88 (0.82, 0.95)	0.33 (0.29, 0.37)	
	30	0.04 (0.03, 0.04)	0.12 (0.11, 0.13)	0.26 (0.24, 0.27)	0.87 (0.81, 0.94)	0.33 (0.29, 0.37)	
	40	0.08 (0.07, 0.09)	0.22 (0.20, 0.24)	0.42 (0.39, 0.45)	0.84 (0.78, 0.91)	0.32 (0.29, 0.37)	
	50	0.14 (0.13, 0.15)	0.34 (0.32, 0.37)	0.53 (0.50, 0.57)	0.77 (0.71, 0.84)	0.31 (0.27, 0.36)	
	60	0.21 (0.19, 0.23)	0.41 (0.37, 0.44)	0.58 (0.54, 0.63)	0.65 (0.59, 0.72)	0.29 (0.25, 0.34)	
	70	0.21 (0.19, 0.24)	0.40 (0.36, 0.45)	- (-, -)	0.48 (0.42, 0.55)	0.25 (0.21, 0.31)	
	80	0.23 (0.20, 0.28)	- (-, -)	- (-, -)	0.33 (0.27, 0.41)	0.21 (0.16, 0.27)	
American	0	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.61 (0.47, 0.91)	0.24 (0.18, 0.34)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.01 (0.00, 0.03)	0.62 (0.48, 0.93)	0.25 (0.19, 0.35)	
	20	0.00 (0.00, 0.01)	0.01 (0.00, 0.03)	0.08 (0.05, 0.12)	0.62 (0.48, 0.94)	0.25 (0.19, 0.35)	
	30	0.01 (0.00, 0.03)	0.08 (0.05, 0.12)	0.23 (0.17, 0.31)	0.63 (0.49, 0.95)	0.25 (0.19, 0.36)	
	40	0.07 (0.04, 0.10)	0.23 (0.17, 0.30)	0.44 (0.34, 0.56)	0.64 (0.49, 0.96)	0.26 (0.20, 0.37)	
	50	0.16 (0.11, 0.23)	0.38 (0.29, 0.51)	0.53 (0.40, 0.70)	0.59 (0.44, 0.93)	0.27 (0.20, 0.38)	
	60	0.24 (0.16, 0.35)	0.40 (0.28, 0.56)	0.43 (0.30, 0.62)	0.46 (0.31, 0.82)	0.24 (0.17, 0.36)	
	70	0.19 (0.10, 0.32)	0.22 (0.12, 0.40)	- (-, -)	0.26 (0.14, 0.67)	0.20 (0.13, 0.34)	
	80	0.05 (0.01, 0.22)	- (-, -)	- (-, -)	0.10 (0.01, 0.67)	0.12 (0.05, 0.31)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.68 (0.64, 0.74)	0.22 (0.20, 0.24)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.02, 0.02)	0.69 (0.64, 0.75)	0.22 (0.20, 0.24)	
	20	0.00 (0.00, 0.01)	0.02 (0.02, 0.02)	0.06 (0.06, 0.07)	0.69 (0.64, 0.75)	0.22 (0.20, 0.24)	
	30	0.01 (0.01, 0.02)	0.06 (0.05, 0.07)	0.16 (0.15, 0.17)	0.69 (0.64, 0.75)	0.22 (0.20, 0.24)	
	40	0.05 (0.04, 0.05)	0.15 (0.14, 0.16)	0.31 (0.29, 0.33)	0.69 (0.64, 0.74)	0.22 (0.20, 0.24)	
	50	0.10 (0.09, 0.11)	0.27 (0.25, 0.29)	0.45 (0.42, 0.49)	0.66 (0.60, 0.71)	0.22 (0.20, 0.24)	
	60	0.17 (0.16, 0.19)	0.37 (0.34, 0.40)	0.50 (0.46, 0.54)	0.58 (0.53, 0.64)	0.21 (0.19, 0.23)	
	70	0.22 (0.19, 0.24)	0.37 (0.33, 0.41)	- (-, -)	0.45 (0.40, 0.52)	0.18 (0.16, 0.20)	
	80	0.20 (0.16, 0.23)	- (-, -)	- (-, -)	0.31 (0.25, 0.39)	0.15 (0.12, 0.18)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XX-9 - continued

CANCER OF THE ORAL CAVITY AND PHARYNX (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	1.40 (1.38, 1.43)	0.39 (0.38, 0.39)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.04, 0.04)	1.42 (1.40, 1.45)	0.39 (0.39, 0.40)	
	20	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	0.16 (0.16, 0.17)	1.43 (1.40, 1.45)	0.39 (0.39, 0.40)	
	30	0.03 (0.03, 0.03)	0.16 (0.15, 0.16)	0.46 (0.45, 0.47)	1.44 (1.41, 1.46)	0.40 (0.39, 0.40)	
	40	0.13 (0.13, 0.14)	0.44 (0.43, 0.45)	0.84 (0.83, 0.86)	1.43 (1.41, 1.46)	0.40 (0.39, 0.41)	
	50	0.32 (0.31, 0.33)	0.74 (0.72, 0.76)	1.12 (1.09, 1.14)	1.35 (1.33, 1.38)	0.39 (0.39, 0.40)	
	60	0.45 (0.44, 0.47)	0.86 (0.84, 0.88)	1.06 (1.04, 1.09)	1.12 (1.09, 1.14)	0.35 (0.34, 0.35)	
	70	0.49 (0.47, 0.50)	0.73 (0.71, 0.75)	- (- , -)	0.79 (0.77, 0.82)	0.27 (0.26, 0.28)	
	80	0.38 (0.36, 0.40)	- (- , -)	- (- , -)	0.47 (0.45, 0.50)	0.19 (0.18, 0.20)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.43 (1.40, 1.46)	0.37 (0.36, 0.38)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	1.45 (1.42, 1.47)	0.38 (0.37, 0.38)	
	20	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	0.16 (0.15, 0.16)	1.45 (1.42, 1.48)	0.38 (0.37, 0.38)	
	30	0.03 (0.02, 0.03)	0.15 (0.15, 0.16)	0.46 (0.44, 0.47)	1.46 (1.43, 1.49)	0.38 (0.37, 0.39)	
	40	0.13 (0.12, 0.13)	0.44 (0.42, 0.45)	0.85 (0.83, 0.87)	1.46 (1.43, 1.49)	0.38 (0.38, 0.39)	
	50	0.32 (0.31, 0.33)	0.75 (0.73, 0.77)	1.13 (1.11, 1.16)	1.38 (1.35, 1.40)	0.38 (0.37, 0.38)	
	60	0.46 (0.45, 0.48)	0.88 (0.86, 0.90)	1.09 (1.06, 1.12)	1.14 (1.11, 1.17)	0.34 (0.33, 0.34)	
	70	0.50 (0.48, 0.52)	0.75 (0.72, 0.77)	- (- , -)	0.81 (0.78, 0.83)	0.27 (0.26, 0.27)	
	80	0.39 (0.37, 0.41)	- (- , -)	- (- , -)	0.48 (0.46, 0.51)	0.19 (0.18, 0.20)	
Black	0	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.01, 0.02)	1.30 (1.23, 1.37)	0.51 (0.49, 0.53)	
	10	0.01 (0.00, 0.01)	0.02 (0.01, 0.02)	0.05 (0.04, 0.06)	1.33 (1.26, 1.41)	0.52 (0.50, 0.55)	
	20	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.18 (0.16, 0.20)	1.34 (1.26, 1.41)	0.53 (0.51, 0.55)	
	30	0.03 (0.02, 0.04)	0.17 (0.16, 0.19)	0.55 (0.51, 0.59)	1.36 (1.29, 1.44)	0.54 (0.52, 0.56)	
	40	0.15 (0.14, 0.17)	0.54 (0.50, 0.57)	0.94 (0.89, 1.00)	1.37 (1.30, 1.45)	0.55 (0.53, 0.57)	
	50	0.41 (0.38, 0.45)	0.85 (0.79, 0.90)	1.17 (1.10, 1.25)	1.30 (1.23, 1.39)	0.53 (0.51, 0.56)	
	60	0.50 (0.46, 0.55)	0.88 (0.82, 0.96)	1.02 (0.94, 1.10)	1.03 (0.96, 1.12)	0.46 (0.43, 0.48)	
	70	0.51 (0.45, 0.57)	0.69 (0.61, 0.76)	- (- , -)	0.71 (0.63, 0.80)	0.31 (0.29, 0.34)	
	80	0.32 (0.26, 0.40)	- (- , -)	- (- , -)	0.37 (0.29, 0.47)	0.21 (0.18, 0.24)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XX-9 - continued

CANCER OF THE ORAL CAVITY AND PHARYNX (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	1.12 (1.03, 1.23)	0.43 (0.37, 0.51)	
	10	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.05 (0.05, 0.07)	1.13 (1.04, 1.25)	0.43 (0.37, 0.51)	
Islander	20	0.01 (0.01, 0.02)	0.05 (0.04, 0.06)	0.17 (0.16, 0.19)	1.14 (1.04, 1.25)	0.43 (0.38, 0.51)	
	30	0.04 (0.03, 0.05)	0.16 (0.15, 0.18)	0.37 (0.34, 0.40)	1.13 (1.04, 1.24)	0.44 (0.38, 0.52)	
	40	0.12 (0.11, 0.14)	0.33 (0.30, 0.36)	0.61 (0.56, 0.66)	1.10 (1.01, 1.21)	0.43 (0.37, 0.52)	
	50	0.21 (0.19, 0.24)	0.49 (0.45, 0.54)	0.76 (0.69, 0.82)	0.99 (0.90, 1.11)	0.42 (0.36, 0.50)	
	60	0.29 (0.26, 0.33)	0.57 (0.51, 0.63)	0.75 (0.68, 0.83)	0.82 (0.73, 0.93)	0.38 (0.32, 0.46)	
	70	0.31 (0.26, 0.36)	0.51 (0.44, 0.59)	- (-, -)	0.58 (0.49, 0.70)	0.33 (0.26, 0.42)	
	80	0.27 (0.21, 0.34)	- (-, -)	- (-, -)	0.37 (0.27, 0.50)	0.25 (0.17, 0.36)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.84 (0.59, 1.53)	0.32 (0.22, 0.53)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.02 (0.00, 0.05)	0.86 (0.61, 1.56)	0.32 (0.22, 0.54)	
	20	0.00 (0.00, 0.02)	0.02 (0.00, 0.05)	0.10 (0.06, 0.16)	0.87 (0.61, 1.57)	0.33 (0.22, 0.55)	
	30	0.02 (0.00, 0.05)	0.10 (0.06, 0.17)	0.34 (0.24, 0.47)	0.89 (0.63, 1.61)	0.33 (0.23, 0.56)	
	40	0.09 (0.05, 0.15)	0.34 (0.23, 0.47)	0.63 (0.47, 0.85)	0.90 (0.63, 1.65)	0.34 (0.23, 0.58)	
	50	0.26 (0.17, 0.38)	0.58 (0.41, 0.80)	0.76 (0.54, 1.06)	0.86 (0.58, 1.65)	0.36 (0.24, 0.61)	
	60	0.34 (0.21, 0.54)	0.55 (0.35, 0.84)	0.57 (0.36, 0.91)	0.65 (0.39, 1.52)	0.33 (0.21, 0.61)	
	70	0.25 (0.11, 0.49)	0.27 (0.13, 0.59)	- (-, -)	0.37 (0.14, 1.41)	0.28 (0.15, 0.62)	
	80	0.03 (0.00, 0.42)	- (-, -)	- (-, -)	0.19 (0.00, 1.80)	0.21 (0.06, 0.74)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.97 (0.88, 1.08)	0.32 (0.29, 0.36)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.02, 0.03)	0.98 (0.90, 1.09)	0.32 (0.29, 0.37)	
	20	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.09 (0.08, 0.10)	0.99 (0.90, 1.10)	0.33 (0.29, 0.37)	
	30	0.02 (0.01, 0.02)	0.08 (0.07, 0.09)	0.24 (0.22, 0.26)	0.99 (0.91, 1.10)	0.33 (0.29, 0.37)	
	40	0.07 (0.06, 0.07)	0.22 (0.20, 0.24)	0.48 (0.44, 0.52)	0.99 (0.90, 1.10)	0.33 (0.30, 0.37)	
	50	0.16 (0.14, 0.18)	0.43 (0.39, 0.47)	0.68 (0.63, 0.74)	0.95 (0.86, 1.07)	0.33 (0.29, 0.37)	
	60	0.28 (0.25, 0.32)	0.56 (0.51, 0.61)	0.72 (0.65, 0.79)	0.84 (0.75, 0.97)	0.31 (0.27, 0.36)	
	70	0.32 (0.27, 0.36)	0.50 (0.44, 0.58)	- (-, -)	0.65 (0.55, 0.78)	0.27 (0.23, 0.32)	
	80	0.27 (0.21, 0.34)	- (-, -)	- (-, -)	0.47 (0.35, 0.64)	0.22 (0.17, 0.29)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XX-9 - continued

CANCER OF THE ORAL CAVITY AND PHARYNX (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.65 (0.64, 0.67)	0.19 (0.18, 0.19)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.66 (0.64, 0.68)	0.19 (0.19, 0.19)	
	20	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.07 (0.07, 0.08)	0.66 (0.64, 0.67)	0.19 (0.19, 0.19)	
	30	0.02 (0.02, 0.02)	0.07 (0.06, 0.07)	0.16 (0.15, 0.16)	0.65 (0.64, 0.67)	0.19 (0.19, 0.19)	
	40	0.05 (0.04, 0.05)	0.14 (0.13, 0.15)	0.30 (0.29, 0.30)	0.64 (0.62, 0.66)	0.19 (0.19, 0.19)	
	50	0.10 (0.09, 0.10)	0.25 (0.24, 0.26)	0.44 (0.42, 0.45)	0.60 (0.59, 0.62)	0.19 (0.18, 0.19)	
	60	0.17 (0.16, 0.17)	0.36 (0.34, 0.37)	0.49 (0.48, 0.50)	0.53 (0.52, 0.55)	0.18 (0.17, 0.18)	
	70	0.21 (0.20, 0.22)	0.36 (0.35, 0.38)	- (-, -)	0.41 (0.40, 0.43)	0.16 (0.15, 0.16)	
	80	0.20 (0.19, 0.21)	- (-, -)	- (-, -)	0.27 (0.25, 0.28)	0.12 (0.12, 0.13)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.66 (0.64, 0.68)	0.19 (0.18, 0.19)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.02, 0.03)	0.67 (0.65, 0.68)	0.19 (0.19, 0.20)	
	20	0.01 (0.01, 0.01)	0.03 (0.02, 0.03)	0.07 (0.06, 0.07)	0.66 (0.65, 0.68)	0.19 (0.19, 0.20)	
	30	0.02 (0.01, 0.02)	0.06 (0.06, 0.06)	0.15 (0.14, 0.16)	0.66 (0.64, 0.68)	0.19 (0.19, 0.20)	
	40	0.04 (0.04, 0.05)	0.14 (0.13, 0.14)	0.29 (0.28, 0.31)	0.65 (0.63, 0.67)	0.19 (0.19, 0.20)	
	50	0.09 (0.09, 0.10)	0.26 (0.25, 0.27)	0.45 (0.43, 0.46)	0.62 (0.60, 0.63)	0.19 (0.18, 0.19)	
	60	0.17 (0.16, 0.18)	0.37 (0.36, 0.38)	0.50 (0.49, 0.52)	0.55 (0.53, 0.56)	0.18 (0.18, 0.19)	
	70	0.22 (0.21, 0.23)	0.37 (0.36, 0.39)	- (-, -)	0.42 (0.41, 0.44)	0.16 (0.15, 0.16)	
	80	0.20 (0.19, 0.21)	- (-, -)	- (-, -)	0.27 (0.25, 0.28)	0.12 (0.12, 0.13)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.52 (0.48, 0.57)	0.16 (0.15, 0.18)	
	10	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.03 (0.02, 0.04)	0.53 (0.49, 0.58)	0.17 (0.16, 0.18)	
	20	0.01 (0.01, 0.01)	0.03 (0.02, 0.03)	0.09 (0.08, 0.10)	0.53 (0.49, 0.58)	0.17 (0.16, 0.18)	
	30	0.02 (0.01, 0.02)	0.08 (0.07, 0.09)	0.19 (0.17, 0.21)	0.53 (0.48, 0.57)	0.17 (0.16, 0.18)	
	40	0.06 (0.05, 0.07)	0.17 (0.15, 0.19)	0.30 (0.27, 0.33)	0.52 (0.47, 0.56)	0.17 (0.16, 0.18)	
	50	0.11 (0.10, 0.13)	0.24 (0.22, 0.27)	0.37 (0.33, 0.40)	0.47 (0.43, 0.52)	0.16 (0.15, 0.17)	
	60	0.14 (0.12, 0.17)	0.28 (0.24, 0.31)	0.36 (0.32, 0.41)	0.39 (0.35, 0.44)	0.15 (0.13, 0.16)	
	70	0.16 (0.13, 0.19)	0.26 (0.22, 0.30)	- (-, -)	0.30 (0.25, 0.35)	0.12 (0.11, 0.13)	
	80	0.15 (0.12, 0.19)	- (-, -)	- (-, -)	0.20 (0.16, 0.26)	0.09 (0.08, 0.11)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XX-9 - continued

CANCER OF THE ORAL CAVITY AND PHARYNX (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.01, 0.03)	0.66 (0.59, 0.76)	0.23 (0.18, 0.30)	
	10	0.01 (0.00, 0.01)	0.02 (0.01, 0.03)	0.05 (0.04, 0.06)	0.67 (0.59, 0.76)	0.23 (0.18, 0.30)	
Islander	20	0.02 (0.01, 0.02)	0.05 (0.04, 0.06)	0.09 (0.08, 0.11)	0.66 (0.59, 0.76)	0.23 (0.18, 0.30)	
	30	0.03 (0.02, 0.04)	0.08 (0.07, 0.09)	0.15 (0.13, 0.17)	0.65 (0.57, 0.75)	0.23 (0.18, 0.30)	
	40	0.05 (0.04, 0.06)	0.12 (0.11, 0.14)	0.26 (0.23, 0.29)	0.62 (0.55, 0.72)	0.23 (0.18, 0.30)	
	50	0.08 (0.06, 0.09)	0.21 (0.18, 0.24)	0.35 (0.31, 0.39)	0.58 (0.51, 0.68)	0.23 (0.18, 0.30)	
	60	0.14 (0.11, 0.16)	0.27 (0.24, 0.31)	0.44 (0.39, 0.50)	0.52 (0.44, 0.62)	0.21 (0.16, 0.28)	
	70	0.15 (0.12, 0.18)	0.32 (0.27, 0.38)	- (-, -)	0.40 (0.33, 0.51)	0.19 (0.14, 0.27)	
	80	0.21 (0.16, 0.27)	- (-, -)	- (-, -)	0.31 (0.23, 0.42)	0.18 (0.12, 0.27)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.39 (0.24, 0.89)	0.18 (0.12, 0.33)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.40 (0.25, 0.90)	0.18 (0.12, 0.33)	
	20	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.06 (0.03, 0.11)	0.40 (0.25, 0.91)	0.18 (0.12, 0.34)	
	30	0.01 (0.00, 0.04)	0.06 (0.03, 0.11)	0.13 (0.07, 0.21)	0.41 (0.25, 0.92)	0.18 (0.12, 0.34)	
	40	0.05 (0.02, 0.10)	0.12 (0.07, 0.21)	0.25 (0.15, 0.40)	0.40 (0.25, 0.92)	0.19 (0.12, 0.35)	
	50	0.07 (0.03, 0.14)	0.21 (0.11, 0.35)	0.32 (0.19, 0.53)	0.36 (0.21, 0.90)	0.19 (0.12, 0.36)	
	60	0.14 (0.06, 0.28)	0.26 (0.14, 0.47)	0.31 (0.16, 0.57)	0.31 (0.16, 0.87)	0.17 (0.10, 0.35)	
	70	0.14 (0.04, 0.33)	0.19 (0.06, 0.46)	- (-, -)	0.19 (0.06, 0.84)	0.14 (0.07, 0.34)	
	80	0.07 (0.01, 0.35)	- (-, -)	- (-, -)	0.07 (0.01, 0.93)	0.07 (0.02, 0.33)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.43 (0.38, 0.50)	0.13 (0.11, 0.15)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.02 (0.01, 0.02)	0.44 (0.38, 0.51)	0.13 (0.11, 0.16)	
	20	0.01 (0.00, 0.01)	0.02 (0.01, 0.02)	0.04 (0.03, 0.05)	0.44 (0.38, 0.51)	0.13 (0.11, 0.16)	
	30	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	0.09 (0.07, 0.10)	0.43 (0.38, 0.50)	0.13 (0.11, 0.16)	
	40	0.03 (0.02, 0.03)	0.08 (0.07, 0.09)	0.15 (0.13, 0.17)	0.43 (0.37, 0.50)	0.13 (0.11, 0.16)	
	50	0.05 (0.04, 0.06)	0.13 (0.11, 0.15)	0.25 (0.22, 0.29)	0.41 (0.35, 0.48)	0.13 (0.11, 0.15)	
	60	0.08 (0.06, 0.10)	0.21 (0.18, 0.24)	0.32 (0.28, 0.37)	0.37 (0.31, 0.44)	0.13 (0.10, 0.15)	
	70	0.14 (0.12, 0.17)	0.26 (0.22, 0.31)	- (-, -)	0.32 (0.26, 0.39)	0.11 (0.09, 0.14)	
	80	0.15 (0.11, 0.20)	- (-, -)	- (-, -)	0.22 (0.16, 0.31)	0.10 (0.08, 0.13)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXI-7
OVARY CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	1.42 (1.39, 1.44)	1.05 (1.04, 1.06)	
	10	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.07 (0.07, 0.08)	1.43 (1.41, 1.46)	1.07 (1.06, 1.08)	
	20	0.02 (0.02, 0.02)	0.06 (0.06, 0.06)	0.18 (0.17, 0.18)	1.43 (1.40, 1.45)	1.07 (1.06, 1.08)	
	30	0.04 (0.04, 0.04)	0.16 (0.15, 0.16)	0.39 (0.38, 0.40)	1.41 (1.39, 1.44)	1.07 (1.06, 1.08)	
	40	0.12 (0.12, 0.12)	0.36 (0.35, 0.37)	0.70 (0.69, 0.72)	1.38 (1.36, 1.41)	1.07 (1.06, 1.08)	
	50	0.24 (0.23, 0.25)	0.60 (0.58, 0.61)	0.98 (0.96, 1.00)	1.29 (1.27, 1.32)	1.05 (1.04, 1.06)	
	60	0.37 (0.36, 0.38)	0.77 (0.75, 0.79)	1.04 (1.01, 1.06)	1.10 (1.08, 1.12)	0.97 (0.95, 0.98)	
	70	0.45 (0.43, 0.46)	0.74 (0.72, 0.76)	- (- , -)	0.82 (0.80, 0.84)	0.79 (0.78, 0.80)	
	80	0.40 (0.38, 0.41)	- (- , -)	- (- , -)	0.49 (0.48, 0.51)	0.53 (0.52, 0.54)	
White	0	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	1.50 (1.47, 1.52)	1.10 (1.09, 1.11)	
	10	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.07 (0.07, 0.08)	1.51 (1.48, 1.54)	1.11 (1.10, 1.13)	
	20	0.02 (0.02, 0.02)	0.06 (0.06, 0.06)	0.18 (0.18, 0.19)	1.50 (1.48, 1.53)	1.12 (1.11, 1.13)	
	30	0.04 (0.04, 0.04)	0.16 (0.16, 0.17)	0.42 (0.40, 0.43)	1.49 (1.46, 1.52)	1.12 (1.11, 1.13)	
	40	0.13 (0.12, 0.13)	0.38 (0.37, 0.39)	0.75 (0.74, 0.77)	1.46 (1.43, 1.49)	1.12 (1.11, 1.13)	
	50	0.26 (0.25, 0.27)	0.64 (0.62, 0.66)	1.04 (1.02, 1.07)	1.36 (1.34, 1.39)	1.09 (1.08, 1.11)	
	60	0.40 (0.39, 0.41)	0.82 (0.80, 0.84)	1.09 (1.07, 1.12)	1.15 (1.13, 1.18)	1.00 (0.99, 1.01)	
	70	0.47 (0.46, 0.49)	0.78 (0.76, 0.80)	- (- , -)	0.85 (0.82, 0.87)	0.81 (0.80, 0.82)	
	80	0.41 (0.39, 0.42)	- (- , -)	- (- , -)	0.50 (0.48, 0.52)	0.54 (0.53, 0.55)	
Black	0	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.02 (0.02, 0.03)	0.99 (0.92, 1.05)	0.77 (0.74, 0.80)	
	10	0.01 (0.01, 0.01)	0.02 (0.02, 0.03)	0.06 (0.05, 0.07)	1.01 (0.94, 1.07)	0.79 (0.76, 0.81)	
	20	0.01 (0.01, 0.02)	0.05 (0.04, 0.06)	0.13 (0.12, 0.15)	1.00 (0.94, 1.07)	0.79 (0.76, 0.82)	
	30	0.04 (0.03, 0.05)	0.12 (0.10, 0.13)	0.27 (0.25, 0.29)	0.99 (0.93, 1.06)	0.79 (0.76, 0.82)	
	40	0.08 (0.07, 0.09)	0.24 (0.21, 0.26)	0.47 (0.43, 0.51)	0.97 (0.91, 1.04)	0.79 (0.77, 0.82)	
	50	0.16 (0.14, 0.18)	0.41 (0.37, 0.44)	0.66 (0.61, 0.71)	0.93 (0.86, 1.00)	0.79 (0.76, 0.82)	
	60	0.27 (0.24, 0.30)	0.55 (0.50, 0.60)	0.77 (0.71, 0.83)	0.84 (0.77, 0.91)	0.74 (0.71, 0.77)	
	70	0.33 (0.29, 0.38)	0.60 (0.54, 0.66)	- (- , -)	0.68 (0.61, 0.75)	0.62 (0.59, 0.65)	
	80	0.39 (0.34, 0.45)	- (- , -)	- (- , -)	0.51 (0.44, 0.59)	0.44 (0.41, 0.47)	

Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXI-7 - continued
OVARY CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/Pacific Islander	0	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	1.13 (1.04, 1.23)	0.68 (0.62, 0.76)	
	10	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.08 (0.07, 0.10)	1.13 (1.05, 1.24)	0.68 (0.62, 0.77)	
	20	0.03 (0.02, 0.04)	0.07 (0.06, 0.09)	0.19 (0.17, 0.21)	1.13 (1.04, 1.23)	0.69 (0.62, 0.77)	
	30	0.05 (0.04, 0.06)	0.16 (0.15, 0.18)	0.34 (0.32, 0.37)	1.10 (1.02, 1.21)	0.69 (0.62, 0.77)	
	40	0.12 (0.10, 0.13)	0.30 (0.27, 0.33)	0.54 (0.50, 0.59)	1.06 (0.97, 1.17)	0.68 (0.62, 0.76)	
	50	0.18 (0.16, 0.21)	0.43 (0.39, 0.47)	0.70 (0.64, 0.76)	0.95 (0.87, 1.06)	0.65 (0.59, 0.73)	
	60	0.25 (0.22, 0.29)	0.53 (0.48, 0.58)	0.72 (0.65, 0.79)	0.79 (0.71, 0.90)	0.58 (0.52, 0.66)	
	70	0.30 (0.26, 0.34)	0.50 (0.44, 0.57)	- (-, -)	0.57 (0.49, 0.68)	0.46 (0.40, 0.54)	
	80	0.24 (0.20, 0.30)	- (-, -)	- (-, -)	0.33 (0.25, 0.45)	0.30 (0.24, 0.39)	
American Indian/Alaska Native ^a	0	0.00 (0.00, 0.04)	0.01 (0.00, 0.05)	0.01 (0.00, 0.06)	0.94 (0.69, 1.49)	0.77 (0.60, 1.00)	
	10	0.01 (0.00, 0.03)	0.01 (0.00, 0.05)	0.03 (0.01, 0.07)	0.95 (0.70, 1.51)	0.78 (0.61, 1.02)	
	20	0.01 (0.00, 0.04)	0.03 (0.01, 0.06)	0.13 (0.08, 0.20)	0.95 (0.70, 1.51)	0.78 (0.62, 1.02)	
	30	0.02 (0.01, 0.05)	0.12 (0.07, 0.19)	0.35 (0.25, 0.48)	0.96 (0.70, 1.52)	0.79 (0.62, 1.03)	
	40	0.10 (0.06, 0.16)	0.34 (0.24, 0.46)	0.59 (0.44, 0.80)	0.95 (0.69, 1.52)	0.79 (0.62, 1.04)	
	50	0.24 (0.16, 0.35)	0.51 (0.35, 0.71)	0.68 (0.48, 0.94)	0.87 (0.62, 1.46)	0.77 (0.59, 1.03)	
	60	0.28 (0.17, 0.45)	0.46 (0.29, 0.71)	0.66 (0.43, 1.02)	0.67 (0.43, 1.29)	0.69 (0.51, 0.96)	
	70	0.21 (0.09, 0.42)	0.44 (0.22, 0.80)	- (-, -)	0.45 (0.22, 1.14)	0.62 (0.42, 0.93)	
	80	0.30 (0.10, 0.71)	- (-, -)	- (-, -)	0.32 (0.11, 1.21)	0.50 (0.27, 0.90)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.04 (0.03, 0.04)	1.34 (1.25, 1.43)	0.82 (0.77, 0.86)	
	10	0.01 (0.01, 0.02)	0.04 (0.03, 0.04)	0.07 (0.06, 0.07)	1.35 (1.26, 1.44)	0.82 (0.78, 0.87)	
	20	0.02 (0.02, 0.03)	0.05 (0.04, 0.06)	0.15 (0.14, 0.17)	1.34 (1.25, 1.43)	0.82 (0.78, 0.87)	
	30	0.03 (0.03, 0.04)	0.13 (0.12, 0.14)	0.33 (0.31, 0.36)	1.32 (1.24, 1.42)	0.83 (0.78, 0.88)	
	40	0.10 (0.09, 0.11)	0.30 (0.28, 0.33)	0.59 (0.55, 0.63)	1.30 (1.21, 1.40)	0.82 (0.78, 0.87)	
	50	0.21 (0.19, 0.23)	0.49 (0.45, 0.53)	0.87 (0.82, 0.93)	1.21 (1.13, 1.31)	0.80 (0.75, 0.85)	
	60	0.29 (0.27, 0.33)	0.69 (0.64, 0.75)	0.97 (0.90, 1.05)	1.04 (0.96, 1.14)	0.74 (0.69, 0.79)	
	70	0.43 (0.39, 0.48)	0.74 (0.67, 0.82)	- (-, -)	0.82 (0.74, 0.92)	0.63 (0.58, 0.68)	
	80	0.39 (0.33, 0.46)	- (-, -)	- (-, -)	0.49 (0.41, 0.59)	0.44 (0.39, 0.49)	

Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXII-9
PANCREAS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.31 (1.29, 1.33)	1.20 (1.19, 1.21)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.32 (1.31, 1.34)	1.22 (1.21, 1.22)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.05, 0.05)	1.33 (1.31, 1.35)	1.22 (1.21, 1.23)	
	30	0.01 (0.01, 0.01)	0.05 (0.04, 0.05)	0.18 (0.17, 0.18)	1.34 (1.32, 1.36)	1.23 (1.22, 1.24)	
	40	0.04 (0.04, 0.04)	0.17 (0.17, 0.18)	0.47 (0.46, 0.48)	1.35 (1.33, 1.37)	1.24 (1.24, 1.25)	
	50	0.14 (0.13, 0.14)	0.45 (0.44, 0.46)	0.90 (0.89, 0.92)	1.35 (1.33, 1.37)	1.25 (1.24, 1.26)	
	60	0.33 (0.32, 0.34)	0.82 (0.80, 0.83)	1.18 (1.16, 1.20)	1.29 (1.27, 1.31)	1.21 (1.20, 1.22)	
	70	0.56 (0.55, 0.57)	0.98 (0.96, 1.00)	- (- , -)	1.10 (1.09, 1.12)	1.07 (1.06, 1.08)	
	80	0.60 (0.58, 0.61)	- (- , -)	- (- , -)	0.77 (0.76, 0.79)	0.79 (0.79, 0.80)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.30 (1.28, 1.32)	1.20 (1.19, 1.20)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.31 (1.29, 1.33)	1.21 (1.20, 1.22)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.04, 0.05)	1.32 (1.30, 1.34)	1.22 (1.21, 1.22)	
	30	0.01 (0.01, 0.01)	0.05 (0.04, 0.05)	0.17 (0.17, 0.18)	1.33 (1.31, 1.35)	1.23 (1.22, 1.23)	
	40	0.04 (0.04, 0.04)	0.17 (0.16, 0.17)	0.47 (0.46, 0.48)	1.34 (1.32, 1.35)	1.24 (1.23, 1.24)	
	50	0.13 (0.13, 0.14)	0.44 (0.43, 0.45)	0.90 (0.88, 0.91)	1.33 (1.31, 1.35)	1.24 (1.23, 1.25)	
	60	0.33 (0.32, 0.33)	0.81 (0.80, 0.83)	1.17 (1.15, 1.19)	1.27 (1.25, 1.29)	1.20 (1.19, 1.21)	
	70	0.56 (0.55, 0.57)	0.97 (0.95, 0.99)	- (- , -)	1.09 (1.07, 1.11)	1.06 (1.05, 1.06)	
	80	0.59 (0.57, 0.60)	- (- , -)	- (- , -)	0.75 (0.73, 0.77)	0.78 (0.77, 0.79)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.40 (1.34, 1.46)	1.26 (1.24, 1.29)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.43 (1.38, 1.49)	1.29 (1.27, 1.32)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.06 (0.06, 0.07)	1.44 (1.38, 1.50)	1.30 (1.28, 1.33)	
	30	0.01 (0.01, 0.01)	0.06 (0.06, 0.07)	0.24 (0.22, 0.25)	1.46 (1.41, 1.53)	1.32 (1.29, 1.35)	
	40	0.06 (0.05, 0.06)	0.23 (0.22, 0.25)	0.61 (0.58, 0.64)	1.49 (1.43, 1.55)	1.35 (1.32, 1.37)	
	50	0.19 (0.17, 0.20)	0.58 (0.55, 0.62)	1.08 (1.03, 1.13)	1.51 (1.45, 1.58)	1.37 (1.34, 1.40)	
	60	0.44 (0.41, 0.47)	0.99 (0.95, 1.05)	1.35 (1.29, 1.41)	1.48 (1.42, 1.55)	1.34 (1.31, 1.37)	
	70	0.69 (0.65, 0.74)	1.14 (1.07, 1.20)	- (- , -)	1.30 (1.23, 1.38)	1.21 (1.17, 1.24)	
	80	0.71 (0.65, 0.78)	- (- , -)	- (- , -)	0.98 (0.90, 1.07)	0.92 (0.88, 0.96)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXII-9 - continued
PANCREAS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.34 (1.26, 1.44)	1.21 (1.14, 1.29)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.35 (1.27, 1.45)	1.22 (1.15, 1.30)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	1.36 (1.27, 1.45)	1.23 (1.15, 1.31)	
	30	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	0.13 (0.12, 0.15)	1.36 (1.28, 1.46)	1.23 (1.16, 1.31)	
	40	0.03 (0.02, 0.03)	0.13 (0.11, 0.14)	0.36 (0.34, 0.39)	1.36 (1.28, 1.46)	1.23 (1.16, 1.31)	
	50	0.10 (0.09, 0.11)	0.34 (0.31, 0.37)	0.74 (0.69, 0.78)	1.35 (1.27, 1.45)	1.23 (1.15, 1.31)	
	60	0.25 (0.23, 0.27)	0.66 (0.62, 0.70)	1.10 (1.03, 1.17)	1.30 (1.21, 1.40)	1.19 (1.11, 1.27)	
	70	0.45 (0.41, 0.48)	0.92 (0.86, 0.99)	- (-, -)	1.14 (1.05, 1.24)	1.07 (1.00, 1.16)	
	80	0.59 (0.53, 0.66)	- (-, -)	- (-, -)	0.86 (0.76, 0.98)	0.84 (0.76, 0.95)	
American	0	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.96 (0.75, 1.32)	0.80 (0.68, 0.96)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.01)	0.00 (0.00, 0.02)	0.02 (0.01, 0.04)	0.98 (0.76, 1.34)	0.81 (0.69, 0.98)	
	20	0.00 (0.00, 0.01)	0.02 (0.01, 0.04)	0.06 (0.03, 0.09)	0.98 (0.76, 1.35)	0.82 (0.70, 0.99)	
	30	0.01 (0.00, 0.04)	0.06 (0.03, 0.09)	0.13 (0.09, 0.18)	1.00 (0.78, 1.37)	0.84 (0.71, 1.00)	
	40	0.04 (0.02, 0.07)	0.12 (0.08, 0.17)	0.38 (0.28, 0.50)	1.01 (0.78, 1.40)	0.86 (0.72, 1.03)	
	50	0.08 (0.04, 0.12)	0.35 (0.25, 0.47)	0.71 (0.54, 0.91)	1.00 (0.77, 1.41)	0.87 (0.74, 1.05)	
	60	0.29 (0.20, 0.41)	0.68 (0.51, 0.89)	0.95 (0.72, 1.24)	1.00 (0.75, 1.43)	0.85 (0.70, 1.05)	
	70	0.45 (0.31, 0.65)	0.77 (0.54, 1.07)	- (-, -)	0.83 (0.58, 1.31)	0.74 (0.57, 0.96)	
	80	0.44 (0.24, 0.77)	- (-, -)	- (-, -)	0.52 (0.28, 1.14)	0.58 (0.38, 0.87)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.46 (1.38, 1.55)	1.13 (1.09, 1.18)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.48 (1.40, 1.57)	1.15 (1.11, 1.19)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	1.48 (1.40, 1.57)	1.15 (1.11, 1.20)	
	30	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	0.15 (0.14, 0.16)	1.49 (1.41, 1.58)	1.16 (1.12, 1.21)	
	40	0.03 (0.02, 0.03)	0.14 (0.13, 0.15)	0.43 (0.40, 0.45)	1.50 (1.42, 1.59)	1.17 (1.13, 1.22)	
	50	0.12 (0.11, 0.13)	0.41 (0.38, 0.43)	0.89 (0.84, 0.93)	1.51 (1.42, 1.60)	1.18 (1.13, 1.22)	
	60	0.31 (0.28, 0.33)	0.81 (0.76, 0.85)	1.22 (1.16, 1.29)	1.46 (1.37, 1.56)	1.15 (1.10, 1.19)	
	70	0.56 (0.52, 0.60)	1.02 (0.96, 1.09)	- (-, -)	1.29 (1.20, 1.40)	1.03 (0.98, 1.08)	
	80	0.61 (0.55, 0.68)	- (-, -)	- (-, -)	0.97 (0.86, 1.09)	0.79 (0.74, 0.85)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXII-9 - continued
PANCREAS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.30 (1.27, 1.32)	1.20 (1.19, 1.21)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.31 (1.29, 1.34)	1.22 (1.21, 1.23)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.05, 0.06)	1.32 (1.30, 1.35)	1.23 (1.22, 1.24)	
	30	0.01 (0.01, 0.01)	0.05 (0.05, 0.06)	0.21 (0.20, 0.22)	1.34 (1.31, 1.36)	1.24 (1.23, 1.25)	
	40	0.05 (0.04, 0.05)	0.20 (0.20, 0.21)	0.54 (0.52, 0.55)	1.35 (1.33, 1.38)	1.26 (1.25, 1.27)	
	50	0.16 (0.16, 0.17)	0.51 (0.50, 0.52)	0.99 (0.97, 1.01)	1.35 (1.33, 1.38)	1.27 (1.26, 1.28)	
	60	0.37 (0.36, 0.39)	0.89 (0.87, 0.91)	1.21 (1.18, 1.24)	1.28 (1.26, 1.31)	1.22 (1.21, 1.23)	
	70	0.61 (0.60, 0.63)	1.00 (0.97, 1.02)	- (-, -)	1.08 (1.06, 1.11)	1.06 (1.05, 1.07)	
	80	0.59 (0.57, 0.61)	- (-, -)	- (-, -)	0.73 (0.70, 0.76)	0.77 (0.76, 0.79)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.30 (1.28, 1.33)	1.21 (1.20, 1.22)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.32 (1.29, 1.35)	1.23 (1.21, 1.24)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.05, 0.06)	1.32 (1.30, 1.35)	1.23 (1.22, 1.25)	
	30	0.01 (0.01, 0.01)	0.05 (0.05, 0.06)	0.21 (0.20, 0.22)	1.34 (1.31, 1.37)	1.25 (1.24, 1.26)	
	40	0.05 (0.04, 0.05)	0.20 (0.19, 0.21)	0.53 (0.52, 0.55)	1.35 (1.32, 1.38)	1.26 (1.25, 1.28)	
	50	0.16 (0.16, 0.17)	0.51 (0.49, 0.52)	0.99 (0.97, 1.01)	1.35 (1.32, 1.38)	1.27 (1.26, 1.28)	
	60	0.37 (0.36, 0.38)	0.89 (0.87, 0.91)	1.21 (1.18, 1.24)	1.28 (1.25, 1.31)	1.22 (1.21, 1.23)	
	70	0.62 (0.60, 0.64)	1.00 (0.97, 1.02)	- (-, -)	1.08 (1.05, 1.11)	1.06 (1.04, 1.07)	
	80	0.59 (0.57, 0.61)	- (-, -)	- (-, -)	0.72 (0.69, 0.75)	0.77 (0.75, 0.79)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.25 (1.17, 1.33)	1.18 (1.15, 1.21)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	1.28 (1.20, 1.36)	1.21 (1.18, 1.25)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.07 (0.06, 0.08)	1.29 (1.21, 1.37)	1.22 (1.19, 1.26)	
	30	0.01 (0.01, 0.02)	0.07 (0.06, 0.08)	0.28 (0.26, 0.31)	1.32 (1.24, 1.41)	1.25 (1.21, 1.29)	
	40	0.06 (0.05, 0.07)	0.28 (0.25, 0.31)	0.68 (0.63, 0.73)	1.35 (1.27, 1.44)	1.28 (1.24, 1.32)	
	50	0.23 (0.21, 0.26)	0.66 (0.61, 0.71)	1.12 (1.05, 1.20)	1.38 (1.30, 1.47)	1.31 (1.27, 1.35)	
	60	0.50 (0.45, 0.55)	1.03 (0.95, 1.11)	1.27 (1.18, 1.36)	1.33 (1.23, 1.43)	1.26 (1.22, 1.31)	
	70	0.71 (0.64, 0.78)	1.02 (0.93, 1.12)	- (-, -)	1.11 (1.01, 1.22)	1.11 (1.06, 1.16)	
	80	0.58 (0.49, 0.68)	- (-, -)	- (-, -)	0.73 (0.61, 0.87)	0.81 (0.75, 0.87)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXII-9 - continued
PANCREAS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	1.33 (1.21, 1.46)	1.14 (1.05, 1.25)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	1.34 (1.22, 1.48)	1.15 (1.06, 1.27)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	1.34 (1.23, 1.48)	1.16 (1.07, 1.27)	
	30	0.01 (0.01, 0.01)	0.04 (0.03, 0.05)	0.14 (0.12, 0.16)	1.35 (1.24, 1.49)	1.16 (1.07, 1.28)	
	40	0.03 (0.02, 0.04)	0.13 (0.12, 0.16)	0.40 (0.36, 0.44)	1.35 (1.24, 1.49)	1.17 (1.07, 1.28)	
	50	0.11 (0.09, 0.12)	0.37 (0.33, 0.42)	0.81 (0.74, 0.88)	1.35 (1.23, 1.49)	1.16 (1.07, 1.28)	
	60	0.28 (0.24, 0.32)	0.73 (0.66, 0.80)	1.13 (1.04, 1.24)	1.29 (1.17, 1.44)	1.12 (1.02, 1.24)	
	70	0.50 (0.44, 0.56)	0.95 (0.85, 1.06)	- (-, -)	1.13 (1.00, 1.28)	1.00 (0.90, 1.13)	
	80	0.59 (0.50, 0.70)	- (-, -)	- (-, -)	0.83 (0.69, 1.01)	0.77 (0.65, 0.92)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.93 (0.66, 1.64)	0.79 (0.60, 1.07)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.02 (0.01, 0.06)	0.95 (0.67, 1.67)	0.80 (0.61, 1.09)	
	20	0.00 (0.00, 0.02)	0.02 (0.01, 0.06)	0.08 (0.04, 0.14)	0.96 (0.68, 1.69)	0.81 (0.62, 1.11)	
	30	0.02 (0.01, 0.06)	0.08 (0.04, 0.14)	0.16 (0.09, 0.25)	0.99 (0.70, 1.73)	0.83 (0.63, 1.14)	
	40	0.06 (0.03, 0.11)	0.14 (0.08, 0.23)	0.45 (0.31, 0.66)	1.00 (0.70, 1.77)	0.86 (0.66, 1.18)	
	50	0.09 (0.04, 0.17)	0.42 (0.27, 0.62)	0.81 (0.56, 1.14)	0.99 (0.68, 1.81)	0.88 (0.66, 1.22)	
	60	0.36 (0.22, 0.57)	0.79 (0.53, 1.15)	0.98 (0.65, 1.45)	0.99 (0.66, 1.87)	0.86 (0.63, 1.23)	
	70	0.51 (0.29, 0.85)	0.74 (0.42, 1.24)	- (-, -)	0.75 (0.43, 1.79)	0.76 (0.49, 1.21)	
	80	0.34 (0.09, 0.92)	- (-, -)	- (-, -)	0.36 (0.09, 1.91)	0.69 (0.34, 1.37)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.26 (1.16, 1.38)	1.06 (1.00, 1.12)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.27 (1.18, 1.39)	1.07 (1.01, 1.13)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.03, 0.05)	1.28 (1.18, 1.40)	1.08 (1.02, 1.14)	
	30	0.01 (0.01, 0.01)	0.04 (0.03, 0.05)	0.16 (0.14, 0.18)	1.29 (1.19, 1.42)	1.09 (1.03, 1.15)	
	40	0.03 (0.03, 0.04)	0.16 (0.14, 0.17)	0.45 (0.41, 0.49)	1.30 (1.20, 1.43)	1.10 (1.04, 1.16)	
	50	0.13 (0.11, 0.14)	0.43 (0.39, 0.47)	0.91 (0.84, 0.97)	1.31 (1.21, 1.44)	1.11 (1.05, 1.17)	
	60	0.32 (0.29, 0.36)	0.83 (0.76, 0.90)	1.15 (1.06, 1.25)	1.26 (1.15, 1.39)	1.07 (1.01, 1.14)	
	70	0.58 (0.53, 0.65)	0.96 (0.87, 1.06)	- (-, -)	1.08 (0.97, 1.23)	0.93 (0.86, 1.00)	
	80	0.53 (0.44, 0.62)	- (-, -)	- (-, -)	0.70 (0.58, 0.88)	0.66 (0.58, 0.75)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXII-9 - continued
PANCREAS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.32 (1.29, 1.34)	1.19 (1.18, 1.20)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.33 (1.31, 1.35)	1.21 (1.20, 1.22)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.04, 0.04)	1.33 (1.31, 1.36)	1.21 (1.20, 1.22)	
	30	0.01 (0.01, 0.01)	0.04 (0.04, 0.04)	0.15 (0.14, 0.15)	1.34 (1.31, 1.36)	1.22 (1.21, 1.23)	
	40	0.03 (0.03, 0.03)	0.14 (0.13, 0.15)	0.41 (0.40, 0.42)	1.34 (1.32, 1.37)	1.23 (1.21, 1.24)	
	50	0.11 (0.10, 0.11)	0.39 (0.38, 0.40)	0.83 (0.81, 0.85)	1.34 (1.31, 1.36)	1.23 (1.22, 1.24)	
	60	0.29 (0.28, 0.30)	0.75 (0.74, 0.77)	1.15 (1.13, 1.18)	1.29 (1.26, 1.31)	1.19 (1.18, 1.21)	
	70	0.52 (0.50, 0.53)	0.97 (0.95, 0.99)	- (-, -)	1.12 (1.09, 1.14)	1.07 (1.06, 1.08)	
	80	0.60 (0.58, 0.62)	- (-, -)	- (-, -)	0.80 (0.78, 0.83)	0.81 (0.79, 0.82)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.29 (1.27, 1.32)	1.18 (1.17, 1.19)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.31 (1.28, 1.33)	1.19 (1.18, 1.20)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.04, 0.04)	1.31 (1.28, 1.33)	1.20 (1.18, 1.21)	
	30	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	0.14 (0.14, 0.15)	1.31 (1.29, 1.34)	1.20 (1.19, 1.21)	
	40	0.03 (0.03, 0.03)	0.13 (0.13, 0.14)	0.40 (0.39, 0.42)	1.32 (1.29, 1.34)	1.21 (1.20, 1.22)	
	50	0.11 (0.10, 0.11)	0.38 (0.37, 0.39)	0.82 (0.80, 0.84)	1.31 (1.28, 1.34)	1.21 (1.20, 1.22)	
	60	0.29 (0.27, 0.30)	0.74 (0.72, 0.76)	1.13 (1.11, 1.16)	1.26 (1.23, 1.28)	1.17 (1.16, 1.19)	
	70	0.51 (0.50, 0.53)	0.95 (0.93, 0.97)	- (-, -)	1.09 (1.06, 1.11)	1.05 (1.04, 1.06)	
	80	0.59 (0.57, 0.61)	- (-, -)	- (-, -)	0.77 (0.75, 0.80)	0.79 (0.78, 0.80)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.53 (1.45, 1.62)	1.33 (1.30, 1.37)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	1.57 (1.48, 1.65)	1.37 (1.33, 1.40)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.06 (0.05, 0.07)	1.57 (1.49, 1.66)	1.37 (1.33, 1.41)	
	30	0.01 (0.00, 0.01)	0.06 (0.05, 0.07)	0.20 (0.18, 0.22)	1.58 (1.50, 1.67)	1.38 (1.34, 1.42)	
	40	0.05 (0.04, 0.06)	0.20 (0.18, 0.22)	0.55 (0.51, 0.59)	1.60 (1.52, 1.69)	1.40 (1.36, 1.44)	
	50	0.15 (0.13, 0.17)	0.52 (0.48, 0.56)	1.04 (0.97, 1.10)	1.62 (1.53, 1.71)	1.42 (1.39, 1.46)	
	60	0.40 (0.36, 0.44)	0.97 (0.90, 1.04)	1.42 (1.33, 1.50)	1.60 (1.50, 1.70)	1.40 (1.36, 1.45)	
	70	0.68 (0.62, 0.74)	1.21 (1.13, 1.30)	- (-, -)	1.43 (1.33, 1.54)	1.27 (1.23, 1.31)	
	80	0.79 (0.71, 0.88)	- (-, -)	- (-, -)	1.11 (1.00, 1.23)	0.98 (0.93, 1.03)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXII-9 - continued
PANCREAS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	1.35 (1.23, 1.50)	1.27 (1.16, 1.40)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	1.37 (1.24, 1.51)	1.28 (1.17, 1.41)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.03, 0.04)	1.37 (1.25, 1.51)	1.29 (1.18, 1.41)	
	30	0.01 (0.00, 0.01)	0.03 (0.02, 0.04)	0.12 (0.11, 0.14)	1.37 (1.25, 1.52)	1.29 (1.18, 1.42)	
	40	0.03 (0.02, 0.03)	0.12 (0.10, 0.14)	0.33 (0.30, 0.37)	1.37 (1.25, 1.52)	1.29 (1.18, 1.42)	
	50	0.09 (0.08, 0.11)	0.31 (0.28, 0.34)	0.68 (0.62, 0.74)	1.36 (1.23, 1.51)	1.29 (1.18, 1.42)	
	60	0.22 (0.19, 0.25)	0.60 (0.55, 0.66)	1.07 (0.98, 1.16)	1.30 (1.17, 1.45)	1.25 (1.14, 1.38)	
	70	0.40 (0.36, 0.45)	0.90 (0.82, 0.99)	- (-, -)	1.15 (1.02, 1.31)	1.13 (1.02, 1.27)	
	80	0.59 (0.51, 0.68)	- (-, -)	- (-, -)	0.89 (0.75, 1.06)	0.90 (0.78, 1.06)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.05)	0.01 (0.00, 0.05)	0.97 (0.67, 1.57)	0.83 (0.66, 1.06)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.03)	0.01 (0.00, 0.03)	0.01 (0.00, 0.05)	0.99 (0.68, 1.59)	0.84 (0.67, 1.08)	
	20	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.04 (0.01, 0.08)	0.99 (0.68, 1.59)	0.84 (0.68, 1.08)	
	30	0.01 (0.00, 0.04)	0.04 (0.01, 0.08)	0.10 (0.05, 0.18)	0.99 (0.69, 1.61)	0.85 (0.68, 1.09)	
	40	0.03 (0.01, 0.07)	0.09 (0.05, 0.17)	0.30 (0.19, 0.46)	1.00 (0.69, 1.63)	0.86 (0.69, 1.11)	
	50	0.06 (0.03, 0.13)	0.28 (0.17, 0.44)	0.62 (0.41, 0.89)	1.00 (0.68, 1.65)	0.88 (0.70, 1.13)	
	60	0.23 (0.13, 0.38)	0.58 (0.38, 0.87)	0.92 (0.62, 1.33)	0.99 (0.66, 1.67)	0.85 (0.67, 1.12)	
	70	0.40 (0.23, 0.67)	0.79 (0.49, 1.23)	- (-, -)	0.87 (0.54, 1.63)	0.73 (0.53, 1.03)	
	80	0.51 (0.24, 0.99)	- (-, -)	- (-, -)	0.63 (0.29, 1.59)	0.52 (0.31, 0.89)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.63 (1.51, 1.76)	1.19 (1.13, 1.26)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.65 (1.52, 1.78)	1.21 (1.15, 1.27)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	1.65 (1.53, 1.79)	1.21 (1.15, 1.28)	
	30	0.01 (0.00, 0.01)	0.03 (0.03, 0.04)	0.14 (0.12, 0.15)	1.65 (1.53, 1.79)	1.21 (1.15, 1.28)	
	40	0.03 (0.02, 0.03)	0.13 (0.11, 0.15)	0.41 (0.37, 0.44)	1.66 (1.53, 1.80)	1.22 (1.16, 1.29)	
	50	0.11 (0.09, 0.12)	0.39 (0.35, 0.42)	0.87 (0.81, 0.93)	1.66 (1.53, 1.80)	1.22 (1.16, 1.29)	
	60	0.29 (0.26, 0.32)	0.79 (0.73, 0.85)	1.27 (1.19, 1.37)	1.60 (1.48, 1.75)	1.19 (1.13, 1.26)	
	70	0.54 (0.49, 0.60)	1.07 (0.99, 1.17)	- (-, -)	1.43 (1.30, 1.58)	1.08 (1.02, 1.16)	
	80	0.67 (0.59, 0.75)	- (-, -)	- (-, -)	1.12 (0.97, 1.30)	0.87 (0.79, 0.95)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXIII-7
PROSTATE CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	16.72 (16.63, 16.81)	2.87 (2.85, 2.89)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	16.95 (16.86, 17.04)	2.92 (2.90, 2.94)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.29 (0.29, 0.30)	17.04 (16.95, 17.13)	2.94 (2.92, 2.96)	
	30	0.01 (0.01, 0.01)	0.30 (0.29, 0.30)	2.51 (2.49, 2.53)	17.27 (17.18, 17.36)	2.98 (2.96, 3.00)	
	40	0.29 (0.29, 0.30)	2.54 (2.52, 2.57)	8.49 (8.43, 8.54)	17.55 (17.46, 17.65)	3.03 (3.01, 3.05)	
	50	2.34 (2.32, 2.36)	8.52 (8.46, 8.58)	14.86 (14.78, 14.94)	17.94 (17.85, 18.04)	3.14 (3.12, 3.16)	
	60	6.83 (6.78, 6.88)	13.84 (13.76, 13.92)	16.71 (16.62, 16.81)	17.25 (17.15, 17.35)	3.34 (3.31, 3.36)	
	70	8.98 (8.91, 9.05)	12.66 (12.57, 12.76)	- (- , -)	13.36 (13.25, 13.46)	3.60 (3.57, 3.63)	
	80	6.26 (6.18, 6.34)	- (- , -)	- (- , -)	7.44 (7.34, 7.54)	3.82 (3.78, 3.85)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	16.15 (16.06, 16.25)	2.70 (2.68, 2.72)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	16.34 (16.25, 16.44)	2.73 (2.71, 2.75)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.27 (0.26, 0.27)	16.43 (16.34, 16.53)	2.75 (2.73, 2.77)	
	30	0.01 (0.01, 0.01)	0.27 (0.26, 0.28)	2.39 (2.37, 2.42)	16.63 (16.53, 16.73)	2.78 (2.76, 2.80)	
	40	0.27 (0.26, 0.27)	2.42 (2.40, 2.45)	8.23 (8.17, 8.29)	16.89 (16.79, 16.99)	2.83 (2.81, 2.85)	
	50	2.24 (2.21, 2.26)	8.27 (8.20, 8.33)	14.39 (14.30, 14.47)	17.25 (17.15, 17.35)	2.93 (2.90, 2.95)	
	60	6.63 (6.57, 6.69)	13.37 (13.27, 13.46)	16.03 (15.92, 16.13)	16.52 (16.41, 16.62)	3.10 (3.08, 3.12)	
	70	8.58 (8.50, 8.66)	11.97 (11.87, 12.07)	- (- , -)	12.59 (12.49, 12.70)	3.34 (3.32, 3.37)	
	80	5.75 (5.67, 5.84)	- (- , -)	- (- , -)	6.81 (6.71, 6.92)	3.57 (3.54, 3.61)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	19.43 (19.13, 19.74)	4.50 (4.43, 4.58)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.03)	19.95 (19.64, 20.27)	4.63 (4.55, 4.71)	
	20	0.00 (0.00, 0.00)	0.02 (0.02, 0.03)	0.61 (0.58, 0.63)	20.13 (19.82, 20.45)	4.67 (4.59, 4.75)	
	30	0.02 (0.02, 0.03)	0.62 (0.59, 0.65)	4.30 (4.20, 4.40)	20.63 (20.31, 20.96)	4.78 (4.70, 4.86)	
	40	0.62 (0.59, 0.65)	4.41 (4.31, 4.52)	12.23 (12.02, 12.44)	21.28 (20.95, 21.61)	4.93 (4.85, 5.01)	
	50	4.08 (3.97, 4.18)	12.46 (12.24, 12.68)	19.30 (18.99, 19.61)	22.18 (21.83, 22.53)	5.24 (5.16, 5.33)	
	60	10.13 (9.91, 10.35)	18.39 (18.05, 18.72)	21.31 (20.94, 21.69)	21.86 (21.47, 22.26)	5.87 (5.77, 5.97)	
	70	12.32 (12.00, 12.66)	16.69 (16.28, 17.11)	- (- , -)	17.52 (17.07, 17.97)	6.69 (6.57, 6.82)	
	80	9.07 (8.64, 9.52)	- (- , -)	- (- , -)	10.78 (10.25, 11.35)	7.33 (7.15, 7.52)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXIII-7 - continued
PROSTATE CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific Islander	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	12.52 (12.18, 12.88)	2.29 (2.10, 2.50)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	12.65 (12.30, 13.01)	2.31 (2.12, 2.52)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.09 (0.08, 0.11)	12.69 (12.35, 13.05)	2.32 (2.13, 2.53)	
	30	0.00 (0.00, 0.01)	0.09 (0.08, 0.11)	0.93 (0.89, 0.98)	12.77 (12.43, 13.13)	2.33 (2.14, 2.54)	
	40	0.09 (0.08, 0.10)	0.94 (0.89, 0.99)	4.33 (4.19, 4.47)	12.87 (12.52, 13.23)	2.35 (2.15, 2.56)	
	50	0.86 (0.82, 0.91)	4.32 (4.18, 4.47)	9.35 (9.11, 9.59)	13.02 (12.66, 13.39)	2.38 (2.19, 2.60)	
	60	3.64 (3.51, 3.77)	8.93 (8.69, 9.18)	11.90 (11.59, 12.22)	12.79 (12.43, 13.18)	2.45 (2.25, 2.68)	
	70	6.10 (5.89, 6.32)	9.53 (9.22, 9.85)	- (- , -)	10.55 (10.17, 10.96)	2.59 (2.36, 2.83)	
American Indian/ Alaska Native ^a	80	4.82 (4.55, 5.11)	- (- , -)	- (- , -)	6.27 (5.86, 6.72)	2.72 (2.44, 3.03)	
	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	6.99 (6.14, 8.15)	2.18 (1.82, 2.62)	
	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	7.13 (6.26, 8.32)	2.22 (1.86, 2.68)	
	20	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.09 (0.05, 0.14)	7.20 (6.33, 8.41)	2.25 (1.88, 2.71)	
	30	0.00 (0.00, 0.03)	0.09 (0.05, 0.14)	0.93 (0.76, 1.13)	7.38 (6.48, 8.61)	2.31 (1.93, 2.78)	
	40	0.09 (0.05, 0.15)	0.96 (0.78, 1.17)	3.40 (2.96, 3.89)	7.65 (6.72, 8.92)	2.40 (2.00, 2.89)	
	50	0.92 (0.74, 1.12)	3.49 (3.03, 4.00)	6.03 (5.33, 6.81)	7.96 (6.98, 9.30)	2.55 (2.13, 3.07)	
	60	2.83 (2.40, 3.33)	5.63 (4.90, 6.45)	7.28 (6.34, 8.34)	7.76 (6.71, 9.21)	2.79 (2.33, 3.38)	
Hispanic ^b	70	3.45 (2.82, 4.18)	5.47 (4.53, 6.59)	- (- , -)	6.07 (4.97, 7.71)	3.14 (2.58, 3.86)	
	80	3.16 (2.24, 4.37)	- (- , -)	- (- , -)	4.08 (2.83, 6.30)	3.41 (2.62, 4.47)	
	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	16.25 (15.90, 16.62)	3.19 (3.05, 3.33)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	16.45 (16.10, 16.82)	3.23 (3.09, 3.37)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.17 (0.16, 0.19)	16.55 (16.19, 16.92)	3.25 (3.11, 3.39)	
	30	0.00 (0.00, 0.01)	0.18 (0.16, 0.19)	1.67 (1.61, 1.72)	16.74 (16.38, 17.12)	3.28 (3.15, 3.43)	
	40	0.17 (0.16, 0.19)	1.68 (1.63, 1.74)	6.76 (6.61, 6.92)	16.97 (16.61, 17.36)	3.33 (3.19, 3.48)	
	50	1.56 (1.50, 1.61)	6.80 (6.65, 6.96)	13.20 (12.95, 13.46)	17.34 (16.96, 17.73)	3.43 (3.28, 3.58)	
-	60	5.68 (5.53, 5.83)	12.61 (12.35, 12.88)	16.01 (15.68, 16.35)	17.09 (16.69, 17.51)	3.61 (3.45, 3.77)	
	70	8.46 (8.22, 8.70)	12.61 (12.27, 12.97)	- (- , -)	13.93 (13.49, 14.39)	3.85 (3.68, 4.04)	
	80	6.40 (6.07, 6.75)	- (- , -)	- (- , -)	8.42 (7.90, 8.99)	4.07 (3.84, 4.32)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXIV-9
STOMACH CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.90 (0.89, 0.91)	0.46 (0.46, 0.47)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.91 (0.90, 0.92)	0.47 (0.47, 0.48)	
	20	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.05 (0.05, 0.06)	0.91 (0.90, 0.93)	0.47 (0.47, 0.48)	
	30	0.01 (0.01, 0.02)	0.05 (0.05, 0.05)	0.15 (0.14, 0.15)	0.92 (0.90, 0.93)	0.48 (0.47, 0.48)	
	40	0.04 (0.04, 0.04)	0.13 (0.13, 0.14)	0.33 (0.32, 0.34)	0.92 (0.90, 0.93)	0.48 (0.47, 0.48)	
	50	0.10 (0.09, 0.10)	0.30 (0.29, 0.31)	0.60 (0.59, 0.61)	0.90 (0.89, 0.92)	0.47 (0.47, 0.48)	
	60	0.22 (0.21, 0.22)	0.53 (0.52, 0.54)	0.78 (0.77, 0.80)	0.86 (0.84, 0.87)	0.46 (0.45, 0.46)	
	70	0.36 (0.36, 0.37)	0.65 (0.64, 0.67)	- (- , -)	0.74 (0.72, 0.75)	0.42 (0.41, 0.42)	
	80	0.41 (0.40, 0.42)	- (- , -)	- (- , -)	0.53 (0.52, 0.55)	0.34 (0.33, 0.35)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.79 (0.78, 0.81)	0.41 (0.41, 0.42)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.80 (0.79, 0.81)	0.42 (0.41, 0.42)	
	20	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.05 (0.05, 0.05)	0.80 (0.79, 0.82)	0.42 (0.42, 0.43)	
	30	0.01 (0.01, 0.01)	0.05 (0.04, 0.05)	0.13 (0.12, 0.13)	0.81 (0.79, 0.82)	0.42 (0.42, 0.43)	
	40	0.03 (0.03, 0.03)	0.12 (0.11, 0.12)	0.29 (0.29, 0.30)	0.80 (0.79, 0.82)	0.42 (0.42, 0.43)	
	50	0.09 (0.08, 0.09)	0.27 (0.26, 0.28)	0.53 (0.52, 0.54)	0.79 (0.78, 0.81)	0.42 (0.41, 0.42)	
	60	0.19 (0.19, 0.20)	0.47 (0.46, 0.48)	0.69 (0.67, 0.70)	0.75 (0.73, 0.76)	0.41 (0.40, 0.41)	
	70	0.32 (0.31, 0.33)	0.57 (0.56, 0.58)	- (- , -)	0.64 (0.62, 0.65)	0.37 (0.37, 0.38)	
	80	0.36 (0.35, 0.37)	- (- , -)	- (- , -)	0.46 (0.44, 0.47)	0.30 (0.30, 0.31)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	1.12 (1.07, 1.17)	0.74 (0.72, 0.76)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	1.14 (1.09, 1.20)	0.76 (0.74, 0.78)	
	20	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.07 (0.06, 0.08)	1.15 (1.10, 1.20)	0.76 (0.74, 0.78)	
	30	0.01 (0.01, 0.02)	0.06 (0.06, 0.07)	0.19 (0.18, 0.21)	1.17 (1.11, 1.22)	0.77 (0.75, 0.79)	
	40	0.05 (0.05, 0.06)	0.19 (0.17, 0.20)	0.45 (0.42, 0.48)	1.18 (1.13, 1.24)	0.78 (0.76, 0.80)	
	50	0.14 (0.13, 0.15)	0.42 (0.39, 0.44)	0.80 (0.75, 0.84)	1.19 (1.13, 1.25)	0.79 (0.77, 0.81)	
	60	0.31 (0.29, 0.34)	0.73 (0.69, 0.78)	1.05 (0.99, 1.10)	1.17 (1.11, 1.24)	0.79 (0.76, 0.81)	
	70	0.53 (0.49, 0.57)	0.92 (0.87, 0.98)	- (- , -)	1.08 (1.02, 1.15)	0.75 (0.73, 0.78)	
	80	0.63 (0.57, 0.69)	- (- , -)	- (- , -)	0.88 (0.80, 0.97)	0.66 (0.63, 0.69)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXIV-9 - continued
STOMACH CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	2.06 (1.95, 2.17)	1.32 (1.24, 1.42)	
Islander	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	2.07 (1.97, 2.19)	1.33 (1.25, 1.43)	
	20	0.00 (0.00, 0.01)	0.03 (0.02, 0.03)	0.09 (0.08, 0.10)	2.08 (1.97, 2.20)	1.34 (1.25, 1.43)	
	30	0.02 (0.02, 0.03)	0.09 (0.08, 0.09)	0.23 (0.21, 0.25)	2.08 (1.98, 2.20)	1.34 (1.25, 1.44)	
	40	0.06 (0.06, 0.07)	0.21 (0.19, 0.23)	0.53 (0.50, 0.56)	2.07 (1.97, 2.19)	1.34 (1.25, 1.43)	
	50	0.15 (0.13, 0.16)	0.47 (0.44, 0.50)	1.07 (1.02, 1.13)	2.04 (1.93, 2.16)	1.32 (1.23, 1.42)	
	60	0.34 (0.31, 0.36)	0.96 (0.91, 1.01)	1.64 (1.56, 1.72)	1.96 (1.85, 2.08)	1.28 (1.19, 1.38)	
	70	0.68 (0.63, 0.72)	1.42 (1.33, 1.50)	- (- , -)	1.76 (1.65, 1.89)	1.22 (1.12, 1.32)	
	80	0.92 (0.84, 1.00)	- (- , -)	- (- , -)	1.36 (1.23, 1.50)	1.07 (0.96, 1.20)	
American	0	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	1.30 (1.04, 1.70)	0.73 (0.62, 0.89)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.01)	0.00 (0.00, 0.02)	0.03 (0.02, 0.06)	1.32 (1.06, 1.73)	0.75 (0.63, 0.90)	
	20	0.00 (0.00, 0.02)	0.03 (0.02, 0.06)	0.09 (0.06, 0.13)	1.33 (1.06, 1.75)	0.75 (0.63, 0.91)	
	30	0.03 (0.01, 0.05)	0.08 (0.05, 0.12)	0.23 (0.17, 0.30)	1.35 (1.08, 1.77)	0.76 (0.64, 0.92)	
	40	0.06 (0.03, 0.09)	0.20 (0.15, 0.27)	0.47 (0.36, 0.59)	1.36 (1.08, 1.79)	0.77 (0.65, 0.94)	
	50	0.15 (0.10, 0.22)	0.43 (0.32, 0.56)	0.83 (0.65, 1.03)	1.35 (1.07, 1.80)	0.77 (0.64, 0.94)	
	60	0.30 (0.21, 0.41)	0.72 (0.55, 0.94)	1.14 (0.89, 1.45)	1.29 (0.99, 1.77)	0.77 (0.64, 0.96)	
	70	0.50 (0.35, 0.70)	0.99 (0.72, 1.33)	- (- , -)	1.16 (0.85, 1.71)	0.73 (0.57, 0.94)	
	80	0.69 (0.42, 1.06)	- (- , -)	- (- , -)	0.93 (0.58, 1.64)	0.54 (0.37, 0.82)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	1.62 (1.54, 1.72)	0.91 (0.87, 0.95)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	1.64 (1.56, 1.73)	0.92 (0.88, 0.96)	
	20	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	0.08 (0.08, 0.09)	1.65 (1.56, 1.74)	0.92 (0.88, 0.96)	
	30	0.02 (0.02, 0.02)	0.08 (0.07, 0.08)	0.22 (0.21, 0.24)	1.66 (1.57, 1.75)	0.93 (0.89, 0.97)	
	40	0.06 (0.05, 0.06)	0.20 (0.19, 0.22)	0.51 (0.49, 0.54)	1.65 (1.57, 1.75)	0.93 (0.89, 0.97)	
	50	0.15 (0.14, 0.16)	0.47 (0.44, 0.49)	0.97 (0.92, 1.01)	1.63 (1.54, 1.73)	0.92 (0.88, 0.96)	
	60	0.33 (0.31, 0.36)	0.86 (0.81, 0.91)	1.32 (1.26, 1.39)	1.56 (1.47, 1.66)	0.88 (0.84, 0.93)	
	70	0.59 (0.55, 0.63)	1.11 (1.04, 1.18)	- (- , -)	1.37 (1.28, 1.48)	0.81 (0.77, 0.86)	
	80	0.69 (0.62, 0.75)	- (- , -)	- (- , -)	1.03 (0.92, 1.16)	0.66 (0.61, 0.72)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXIV-9 - continued
STOMACH CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.13 (1.11, 1.15)	0.56 (0.56, 0.57)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	1.15 (1.12, 1.17)	0.57 (0.57, 0.58)	
	20	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.06 (0.06, 0.07)	1.15 (1.13, 1.18)	0.58 (0.57, 0.59)	
	30	0.01 (0.01, 0.02)	0.06 (0.06, 0.06)	0.18 (0.18, 0.19)	1.16 (1.14, 1.19)	0.58 (0.57, 0.59)	
	40	0.05 (0.04, 0.05)	0.17 (0.17, 0.18)	0.45 (0.44, 0.46)	1.17 (1.14, 1.19)	0.59 (0.58, 0.60)	
	50	0.13 (0.12, 0.14)	0.42 (0.40, 0.43)	0.81 (0.79, 0.83)	1.16 (1.14, 1.19)	0.59 (0.58, 0.60)	
	60	0.31 (0.30, 0.32)	0.74 (0.72, 0.76)	1.04 (1.02, 1.06)	1.12 (1.09, 1.14)	0.57 (0.56, 0.58)	
	70	0.51 (0.49, 0.53)	0.87 (0.85, 0.89)	- (-, -)	0.96 (0.93, 0.99)	0.52 (0.51, 0.53)	
	80	0.56 (0.53, 0.58)	- (-, -)	- (-, -)	0.70 (0.67, 0.73)	0.43 (0.42, 0.44)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.01 (0.99, 1.04)	0.51 (0.50, 0.52)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	1.02 (1.00, 1.05)	0.51 (0.51, 0.52)	
	20	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.06 (0.05, 0.06)	1.03 (1.01, 1.05)	0.52 (0.51, 0.52)	
	30	0.01 (0.01, 0.02)	0.05 (0.05, 0.06)	0.16 (0.16, 0.17)	1.04 (1.01, 1.06)	0.52 (0.51, 0.53)	
	40	0.04 (0.04, 0.04)	0.15 (0.15, 0.16)	0.40 (0.39, 0.42)	1.04 (1.02, 1.07)	0.52 (0.52, 0.53)	
	50	0.12 (0.11, 0.12)	0.38 (0.36, 0.39)	0.73 (0.71, 0.75)	1.04 (1.01, 1.06)	0.52 (0.52, 0.53)	
	60	0.28 (0.27, 0.29)	0.66 (0.64, 0.68)	0.93 (0.90, 0.95)	0.99 (0.96, 1.02)	0.51 (0.50, 0.52)	
	70	0.45 (0.44, 0.47)	0.77 (0.75, 0.79)	- (-, -)	0.84 (0.82, 0.87)	0.47 (0.46, 0.48)	
	80	0.49 (0.47, 0.51)	- (-, -)	- (-, -)	0.60 (0.57, 0.63)	0.39 (0.37, 0.40)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	1.30 (1.22, 1.38)	0.87 (0.84, 0.90)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.03)	1.33 (1.25, 1.42)	0.90 (0.87, 0.93)	
	20	0.00 (0.00, 0.01)	0.02 (0.01, 0.03)	0.08 (0.07, 0.09)	1.34 (1.26, 1.43)	0.90 (0.87, 0.94)	
	30	0.02 (0.01, 0.02)	0.08 (0.07, 0.09)	0.25 (0.23, 0.28)	1.37 (1.29, 1.46)	0.92 (0.89, 0.95)	
	40	0.07 (0.06, 0.08)	0.24 (0.22, 0.27)	0.60 (0.56, 0.65)	1.40 (1.31, 1.49)	0.94 (0.91, 0.98)	
	50	0.19 (0.17, 0.21)	0.57 (0.53, 0.62)	1.03 (0.97, 1.11)	1.42 (1.33, 1.52)	0.96 (0.93, 1.00)	
	60	0.44 (0.40, 0.49)	0.98 (0.90, 1.05)	1.30 (1.21, 1.39)	1.43 (1.33, 1.54)	0.97 (0.93, 1.01)	
	70	0.71 (0.64, 0.79)	1.14 (1.05, 1.25)	- (-, -)	1.31 (1.20, 1.44)	0.93 (0.89, 0.98)	
	80	0.78 (0.68, 0.90)	- (-, -)	- (-, -)	1.09 (0.94, 1.27)	0.81 (0.75, 0.87)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

- Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXIV-9 - continued
STOMACH CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	2.47 (2.31, 2.65)	1.54 (1.42, 1.69)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.03 (0.02, 0.04)	2.50 (2.34, 2.68)	1.56 (1.43, 1.70)	
	20	0.00 (0.00, 0.01)	0.03 (0.02, 0.04)	0.10 (0.09, 0.11)	2.51 (2.35, 2.69)	1.56 (1.44, 1.71)	
	30	0.02 (0.02, 0.03)	0.10 (0.08, 0.11)	0.27 (0.25, 0.30)	2.52 (2.36, 2.70)	1.57 (1.44, 1.72)	
	40	0.07 (0.06, 0.08)	0.25 (0.23, 0.28)	0.69 (0.64, 0.75)	2.51 (2.35, 2.70)	1.57 (1.44, 1.72)	
	50	0.18 (0.16, 0.21)	0.63 (0.58, 0.68)	1.40 (1.31, 1.50)	2.49 (2.32, 2.67)	1.56 (1.43, 1.71)	
	60	0.47 (0.42, 0.51)	1.27 (1.19, 1.37)	2.11 (1.98, 2.26)	2.41 (2.24, 2.60)	1.53 (1.40, 1.69)	
	70	0.90 (0.82, 0.98)	1.84 (1.70, 1.98)	- (- , -)	2.16 (1.98, 2.36)	1.45 (1.30, 1.62)	
	80	1.24 (1.10, 1.38)	- (- , -)	- (- , -)	1.67 (1.47, 1.91)	1.28 (1.11, 1.49)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	1.65 (1.22, 2.47)	0.85 (0.68, 1.10)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.03 (0.01, 0.07)	1.68 (1.25, 2.52)	0.86 (0.70, 1.13)	
	20	0.00 (0.00, 0.03)	0.03 (0.01, 0.08)	0.09 (0.05, 0.16)	1.70 (1.26, 2.55)	0.87 (0.70, 1.14)	
	30	0.03 (0.01, 0.07)	0.09 (0.05, 0.16)	0.28 (0.19, 0.40)	1.74 (1.29, 2.61)	0.89 (0.72, 1.17)	
	40	0.07 (0.03, 0.12)	0.26 (0.17, 0.38)	0.58 (0.42, 0.78)	1.78 (1.31, 2.67)	0.91 (0.73, 1.20)	
	50	0.21 (0.13, 0.32)	0.54 (0.38, 0.75)	1.11 (0.82, 1.48)	1.80 (1.31, 2.74)	0.93 (0.74, 1.23)	
	60	0.36 (0.23, 0.56)	0.99 (0.69, 1.37)	1.54 (1.11, 2.12)	1.74 (1.23, 2.77)	0.96 (0.75, 1.29)	
	70	0.75 (0.48, 1.13)	1.42 (0.95, 2.06)	- (- , -)	1.66 (1.09, 2.86)	0.92 (0.68, 1.32)	
	80	1.01 (0.51, 1.81)	- (- , -)	- (- , -)	1.37 (0.69, 3.12)	0.70 (0.43, 1.26)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	1.85 (1.73, 1.99)	1.07 (1.01, 1.14)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.02, 0.03)	1.87 (1.75, 2.02)	1.08 (1.02, 1.15)	
	20	0.01 (0.00, 0.01)	0.02 (0.02, 0.03)	0.08 (0.07, 0.09)	1.88 (1.76, 2.03)	1.09 (1.03, 1.16)	
	30	0.02 (0.02, 0.02)	0.08 (0.07, 0.09)	0.24 (0.22, 0.26)	1.90 (1.77, 2.05)	1.10 (1.04, 1.17)	
	40	0.06 (0.05, 0.07)	0.23 (0.21, 0.25)	0.63 (0.59, 0.68)	1.91 (1.78, 2.06)	1.10 (1.04, 1.18)	
	50	0.17 (0.15, 0.19)	0.59 (0.55, 0.64)	1.24 (1.16, 1.32)	1.90 (1.77, 2.06)	1.11 (1.04, 1.18)	
	60	0.45 (0.41, 0.49)	1.14 (1.06, 1.22)	1.64 (1.53, 1.76)	1.85 (1.71, 2.01)	1.08 (1.02, 1.16)	
	70	0.80 (0.73, 0.87)	1.38 (1.27, 1.50)	- (- , -)	1.62 (1.47, 1.79)	1.02 (0.94, 1.10)	
	80	0.82 (0.71, 0.94)	- (- , -)	- (- , -)	1.16 (0.98, 1.38)	0.83 (0.73, 0.94)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXIV-9 - continued
STOMACH CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.69 (0.68, 0.71)	0.38 (0.37, 0.38)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.70 (0.68, 0.72)	0.38 (0.38, 0.39)	
	20	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.05 (0.04, 0.05)	0.70 (0.69, 0.72)	0.38 (0.38, 0.39)	
	30	0.01 (0.01, 0.01)	0.04 (0.04, 0.05)	0.11 (0.10, 0.11)	0.70 (0.69, 0.72)	0.38 (0.38, 0.39)	
	40	0.03 (0.03, 0.03)	0.10 (0.09, 0.10)	0.22 (0.21, 0.23)	0.70 (0.68, 0.71)	0.38 (0.37, 0.39)	
	50	0.07 (0.06, 0.07)	0.19 (0.19, 0.20)	0.41 (0.40, 0.42)	0.68 (0.66, 0.70)	0.38 (0.37, 0.38)	
	60	0.13 (0.13, 0.14)	0.36 (0.35, 0.37)	0.57 (0.55, 0.58)	0.64 (0.62, 0.66)	0.36 (0.36, 0.37)	
	70	0.25 (0.24, 0.26)	0.49 (0.47, 0.50)	- (-, -)	0.57 (0.55, 0.59)	0.34 (0.33, 0.35)	
	80	0.32 (0.30, 0.33)	- (-, -)	- (-, -)	0.43 (0.41, 0.45)	0.29 (0.28, 0.29)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.59 (0.58, 0.61)	0.33 (0.33, 0.34)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.60 (0.58, 0.62)	0.34 (0.33, 0.34)	
	20	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.04 (0.04, 0.04)	0.60 (0.58, 0.62)	0.34 (0.33, 0.34)	
	30	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	0.09 (0.09, 0.10)	0.60 (0.58, 0.62)	0.34 (0.33, 0.34)	
	40	0.03 (0.02, 0.03)	0.08 (0.08, 0.09)	0.19 (0.18, 0.20)	0.59 (0.58, 0.61)	0.33 (0.33, 0.34)	
	50	0.06 (0.05, 0.06)	0.17 (0.16, 0.17)	0.35 (0.33, 0.36)	0.58 (0.56, 0.60)	0.33 (0.32, 0.34)	
	60	0.11 (0.11, 0.12)	0.30 (0.29, 0.31)	0.48 (0.47, 0.50)	0.54 (0.53, 0.56)	0.32 (0.31, 0.33)	
	70	0.21 (0.20, 0.22)	0.41 (0.40, 0.43)	- (-, -)	0.48 (0.46, 0.50)	0.30 (0.29, 0.31)	
	80	0.27 (0.26, 0.29)	- (-, -)	- (-, -)	0.37 (0.35, 0.38)	0.25 (0.24, 0.26)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.96 (0.90, 1.03)	0.62 (0.60, 0.65)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.98 (0.92, 1.05)	0.63 (0.61, 0.66)	
	20	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.06 (0.05, 0.07)	0.99 (0.92, 1.06)	0.64 (0.61, 0.66)	
	30	0.01 (0.01, 0.02)	0.05 (0.04, 0.06)	0.14 (0.13, 0.16)	0.99 (0.92, 1.06)	0.64 (0.61, 0.67)	
	40	0.04 (0.03, 0.05)	0.13 (0.12, 0.15)	0.31 (0.28, 0.34)	0.99 (0.93, 1.07)	0.64 (0.62, 0.67)	
	50	0.10 (0.08, 0.11)	0.28 (0.25, 0.31)	0.59 (0.54, 0.64)	0.99 (0.92, 1.07)	0.64 (0.62, 0.67)	
	60	0.20 (0.18, 0.23)	0.54 (0.49, 0.59)	0.85 (0.78, 0.92)	0.98 (0.91, 1.06)	0.64 (0.62, 0.67)	
	70	0.40 (0.36, 0.45)	0.77 (0.70, 0.84)	- (-, -)	0.93 (0.85, 1.02)	0.63 (0.60, 0.67)	
	80	0.54 (0.47, 0.61)	- (-, -)	- (-, -)	0.78 (0.69, 0.88)	0.59 (0.55, 0.62)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXIV-9 - continued
STOMACH CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific Islander	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	1.69 (1.55, 1.85)	1.13 (1.02, 1.27)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.02, 0.04)	1.70 (1.56, 1.87)	1.14 (1.03, 1.28)	
	20	0.00 (0.00, 0.01)	0.03 (0.02, 0.03)	0.08 (0.07, 0.09)	1.71 (1.57, 1.87)	1.14 (1.03, 1.28)	
	30	0.02 (0.02, 0.03)	0.08 (0.06, 0.09)	0.19 (0.17, 0.21)	1.71 (1.57, 1.87)	1.14 (1.03, 1.28)	
	40	0.05 (0.05, 0.07)	0.17 (0.15, 0.19)	0.39 (0.35, 0.43)	1.69 (1.55, 1.86)	1.14 (1.02, 1.27)	
	50	0.12 (0.10, 0.13)	0.34 (0.30, 0.37)	0.80 (0.74, 0.87)	1.66 (1.51, 1.82)	1.12 (1.00, 1.25)	
	60	0.23 (0.20, 0.26)	0.70 (0.64, 0.77)	1.24 (1.15, 1.34)	1.58 (1.44, 1.75)	1.08 (0.96, 1.22)	
	70	0.51 (0.46, 0.56)	1.08 (0.99, 1.18)	- (-, -)	1.44 (1.29, 1.62)	1.03 (0.91, 1.18)	
American Indian/ Alaska Native ^a	80	0.69 (0.60, 0.78)	- (-, -)	- (-, -)	1.12 (0.95, 1.31)	0.91 (0.77, 1.08)	
	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.01 (0.00, 0.05)	0.99 (0.69, 1.59)	0.63 (0.48, 0.86)	
	10	0.00 (0.00, 0.02)	0.01 (0.00, 0.04)	0.04 (0.01, 0.08)	1.01 (0.70, 1.61)	0.64 (0.48, 0.87)	
	20	0.01 (0.00, 0.04)	0.03 (0.01, 0.08)	0.08 (0.04, 0.14)	1.01 (0.70, 1.62)	0.65 (0.49, 0.88)	
	30	0.03 (0.01, 0.07)	0.07 (0.04, 0.13)	0.17 (0.11, 0.27)	1.01 (0.70, 1.63)	0.65 (0.49, 0.89)	
	40	0.05 (0.02, 0.09)	0.15 (0.08, 0.24)	0.36 (0.24, 0.54)	1.00 (0.69, 1.63)	0.65 (0.49, 0.89)	
	50	0.10 (0.05, 0.18)	0.33 (0.20, 0.51)	0.57 (0.38, 0.83)	0.99 (0.66, 1.63)	0.63 (0.47, 0.88)	
	60	0.24 (0.13, 0.40)	0.50 (0.31, 0.76)	0.81 (0.53, 1.20)	0.93 (0.60, 1.61)	0.62 (0.45, 0.89)	
Hispanic ^b	70	0.30 (0.15, 0.54)	0.66 (0.38, 1.07)	- (-, -)	0.80 (0.46, 1.56)	0.58 (0.39, 0.89)	
	80	0.48 (0.22, 0.93)	- (-, -)	- (-, -)	0.67 (0.31, 1.64)	0.44 (0.23, 0.84)	
	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	1.43 (1.32, 1.56)	0.78 (0.73, 0.83)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	1.45 (1.34, 1.58)	0.78 (0.73, 0.84)	
	20	0.00 (0.00, 0.01)	0.03 (0.02, 0.03)	0.08 (0.07, 0.09)	1.45 (1.34, 1.58)	0.79 (0.73, 0.84)	
	30	0.02 (0.02, 0.03)	0.08 (0.07, 0.09)	0.20 (0.18, 0.22)	1.45 (1.34, 1.58)	0.79 (0.74, 0.84)	
	40	0.05 (0.05, 0.06)	0.18 (0.16, 0.20)	0.40 (0.37, 0.44)	1.44 (1.32, 1.57)	0.78 (0.73, 0.84)	
	50	0.13 (0.11, 0.14)	0.35 (0.32, 0.39)	0.73 (0.68, 0.79)	1.40 (1.29, 1.53)	0.76 (0.71, 0.82)	
-	60	0.24 (0.21, 0.27)	0.63 (0.58, 0.68)	1.06 (0.98, 1.15)	1.32 (1.20, 1.46)	0.72 (0.67, 0.78)	
	70	0.43 (0.38, 0.48)	0.90 (0.82, 0.99)	- (-, -)	1.19 (1.06, 1.33)	0.66 (0.61, 0.72)	
	80	0.60 (0.52, 0.68)	- (-, -)	- (-, -)	0.95 (0.82, 1.12)	0.56 (0.49, 0.63)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXV-7
TESTIS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.02 (0.02, 0.03)	0.12 (0.12, 0.13)	0.36 (0.35, 0.37)	0.02 (0.02, 0.02)	
	10	0.02 (0.02, 0.02)	0.12 (0.12, 0.13)	0.24 (0.23, 0.25)	0.36 (0.35, 0.37)	0.02 (0.02, 0.02)	
	20	0.10 (0.10, 0.11)	0.22 (0.21, 0.23)	0.29 (0.28, 0.30)	0.34 (0.33, 0.35)	0.02 (0.02, 0.02)	
	30	0.12 (0.11, 0.12)	0.19 (0.19, 0.20)	0.22 (0.22, 0.23)	0.24 (0.24, 0.25)	0.02 (0.01, 0.02)	
	40	0.08 (0.07, 0.08)	0.11 (0.10, 0.11)	0.12 (0.11, 0.12)	0.13 (0.12, 0.13)	0.01 (0.01, 0.01)	
	50	0.03 (0.03, 0.04)	0.04 (0.04, 0.05)	0.05 (0.05, 0.06)	0.06 (0.05, 0.06)	0.01 (0.01, 0.01)	
	60	0.01 (0.01, 0.02)	0.02 (0.02, 0.02)	0.02 (0.02, 0.03)	0.03 (0.02, 0.03)	0.01 (0.01, 0.01)	
	70	0.01 (0.01, 0.01)	0.01 (0.01, 0.02)	- (- , -)	0.01 (0.01, 0.02)	0.01 (0.00, 0.01)	
	80	0.01 (0.00, 0.01)	- (- , -)	- (- , -)	0.01 (0.01, 0.01)	0.00 (0.00, 0.01)	
White	0	0.00 (0.00, 0.00)	0.03 (0.03, 0.03)	0.15 (0.14, 0.16)	0.43 (0.42, 0.44)	0.02 (0.02, 0.02)	
	10	0.03 (0.02, 0.03)	0.15 (0.14, 0.16)	0.29 (0.28, 0.30)	0.43 (0.42, 0.44)	0.02 (0.02, 0.02)	
	20	0.12 (0.12, 0.13)	0.26 (0.25, 0.27)	0.35 (0.34, 0.36)	0.41 (0.40, 0.42)	0.02 (0.02, 0.02)	
	30	0.14 (0.14, 0.15)	0.23 (0.22, 0.24)	0.26 (0.25, 0.27)	0.29 (0.28, 0.30)	0.02 (0.02, 0.02)	
	40	0.09 (0.08, 0.09)	0.12 (0.12, 0.13)	0.14 (0.13, 0.14)	0.15 (0.14, 0.16)	0.01 (0.01, 0.01)	
	50	0.04 (0.03, 0.04)	0.05 (0.05, 0.06)	0.06 (0.05, 0.06)	0.06 (0.06, 0.07)	0.01 (0.01, 0.01)	
	60	0.01 (0.01, 0.02)	0.02 (0.02, 0.03)	0.03 (0.02, 0.03)	0.03 (0.02, 0.03)	0.01 (0.01, 0.01)	
	70	0.01 (0.01, 0.01)	0.01 (0.01, 0.02)	- (- , -)	0.02 (0.01, 0.02)	0.01 (0.00, 0.01)	
	80	0.01 (0.01, 0.01)	- (- , -)	- (- , -)	0.01 (0.01, 0.01)	0.00 (0.00, 0.01)	
Black	0	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	0.09 (0.07, 0.11)	0.01 (0.01, 0.02)	
	10	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	0.05 (0.04, 0.06)	0.09 (0.07, 0.11)	0.01 (0.01, 0.02)	
	20	0.02 (0.01, 0.03)	0.05 (0.04, 0.06)	0.07 (0.06, 0.08)	0.08 (0.07, 0.10)	0.01 (0.01, 0.02)	
	30	0.03 (0.02, 0.04)	0.05 (0.04, 0.06)	0.06 (0.05, 0.07)	0.06 (0.05, 0.08)	0.01 (0.01, 0.01)	
	40	0.02 (0.01, 0.03)	0.03 (0.02, 0.04)	0.03 (0.02, 0.04)	0.04 (0.03, 0.05)	0.01 (0.00, 0.01)	
	50	0.01 (0.01, 0.02)	0.01 (0.01, 0.02)	0.02 (0.01, 0.03)	0.02 (0.01, 0.04)	0.01 (0.00, 0.01)	
	60	0.00 (0.00, 0.01)	0.01 (0.00, 0.02)	0.01 (0.00, 0.02)	0.01 (0.00, 0.03)	0.00 (0.00, 0.01)	
	70	0.00 (0.00, 0.01)	0.00 (0.00, 0.02)	- (- , -)	0.00 (0.00, 0.03)	0.01 (0.00, 0.01)	
	80	0.00 (0.00, 0.02)	- (- , -)	- (- , -)	0.00 (0.00, 0.05)	0.00 (0.00, 0.01)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXV-7 - continued
TESTIS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/Pacific Islander	0	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.11 (0.09, 0.15)	0.01 (0.00, 0.05)	
	10	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.07 (0.06, 0.08)	0.11 (0.09, 0.15)	0.01 (0.00, 0.05)	
	20	0.03 (0.02, 0.03)	0.06 (0.05, 0.07)	0.08 (0.07, 0.09)	0.10 (0.08, 0.14)	0.01 (0.00, 0.05)	
	30	0.03 (0.03, 0.04)	0.05 (0.04, 0.06)	0.06 (0.05, 0.07)	0.07 (0.06, 0.11)	0.01 (0.00, 0.05)	
	40	0.02 (0.01, 0.03)	0.03 (0.02, 0.04)	0.03 (0.02, 0.04)	0.04 (0.03, 0.08)	0.00 (0.00, 0.05)	
	50	0.01 (0.00, 0.01)	0.01 (0.01, 0.02)	0.02 (0.01, 0.03)	0.02 (0.01, 0.07)	0.00 (0.00, 0.05)	
	60	0.00 (0.00, 0.01)	0.01 (0.00, 0.02)	0.01 (0.00, 0.03)	0.01 (0.00, 0.06)	0.00 (0.00, 0.05)	
	70	0.01 (0.00, 0.02)	0.01 (0.00, 0.03)	- (-, -)	0.01 (0.00, 0.06)	0.00 (0.00, 0.05)	
	80	0.00 (0.00, 0.02)	- (-, -)	- (-, -)	0.00 (0.00, 0.08)	0.00 (0.00, 0.07)	
American Indian/Alaska Native ^a	0	0.00 (0.00, 0.04)	0.02 (0.01, 0.07)	0.11 (0.07, 0.19)	0.28 (0.18, 0.91)	0.01 (0.00, 0.19)	
	10	0.02 (0.01, 0.05)	0.12 (0.07, 0.18)	0.23 (0.16, 0.32)	0.28 (0.19, 0.93)	0.01 (0.00, 0.20)	
	20	0.10 (0.06, 0.15)	0.21 (0.14, 0.30)	0.22 (0.15, 0.31)	0.26 (0.17, 0.92)	0.01 (0.00, 0.20)	
	30	0.12 (0.07, 0.18)	0.12 (0.07, 0.20)	0.13 (0.08, 0.21)	0.17 (0.09, 0.86)	0.01 (0.00, 0.20)	
	40	0.01 (0.00, 0.03)	0.01 (0.00, 0.05)	0.03 (0.00, 0.11)	0.06 (0.01, 0.80)	0.00 (0.00, 0.21)	
	50	0.00 (0.00, 0.05)	0.02 (0.00, 0.11)	0.05 (0.01, 0.19)	0.05 (0.01, 0.83)	0.00 (0.00, 0.22)	
	60	0.02 (0.00, 0.11)	0.05 (0.01, 0.21)	0.05 (0.01, 0.29)	0.05 (0.01, 0.91)	0.00 (0.00, 0.24)	
	70	0.04 (0.00, 0.22)	0.04 (0.00, 0.33)	- (-, -)	0.04 (0.00, 1.07)	0.00 (0.00, 0.30)	
	80	0.01 (0.00, 0.41)	- (-, -)	- (-, -)	0.01 (0.00, 1.57)	0.00 (0.00, 0.47)	
Hispanic ^b	0	0.00 (0.00, 0.01)	0.03 (0.02, 0.03)	0.12 (0.11, 0.13)	0.27 (0.25, 0.31)	0.02 (0.02, 0.04)	
	10	0.02 (0.02, 0.03)	0.12 (0.11, 0.13)	0.20 (0.19, 0.21)	0.27 (0.25, 0.31)	0.02 (0.02, 0.04)	
	20	0.09 (0.09, 0.10)	0.18 (0.16, 0.19)	0.22 (0.20, 0.23)	0.25 (0.23, 0.29)	0.02 (0.02, 0.04)	
	30	0.08 (0.08, 0.09)	0.12 (0.11, 0.14)	0.14 (0.13, 0.16)	0.16 (0.14, 0.20)	0.02 (0.01, 0.03)	
	40	0.04 (0.03, 0.05)	0.06 (0.05, 0.07)	0.07 (0.06, 0.08)	0.07 (0.06, 0.12)	0.01 (0.01, 0.03)	
	50	0.02 (0.01, 0.03)	0.03 (0.02, 0.04)	0.03 (0.02, 0.05)	0.03 (0.02, 0.08)	0.01 (0.01, 0.03)	
	60	0.01 (0.01, 0.02)	0.02 (0.01, 0.03)	0.02 (0.01, 0.03)	0.02 (0.01, 0.07)	0.01 (0.00, 0.03)	
	70	0.00 (0.00, 0.01)	0.00 (0.00, 0.02)	- (-, -)	0.00 (0.00, 0.07)	0.01 (0.00, 0.03)	
	80	0.00 (0.00, 0.02)	- (-, -)	- (-, -)	0.00 (0.00, 0.09)	0.01 (0.00, 0.04)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVI-9
THYROID CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.07 (0.07, 0.07)	0.73 (0.72, 0.74)	0.05 (0.05, 0.05)	
	10	0.01 (0.01, 0.01)	0.07 (0.07, 0.07)	0.18 (0.18, 0.19)	0.74 (0.73, 0.75)	0.05 (0.05, 0.06)	
	20	0.06 (0.06, 0.06)	0.17 (0.17, 0.17)	0.30 (0.30, 0.31)	0.73 (0.72, 0.74)	0.05 (0.05, 0.06)	
	30	0.11 (0.11, 0.11)	0.25 (0.24, 0.25)	0.39 (0.38, 0.39)	0.68 (0.67, 0.69)	0.05 (0.05, 0.06)	
	40	0.14 (0.13, 0.14)	0.28 (0.28, 0.29)	0.42 (0.41, 0.43)	0.58 (0.57, 0.59)	0.05 (0.05, 0.06)	
	50	0.15 (0.14, 0.15)	0.29 (0.29, 0.30)	0.40 (0.39, 0.41)	0.45 (0.44, 0.46)	0.05 (0.05, 0.06)	
	60	0.15 (0.15, 0.16)	0.27 (0.26, 0.28)	0.32 (0.31, 0.32)	0.32 (0.32, 0.33)	0.05 (0.05, 0.05)	
	70	0.13 (0.13, 0.14)	0.19 (0.18, 0.19)	- (- , -)	0.20 (0.19, 0.20)	0.05 (0.05, 0.05)	
	80	0.07 (0.07, 0.08)	- (- , -)	- (- , -)	0.09 (0.08, 0.10)	0.04 (0.04, 0.04)	
White	0	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.08 (0.08, 0.08)	0.77 (0.76, 0.78)	0.05 (0.05, 0.06)	
	10	0.01 (0.01, 0.02)	0.08 (0.07, 0.08)	0.20 (0.19, 0.20)	0.78 (0.77, 0.79)	0.05 (0.05, 0.06)	
	20	0.06 (0.06, 0.07)	0.18 (0.18, 0.19)	0.33 (0.32, 0.33)	0.77 (0.76, 0.78)	0.05 (0.05, 0.06)	
	30	0.12 (0.12, 0.12)	0.26 (0.26, 0.27)	0.41 (0.40, 0.42)	0.71 (0.70, 0.72)	0.05 (0.05, 0.06)	
	40	0.15 (0.14, 0.15)	0.29 (0.29, 0.30)	0.44 (0.43, 0.45)	0.60 (0.59, 0.61)	0.05 (0.05, 0.06)	
	50	0.15 (0.15, 0.16)	0.30 (0.29, 0.31)	0.41 (0.40, 0.42)	0.46 (0.45, 0.48)	0.05 (0.05, 0.06)	
	60	0.16 (0.15, 0.17)	0.28 (0.27, 0.29)	0.32 (0.31, 0.33)	0.33 (0.32, 0.34)	0.05 (0.05, 0.05)	
	70	0.14 (0.13, 0.14)	0.19 (0.18, 0.20)	- (- , -)	0.20 (0.19, 0.21)	0.05 (0.05, 0.05)	
	80	0.08 (0.07, 0.08)	- (- , -)	- (- , -)	0.09 (0.08, 0.10)	0.04 (0.03, 0.04)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.03 (0.03, 0.04)	0.40 (0.37, 0.42)	0.04 (0.04, 0.04)	
	10	0.00 (0.00, 0.01)	0.03 (0.03, 0.04)	0.09 (0.08, 0.10)	0.41 (0.38, 0.43)	0.04 (0.04, 0.05)	
	20	0.03 (0.02, 0.03)	0.08 (0.08, 0.09)	0.15 (0.14, 0.16)	0.41 (0.38, 0.43)	0.04 (0.04, 0.05)	
	30	0.06 (0.05, 0.06)	0.13 (0.12, 0.14)	0.22 (0.20, 0.23)	0.38 (0.36, 0.41)	0.04 (0.04, 0.05)	
	40	0.07 (0.06, 0.08)	0.16 (0.15, 0.18)	0.25 (0.23, 0.27)	0.33 (0.31, 0.36)	0.04 (0.04, 0.05)	
	50	0.10 (0.09, 0.11)	0.19 (0.17, 0.21)	0.25 (0.23, 0.27)	0.28 (0.25, 0.31)	0.04 (0.04, 0.05)	
	60	0.10 (0.09, 0.12)	0.17 (0.15, 0.19)	0.20 (0.17, 0.22)	0.20 (0.18, 0.23)	0.04 (0.04, 0.05)	
	70	0.08 (0.07, 0.10)	0.12 (0.10, 0.14)	- (- , -)	0.13 (0.11, 0.15)	0.04 (0.04, 0.05)	
	80	0.06 (0.04, 0.08)	- (- , -)	- (- , -)	0.07 (0.05, 0.10)	0.04 (0.03, 0.05)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVI-9 - continued
THYROID CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/Pacific Islander	0	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.07 (0.06, 0.08)	0.79 (0.75, 0.85)	0.09 (0.07, 0.12)	
	10	0.01 (0.01, 0.01)	0.07 (0.06, 0.08)	0.17 (0.15, 0.18)	0.80 (0.76, 0.85)	0.09 (0.07, 0.12)	
	20	0.06 (0.05, 0.06)	0.16 (0.14, 0.17)	0.29 (0.27, 0.31)	0.79 (0.75, 0.85)	0.09 (0.07, 0.12)	
	30	0.10 (0.09, 0.11)	0.23 (0.22, 0.25)	0.38 (0.36, 0.40)	0.74 (0.70, 0.79)	0.09 (0.07, 0.12)	
	40	0.13 (0.12, 0.15)	0.28 (0.26, 0.30)	0.43 (0.41, 0.46)	0.64 (0.60, 0.70)	0.09 (0.07, 0.12)	
	50	0.15 (0.14, 0.16)	0.30 (0.28, 0.33)	0.44 (0.41, 0.47)	0.52 (0.48, 0.57)	0.09 (0.07, 0.12)	
	60	0.16 (0.14, 0.18)	0.30 (0.27, 0.33)	0.36 (0.33, 0.40)	0.38 (0.34, 0.43)	0.09 (0.06, 0.12)	
	70	0.15 (0.13, 0.18)	0.22 (0.19, 0.25)	- (-, -)	0.24 (0.21, 0.29)	0.08 (0.06, 0.11)	
	80	0.09 (0.06, 0.11)	- (-, -)	- (-, -)	0.11 (0.08, 0.16)	0.06 (0.04, 0.10)	
American Indian/Alaska Native ^a	0	0.00 (0.00, 0.02)	0.01 (0.01, 0.04)	0.05 (0.03, 0.08)	0.44 (0.34, 0.72)	0.06 (0.03, 0.14)	
	10	0.01 (0.01, 0.03)	0.05 (0.03, 0.07)	0.15 (0.11, 0.20)	0.45 (0.34, 0.73)	0.06 (0.03, 0.14)	
	20	0.03 (0.02, 0.06)	0.14 (0.10, 0.19)	0.19 (0.14, 0.25)	0.44 (0.33, 0.72)	0.06 (0.03, 0.15)	
	30	0.11 (0.07, 0.15)	0.16 (0.12, 0.21)	0.26 (0.20, 0.33)	0.41 (0.31, 0.70)	0.06 (0.03, 0.15)	
	40	0.05 (0.03, 0.08)	0.16 (0.11, 0.22)	0.23 (0.16, 0.32)	0.31 (0.22, 0.61)	0.06 (0.03, 0.15)	
	50	0.11 (0.07, 0.16)	0.18 (0.12, 0.27)	0.23 (0.16, 0.35)	0.27 (0.18, 0.58)	0.06 (0.03, 0.16)	
	60	0.08 (0.04, 0.15)	0.13 (0.07, 0.24)	0.17 (0.09, 0.32)	0.17 (0.09, 0.51)	0.06 (0.03, 0.17)	
	70	0.06 (0.02, 0.16)	0.11 (0.04, 0.27)	- (-, -)	0.11 (0.04, 0.50)	0.07 (0.03, 0.19)	
	80	0.07 (0.01, 0.25)	- (-, -)	- (-, -)	0.07 (0.01, 0.62)	0.08 (0.03, 0.26)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.05, 0.06)	0.69 (0.65, 0.73)	0.07 (0.06, 0.08)	
	10	0.01 (0.01, 0.01)	0.05 (0.05, 0.06)	0.13 (0.13, 0.14)	0.69 (0.66, 0.74)	0.07 (0.06, 0.09)	
	20	0.04 (0.04, 0.05)	0.12 (0.12, 0.13)	0.23 (0.22, 0.24)	0.69 (0.65, 0.73)	0.07 (0.06, 0.09)	
	30	0.08 (0.08, 0.09)	0.19 (0.18, 0.20)	0.31 (0.30, 0.33)	0.65 (0.61, 0.69)	0.07 (0.06, 0.09)	
	40	0.11 (0.10, 0.11)	0.23 (0.22, 0.25)	0.37 (0.35, 0.39)	0.57 (0.54, 0.62)	0.07 (0.06, 0.09)	
	50	0.13 (0.12, 0.14)	0.27 (0.25, 0.29)	0.39 (0.37, 0.42)	0.48 (0.44, 0.52)	0.07 (0.06, 0.09)	
	60	0.14 (0.13, 0.16)	0.28 (0.25, 0.31)	0.34 (0.31, 0.37)	0.37 (0.33, 0.41)	0.07 (0.06, 0.08)	
	70	0.15 (0.13, 0.17)	0.22 (0.19, 0.25)	- (-, -)	0.25 (0.21, 0.29)	0.06 (0.05, 0.07)	
	80	0.09 (0.07, 0.11)	- (-, -)	- (-, -)	0.13 (0.10, 0.18)	0.04 (0.03, 0.06)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVI-9 - continued
THYROID CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.02, 0.02)	0.39 (0.37, 0.40)	0.04 (0.04, 0.05)	
	10	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.06 (0.06, 0.06)	0.39 (0.38, 0.40)	0.04 (0.04, 0.05)	
	20	0.02 (0.02, 0.02)	0.06 (0.05, 0.06)	0.11 (0.11, 0.12)	0.39 (0.38, 0.40)	0.04 (0.04, 0.05)	
	30	0.04 (0.04, 0.04)	0.10 (0.09, 0.10)	0.17 (0.16, 0.18)	0.38 (0.37, 0.39)	0.05 (0.04, 0.05)	
	40	0.06 (0.05, 0.06)	0.13 (0.13, 0.14)	0.23 (0.22, 0.24)	0.34 (0.33, 0.36)	0.05 (0.04, 0.05)	
	50	0.08 (0.08, 0.08)	0.18 (0.17, 0.19)	0.26 (0.25, 0.27)	0.30 (0.29, 0.31)	0.05 (0.04, 0.05)	
	60	0.11 (0.10, 0.12)	0.20 (0.19, 0.21)	0.23 (0.22, 0.24)	0.23 (0.22, 0.25)	0.04 (0.04, 0.05)	
	70	0.10 (0.09, 0.11)	0.14 (0.13, 0.15)	- (-, -)	0.15 (0.14, 0.16)	0.04 (0.04, 0.04)	
	80	0.06 (0.05, 0.07)	- (-, -)	- (-, -)	0.07 (0.06, 0.08)	0.03 (0.02, 0.03)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.41 (0.40, 0.43)	0.05 (0.04, 0.05)	
	10	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.06 (0.06, 0.07)	0.42 (0.40, 0.43)	0.05 (0.04, 0.05)	
	20	0.02 (0.02, 0.02)	0.06 (0.06, 0.06)	0.12 (0.12, 0.13)	0.41 (0.40, 0.43)	0.05 (0.04, 0.05)	
	30	0.04 (0.04, 0.05)	0.10 (0.10, 0.11)	0.18 (0.18, 0.19)	0.40 (0.39, 0.41)	0.05 (0.04, 0.05)	
	40	0.06 (0.06, 0.07)	0.14 (0.14, 0.15)	0.25 (0.24, 0.26)	0.36 (0.35, 0.38)	0.05 (0.04, 0.05)	
	50	0.08 (0.08, 0.09)	0.19 (0.18, 0.20)	0.28 (0.27, 0.29)	0.31 (0.30, 0.32)	0.05 (0.04, 0.05)	
	60	0.12 (0.11, 0.13)	0.21 (0.20, 0.22)	0.24 (0.23, 0.25)	0.25 (0.23, 0.26)	0.04 (0.04, 0.05)	
	70	0.11 (0.10, 0.11)	0.15 (0.14, 0.16)	- (-, -)	0.15 (0.14, 0.16)	0.04 (0.04, 0.04)	
	80	0.06 (0.05, 0.07)	- (-, -)	- (-, -)	0.07 (0.06, 0.08)	0.03 (0.02, 0.03)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.17 (0.15, 0.20)	0.03 (0.02, 0.03)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.03 (0.02, 0.03)	0.17 (0.15, 0.20)	0.03 (0.02, 0.03)	
	20	0.01 (0.00, 0.01)	0.02 (0.02, 0.03)	0.05 (0.04, 0.06)	0.17 (0.15, 0.20)	0.03 (0.02, 0.03)	
	30	0.02 (0.01, 0.02)	0.04 (0.03, 0.05)	0.09 (0.07, 0.10)	0.17 (0.15, 0.20)	0.03 (0.02, 0.03)	
	40	0.02 (0.02, 0.03)	0.07 (0.06, 0.09)	0.12 (0.10, 0.14)	0.16 (0.13, 0.19)	0.03 (0.02, 0.04)	
	50	0.05 (0.04, 0.07)	0.10 (0.08, 0.12)	0.13 (0.11, 0.15)	0.14 (0.12, 0.18)	0.03 (0.02, 0.04)	
	60	0.05 (0.04, 0.07)	0.09 (0.07, 0.11)	0.10 (0.08, 0.13)	0.10 (0.08, 0.14)	0.03 (0.02, 0.04)	
	70	0.05 (0.03, 0.07)	0.07 (0.05, 0.10)	- (-, -)	0.07 (0.05, 0.11)	0.03 (0.02, 0.04)	
	80	0.04 (0.02, 0.07)	- (-, -)	- (-, -)	0.04 (0.02, 0.09)	0.03 (0.02, 0.05)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVI-9 - continued
THYROID CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/Pacific Islander	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.03)	0.41 (0.36, 0.48)	0.06 (0.04, 0.11)	
	10	0.00 (0.00, 0.00)	0.02 (0.01, 0.03)	0.06 (0.05, 0.07)	0.41 (0.36, 0.49)	0.06 (0.04, 0.12)	
	20	0.02 (0.01, 0.03)	0.06 (0.05, 0.07)	0.10 (0.09, 0.11)	0.41 (0.36, 0.49)	0.06 (0.04, 0.12)	
	30	0.04 (0.03, 0.04)	0.08 (0.07, 0.09)	0.14 (0.12, 0.16)	0.40 (0.34, 0.47)	0.06 (0.04, 0.12)	
	40	0.04 (0.04, 0.05)	0.10 (0.09, 0.12)	0.19 (0.17, 0.22)	0.36 (0.31, 0.44)	0.06 (0.04, 0.12)	
	50	0.06 (0.05, 0.07)	0.15 (0.13, 0.18)	0.24 (0.21, 0.28)	0.32 (0.27, 0.40)	0.06 (0.04, 0.12)	
	60	0.10 (0.08, 0.12)	0.19 (0.16, 0.23)	0.25 (0.21, 0.30)	0.28 (0.22, 0.35)	0.06 (0.03, 0.12)	
	70	0.10 (0.08, 0.13)	0.17 (0.13, 0.22)	- (-, -)	0.20 (0.15, 0.28)	0.06 (0.03, 0.12)	
	80	0.09 (0.05, 0.13)	- (-, -)	- (-, -)	0.13 (0.07, 0.23)	0.04 (0.01, 0.12)	
American Indian/Alaska Native ^a	0	0.00 (0.00, 0.05)	0.01 (0.00, 0.05)	0.02 (0.00, 0.07)	0.26 (0.16, 0.91)	0.02 (0.00, 0.20)	
	10	0.00 (0.00, 0.02)	0.01 (0.00, 0.05)	0.05 (0.02, 0.10)	0.26 (0.15, 0.92)	0.02 (0.00, 0.21)	
	20	0.01 (0.00, 0.04)	0.05 (0.02, 0.10)	0.08 (0.04, 0.14)	0.27 (0.15, 0.93)	0.02 (0.00, 0.21)	
	30	0.03 (0.01, 0.08)	0.07 (0.03, 0.13)	0.13 (0.07, 0.21)	0.26 (0.15, 0.94)	0.02 (0.00, 0.21)	
	40	0.03 (0.01, 0.08)	0.09 (0.05, 0.18)	0.17 (0.09, 0.31)	0.23 (0.12, 0.94)	0.02 (0.00, 0.22)	
	50	0.07 (0.02, 0.14)	0.15 (0.07, 0.28)	0.21 (0.10, 0.40)	0.21 (0.10, 0.96)	0.02 (0.00, 0.23)	
	60	0.09 (0.03, 0.22)	0.15 (0.06, 0.35)	0.16 (0.06, 0.42)	0.16 (0.06, 0.99)	0.02 (0.00, 0.26)	
	70	0.08 (0.01, 0.27)	0.08 (0.01, 0.38)	- (-, -)	0.08 (0.01, 1.10)	0.02 (0.00, 0.32)	
	80	0.01 (0.00, 0.41)	- (-, -)	- (-, -)	0.01 (0.00, 1.57)	0.02 (0.00, 0.49)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.31 (0.27, 0.37)	0.06 (0.05, 0.08)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.04 (0.03, 0.04)	0.31 (0.27, 0.37)	0.06 (0.05, 0.08)	
	20	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	0.07 (0.06, 0.08)	0.31 (0.27, 0.37)	0.06 (0.05, 0.08)	
	30	0.02 (0.02, 0.03)	0.06 (0.05, 0.07)	0.11 (0.10, 0.12)	0.30 (0.27, 0.36)	0.06 (0.05, 0.08)	
	40	0.04 (0.03, 0.04)	0.09 (0.07, 0.10)	0.16 (0.14, 0.19)	0.28 (0.25, 0.35)	0.06 (0.05, 0.08)	
	50	0.05 (0.04, 0.06)	0.13 (0.11, 0.15)	0.21 (0.18, 0.24)	0.26 (0.22, 0.32)	0.06 (0.05, 0.08)	
	60	0.09 (0.07, 0.11)	0.17 (0.14, 0.20)	0.21 (0.17, 0.25)	0.22 (0.18, 0.29)	0.06 (0.05, 0.08)	
	70	0.10 (0.07, 0.12)	0.14 (0.11, 0.18)	- (-, -)	0.15 (0.11, 0.22)	0.05 (0.04, 0.08)	
	80	0.06 (0.03, 0.10)	- (-, -)	- (-, -)	0.08 (0.04, 0.17)	0.03 (0.02, 0.07)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVI-9 - continued
THYROID CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.13 (0.12, 0.13)	1.08 (1.07, 1.10)	0.06 (0.06, 0.06)	
	10	0.02 (0.02, 0.02)	0.13 (0.12, 0.13)	0.31 (0.30, 0.32)	1.10 (1.08, 1.11)	0.06 (0.06, 0.06)	
	20	0.10 (0.10, 0.11)	0.29 (0.28, 0.30)	0.50 (0.49, 0.51)	1.08 (1.06, 1.09)	0.06 (0.06, 0.06)	
	30	0.18 (0.18, 0.19)	0.40 (0.39, 0.41)	0.61 (0.59, 0.62)	0.98 (0.96, 1.00)	0.06 (0.06, 0.06)	
	40	0.22 (0.21, 0.22)	0.43 (0.42, 0.44)	0.61 (0.59, 0.62)	0.80 (0.79, 0.82)	0.06 (0.06, 0.06)	
	50	0.21 (0.21, 0.22)	0.40 (0.39, 0.41)	0.53 (0.52, 0.55)	0.60 (0.58, 0.61)	0.06 (0.06, 0.06)	
	60	0.19 (0.19, 0.20)	0.34 (0.32, 0.35)	0.39 (0.38, 0.41)	0.40 (0.39, 0.42)	0.06 (0.06, 0.06)	
	70	0.16 (0.15, 0.17)	0.22 (0.21, 0.23)	- (-, -)	0.24 (0.22, 0.25)	0.05 (0.05, 0.06)	
	80	0.08 (0.08, 0.09)	- (-, -)	- (-, -)	0.10 (0.09, 0.11)	0.04 (0.04, 0.05)	
White	0	0.00 (0.00, 0.00)	0.03 (0.02, 0.03)	0.14 (0.13, 0.15)	1.14 (1.12, 1.17)	0.06 (0.06, 0.06)	
	10	0.02 (0.02, 0.03)	0.14 (0.13, 0.15)	0.34 (0.33, 0.35)	1.15 (1.13, 1.18)	0.06 (0.06, 0.06)	
	20	0.12 (0.11, 0.12)	0.32 (0.31, 0.33)	0.55 (0.53, 0.56)	1.13 (1.11, 1.15)	0.06 (0.06, 0.06)	
	30	0.20 (0.20, 0.21)	0.43 (0.42, 0.44)	0.65 (0.63, 0.66)	1.02 (1.00, 1.04)	0.06 (0.06, 0.06)	
	40	0.23 (0.23, 0.24)	0.45 (0.44, 0.46)	0.63 (0.62, 0.65)	0.83 (0.81, 0.85)	0.06 (0.06, 0.06)	
	50	0.22 (0.21, 0.23)	0.41 (0.40, 0.42)	0.55 (0.53, 0.56)	0.61 (0.59, 0.63)	0.06 (0.06, 0.06)	
	60	0.20 (0.19, 0.21)	0.34 (0.33, 0.35)	0.40 (0.38, 0.41)	0.41 (0.39, 0.42)	0.06 (0.06, 0.06)	
	70	0.16 (0.15, 0.17)	0.22 (0.21, 0.24)	- (-, -)	0.24 (0.23, 0.25)	0.05 (0.05, 0.06)	
	80	0.08 (0.08, 0.09)	- (-, -)	- (-, -)	0.10 (0.09, 0.11)	0.04 (0.04, 0.05)	
Black	0	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.05 (0.04, 0.06)	0.61 (0.57, 0.66)	0.05 (0.05, 0.06)	
	10	0.01 (0.00, 0.01)	0.05 (0.05, 0.06)	0.15 (0.13, 0.16)	0.62 (0.58, 0.67)	0.05 (0.05, 0.06)	
	20	0.05 (0.04, 0.06)	0.14 (0.13, 0.16)	0.25 (0.23, 0.27)	0.62 (0.58, 0.67)	0.05 (0.05, 0.06)	
	30	0.09 (0.08, 0.11)	0.20 (0.19, 0.22)	0.33 (0.31, 0.36)	0.58 (0.53, 0.62)	0.05 (0.05, 0.06)	
	40	0.11 (0.10, 0.12)	0.24 (0.22, 0.27)	0.37 (0.34, 0.40)	0.49 (0.45, 0.53)	0.05 (0.05, 0.06)	
	50	0.14 (0.12, 0.16)	0.27 (0.24, 0.30)	0.35 (0.32, 0.39)	0.40 (0.36, 0.44)	0.06 (0.05, 0.06)	
	60	0.14 (0.12, 0.16)	0.23 (0.20, 0.27)	0.27 (0.24, 0.31)	0.28 (0.24, 0.32)	0.05 (0.05, 0.06)	
	70	0.11 (0.09, 0.14)	0.16 (0.13, 0.19)	- (-, -)	0.17 (0.14, 0.21)	0.05 (0.04, 0.06)	
	80	0.07 (0.05, 0.10)	- (-, -)	- (-, -)	0.08 (0.06, 0.12)	0.05 (0.04, 0.06)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XVI-9 - continued
THYROID CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.01)	0.02 (0.01, 0.03)	0.11 (0.10, 0.13)	1.14 (1.08, 1.23)	0.11 (0.08, 0.16)	
Islander	10	0.02 (0.01, 0.03)	0.11 (0.10, 0.13)	0.27 (0.25, 0.30)	1.15 (1.09, 1.24)	0.11 (0.08, 0.16)	
	20	0.09 (0.08, 0.11)	0.25 (0.23, 0.27)	0.46 (0.43, 0.50)	1.13 (1.07, 1.22)	0.11 (0.08, 0.16)	
	30	0.16 (0.14, 0.18)	0.37 (0.35, 0.40)	0.60 (0.56, 0.63)	1.05 (0.98, 1.13)	0.11 (0.08, 0.16)	
	40	0.21 (0.20, 0.23)	0.44 (0.41, 0.47)	0.65 (0.60, 0.69)	0.89 (0.83, 0.98)	0.11 (0.08, 0.16)	
	50	0.23 (0.20, 0.25)	0.44 (0.40, 0.48)	0.61 (0.56, 0.66)	0.69 (0.63, 0.77)	0.11 (0.08, 0.16)	
	60	0.21 (0.19, 0.25)	0.39 (0.35, 0.44)	0.46 (0.41, 0.52)	0.47 (0.42, 0.55)	0.11 (0.08, 0.16)	
	70	0.19 (0.16, 0.23)	0.26 (0.22, 0.31)	- (-, -)	0.27 (0.23, 0.35)	0.09 (0.07, 0.15)	
	80	0.08 (0.06, 0.12)	- (-, -)	- (-, -)	0.10 (0.07, 0.18)	0.08 (0.05, 0.14)	
American	0	0.00 (0.00, 0.04)	0.02 (0.01, 0.07)	0.08 (0.04, 0.14)	0.62 (0.46, 1.11)	0.09 (0.04, 0.24)	
Indian/ Alaska Native ^a	10	0.02 (0.01, 0.05)	0.08 (0.04, 0.13)	0.25 (0.18, 0.35)	0.63 (0.46, 1.12)	0.09 (0.04, 0.24)	
	20	0.05 (0.03, 0.10)	0.23 (0.16, 0.32)	0.30 (0.22, 0.41)	0.61 (0.45, 1.10)	0.09 (0.04, 0.25)	
	30	0.18 (0.12, 0.26)	0.25 (0.17, 0.35)	0.39 (0.29, 0.52)	0.56 (0.40, 1.06)	0.10 (0.04, 0.25)	
	40	0.08 (0.04, 0.12)	0.22 (0.14, 0.33)	0.29 (0.19, 0.43)	0.39 (0.25, 0.90)	0.09 (0.04, 0.25)	
	50	0.15 (0.09, 0.24)	0.22 (0.13, 0.36)	0.26 (0.16, 0.43)	0.33 (0.19, 0.86)	0.09 (0.04, 0.26)	
	60	0.07 (0.02, 0.18)	0.12 (0.04, 0.27)	0.19 (0.07, 0.43)	0.19 (0.07, 0.75)	0.10 (0.04, 0.27)	
	70	0.05 (0.01, 0.19)	0.13 (0.03, 0.39)	- (-, -)	0.13 (0.03, 0.79)	0.10 (0.04, 0.31)	
	80	0.11 (0.01, 0.42)	- (-, -)	- (-, -)	0.11 (0.01, 0.98)	0.11 (0.03, 0.39)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.10 (0.10, 0.11)	1.07 (1.01, 1.15)	0.08 (0.07, 0.10)	
	10	0.02 (0.01, 0.02)	0.10 (0.09, 0.11)	0.25 (0.23, 0.26)	1.08 (1.02, 1.16)	0.08 (0.07, 0.10)	
	20	0.09 (0.08, 0.09)	0.23 (0.22, 0.25)	0.41 (0.39, 0.43)	1.07 (1.01, 1.14)	0.08 (0.07, 0.10)	
	30	0.15 (0.14, 0.16)	0.32 (0.31, 0.34)	0.52 (0.50, 0.55)	0.99 (0.93, 1.06)	0.08 (0.07, 0.10)	
	40	0.18 (0.17, 0.19)	0.38 (0.36, 0.41)	0.57 (0.53, 0.60)	0.85 (0.79, 0.92)	0.08 (0.07, 0.10)	
	50	0.20 (0.19, 0.22)	0.39 (0.36, 0.43)	0.56 (0.52, 0.61)	0.68 (0.62, 0.75)	0.08 (0.07, 0.10)	
	60	0.20 (0.17, 0.22)	0.37 (0.33, 0.41)	0.45 (0.40, 0.50)	0.49 (0.44, 0.56)	0.08 (0.06, 0.10)	
	70	0.19 (0.16, 0.23)	0.28 (0.24, 0.32)	- (-, -)	0.32 (0.27, 0.39)	0.07 (0.05, 0.09)	
	80	0.11 (0.08, 0.14)	- (-, -)	- (-, -)	0.16 (0.11, 0.24)	0.04 (0.03, 0.07)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXVII-9

URINARY BLADDER CANCER (Invasive and In Situ)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	2.35 (2.33, 2.37)	0.53 (0.52, 0.53)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	2.38 (2.36, 2.40)	0.54 (0.53, 0.54)	
	20	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.07 (0.07, 0.07)	2.39 (2.37, 2.41)	0.54 (0.53, 0.54)	
	30	0.01 (0.01, 0.01)	0.07 (0.07, 0.07)	0.28 (0.27, 0.28)	2.41 (2.38, 2.43)	0.54 (0.54, 0.55)	
	40	0.06 (0.05, 0.06)	0.27 (0.26, 0.27)	0.81 (0.79, 0.82)	2.43 (2.40, 2.45)	0.55 (0.54, 0.56)	
	50	0.22 (0.21, 0.22)	0.77 (0.76, 0.78)	1.65 (1.63, 1.66)	2.44 (2.41, 2.46)	0.56 (0.55, 0.57)	
	60	0.59 (0.58, 0.60)	1.52 (1.50, 1.54)	2.19 (2.17, 2.22)	2.37 (2.34, 2.39)	0.57 (0.56, 0.57)	
	70	1.08 (1.06, 1.09)	1.86 (1.83, 1.88)	- (- , -)	2.06 (2.03, 2.08)	0.56 (0.55, 0.57)	
	80	1.12 (1.10, 1.14)	- (- , -)	- (- , -)	1.41 (1.38, 1.43)	0.50 (0.49, 0.51)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	2.55 (2.53, 2.58)	0.56 (0.55, 0.56)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.02, 0.02)	2.58 (2.56, 2.61)	0.56 (0.56, 0.57)	
	20	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.08 (0.08, 0.08)	2.59 (2.56, 2.62)	0.56 (0.56, 0.57)	
	30	0.01 (0.01, 0.02)	0.07 (0.07, 0.08)	0.30 (0.30, 0.31)	2.61 (2.58, 2.64)	0.57 (0.56, 0.57)	
	40	0.06 (0.06, 0.06)	0.29 (0.29, 0.30)	0.89 (0.88, 0.91)	2.63 (2.60, 2.65)	0.58 (0.57, 0.58)	
	50	0.24 (0.23, 0.24)	0.85 (0.84, 0.87)	1.81 (1.78, 1.83)	2.64 (2.61, 2.66)	0.58 (0.58, 0.59)	
	60	0.65 (0.64, 0.66)	1.66 (1.64, 1.69)	2.37 (2.35, 2.40)	2.55 (2.52, 2.57)	0.59 (0.58, 0.60)	
	70	1.17 (1.15, 1.19)	1.99 (1.96, 2.01)	- (- , -)	2.19 (2.16, 2.22)	0.58 (0.57, 0.58)	
	80	1.18 (1.16, 1.20)	- (- , -)	- (- , -)	1.47 (1.44, 1.50)	0.51 (0.50, 0.52)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.13 (1.08, 1.19)	0.36 (0.35, 0.38)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.16 (1.11, 1.21)	0.37 (0.36, 0.39)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.04, 0.05)	1.17 (1.12, 1.22)	0.37 (0.36, 0.39)	
	30	0.01 (0.01, 0.01)	0.05 (0.04, 0.05)	0.17 (0.16, 0.19)	1.18 (1.13, 1.24)	0.38 (0.37, 0.40)	
	40	0.04 (0.03, 0.04)	0.17 (0.16, 0.18)	0.45 (0.43, 0.48)	1.20 (1.15, 1.26)	0.39 (0.37, 0.40)	
	50	0.14 (0.13, 0.15)	0.44 (0.41, 0.47)	0.86 (0.82, 0.91)	1.23 (1.17, 1.29)	0.40 (0.38, 0.42)	
	60	0.34 (0.31, 0.36)	0.81 (0.77, 0.86)	1.13 (1.08, 1.19)	1.22 (1.16, 1.29)	0.42 (0.40, 0.44)	
	70	0.60 (0.55, 0.64)	1.00 (0.94, 1.06)	- (- , -)	1.11 (1.05, 1.18)	0.44 (0.42, 0.46)	
	80	0.65 (0.60, 0.72)	- (- , -)	- (- , -)	0.83 (0.76, 0.91)	0.43 (0.40, 0.46)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXVII-9 - continued

URINARY BLADDER CANCER (Invasive and In Situ)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific Islander	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	1.43 (1.34, 1.53)	0.36 (0.31, 0.42)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	1.44 (1.35, 1.54)	0.37 (0.32, 0.43)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	1.44 (1.35, 1.55)	0.37 (0.32, 0.43)	
	30	0.00 (0.00, 0.01)	0.03 (0.02, 0.03)	0.10 (0.09, 0.11)	1.45 (1.36, 1.55)	0.37 (0.32, 0.43)	
	40	0.02 (0.02, 0.03)	0.09 (0.08, 0.10)	0.32 (0.29, 0.34)	1.45 (1.36, 1.56)	0.37 (0.32, 0.43)	
	50	0.07 (0.06, 0.08)	0.30 (0.28, 0.33)	0.75 (0.70, 0.79)	1.45 (1.36, 1.56)	0.37 (0.32, 0.43)	
	60	0.24 (0.21, 0.26)	0.70 (0.65, 0.75)	1.18 (1.11, 1.25)	1.43 (1.33, 1.54)	0.38 (0.32, 0.44)	
	70	0.50 (0.47, 0.54)	1.02 (0.95, 1.09)	- (-, -)	1.30 (1.19, 1.41)	0.38 (0.32, 0.45)	
American Indian/ Alaska Native ^a	80	0.65 (0.58, 0.71)	- (-, -)	- (-, -)	0.99 (0.88, 1.12)	0.36 (0.29, 0.44)	
	0	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.69 (0.52, 1.02)	0.23 (0.15, 0.35)	
	10	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.00 (0.00, 0.02)	0.70 (0.53, 1.04)	0.23 (0.16, 0.36)	
	20	0.00 (0.00, 0.01)	0.00 (0.00, 0.02)	0.01 (0.00, 0.03)	0.71 (0.54, 1.05)	0.23 (0.16, 0.36)	
	30	0.00 (0.00, 0.02)	0.01 (0.00, 0.03)	0.11 (0.07, 0.16)	0.72 (0.54, 1.06)	0.24 (0.16, 0.37)	
	40	0.01 (0.00, 0.03)	0.10 (0.07, 0.16)	0.32 (0.23, 0.43)	0.74 (0.55, 1.09)	0.25 (0.16, 0.38)	
	50	0.10 (0.06, 0.16)	0.32 (0.23, 0.44)	0.56 (0.42, 0.73)	0.76 (0.57, 1.12)	0.25 (0.17, 0.39)	
	60	0.24 (0.16, 0.35)	0.49 (0.35, 0.67)	0.70 (0.51, 0.95)	0.71 (0.51, 1.10)	0.25 (0.16, 0.40)	
Hispanic ^b	70	0.30 (0.18, 0.46)	0.54 (0.35, 0.81)	- (-, -)	0.55 (0.36, 0.99)	0.26 (0.15, 0.44)	
	80	0.35 (0.18, 0.64)	- (-, -)	- (-, -)	0.36 (0.18, 0.93)	0.30 (0.16, 0.56)	
	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.53 (1.45, 1.62)	0.39 (0.36, 0.42)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.55 (1.46, 1.64)	0.40 (0.37, 0.43)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	1.55 (1.47, 1.65)	0.40 (0.37, 0.43)	
	30	0.01 (0.01, 0.01)	0.03 (0.03, 0.04)	0.13 (0.12, 0.14)	1.57 (1.48, 1.66)	0.40 (0.37, 0.43)	
	40	0.03 (0.02, 0.03)	0.13 (0.12, 0.14)	0.40 (0.38, 0.43)	1.57 (1.49, 1.67)	0.41 (0.38, 0.44)	
	50	0.10 (0.09, 0.11)	0.38 (0.36, 0.41)	0.87 (0.82, 0.91)	1.58 (1.49, 1.68)	0.41 (0.38, 0.44)	
-	60	0.30 (0.27, 0.32)	0.80 (0.76, 0.85)	1.32 (1.25, 1.39)	1.56 (1.46, 1.66)	0.42 (0.39, 0.45)	
	70	0.57 (0.53, 0.61)	1.14 (1.07, 1.22)	- (-, -)	1.41 (1.31, 1.52)	0.41 (0.38, 0.45)	
	80	0.76 (0.69, 0.83)	- (-, -)	- (-, -)	1.12 (1.00, 1.25)	0.40 (0.36, 0.44)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXVII-9 - continued

URINARY BLADDER CANCER (Invasive and In Situ)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	3.70 (3.65, 3.74)	0.78 (0.77, 0.79)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.02, 0.02)	3.75 (3.70, 3.79)	0.79 (0.78, 0.80)	
	20	0.00 (0.00, 0.01)	0.02 (0.02, 0.02)	0.10 (0.10, 0.11)	3.77 (3.72, 3.81)	0.80 (0.79, 0.81)	
	30	0.02 (0.02, 0.02)	0.10 (0.10, 0.10)	0.42 (0.41, 0.43)	3.81 (3.77, 3.86)	0.81 (0.80, 0.82)	
	40	0.08 (0.08, 0.09)	0.41 (0.40, 0.42)	1.26 (1.24, 1.28)	3.86 (3.82, 3.91)	0.82 (0.81, 0.83)	
	50	0.34 (0.33, 0.35)	1.22 (1.20, 1.25)	2.66 (2.62, 2.69)	3.92 (3.87, 3.96)	0.84 (0.83, 0.85)	
	60	0.96 (0.94, 0.98)	2.51 (2.48, 2.54)	3.60 (3.55, 3.64)	3.88 (3.83, 3.92)	0.87 (0.86, 0.88)	
	70	1.86 (1.83, 1.90)	3.17 (3.13, 3.22)	- (- , -)	3.50 (3.45, 3.56)	0.88 (0.87, 0.89)	
	80	2.06 (2.01, 2.10)	- (- , -)	- (- , -)	2.58 (2.52, 2.64)	0.84 (0.83, 0.86)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	4.03 (3.99, 4.08)	0.83 (0.82, 0.84)	
	10	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.02, 0.03)	4.08 (4.03, 4.13)	0.84 (0.83, 0.85)	
	20	0.00 (0.00, 0.01)	0.02 (0.02, 0.03)	0.11 (0.11, 0.12)	4.10 (4.05, 4.15)	0.85 (0.84, 0.86)	
	30	0.02 (0.02, 0.02)	0.11 (0.10, 0.11)	0.46 (0.45, 0.47)	4.15 (4.10, 4.20)	0.86 (0.85, 0.87)	
	40	0.09 (0.09, 0.09)	0.45 (0.44, 0.46)	1.39 (1.37, 1.42)	4.20 (4.15, 4.25)	0.87 (0.86, 0.88)	
	50	0.37 (0.36, 0.38)	1.35 (1.32, 1.37)	2.91 (2.87, 2.95)	4.25 (4.20, 4.30)	0.89 (0.88, 0.91)	
	60	1.06 (1.03, 1.08)	2.74 (2.70, 2.78)	3.90 (3.85, 3.95)	4.19 (4.13, 4.24)	0.92 (0.90, 0.93)	
	70	2.02 (1.99, 2.06)	3.41 (3.36, 3.46)	- (- , -)	3.75 (3.70, 3.81)	0.92 (0.91, 0.94)	
	80	2.19 (2.15, 2.24)	- (- , -)	- (- , -)	2.73 (2.67, 2.79)	0.87 (0.86, 0.89)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	1.48 (1.40, 1.57)	0.40 (0.38, 0.42)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	1.52 (1.44, 1.62)	0.41 (0.39, 0.43)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.07 (0.06, 0.08)	1.54 (1.45, 1.63)	0.41 (0.39, 0.44)	
	30	0.01 (0.01, 0.02)	0.07 (0.06, 0.08)	0.26 (0.24, 0.29)	1.57 (1.49, 1.67)	0.42 (0.40, 0.45)	
	40	0.06 (0.05, 0.07)	0.26 (0.23, 0.28)	0.66 (0.62, 0.71)	1.61 (1.52, 1.71)	0.43 (0.41, 0.46)	
	50	0.21 (0.19, 0.24)	0.65 (0.60, 0.70)	1.22 (1.15, 1.30)	1.66 (1.56, 1.76)	0.45 (0.43, 0.48)	
	60	0.50 (0.45, 0.55)	1.17 (1.09, 1.25)	1.58 (1.48, 1.68)	1.67 (1.57, 1.79)	0.48 (0.46, 0.51)	
	70	0.89 (0.82, 0.98)	1.45 (1.34, 1.56)	- (- , -)	1.57 (1.45, 1.71)	0.53 (0.49, 0.56)	
	80	1.01 (0.89, 1.14)	- (- , -)	- (- , -)	1.24 (1.09, 1.42)	0.54 (0.49, 0.60)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXVII-9 - continued

URINARY BLADDER CANCER (Invasive and In Situ)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	2.29 (2.13, 2.48)	0.53 (0.45, 0.64)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	2.32 (2.15, 2.50)	0.54 (0.45, 0.65)	
Islander	20	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.05 (0.04, 0.06)	2.32 (2.16, 2.51)	0.54 (0.45, 0.65)	
	30	0.01 (0.01, 0.01)	0.05 (0.04, 0.06)	0.17 (0.15, 0.19)	2.33 (2.17, 2.52)	0.54 (0.45, 0.65)	
	40	0.04 (0.03, 0.05)	0.16 (0.14, 0.18)	0.55 (0.50, 0.60)	2.34 (2.18, 2.53)	0.55 (0.46, 0.66)	
	50	0.13 (0.11, 0.15)	0.52 (0.47, 0.57)	1.29 (1.20, 1.38)	2.35 (2.18, 2.54)	0.55 (0.46, 0.66)	
	60	0.41 (0.37, 0.45)	1.22 (1.13, 1.31)	1.95 (1.82, 2.09)	2.32 (2.15, 2.52)	0.56 (0.46, 0.68)	
	70	0.90 (0.82, 0.98)	1.72 (1.59, 1.86)	- (- , -)	2.13 (1.94, 2.35)	0.56 (0.46, 0.69)	
	80	1.09 (0.97, 1.23)	- (- , -)	- (- , -)	1.64 (1.42, 1.89)	0.54 (0.42, 0.71)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	1.07 (0.77, 1.80)	0.30 (0.17, 0.54)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	1.10 (0.78, 1.84)	0.31 (0.18, 0.56)	
	20	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	1.11 (0.79, 1.86)	0.31 (0.18, 0.56)	
	30	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.15 (0.09, 0.25)	1.14 (0.81, 1.90)	0.32 (0.18, 0.58)	
	40	0.01 (0.00, 0.04)	0.16 (0.09, 0.26)	0.48 (0.33, 0.69)	1.18 (0.84, 1.97)	0.33 (0.19, 0.60)	
	50	0.16 (0.09, 0.26)	0.50 (0.33, 0.72)	0.87 (0.62, 1.19)	1.23 (0.87, 2.07)	0.35 (0.20, 0.63)	
	60	0.37 (0.23, 0.58)	0.77 (0.53, 1.11)	1.16 (0.79, 1.67)	1.17 (0.80, 2.08)	0.36 (0.20, 0.68)	
	70	0.49 (0.27, 0.81)	0.95 (0.57, 1.51)	- (- , -)	0.96 (0.58, 2.04)	0.38 (0.19, 0.78)	
	80	0.70 (0.32, 1.39)	- (- , -)	- (- , -)	0.72 (0.32, 2.29)	0.47 (0.20, 1.08)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	2.37 (2.22, 2.55)	0.56 (0.51, 0.62)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	2.40 (2.24, 2.58)	0.57 (0.52, 0.63)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.04, 0.06)	2.41 (2.26, 2.59)	0.57 (0.52, 0.64)	
	30	0.01 (0.01, 0.01)	0.05 (0.04, 0.05)	0.19 (0.17, 0.21)	2.44 (2.28, 2.62)	0.58 (0.53, 0.64)	
	40	0.04 (0.03, 0.05)	0.19 (0.17, 0.21)	0.63 (0.58, 0.67)	2.46 (2.30, 2.65)	0.59 (0.54, 0.65)	
	50	0.15 (0.14, 0.17)	0.61 (0.56, 0.65)	1.40 (1.32, 1.49)	2.50 (2.34, 2.69)	0.60 (0.55, 0.67)	
	60	0.49 (0.44, 0.53)	1.33 (1.25, 1.42)	2.16 (2.03, 2.29)	2.51 (2.33, 2.70)	0.62 (0.56, 0.69)	
	70	0.98 (0.90, 1.06)	1.93 (1.79, 2.07)	- (- , -)	2.33 (2.14, 2.56)	0.63 (0.56, 0.71)	
	80	1.35 (1.21, 1.50)	- (- , -)	- (- , -)	1.92 (1.69, 2.21)	0.66 (0.57, 0.76)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXVII-9 - continued

URINARY BLADDER CANCER (Invasive and In Situ)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.17 (1.15, 1.19)	0.32 (0.32, 0.33)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.18 (1.16, 1.21)	0.33 (0.32, 0.33)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.04, 0.04)	1.19 (1.16, 1.21)	0.33 (0.32, 0.33)	
	30	0.01 (0.01, 0.01)	0.04 (0.03, 0.04)	0.13 (0.13, 0.14)	1.19 (1.17, 1.21)	0.33 (0.32, 0.34)	
	40	0.03 (0.03, 0.03)	0.13 (0.12, 0.13)	0.37 (0.36, 0.38)	1.19 (1.17, 1.22)	0.33 (0.33, 0.34)	
	50	0.10 (0.10, 0.10)	0.35 (0.34, 0.36)	0.74 (0.73, 0.76)	1.19 (1.17, 1.21)	0.34 (0.33, 0.34)	
	60	0.26 (0.25, 0.27)	0.68 (0.66, 0.69)	1.03 (1.00, 1.05)	1.14 (1.12, 1.16)	0.34 (0.33, 0.34)	
	70	0.47 (0.45, 0.48)	0.86 (0.84, 0.88)	- (- , -)	0.99 (0.97, 1.01)	0.33 (0.33, 0.34)	
	80	0.53 (0.51, 0.55)	- (- , -)	- (- , -)	0.70 (0.68, 0.73)	0.30 (0.29, 0.31)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	1.25 (1.22, 1.27)	0.33 (0.32, 0.33)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.26 (1.23, 1.29)	0.33 (0.32, 0.34)	
	20	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.04, 0.05)	1.26 (1.24, 1.29)	0.33 (0.32, 0.34)	
	30	0.01 (0.01, 0.01)	0.04 (0.04, 0.04)	0.15 (0.14, 0.15)	1.27 (1.24, 1.29)	0.33 (0.33, 0.34)	
	40	0.03 (0.03, 0.03)	0.14 (0.13, 0.15)	0.41 (0.39, 0.42)	1.27 (1.24, 1.29)	0.33 (0.33, 0.34)	
	50	0.11 (0.10, 0.12)	0.38 (0.37, 0.39)	0.80 (0.78, 0.82)	1.26 (1.23, 1.29)	0.34 (0.33, 0.34)	
	60	0.28 (0.27, 0.30)	0.73 (0.71, 0.75)	1.09 (1.06, 1.11)	1.20 (1.18, 1.23)	0.34 (0.33, 0.35)	
	70	0.50 (0.48, 0.51)	0.90 (0.88, 0.92)	- (- , -)	1.03 (1.00, 1.06)	0.33 (0.33, 0.34)	
	80	0.55 (0.53, 0.57)	- (- , -)	- (- , -)	0.72 (0.70, 0.74)	0.30 (0.29, 0.30)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.82 (0.76, 0.89)	0.33 (0.32, 0.35)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.84 (0.78, 0.91)	0.34 (0.32, 0.36)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	0.84 (0.78, 0.91)	0.34 (0.32, 0.36)	
	30	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	0.10 (0.08, 0.11)	0.85 (0.79, 0.91)	0.35 (0.33, 0.37)	
	40	0.02 (0.02, 0.03)	0.09 (0.08, 0.11)	0.27 (0.24, 0.30)	0.86 (0.79, 0.92)	0.35 (0.33, 0.37)	
	50	0.07 (0.06, 0.09)	0.26 (0.23, 0.29)	0.55 (0.51, 0.60)	0.87 (0.81, 0.94)	0.36 (0.34, 0.38)	
	60	0.20 (0.18, 0.23)	0.52 (0.48, 0.57)	0.78 (0.72, 0.85)	0.87 (0.80, 0.94)	0.37 (0.35, 0.39)	
	70	0.38 (0.34, 0.43)	0.69 (0.63, 0.76)	- (- , -)	0.80 (0.72, 0.88)	0.38 (0.36, 0.41)	
	80	0.46 (0.40, 0.52)	- (- , -)	- (- , -)	0.61 (0.53, 0.70)	0.37 (0.34, 0.40)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table XXVII-9 - continued

URINARY BLADDER CANCER (Invasive and In Situ)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.66 (0.58, 0.76)	0.21 (0.17, 0.28)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.66 (0.58, 0.77)	0.22 (0.17, 0.29)	
Islander	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.66 (0.58, 0.77)	0.22 (0.17, 0.29)	
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.02, 0.04)	0.66 (0.58, 0.77)	0.22 (0.17, 0.29)	
	40	0.01 (0.01, 0.01)	0.03 (0.02, 0.04)	0.12 (0.10, 0.14)	0.67 (0.58, 0.77)	0.22 (0.17, 0.29)	
	50	0.02 (0.02, 0.03)	0.11 (0.09, 0.14)	0.30 (0.26, 0.34)	0.67 (0.58, 0.77)	0.22 (0.17, 0.29)	
	60	0.09 (0.07, 0.11)	0.29 (0.25, 0.33)	0.54 (0.48, 0.61)	0.66 (0.57, 0.77)	0.22 (0.17, 0.29)	
	70	0.21 (0.18, 0.24)	0.48 (0.41, 0.55)	- (-, -)	0.60 (0.52, 0.72)	0.22 (0.17, 0.30)	
	80	0.32 (0.26, 0.39)	- (-, -)	- (-, -)	0.48 (0.38, 0.60)	0.21 (0.15, 0.30)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.34 (0.19, 0.84)	0.17 (0.09, 0.35)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.34 (0.19, 0.85)	0.17 (0.09, 0.35)	
	20	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.02 (0.00, 0.06)	0.34 (0.19, 0.85)	0.18 (0.09, 0.36)	
	30	0.01 (0.00, 0.04)	0.02 (0.00, 0.05)	0.06 (0.02, 0.13)	0.34 (0.19, 0.86)	0.18 (0.09, 0.36)	
	40	0.01 (0.00, 0.04)	0.05 (0.02, 0.12)	0.16 (0.08, 0.30)	0.34 (0.19, 0.87)	0.18 (0.09, 0.36)	
	50	0.05 (0.01, 0.11)	0.16 (0.08, 0.30)	0.28 (0.15, 0.48)	0.35 (0.19, 0.89)	0.18 (0.09, 0.37)	
	60	0.12 (0.05, 0.25)	0.24 (0.12, 0.45)	0.32 (0.16, 0.59)	0.32 (0.16, 0.89)	0.17 (0.08, 0.38)	
	70	0.14 (0.05, 0.33)	0.22 (0.09, 0.51)	- (-, -)	0.22 (0.09, 0.88)	0.17 (0.07, 0.41)	
	80	0.11 (0.02, 0.42)	- (-, -)	- (-, -)	0.11 (0.02, 0.98)	0.20 (0.07, 0.54)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.83 (0.75, 0.94)	0.26 (0.23, 0.30)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.84 (0.75, 0.95)	0.26 (0.23, 0.30)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.02, 0.03)	0.84 (0.75, 0.95)	0.27 (0.23, 0.30)	
	30	0.01 (0.00, 0.01)	0.02 (0.02, 0.03)	0.07 (0.06, 0.08)	0.84 (0.76, 0.95)	0.27 (0.24, 0.30)	
	40	0.02 (0.01, 0.02)	0.07 (0.06, 0.08)	0.20 (0.18, 0.22)	0.84 (0.75, 0.95)	0.27 (0.24, 0.30)	
	50	0.05 (0.04, 0.06)	0.19 (0.16, 0.21)	0.41 (0.37, 0.45)	0.84 (0.75, 0.95)	0.27 (0.24, 0.31)	
	60	0.14 (0.12, 0.16)	0.37 (0.33, 0.41)	0.65 (0.58, 0.71)	0.82 (0.72, 0.93)	0.27 (0.24, 0.31)	
	70	0.25 (0.22, 0.29)	0.56 (0.49, 0.63)	- (-, -)	0.74 (0.64, 0.86)	0.27 (0.24, 0.31)	
	80	0.38 (0.32, 0.45)	- (-, -)	- (-, -)	0.62 (0.50, 0.75)	0.25 (0.21, 0.30)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.