

Table 1.7
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
All Races, 1998-2007

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
All Sites	-0.9*	-1.2*	-0.8*	-1.4*	-1.8*	-1.2*
Oral Cavity & Pharynx:						
Lip	-1.1*	-1.3*	-1.1*	-1.6*	-1.5*	-2.1*
Tongue	-6.3*	-7.0*	-4.3*	-4.5*	-5.1*	-2.9
Salivary gland	1.2*	1.5*	0.6	0.4*	0.4	-0.7*
Floor of mouth	0.5	0.3	0.6	-1.0*	-0.7	-2.0*
Gum & other oral cavity	-6.2*	-6.1*	-6.8*	-8.0*	-7.6*	-9.1*
Nasopharynx	-2.5*	-2.9*	-2.1*	-3.7*	-3.8*	-3.7*
Tonsil	-0.2	-0.3	-0.1	-2.2*	-2.2*	-2.4*
Oropharynx	1.9*	1.9*	0.8	0.3	0.7	-1.5
Hypopharynx	0.3	0.5	-0.7	0.3	0.1	0.3
Other oral cavity & pharynx	-4.8*	-4.7*	-5.1*	-4.5*	-4.1*	-6.7*
Digestive System:	-5.3*	-6.3*	-4.0*	-1.5*	-1.7*	-1.6*
Esophagus	-1.3*	-1.5*	-1.2*	-1.3*	-1.3*	-1.5*
Stomach	-0.9	-0.9	-1.3	-0.1	0.2	-1.4*
Small intestine	-1.7*	-1.9*	-1.7*	-3.2*	-3.5*	-3.0*
Colon & Rectum:	1.1	1.4	0.5	-1.2*	-1.3*	-1.1
Colon	-2.6*	-3.0*	-2.4*	-2.9*	-3.0*	-2.9*
Rectum	-2.6*	-3.1*	-2.3*	-	-	-
Anus, anal canal & anorectum	-2.5*	-2.8*	-2.5*	-	-	-
Liver & intrahep. bile duct:	1.8*	1.6*	2.1*	2.3*	2.3*	2.6*
Liver	2.5*	2.6*	1.5*	2.1*	2.2*	1.4*
Intrahepatic bile duct	3.4*	3.5*	2.4*	1.7*	2.0*	0.5*
Gallbladder	-4.4*	-6.0*	-2.6	3.5*	3.3*	3.6*
Other biliary	-0.6	-0.4	-0.7	-2.1*	-1.9*	-2.1*
Pancreas	2.3*	1.8	2.6*	-3.0*	-3.3*	-2.8*
Retroperitoneum	0.7*	0.5	0.9*	0.3*	0.2	0.3*
Peritoneum, omentum & mesentery	-1.4	-2.8*	0.1	-4.5*	-4.2*	-5.3*
Other digestive system	2.4*	1.5	2.5*	3.7*	0.5	4.3*
Respiratory System:	4.6*	4.8*	4.3	-1.4	-1.0	-2.0
Nose, nasal cavity & middle ear	-1.6*	-2.4*	-0.6*	-1.2*	-2.1*	-0.2
Larynx	-0.6	-1.1	-0.2	-1.9*	-2.2*	-1.6
Lung & bronchus	-3.1*	-2.8*	-4.4*	-2.2*	-2.4*	-2.4*
Pleura	-1.5*	-2.4*	-0.5*	-1.2*	-2.1*	-0.2
Trachea & other respiratory organs	-	-	-	-7.4*	-8.0*	-6.1*
Bones & joints	0.7	0.6	1.0	-0.1	-0.4	0.0
Soft tissue (including heart)	1.5*	1.3*	1.5*	-1.2*	-0.8	-1.5*
Skin (excl. basal & squamous):	1.5*	1.6*	1.5*	0.2	0.4	-0.4
Melanoma of the skin	1.6*	1.7*	1.6*	0.1	0.3	-0.4
Other non-epithelial skin	0.6	0.5	0.8	0.3	0.4	-0.3
Breast	-1.9*	0.0	-1.7*	-2.2*	-2.6*	-2.0*
Breast (<i>in situ</i>)	0.0	1.0	0.2	-	-	-

^a The APC is the Annual Percent Change over the time interval.

SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

* The APC is significantly different from zero ($p<.05$).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.7 - continued
 SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
 All Races, 1998-2007

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
Female Genital System:	-1.4*	-	-1.3*	-0.9*	-	-0.7*
Cervix uteri	-3.4*	-	-3.2*	-2.6*	-	-2.4*
Corpus uteri	-0.8*	-	-0.5*	-0.8*	-	-0.5*
Uterus, NOS	1.5	-	2.0	0.4*	-	0.8*
Ovary ^c	-1.8*	-	-1.6*	-0.8*	-	-0.6*
Vagina	-0.7	-	-0.6	-2.8*	-	-2.5*
Vulva	-1.4*	-	-1.2*	-0.3	-	0.2
Other female genital system	1.5	-	1.7	-2.2	-	-2.0
Male Genital System:	-1.2*	-1.6*	-	-3.1*	-3.8*	-
Prostate	-1.2*	-1.7*	-	-3.2*	-3.9*	-
Testis	1.1*	0.9*	-	-1.6*	-1.8*	-
Penis	-1.1	-1.4	-	0.7	0.1	-
Other male genital system	-3.7*	-3.9*	-	-2.6	-3.4	-
Urinary System:	0.7*	0.3	1.0*	-0.3*	-0.4*	-0.5*
Urinary bladder	-0.4*	-0.5*	-0.8*	0.0	-0.1	-0.6*
Kidney & renal pelvis	2.4*	2.0*	2.8*	-0.6*	-0.7*	-0.6*
Ureter	1.7	0.9	2.5	-0.6	-1.5	0.0
Other urinary system	0.3	-0.2	1.6	3.5	5.1	1.2
Eye & Orbit	-0.3	-0.2	0.0	-2.4*	-2.2	-2.8*
Brain & Nervous System: ^d	-0.6*	-0.7*	-0.6	-1.2*	-1.3*	-1.2*
Brain	-0.7*	-0.8*	-0.6	-	-	-
Cranial nerves & other nervous system	-0.3	-0.1	-0.7	-	-	-
Endocrine System:	5.8*	4.7*	6.2*	0.0	-0.1	0.1
Thyroid	6.1*	5.3*	6.5*	0.9*	1.6*	0.6
Other endocrine & thymus	1.7	1.7	1.5*	-1.5*	-2.3*	-0.7
Lymphoma:	0.2	0.1	0.3	-3.1*	-2.9*	-3.4*
Hodgkin lymphoma	0.8*	0.7	0.8	-1.9*	-2.0*	-1.8*
Non-Hodgkin lymphoma	0.1	0.1	0.2	-3.2*	-3.0*	-3.5*
Myeloma	-0.6	-0.4	-1.0	-1.3*	-1.0*	-1.8*
Leukemia:	-0.8*	-1.2*	-0.5	-1.0*	-0.9*	-1.4*
Lymphocytic:	0.5	0.2	0.9	-1.4*	-1.4*	-1.7*
Acute lymphocytic	1.1	1.5	0.6	-1.1*	-1.3*	-1.0
Chronic lymphocytic	0.7	0.2	1.5	-1.4*	-1.3*	-1.9*
Other lymphocytic	-3.7*	-3.2*	-5.0*	-2.2*	-2.1*	-2.9*
Myeloid & Monocytic:	-1.6*	-1.9*	-1.4*	-0.6*	-0.4	-1.1*
Acute myeloid	-1.8*	-2.2*	-1.4*	1.2*	1.5*	0.6
Chronic myeloid	-1.4*	-1.0	-2.2*	-10.5*	-10.6*	-10.5*
Acute monocytic	0.7	-1.6	3.3*	-6.2*	-6.5*	-5.7*
Other myeloid & monocytic	-2.8	-2.9	-2.6	1.1	1.2	0.3
Other leukemia:	-5.8*	-7.9*	-3.9*	-1.4*	-1.4*	-1.6*
Other acute leukemia	-10.2*	-10.7*	-9.8*	-4.1*	-4.1*	-4.2*
Aleukemic, subleukemic & NOS	-2.1	-5.5*	0.9	0.8*	0.9*	0.5
Kaposi Sarcoma ^e	-4.1*	-3.9*	-6.3*	-	-	-
Mesothelioma ^e	-1.4*	-2.2*	1.0	-	-	-
Ill-defined & unspecified	-3.5*	-3.7*	-3.4*	-1.5	-1.4	-1.8*

^a The APC is the Annual Percent Change over the time interval.

SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^d Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

^e Trend not shown for mortality. Category did not exist in mortality coding until 1999.

* The APC is significantly different from zero ($p<.05$).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.