Table 4.13 Cancer of the Female Breast(In Situ)

SEER Incidence^a Rates, Age-Adjusted and Age-Specific Rates, by Race

_	All Races, Females	White Females	Black Females
Age-Adjusted Rates, 2004-2008			
Age at Diagnosis:			
All ages	30.9	31.4	28.8
Under 65	23.2	23.6	19.9
65 and over	84.2	85.1	90.3
All ages (IARC world std) ^b	23.3	23.7	21.4
Age-Specific Rates, 2004-2008			
Age at Diagnosis:			
<1	-	-	-
1-4	-	-	-
5-9	-	-	-
10-14	-	-	-
15-19	-	-	=
20-24	0.1	-	=
25-29	0.7	0.7	1.0
30-34	2.7	2.6	2.3
35-39	10.8	10.7	10.6
40-44	37.9	38.4	29.2
45-49	60.5	62.3	43.3
50-54	71.2	72.4	58.1
55-59	77.4	78.1	75.8
60-64	90.1	91.2	90.1
65-69	99.4	100.5	101.2
70-74	93.8	95.0	103.7
75-79	90.8	91.8	95.5
80-84	70.4	71.9	70.0
85+	35.1	34.2	53.3

SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

Rates are per 100,000 and are age-adjusted to the IARC world standard population. Statistic not shown. Rate based on less than 16 cases for the time interval.