## Table 27.3 Cancer of the Urinary Bladder (Invasive and In Situ)

Trends in SEER Incidence<sup>ab</sup> and U.S. Mortality<sup>c</sup> Using the Joinpoint Regression Program, 1975-2009 With up to Five Joinpoints, 1992-2009 With up to Three Joinpoints, Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC <sup>d</sup>	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2000-09	2005-09
SEER 9 Delay-Adjusted Incidence <sup>a</sup> , 1975-2009														
All Races	1975-03	0.2*	2003-09	-1.4*									-0.9*	-1.4*
White	1975-03	0.4*	2003-09	-1.7*									-1.0*	-1.7*
Black	1975-09	0.3											0.3	0.3
SEER 13 Delay-Adjusted Incidence <sup>b</sup> , 1992-2009														
All Races	1992-04	-0.1	2004-09	-1.3*									-0.8*	-1.3*
White	1992-03	0.0	2003-09	-1.2*									-0.8*	-1.2*
Black	1992-09	0.1											0.1	0.1
SEER 9 Observed In														
All Races	1975-03	0.2*	2003-09										-1.1*	-1.7*
White	1975-03	0.4*	2003-09	-2.0*									-1.2*	-2.0*
Black	1975-09	0.2											0.2	0.2
SEER 13 Observed Incidence <sup>b</sup> , 1992-2009														
All Races	1992-04	-0.2	2004-09										-1.0*	-1.7*
White	1992-04	-0.1	2004-09										-1.0*	-1.8*
White NH <sup>ef</sup>	1992-03	0.2	2003-09	-1.4*									-0.8*	-1.4*
Black	1992-09	0.0											0.0	0.0
Black NH <sup>ef</sup>	1992-06	1.0	2006-09	-8.5									-2.3	-6.2
API <sup>e</sup>	1992-09	-0.1											-0.1	-0.1
AI/AN <sup>eg</sup>	1992-09	4.7											4.7	4.7
Hispanic <sup>f</sup>	1992-09	-0.6											-0.6	-0.6
U.S. Cancer Mortality <sup>c</sup> , 1975-2009														
All Races	1975-86	-1.6*	1986-09	-0.4*									-0.4*	-0.4*
White	1975-86	-1.7*	1986-09	-0.3*									-0.3*	-0.3*
Black	1975-09	-0.9*											-0.9*	-0.9*
U.S. Cancer Mortal														
All Races	1992-09	-0.6*											-0.6*	-0.6*
White	1992-09	-0.4*											-0.4*	-0.4*
White NH <sup>ef</sup>	1992-09	-0.2											-0.2	-0.2
Black	1992-09	-1.2*											-1.2*	-1.2*
Black NH <sup>ef</sup>	1992-09	-0.9*											-0.9*	-0.9*
API <sup>e</sup>	1992-09	-1.9*											-1.9*	-1.9*
AI/AN <sup>eg</sup>	1992-09	-1.0											-1.0	-1.0
Hispanic <sup>f</sup>	1992-09	-0.6											-0.6	-0.6

Joinpoint Regression Program Version 3.5, April 2011, National Cancer Institute. (<a href="http://surveillance.cancer.gov/joinpoint/">http://surveillance.cancer.gov/joinpoint/</a>).
The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

d The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The Hispanic and Non-Hispanic mortality trends exclude deaths from CT, DC, ME, MD, MN, NH, NY, ND, SC, OK and VT.

The Hispanic and Non-Hispanic mortality trends exclude deaths from CT, DC, ME, MD, MN, NH, NY, ND, SC, OK and V Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

<sup>\*</sup> The APC/AAPC is significantly different from zero (p<.05).</p>

Joinpoint regression line analysis could not be performed on data series.