

Table 10.1

## Kaposi Sarcoma

## Number of Cases

By SEER Location and Year of Diagnosis, 1975-2010

## All Races, Males, All Ages

## Year of Diagnosis

SEER Location	1975-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001	2002-2003	2004-2005	2006-2007	2008-2010
SEER 9 <sup>b</sup>	134	79	326	889	1,574	2,131	2,230	2,057	1,800	841	392	401	336	372	334	508
SEER 9 except SF-Oakland	102	44	85	261	428	776	840	976	874	464	245	265	223	245	237	371
San Francisco-Oakland SMSA <sup>c</sup>	32	35	241	628	1,146	1,355	1,390	1,081	926	377	147	136	113	127	97	137
San Francisco City/County	15	21	201	540	955	1,076	1,089	792	689	254	85	81	72	83	62	90

Age-Adjusted Cancer Incidence Rates<sup>a</sup>

By SEER Location and Year of Diagnosis, 1975-2010

## All Races, Males, All Ages

## Year of Diagnosis

SEER Location	1975-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001	2002-2003	2004-2005	2006-2007	2008-2010
SEER 9 <sup>b</sup>	0.4	0.5	1.7	4.2	7.1	9.2	9.3	8.2	7.0	3.2	1.5	1.5	1.3	1.4	1.2	1.2
SEER 9 except SF-Oakland	0.4	0.4	0.6	1.6	2.3	4.0	4.2	4.6	4.0	2.1	1.1	1.2	1.0	1.1	1.1	1.1
San Francisco-Oakland SMSA <sup>c</sup>	0.7	1.3	7.0	17.2	30.1	34.0	34.0	25.6	22.0	9.0	3.5	3.1	2.7	3.0	2.3	2.1
San Francisco City/County	1.1	3.1	25.1	65.4	111.4	124.9	125.9	89.1	77.4	28.3	9.3	8.8	8.0	9.2	6.8	6.4

<sup>a</sup> Rates are per 100,000 males and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).<sup>b</sup> The SEER 9 areas are San Francisco-Oakland SMSA, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.<sup>c</sup> The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi Sarcoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.

Table 10.2

## Kaposi Sarcoma

## Number of Cases

By SEER Location and Year of Diagnosis, 1975-2010

All Races, Males, Ages 20-54

## Year of Diagnosis

SEER Location	1975-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001	2002-2003	2004-2005	2006-2007	2008-2010
SEER 9 <sup>b</sup>	13	26	254	806	1,480	1,994	2,101	1,922	1,674	756	319	347	262	310	269	379
SEER 9 except SF-Oakland	11	5	35	207	379	695	771	910	809	411	198	227	184	202	196	278
San Francisco-Oakland SMSA <sup>c</sup>	<4	21	219	599	1,101	1,299	1,330	1,012	865	345	121	120	78	108	73	101
San Francisco City/County	<4	18	193	520	924	1,040	1,051	748	647	237	75	74	54	73	53	73

Age-Adjusted Cancer Incidence Rates<sup>a</sup>

By SEER Location and Year of Diagnosis, 1975-2010

All Races, Males, Ages 20-54

## Year of Diagnosis

SEER Location	1975-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001	2002-2003	2004-2005	2006-2007	2008-2010
SEER 9 <sup>b</sup>	0.1	0.3	2.4	7.4	13.1	16.9	17.1	14.9	12.6	5.6	2.3	2.5	1.9	2.3	2.0	1.8
SEER 9 except SF-Oakland	0.1	0.1	0.4	2.2	3.9	6.9	7.4	8.4	7.2	3.6	1.7	2.0	1.6	1.8	1.7	1.6
San Francisco-Oakland SMSA <sup>c</sup>	-	1.3	12.1	32.4	57.3	64.5	64.0	46.9	40.1	15.9	5.4	5.2	3.4	4.9	3.4	3.0
San Francisco City/County	-	5.3	47.8	125.4	214.3	239.5	240.8	165.5	142.3	51.8	15.9	15.7	11.4	16.1	11.5	10.3

<sup>a</sup> Rates are per 100,000 males aged 20-54 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).<sup>b</sup> The SEER 9 areas are San Francisco-Oakland SMSA, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.<sup>c</sup> The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi Sarcoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.

Table 10.3

## Kaposi Sarcoma

## Number of Cases

By SEER Location and Year of Diagnosis, 1975-2010

## All Races, Males, Ages 55+

## Year of Diagnosis

SEER Location	1975-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001	2002-2003	2004-2005	2006-2007	2008-2010
SEER 9 <sup>b</sup>	121	53	71	82	93	135	129	134	126	85	73	53	74	61	65	126
SEER 9 except SF-Oakland	91	39	49	54	49	81	69	65	65	53	47	37	39	42	41	90
San Francisco-Oakland SMSA <sup>c</sup>	30	14	22	28	44	54	60	69	61	32	26	16	35	19	24	36
San Francisco City/County	14	<4	8	20	31	35	38	44	42	17	10	7	18	10	9	17

Age-Adjusted Cancer Incidence Rates<sup>a</sup>

By SEER Location and Year of Diagnosis, 1975-2010

## All Races, Males, Ages 55+

## Year of Diagnosis

SEER Location	1975-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001	2002-2003	2004-2005	2006-2007	2008-2010
SEER 9 <sup>b</sup>	1.9	1.9	2.4	2.5	2.5	3.6	3.6	3.3	3.0	2.1	1.7	1.3	1.5	1.2	1.2	1.4
SEER 9 except SF-Oakland	1.7	1.7	2.0	2.0	1.8	2.7	2.4	2.0	1.9	1.6	1.3	1.1	1.0	1.0	0.9	1.2
San Francisco-Oakland SMSA <sup>c</sup>	2.9	3.1	4.6	4.7	6.6	8.1	9.6	10.4	9.0	4.8	3.7	2.3	4.5	2.3	2.8	2.6
San Francisco City/County	4.7	-	5.6	12.9	20.5	23.5	26.4	30.4	29.8	11.5	6.7	4.4	10.8	5.7	4.8	5.7

<sup>a</sup> Rates are per 100,000 males aged 55+ and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).<sup>b</sup> The SEER 9 areas are San Francisco-Oakland SMSA, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.<sup>c</sup> The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi Sarcoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.

Table 10.4  
Kaposi Sarcoma

Age-adjusted SEER Incidence<sup>a</sup> Rates by Year, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<b>Year of Diagnosis:</b>									
1975-2010	1.74	3.42	0.13	1.80	3.51	0.11	2.18	4.45	0.24
1975	0.23	0.56	-	0.22	0.56	-	-	-	-
1976	0.25	0.47	-	0.24	0.48	-	-	-	-
1977	0.29	0.54	-	0.30	0.55	-	-	-	-
1978	0.24	0.35	0.17	0.23	0.34	-	-	-	-
1979	0.18	0.30	-	0.18	0.30	-	-	-	-
1980	0.28	0.56	-	0.30	0.61	-	-	-	-
1981	0.32	0.54	-	0.34	0.61	-	-	-	-
1982	0.65	1.20	0.18	0.71	1.32	0.18	-	-	-
1983	1.07	2.19	-	1.19	2.40	-	-	-	-
1984	1.81	3.59	-	1.99	3.91	-	1.27	2.89	-
1985	2.47	4.87	0.16	2.80	5.49	0.18	1.09	2.24	-
1986	3.27	6.54	-	3.71	7.34	-	1.65	3.59	-
1987	3.81	7.61	-	4.26	8.45	-	2.09	4.29	-
1988	4.43	8.87	0.15	4.82	9.56	0.15	3.40	7.28	-
1989	4.76	9.53	0.14	5.13	10.18	-	4.57	9.69	-
1990	4.61	9.21	0.18	5.03	9.99	0.15	3.65	7.54	-
1991	4.70	9.40	0.19	5.04	10.00	0.18	4.59	9.53	-
1992	4.38	8.69	0.18	4.60	9.02	0.18	5.02	10.59	-
1993	3.86	7.64	0.17	4.05	7.96	0.15	4.28	8.60	-
1994	3.78	7.46	0.17	3.88	7.58	0.13	4.15	8.42	-
1995	3.29	6.45	0.20	3.28	6.37	0.17	4.94	10.27	-
1996	2.24	4.40	0.16	2.01	3.99	-	4.61	9.09	-
1997	1.08	2.09	0.14	0.92	1.78	-	2.70	5.36	-
1998	0.81	1.58	-	0.70	1.34	-	1.79	3.78	-
1999	0.78	1.47	0.14	0.66	1.24	-	1.94	3.72	-
2000	0.80	1.55	-	0.67	1.30	-	1.99	4.00	-
2001	0.83	1.51	0.17	0.67	1.20	0.15	1.95	3.89	-
2002	0.68	1.28	0.12	0.57	1.07	-	1.51	3.05	-
2003	0.67	1.29	-	0.60	1.12	-	1.52	3.21	-
2004	0.70	1.36	-	0.60	1.16	-	1.69	3.41	-
2005	0.78	1.47	0.12	0.61	1.15	-	2.03	4.01	-
2006	0.61	1.18	-	0.50	0.96	-	1.42	2.88	-
2007	0.69	1.32	-	0.58	1.07	-	1.66	3.48	-
2008	0.67	1.33	-	0.56	1.10	-	1.63	3.15	-
2009	0.63	1.19	0.10	0.54	1.02	-	1.38	2.78	-
2010	0.63	1.19	0.09	0.43	0.80	-	1.77	3.59	-

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 10.5  
Kaposi Sarcoma

SEER Incidence<sup>a</sup> Rates, Age-Adjusted and Age-Specific Rates, by Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Incidence</u>									
<u>Age at Diagnosis</u>									
Age-Adjusted Rates, 2006-2010									
All ages	0.6	1.0	0.1	0.5	0.9	0.1	1.0	2.0	0.2
Under 65	0.5	1.0	0.0	0.4	0.8	0.0	1.1	2.1	0.2
65 and over	1.0	1.7	0.5	1.0	1.7	0.5	0.8	1.4	-
All ages (IARC world std) <sup>b</sup>	0.4	0.8	0.1	0.4	0.7	0.0	0.9	1.7	0.1
Age-Specific Rates, 2006-2010									
<1	-	-	-	-	-	-	-	-	-
1-4	-	-	-	-	-	-	-	-	-
5-9	-	-	-	-	-	-	-	-	-
10-14	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-
20-24	0.2	0.4	-	0.1	0.2	-	0.9	1.6	-
25-29	0.5	1.0	-	0.4	0.7	-	1.9	3.7	-
30-34	0.8	1.4	-	0.6	1.2	-	2.0	3.6	-
35-39	1.0	1.9	-	0.8	1.6	-	2.3	4.5	-
40-44	1.2	2.4	-	1.1	2.1	-	2.6	5.1	-
45-49	1.0	1.9	-	0.9	1.7	-	1.7	3.2	-
50-54	0.6	1.2	-	0.6	1.1	-	1.0	1.9	-
55-59	0.5	0.9	-	0.4	0.8	-	0.6	-	-
60-64	0.5	0.9	-	0.4	0.8	-	-	-	-
65-69	0.6	1.0	0.2	0.6	1.0	-	-	-	-
70-74	0.7	1.1	0.4	0.7	1.1	0.4	-	-	-
75-79	0.9	1.7	0.4	1.0	1.8	0.4	-	-	-
80-84	1.2	2.0	0.7	1.2	2.0	0.7	-	-	-
85+	2.1	3.6	1.3	2.1	3.7	1.3	-	-	-

<sup>a</sup> SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

<sup>b</sup> Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 10.6  
Kaposi Sarcoma

5-Year Relative and Period Survival (Percent) by Race, Sex, Diagnosis Year, Stage and Age

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<b>5-Year Relative Survival (Percent)</b>									
<b>Year of Diagnosis:</b>									
1960-1963 <sup>a</sup>	-	-	-	-	-	-	-	-	-
1970-1973 <sup>a</sup>	-	-	-	-	-	-	-	-	-
1975-1977 <sup>b</sup>	80.7 <sup>f</sup>	80.1 <sup>f</sup>	-	79.9 <sup>f</sup>	79.0 <sup>f</sup>	-	-	-	-
1978-1980 <sup>b</sup>	78.1 <sup>f</sup>	79.0 <sup>f</sup>	73.1 <sup>g</sup>	77.7 <sup>f</sup>	78.0 <sup>f</sup>	72.5 <sup>g</sup>	-	-	-
1981-1983 <sup>b</sup>	35.2	32.9	64.3 <sup>g</sup>	33.7	31.3	64.2 <sup>g</sup>	-	-	-
1984-1986 <sup>b</sup>	12.9	11.6	70.5 <sup>g</sup>	12.6	11.3	70.2 <sup>g</sup>	16.8	15.7	-
1987-1989 <sup>b</sup>	9.8	9.2	68.5 <sup>g</sup>	9.4	8.8	72.3 <sup>g</sup>	12.0	10.8	-
1990-1992 <sup>b</sup>	9.2	8.5	52.6 <sup>f</sup>	8.8	8.1	52.3 <sup>f</sup>	11.5	10.5	-
1993-1995 <sup>b</sup>	21.7	20.9	52.3 <sup>f</sup>	22.2	21.4	60.0 <sup>g</sup>	16.5	15.6	-
1996-1998 <sup>b</sup>	52.3	51.4	72.1 <sup>g</sup>	57.0	55.7	86.5 <sup>f</sup>	39.1	38.5	-
1999-2002 <sup>b</sup>	58.3	57.7	64.0 <sup>f</sup>	66.3	64.5	81.1 <sup>f</sup>	41.7	42.9	-
2003-2009 <sup>b</sup>	70.7 <sup>e</sup>	70.2 <sup>e</sup>	72.5 <sup>f</sup>	78.9	78.3	81.3 <sup>f</sup>	51.7	52.0	-
<b>5-Year Period Survival (Percent)<sup>cd</sup></b>									
2009	72.9	72.1	81.0	78.8	77.4	88.2	55.3	54.6	-
<b>5-Year Relative Survival (Percent), 2003-2009<sup>c</sup></b>									
<b>Age at Diagnosis:</b>									
Ages <45	64.5	64.7	57.0 <sup>f</sup>	70.7	70.7	-	50.1	50.5	45.2 <sup>f</sup>
Ages 45-54	68.8	69.3	-	71.8	71.9	-	56.4 <sup>f</sup>	57.4 <sup>f</sup>	-
Ages 55-64	80.6	80.4	-	85.4	84.5	-	34.0 <sup>g</sup>	23.4 <sup>g</sup>	-
Ages 65-74	80.5	80.7 <sup>f</sup>	78.1 <sup>g</sup>	79.6	80.3 <sup>f</sup>	75.6 <sup>g</sup>	-	-	-
Ages 75+	88.1	90.7	81.7 <sup>f</sup>	88.2	90.6	82.6	-	-	-
Ages <65	67.1	67.3	62.0 <sup>f</sup>	72.9	72.7	78.5 <sup>f</sup>	50.6	50.8	48.1 <sup>f</sup>
Ages 65+	88.2	90.1	82.1 <sup>f</sup>	88.4	89.7	83.9	82.1 <sup>g</sup>	-	-

<sup>a</sup> Based on End Results data from a series of hospital registries and one population-based registry.

<sup>b</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

<sup>c</sup> Based on follow-up of patients into 2010.

<sup>d</sup> SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

<sup>e</sup> Based on follow-up of patients into 2010.

<sup>f</sup> Period survival provides a 2009 estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three year calendar blocks (2007-2009: 0-1 year survival), (2006-2008: 1-2 year survival), (2005-2007: 2-3 year survival), (2004-2006: 3-4 year survival), (2003-2005: 4-5 years survival).

<sup>g</sup> The difference between 1975-1977 and 2003-2009 is statistically significant ( $p<.05$ ).

<sup>h</sup> The standard error is between 5 and 10 percentage points.

<sup>i</sup> The standard error is greater than 10 percentage points.

<sup>j</sup> Statistic could not be calculated due to fewer than 25 cases during the time period.

Table 10.7  
Kaposi Sarcoma

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975-1979	1980-1984	1985-1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
<b>Survival Time</b>																							
1-year	95.0	65.4	59.0	59.8	60.8	61.9	57.7	54.2	61.8	69.0	72.9	73.8	76.6	74.5	80.2	77.7	75.3	82.6	81.8	80.8	82.6	79.4	81.5
2-year	89.6	44.3	32.8	32.1	29.6	27.8	29.9	26.9	46.7	60.4	65.1	65.5	65.0	66.9	72.9	70.8	72.5	73.5	78.4	77.9	80.1	75.3	
3-year	88.7	35.8	19.7	19.8	15.1	16.2	19.5	20.9	42.8	54.0	62.1	61.1	61.3	60.9	66.4	67.5	67.0	70.9	74.5	75.5	78.4		
4-year	83.8	31.7	13.8	14.3	11.0	10.8	15.6	18.5	38.4	51.5	58.8	59.0	56.5	57.4	64.4	66.4	65.1	69.3	73.7	74.7			
5-year	83.8	27.7	10.5	9.8	8.6	9.2	14.0	17.4	35.6	48.3	56.5	58.5	55.6	54.4	62.8	60.1	64.8	68.2	71.7				
6-year	76.9	24.2	8.8	8.0	7.8	8.5	13.8	16.2	35.0	45.5	54.5	55.7	53.1	52.9	62.5	58.3	64.6	66.9					
7-year	76.8	23.2	7.6	7.6	7.5	7.9	13.2	15.1	33.2	44.8	50.0	53.1	52.6	52.9	61.7	58.3	60.8						
8-year	76.8	20.3	6.7	7.0	7.3	7.6	12.6	14.7	32.0	44.4	47.4	50.4	52.0	52.4	60.2	56.4							
9-year	76.8	18.9	6.0	6.9	6.8	7.6	12.5	13.9	30.9	42.6	44.3	49.8	52.0	50.8	58.1								
10-year	76.8	16.9	5.8	6.8	6.8	7.3	12.4	13.7	29.7	42.5	43.1	49.8	51.2	48.8									
11-year	74.7	15.6	5.6	6.7	6.7	7.3	11.8	13.3	28.6	42.0	41.7	49.1	50.0										
12-year	74.7	15.2	5.4	6.7	6.7	7.2	11.7	13.3	27.8	41.7	40.5	49.1											
13-year	74.7	14.6	5.0	6.7	6.5	7.2	11.7	13.3	27.2	41.6	39.4												
14-year	74.7	14.6	4.8	6.3	6.5	6.7	11.7	13.1	26.7	40.3													
15-year	74.7	14.6	4.7	5.7	6.5	6.6	11.6	12.9	26.2														
16-year	74.7	14.6	4.7	5.7	6.3	6.4	11.3	12.8															
17-year	74.7	14.3	4.7	5.6	6.3	6.2	11.1																
18-year	74.7	13.0	4.7	5.3	6.3	6.0																	
19-year	74.7	13.0	4.5	5.3	6.2																		
20-year	74.7	13.0	4.4	5.2																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 10.8  
Kaposi Sarcoma

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975-1979	1980-1984	1985-1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
<b>Survival Time</b>																							
1-year	95.2	63.7	58.6	59.3	60.3	61.6	57.4	53.7	60.9	68.7	72.7	73.0	76.6	73.8	79.5	76.4	75.6	81.4	82.8	80.1	82.6	80.7	81.0
2-year	89.1	41.6	32.1	31.6	28.9	27.3	28.9	26.4	45.4	59.7	64.2	64.0	64.4	65.6	73.2	69.6	72.5	72.2	79.1	77.0	80.3	76.6	
3-year	86.8	33.4	18.9	19.0	14.1	15.8	18.8	20.4	41.8	53.7	60.7	59.6	60.8	59.4	67.5	66.7	67.2	69.2	75.5	74.8	78.2		
4-year	83.7	29.2	12.9	13.6	9.9	10.5	14.8	17.9	37.5	51.1	57.0	57.0	55.5	56.6	64.9	66.2	65.0	67.3	74.6	74.2			
5-year	83.6	25.6	9.6	9.1	7.6	8.8	13.2	16.8	35.0	48.0	55.6	56.1	54.4	54.1	62.6	59.8	64.6	65.8	72.4				
6-year	74.2	21.8	7.9	7.3	6.9	8.2	12.9	15.6	34.3	45.3	53.4	54.0	51.5	53.0	62.6	58.0	64.2	64.1					
7-year	74.1	20.5	6.7	6.8	6.6	7.5	12.5	14.5	32.4	44.5	49.4	52.4	50.7	53.0	62.4	58.0	60.0						
8-year	74.1	18.1	5.9	6.3	6.3	7.3	11.9	14.1	31.2	44.0	46.9	50.7	50.6	52.3	61.1	57.0							
9-year	74.1	16.9	5.2	6.1	5.8	7.3	11.8	13.3	30.2	42.1	43.6	50.1	50.6	50.6	50.6	59.2							
10-year	74.1	15.1	5.1	6.1	5.7	7.0	11.6	13.2	29.1	42.0	42.2	50.1	50.1	49.1									
11-year	69.0	13.8	4.9	6.1	5.5	7.0	11.0	12.7	28.2	41.7	41.2	49.3	48.7										
12-year	68.9	13.6	4.8	6.1	5.4	6.9	10.9	12.7	27.2	41.2	40.0	49.3											
13-year	68.9	13.0	4.5	6.1	5.2	6.9	10.9	12.6	26.6	41.2	38.8												
14-year	68.9	13.0	4.3	5.6	5.2	6.4	10.9	12.3	26.2	40.1													
15-year	68.9	12.9	4.3	5.3	5.1	6.3	10.8	12.1	25.8														
16-year	68.9	12.7	4.2	5.3	4.9	6.1	10.6	12.1															
17-year	68.9	12.7	4.2	5.2	4.9	5.9	10.4																
18-year	68.9	12.3	4.1	4.9	4.8	5.7																	
19-year	68.9	12.3	4.0	4.8	4.6																		
20-year	68.9	12.3	3.9	4.8																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 10.9  
Kaposi Sarcoma

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
<b>Survival Time</b>																							
1-year	92.6	90.0	80.4	79.5	90.0	74.9	71.1	81.7	87.2	78.4	74.9	85.4	75.8	81.8	86.3	88.7	68.6	100.0	64.3	89.8	82.0	44.8	86.5
2-year	90.4	84.0	79.0	56.3	75.2	53.8	71.1	52.7	85.9	78.4	74.9	85.4	69.6	81.8	67.6	80.5	68.6	97.0	64.3	89.8	75.7	44.8	
3-year	90.4	73.4	74.3	56.3	75.2	44.3	52.8	49.8	74.5	64.5	74.9	84.9	65.9	76.9	52.2	71.2	61.6	97.0	57.1	80.0	75.7		
4-year	83.5	69.5	73.6	52.0	75.2	33.0	52.8	46.0	67.1	64.5	74.9	84.9	65.9	66.9	52.2	63.1	61.6	97.0	57.1	80.0			
5-year	81.4	61.0	73.6	49.9	73.1	33.0	52.8	46.0	50.5	55.2	69.1	84.9	65.9	57.2	52.2	58.3	61.6	97.0	57.1				
6-year	81.4	61.0	73.6	44.8	69.1	28.1	52.8	45.6	50.5	49.2	69.1	80.6	65.9	47.2	52.2	56.1	61.6	97.0					
7-year	81.4	61.0	73.6	44.8	69.1	28.1	47.4	40.6	50.5	49.2	58.0	54.3	65.9	47.2	50.9	56.1	61.6						
8-year	81.4	54.1	66.7	44.8	69.1	24.3	43.9	40.6	50.5	49.2	51.8	19.5	65.9	47.2	45.8	40.8							
9-year	81.4	51.0	66.1	44.8	69.1	24.3	43.9	38.9	47.3	49.2	51.8	19.5	65.9	47.2	38.6								
10-year	81.4	45.4	59.7	44.8	69.1	24.3	43.9	38.9	44.8	49.2	51.8	19.5	63.4	38.8									
11-year	81.4	42.7	57.6	40.6	69.1	24.3	43.9	38.9	37.6	49.2	46.5	19.5	63.4										
12-year	81.4	41.9	49.4	40.6	69.1	24.3	43.9	38.9	37.6	49.2	46.5	19.5											
13-year	81.4	41.0	44.1	40.6	69.1	24.3	43.9	38.9	37.6	37.9	46.5												
14-year	81.4	41.0	32.9	40.6	69.1	24.3	43.9	38.9	36.3	37.9													
15-year	81.4	41.0	27.9	19.2	69.1	24.3	43.9	38.9	19.4														
16-year	81.4	41.0	27.9	19.2	69.1	24.3	43.9	38.9															
17-year	81.4	29.3	27.9	19.2	69.1	24.3	43.9	38.9															
18-year	80.2	16.6	27.9	19.2	69.1	24.3	43.9																
19-year	80.2	16.6	27.9	19.2	69.1																		
20-year	80.2	16.6	27.9	0.0																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 10.10  
Kaposi Sarcoma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,  
 Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and  
 Lifetime Risk of Dying from Cancer Given Alive at Current Age  
 Both Sexes, 2008-2010 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer					Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer					Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever	from Cancer			+10 yrs	+20 yrs	+30 yrs	Ever	from Cancer	
All Races	0	0.00	0.00	0.00	0.05	0.00	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.02	0.00	
	10	0.00	0.00	0.01	0.05	0.00		10	0.00	0.00	0.00	0.02	0.00	
	20	0.00	0.01	0.02	0.05	0.00		20	0.00	0.00	0.01	0.02	0.00	
	30	0.01	0.02	0.02	0.04	0.00		30	0.00	0.01	0.01	0.02	0.00	
	40	0.01	0.02	0.02	0.04	0.00		40	0.00	0.01	0.01	0.01	0.00	
	50	0.01	0.01	0.02	0.03	0.00		50	0.00	0.00	0.01	0.01	0.00	
	60	0.01	0.01	0.02	0.02	0.00		60	0.00	0.00	0.01	0.01	0.00	
	70	0.01	0.02	-	0.02	0.00		70	0.00	0.01	-	0.01	0.00	
	80	0.01	-	-	0.02	0.00		80	0.01	-	-	0.01	0.00	
White	0	0.00	0.00	0.00	0.04	0.00	American Indian/ Alaska Native <sup>a</sup>	0	0.00	0.00	0.00	0.02	0.00	
	10	0.00	0.00	0.01	0.04	0.00		10	0.00	0.00	0.00	0.02	0.00	
	20	0.00	0.01	0.02	0.04	0.00		20	0.00	0.00	0.01	0.02	0.00	
	30	0.01	0.02	0.02	0.04	0.00		30	0.00	0.01	0.01	0.02	0.00	
	40	0.01	0.01	0.02	0.03	0.00		40	0.01	0.01	0.01	0.02	0.00	
	50	0.01	0.01	0.02	0.03	0.00		50	0.00	0.00	0.00	0.02	0.00	
	60	0.01	0.01	0.02	0.02	0.00		60	0.00	0.00	0.02	0.02	0.00	
	70	0.01	0.02	-	0.02	0.00		70	0.00	0.02	-	0.02	0.00	
	80	0.01	-	-	0.02	0.00		80	0.02	-	-	0.02	0.00	
Black	0	0.00	0.00	0.01	0.07	0.01	Hispanic <sup>b</sup>	0	0.00	0.00	0.00	0.08	0.01	
	10	0.00	0.01	0.03	0.07	0.01		10	0.00	0.00	0.01	0.08	0.01	
	20	0.01	0.03	0.05	0.07	0.01		20	0.00	0.01	0.02	0.08	0.01	
	30	0.02	0.04	0.05	0.06	0.01		30	0.01	0.02	0.03	0.08	0.01	
	40	0.02	0.03	0.03	0.04	0.00		40	0.01	0.02	0.03	0.07	0.00	
	50	0.01	0.01	0.02	0.02	0.00		50	0.01	0.02	0.03	0.06	0.00	
	60	0.00	0.01	0.01	0.01	0.00		60	0.01	0.03	0.04	0.05	0.00	
	70	0.01	0.01	-	0.01	0.00		70	0.02	0.04	-	0.05	0.00	
	80	0.01	-	-	0.01	0.00		80	0.02	-	-	0.04	0.00	

Devcan 6.7.0, June 2013, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

<sup>a</sup> Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

<sup>b</sup> Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Underlying mortality data for Hispanics exclude deaths from the District of Columbia, Minnesota, New Hampshire, North Dakota and South Carolina.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.

Table 10.11  
Kaposi Sarcoma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,  
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and  
Lifetime Risk of Dying from Cancer Given Alive at Current Age  
Males, 2008-2010 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer					Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer					Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever	from Cancer			+10 yrs	+20 yrs	+30 yrs	Ever	from Cancer	
All Races	0	0.00	0.00	0.01	0.08	0.01	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.03	0.00	
	10	0.00	0.01	0.02	0.08	0.01		10	0.00	0.00	0.01	0.03	0.00	
	20	0.01	0.02	0.04	0.08	0.01		20	0.00	0.01	0.01	0.03	0.00	
	30	0.02	0.04	0.05	0.08	0.00		30	0.01	0.01	0.02	0.03	0.00	
	40	0.02	0.03	0.04	0.06	0.00		40	0.01	0.01	0.01	0.03	0.00	
	50	0.01	0.02	0.03	0.04	0.00		50	0.00	0.01	0.01	0.02	0.00	
	60	0.01	0.02	0.03	0.03	0.00		60	0.00	0.01	0.02	0.02	0.00	
	70	0.01	0.02	-	0.03	0.00		70	0.00	0.01	-	0.02	0.00	
	80	0.02	-	-	0.02	0.00		80	0.01	-	-	0.01	0.00	
White	0	0.00	0.00	0.00	0.07	0.00	American Indian/ Alaska Native <sup>a</sup>	0	0.00	0.00	0.00	0.04	0.00	
	10	0.00	0.00	0.02	0.07	0.00		10	0.00	0.00	0.00	0.04	0.00	
	20	0.00	0.02	0.03	0.07	0.00		20	0.00	0.00	0.01	0.04	0.00	
	30	0.01	0.03	0.04	0.07	0.00		30	0.00	0.01	0.01	0.04	0.00	
	40	0.02	0.03	0.04	0.06	0.00		40	0.01	0.01	0.01	0.04	0.00	
	50	0.01	0.02	0.03	0.04	0.00		50	0.00	0.00	0.01	0.04	0.00	
	60	0.01	0.02	0.03	0.03	0.00		60	0.00	0.01	0.02	0.04	0.00	
	70	0.01	0.03	-	0.03	0.00		70	0.01	0.05	-	0.05	0.00	
	80	0.02	-	-	0.03	0.00		80	0.06	-	-	0.06	0.00	
Black	0	0.00	0.00	0.03	0.13	0.01	Hispanic <sup>b</sup>	0	0.00	0.00	0.01	0.12	0.01	
	10	0.00	0.03	0.07	0.13	0.01		10	0.00	0.01	0.02	0.12	0.01	
	20	0.03	0.07	0.10	0.13	0.01		20	0.01	0.02	0.04	0.12	0.01	
	30	0.04	0.08	0.09	0.11	0.01		30	0.02	0.04	0.05	0.12	0.01	
	40	0.04	0.05	0.06	0.07	0.01		40	0.02	0.04	0.05	0.10	0.01	
	50	0.01	0.02	0.03	0.03	0.01		50	0.01	0.03	0.06	0.08	0.01	
	60	0.01	0.02	0.02	0.02	0.00		60	0.02	0.04	0.07	0.07	0.01	
	70	0.01	0.01	-	0.01	0.00		70	0.03	0.06	-	0.06	0.01	
	80	0.01	-	-	0.01	0.00		80	0.04	-	-	0.04	0.01	

Devcan 6.7.0, June 2013, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

<sup>a</sup> Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

<sup>b</sup> Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Underlying mortality data for Hispanics exclude deaths from the District of Columbia, Minnesota, New Hampshire, North Dakota and South Carolina.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.

Table 10.12  
Kaposi Sarcoma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,  
 Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and  
 Lifetime Risk of Dying from Cancer Given Alive at Current Age  
 Females, 2008-2010 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer					Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer					Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever	from Cancer			+10 yrs	+20 yrs	+30 yrs	Ever	from Cancer	
All Races	0	0.00	0.00	0.00	0.01	0.00	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.00	0.00	0.00
	10	0.00	0.00	0.00	0.01	0.00		10	0.00	0.00	0.00	0.00	0.00	0.00
	20	0.00	0.00	0.00	0.01	0.00		20	0.00	0.00	0.00	0.00	0.00	0.00
	30	0.00	0.00	0.00	0.01	0.00		30	0.00	0.00	0.00	0.00	0.00	0.00
	40	0.00	0.00	0.00	0.01	0.00		40	0.00	0.00	0.00	0.00	0.00	0.00
	50	0.00	0.00	0.00	0.01	0.00		50	0.00	0.00	0.00	0.00	0.00	0.00
	60	0.00	0.00	0.01	0.01	0.00		60	0.00	0.00	0.00	0.00	0.00	0.00
	70	0.00	0.01	-	0.01	0.00		70	0.00	0.00	0.00	-	0.00	0.00
	80	0.01	-	-	0.01	0.00		80	0.00	-	-	0.00	0.00	0.00
White	0	0.00	0.00	0.00	0.01	0.00	American Indian/ Alaska Native <sup>a</sup>	0	0.00	0.00	0.00	0.01	0.01	0.00
	10	0.00	0.00	0.00	0.01	0.00		10	0.00	0.00	0.00	0.01	0.01	0.00
	20	0.00	0.00	0.00	0.01	0.00		20	0.00	0.00	0.01	0.01	0.01	0.00
	30	0.00	0.00	0.00	0.01	0.00		30	0.00	0.01	0.01	0.01	0.01	0.00
	40	0.00	0.00	0.00	0.01	0.00		40	0.01	0.01	0.01	0.01	0.01	0.00
	50	0.00	0.00	0.01	0.01	0.00		50	0.00	0.00	0.00	0.00	0.00	0.00
	60	0.00	0.00	0.01	0.01	0.00		60	0.00	0.00	0.00	0.00	0.00	0.00
	70	0.00	0.01	-	0.01	0.00		70	0.00	0.00	0.00	-	0.00	0.00
	80	0.01	-	-	0.01	0.00		80	0.00	-	-	0.00	0.00	0.00
Black	0	0.00	0.00	0.00	0.01	0.00	Hispanic <sup>b</sup>	0	0.00	0.00	0.00	0.04	0.04	0.00
	10	0.00	0.00	0.00	0.01	0.00		10	0.00	0.00	0.00	0.04	0.04	0.00
	20	0.00	0.00	0.01	0.01	0.00		20	0.00	0.00	0.00	0.04	0.04	0.00
	30	0.00	0.01	0.01	0.01	0.00		30	0.00	0.00	0.00	0.04	0.04	0.00
	40	0.00	0.00	0.00	0.01	0.00		40	0.00	0.00	0.00	0.04	0.04	0.00
	50	0.00	0.00	0.00	0.01	0.00		50	0.00	0.00	0.01	0.04	0.04	0.00
	60	0.00	0.00	0.01	0.01	0.00		60	0.00	0.01	0.03	0.04	0.04	0.00
	70	0.00	0.01	-	0.01	0.00		70	0.01	0.03	-	0.04	0.04	0.00
	80	0.00	-	-	0.01	0.00		80	0.02	-	-	0.03	0.03	0.00

Devcan 6.7.0, June 2013, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

<sup>a</sup> Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

<sup>b</sup> Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Underlying mortality data for Hispanics exclude deaths from the District of Columbia, Minnesota, New Hampshire, North Dakota and South Carolina.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.

Table 10.13  
Kaposi Sarcoma

SEER Incidence  
Age-Adjusted Rates and Trends<sup>a</sup>  
By Race/Ethnicity and Sex

RACE/ETHNICITY	SEER Incidence			SEER 18 Areas <sup>b</sup>			SEER 13 Areas <sup>b</sup>			SEER 9 Areas <sup>b</sup>				
				Rate 2006-2010		Trend 2001-2010 <sup>c</sup>	Rate per 100,000 persons		AAPC (%)		Trend 2001-2010 <sup>d</sup>		AAPC (%)	
	Total	Males	Females	Total	Males	Females	Total	Males	Females	Total	Males	Females		
All Races	0.6	1.0	0.1		-2.5*	-2.4*	-5.3*		-1.8*	-1.8*	-4.9*			
White	0.5	0.9	0.1		-2.5*	-2.6*	-		-2.1*	-2.1*	-4.9*			
White Hispanic <sup>e</sup>	0.8	1.4	0.2		-2.3	-2.7	-		-	-	-			
White Non-Hispanic <sup>e</sup>	0.4	0.8	0.1		-2.6*	-2.7*	-		-	-	-			
Black	1.0	2.0	0.2		-2.7*	-2.5*	-		-2.1	-2.0	-			
Asian/Pacific Islander	0.2	0.4	-		-	-	-		-	-	-			
Amer Ind/Alaska Nat <sup>f</sup>	-	-	-		-	-	-		-	-	-			
Hispanic <sup>e</sup>	0.8	1.4	0.2		-2.5	-3.0	-		-	-	-			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.0.3, April 2013, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

<sup>a</sup> Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

<sup>c</sup> The 2001-2010 AAPC estimates are based on a Joinpoint analysis with up to 3 Joinpoints over diagnosis years 1992-2010.

<sup>d</sup> The 2001-2010 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2010.

<sup>e</sup> Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

<sup>f</sup> Incidence data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

\* The APC is significantly different from zero ( $p < .05$ ).

Table 10.14  
Kaposi Sarcoma

Age-Adjusted SEER Incidence Rates<sup>a</sup>  
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<b>SEER Incidence Rates<sup>a</sup>, 2006-2010</b>									
California	0.67	1.26	0.10	0.68	1.28	0.09	1.19	2.26	-
Greater Bay Area	0.89	1.68	0.13	1.08	2.00	0.15	1.58	3.03	-
San Francisco-Oakland	1.12	2.14	0.14	1.38	2.58	-	1.73	3.39	-
San Jose-Monterey	0.48	0.86	-	0.58	1.03	-	-	-	-
Los Angeles	0.96	1.89	0.09	1.01	1.93	0.11	1.35	2.90	-
Greater California	0.44	0.80	0.09	0.41	0.78	0.06	0.87	1.43	-
Connecticut	0.35	0.63	-	0.30	0.58	-	-	-	-
Detroit	0.51	0.96	-	0.30	0.52	-	1.06	2.28	-
Georgia	0.73	1.44	0.07	0.43	0.83	-	1.30	2.73	-
Atlanta	1.53	3.04	-	1.00	1.90	-	2.28	4.90	-
Rural Georgia	-	-	-	-	-	-	-	-	-
Greater Georgia	0.26	0.50	-	0.19	0.37	-	0.41	0.81	-
Hawaii	0.36	0.73	-	-	-	-	-	-	-
Iowa	-	-	-	-	-	-	-	-	-
Kentucky	0.20	0.36	-	0.19	0.34	-	-	-	-
Louisiana	0.54	0.95	0.15	0.45	0.81	-	0.72	1.26	-
New Jersey	0.46	0.81	0.17	0.39	0.69	0.12	0.79	1.35	-
New Mexico	0.56	1.02	-	0.49	0.90	-	-	-	-
Seattle-Puget Sound	0.52	1.01	-	0.51	0.98	-	-	-	-
Utah	0.17	0.33	-	0.16	0.32	-	-	-	-
9 SEER Areas <sup>b</sup>	0.65	1.24	0.08	0.52	0.99	0.07	1.57	3.18	-
11 SEER Areas <sup>b</sup>	0.71	1.37	0.09	0.64	1.21	0.08	1.52	3.11	0.15
13 SEER Areas <sup>b</sup>	0.70	1.36	0.09	0.64	1.21	0.08	1.50	3.08	0.15
18 SEER Areas <sup>b</sup>	0.55	1.04	0.09	0.49	0.92	0.08	1.05	2.02	0.19

<sup>a</sup> Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130)  
<sup>b</sup> The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.  
The SEER 11 areas comprise the SEER 9 areas plus San Jose-Monterey and Los Angeles.  
The SEER 13 areas comprise the SEER 11 areas plus the Alaska Native Registry and Rural Georgia.  
The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 10.15  
Kaposi Sarcoma

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2010  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 <sup>e</sup>	0 to <35 <sup>e</sup>	>=35 <sup>g</sup>	Complete <sup>h</sup>
Race	Sex										
All Races <sup>b</sup>	Both Sexes	7,319	5,335	6,995	3,499	1,165	334	23,149	24,815	+	+
	Males	6,865	5,044	6,752	3,293	1,143	311	21,954	23,533	+	+
	Females	454	291	243	206	22	23	1,195	1,282	+	+
White <sup>b</sup>	Both Sexes	5,002	3,946	5,497	2,916	963	258	17,363	18,729	+	+
	Males	4,685	3,683	5,346	2,755	941	247	16,470	17,761	+	+
	Females	317	263	151	161	22	11	893	968	+	+
Black <sup>b</sup>	Both Sexes	1,915	1,174	1,068	374	93	40	4,531	4,676	+	+
	Males	1,814	1,151	1,022	352	93	40	4,339	4,484	+	+
	Females	101	23	46	22	0	0	192	192	+	+
Asian/ Pacific	Both Sexes	101	83	99	+	+	+	302	+	+	+
	Males	101	77	99	+	+	+	296	+	+	+
Islander <sup>c</sup>	Females	0	6	0	+	+	+	6	+	+	+
Hispanic <sup>d</sup>	Both Sexes	1,279	974	1,300	+	+	+	3,553	+	+	+
	Males	1,166	929	1,259	+	+	+	3,354	+	+	+
	Females	113	45	41	+	+	+	199	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)								Age-Adjusted <sup>f</sup> All Ages		
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79			
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.0071%	-	-	0.0010%	0.0062%	0.0191%	0.0138%	0.0081%	0.0080%	0.0127%	0.0070%
	Males	0.0138%	-	-	0.0019%	0.0122%	0.0377%	0.0281%	0.0164%	0.0152%	0.0249%	0.0139%
	Females	0.0006%	-	-	0.0002%	0.0007%	0.0003%	0.0007%	0.0022%	0.0057%	0.0006%	
White <sup>c</sup>	Both Sexes	0.0072%	-	-	0.0007%	0.0055%	0.0187%	0.0144%	0.0085%	0.0086%	0.0133%	0.0070%
	Males	0.0138%	-	-	0.0014%	0.0105%	0.0361%	0.0288%	0.0169%	0.0163%	0.0260%	0.0136%
	Females	0.0006%	-	-	-	0.0007%	-	0.0007%	0.0024%	0.0060%	0.0006%	
Black <sup>c</sup>	Both Sexes	0.0103%	-	-	0.0032%	0.0148%	0.0322%	0.0173%	0.0102%	0.0042%	0.0098%	0.0109%
	Males	0.0209%	-	-	0.0065%	0.0313%	0.0674%	0.0365%	0.0211%	0.0088%	-	0.0227%
	Females	0.0007%	-	-	-	-	-	-	-	0.0082%	0.0008%	
Asian/ Pacific	Both Sexes	0.0018%	-	-	0.0016%	0.0043%	0.0035%	0.0015%	0.0035%	0.0045%	0.0018%	
	Males	0.0037%	-	-	0.0034%	0.0089%	0.0076%	0.0034%	0.0080%	0.0097%	0.0038%	
Islander <sup>c</sup>	Females	-	-	-	-	-	-	-	-	-	-	
Hispanic <sup>d</sup>	Both Sexes	0.0073%	-	-	0.0012%	0.0073%	0.0229%	0.0178%	0.0110%	0.0168%	0.0327%	0.0098%
	Males	0.0135%	-	-	0.0022%	0.0141%	0.0438%	0.0361%	0.0230%	0.0288%	0.0570%	0.0187%
	Females	0.0008%	-	-	-	0.0013%	-	-	0.0080%	0.0183%	0.0015%	

<sup>a</sup> US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census.

<sup>b c d</sup> Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.


<sup>g h i</sup> (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et al. 1997, Merrill et al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Figure 10.1

# All Cancer Sites vs Kaposi Sarcoma SEER Incidence Rates, 1975-2010 Males, Ages 20-54




Source: SEER 9 areas.

Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines are calculated using the Joinpoint Regression Program Version 4.0.3, April 2013,  
National Cancer Institute.

Figure 10.2


All SEER excluding  
San Francisco/Oakland  
Males, Ages 20-54  
1975-2010



San Francisco/Oakland  
Males, Ages 20-54  
1975-2010



San Francisco County/City  
Males, Ages 20-54  
1975-2010



Source: SEER 9 areas. Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

<sup>a</sup> Kaposi Sarcoma cases are excluded from Colon & Rectum, Lung & Bronchus, and Non-Hodgkin Lymphoma cases.