

Table 1.22
U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2011^a
Using Different Tumor Inclusion Criteria^b

Site	Sex	5-Year Limited Duration			36-year Limited Duration	
		1st Invasive Tumor Ever ^c	1st Per Site in Previous 36 Years ^d	1st Per Site in Previous 5 Years ^e	1st Invasive Tumor Ever ^c	1st Per Site in Previous 36 Years ^d
All Sites	Both Sexes	4,594,732	4,684,591	5,152,908	12,998,655	13,213,910
	Male	2,373,342	2,409,199	2,638,979	6,188,275	6,261,129
	Female	2,221,390	2,275,392	2,513,929	6,810,380	6,952,781
Oral Cavity & Pharynx	Both Sexes	104,979	121,403	125,889	272,072	300,609
	Male	73,503	84,311	87,116	181,087	198,337
	Female	31,476	37,092	38,773	90,985	102,272
Esophagus	Both Sexes	21,165	26,055	26,172	34,428	40,593
	Male	16,662	20,345	20,416	26,846	31,461
	Female	4,503	5,710	5,756	7,582	9,132
Stomach	Both Sexes	36,380	43,398	43,779	72,734	83,348
	Male	21,307	26,023	26,197	41,394	47,992
	Female	15,073	17,375	17,582	31,340	35,356
Colon & Rectum	Both Sexes	403,434	465,899	475,547	1,140,625	1,270,460
	Male	206,233	237,736	242,463	567,872	629,107
	Female	197,201	228,163	233,084	572,753	641,353
Liver & Intrahepatic Bile Duct	Both Sexes	33,682	38,427	38,461	45,468	50,961
	Male	24,703	27,797	27,819	32,312	35,854
	Female	8,979	10,630	10,642	13,156	15,107
Pancreas	Both Sexes	32,756	39,642	39,654	43,238	51,264
	Male	16,110	19,950	19,962	20,679	25,032
	Female	16,646	19,692	19,692	22,559	26,232
Larynx	Both Sexes	31,318	37,292	37,636	87,652	98,482
	Male	25,568	30,391	30,700	70,687	79,142
	Female	5,750	6,901	6,936	16,965	19,340
Lung & Bronchus	Both Sexes	231,126	297,286	306,343	395,186	487,438
	Male	107,276	140,106	143,631	179,129	222,121
	Female	123,850	157,180	162,712	216,057	265,317
Melanoma of the Skin	Both Sexes	282,028	320,618	335,149	924,397	998,829
	Male	151,313	176,523	185,633	459,686	503,903
	Female	130,715	144,095	149,516	464,711	494,926
Breast	Female	855,411	924,344	982,776	2,847,146	3,016,451
Cervix	Female	39,578	41,703	41,825	211,309	217,336
Corpus & Uterus, NOS	Female	177,109	198,883	199,045	584,180	637,931
Ovary ^f	Female	58,762	67,726	67,789	173,192	194,691

^a U.S. 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER 9 registries and 1/1/2011 U.S. population estimates based on the average of 2010 and 2011 population estimates from the U.S. Bureau of the Census.

^b Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

^c (c) First invasive tumor ever

^d (d) First invasive tumor for each cancer site diagnosed during the previous 36 years (1975-2010)

^e (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2006-2010)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2006; Melanoma in 2007.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 36-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2011.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 36-year limited duration prevalence. The 2006 breast cancer is counted for the breast 5-year and 36-year limited duration prevalence.

In method (e) the 2006 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2007 melanoma is counted for 5-year limited duration prevalence for melanoma.

^f Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table 1.22 - continued
 U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2011^a
 Using Different Tumor Inclusion Criteria^b

Site	Sex	5-Year Limited Duration			36-year Limited Duration	
		1st Invasive Tumor Ever ^c	1st Per Site in Previous 36 Years ^d	1st Per Site in Previous 5 Years ^e	1st Invasive Tumor Ever ^c	1st Per Site in Previous 36 Years ^d
Prostate	Male	1,047,674	1,131,608	1,131,634	2,706,583	2,883,365
Testis	Male	42,114	42,942	43,546	212,079	214,995
Urinary Bladder	Both Sexes	200,849	251,314	256,779	560,445	653,826
	Male	152,753	192,335	196,676	418,562	489,001
	Female	48,096	58,979	60,103	141,883	164,825
Kidney & Renal Pelvis	Both Sexes	149,969	182,913	185,010	347,916	406,298
	Male	91,313	113,044	114,471	206,735	243,752
	Female	58,656	69,869	70,539	141,181	162,546
Brain & Nervous System	Both Sexes	44,217	47,403	47,827	131,408	135,930
	Male	24,099	25,776	26,028	69,809	72,101
	Female	20,118	21,627	21,799	61,599	63,829
Thyroid	Both Sexes	174,786	194,868	195,449	527,919	564,773
	Male	38,665	45,067	45,197	114,267	125,269
	Female	136,121	149,801	150,252	413,652	439,504
Hodgkin Lymphoma	Both Sexes	38,529	41,043	41,065	172,689	177,526
	Male	20,912	22,396	22,407	89,550	92,081
	Female	17,617	18,647	18,658	83,139	85,445
Non-Hodgkin Lymphoma	Both Sexes	203,992	239,804	242,418	522,113	584,133
	Male	109,031	129,460	130,668	274,206	307,672
	Female	94,961	110,344	111,750	247,907	276,461
Myeloma	Both Sexes	52,505	62,629	62,743	83,118	95,874
	Male	28,573	34,693	34,761	45,299	52,835
	Female	23,932	27,936	27,982	37,819	43,039
Leukemia	Both Sexes	115,823	134,637	135,040	297,129	327,520
	Male	66,545	78,337	78,566	168,443	186,907
	Female	49,278	56,300	56,474	128,686	140,613
Acute Lymphocytic Leukemia	Both Sexes	16,260	16,720	16,720	68,728	69,433
	Male	9,051	9,286	9,286	37,729	38,032
	Female	7,209	7,434	7,434	30,999	31,401
Childhood (Ages 0-19)	Both Sexes	63,918	63,993	64,508	314,920	315,399
	Male	33,711	33,746	33,991	162,108	162,335
	Female	30,207	30,247	30,517	152,812	153,064
Kaposi Sarcoma	Both Sexes	7,194	7,680	7,680	25,504	26,815
	Male	6,655	7,078	7,078	24,134	25,255
	Female	539	602	602	1,370	1,560
Mesothelioma	Both Sexes	3,004	3,922	3,922	4,663	5,734
	Male	2,172	2,854	2,854	2,829	3,595
	Female	832	1,068	1,068	1,834	2,139

^a U.S. 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER 9 registries and 1/1/2011 U.S. population estimates based on the average of 2010 and 2011 population estimates from the U.S. Bureau of the Census.

^b Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

^c (c) First invasive tumor ever

^d (d) First invasive tumor for each cancer site diagnosed during the previous 36 years (1975-2010)

^e (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2006-2010)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2006; Melanoma in 2007.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 36-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2011.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 36-year limited duration prevalence. The 2006 breast cancer is counted for the breast 5-year and 36-year limited duration prevalence.

In method (e) the 2006 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2007 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table 1.23
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2011^a
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	6,271,036	19,001	41,846	85,407	155,919	356,211	883,826	1,658,545	3,070,283
Females	7,126,123	16,447	36,376	88,223	226,162	638,879	1,316,044	1,758,236	3,045,756
Oral Cavity & Pharynx									
Males	186,491	62	423	1,548	3,544	14,256	45,288	58,097	63,273
Females	95,100	90	522	1,483	3,689	8,554	18,160	23,692	38,911
Esophagus									
Males	26,909	0	0	22	144	1,109	4,697	9,747	11,190
Females	7,642	0	0	11	11	240	1,378	1,832	4,170
Stomach									
Males	42,127	0	29	62	603	2,502	7,119	11,218	20,593
Females	31,908	4	23	154	618	2,252	4,650	6,664	17,543
Colon & Rectum									
Males	575,457	11	69	1,276	5,758	25,233	82,732	139,236	321,141
Females	586,969	0	67	1,472	5,551	23,983	70,097	112,784	373,015
Liver & Intrahep									
Males	32,389	556	508	616	547	1,590	10,492	11,364	6,716
Females	13,553	396	521	359	403	889	3,016	3,463	4,506
Pancreas									
Males	20,801	23	12	130	276	1,653	4,466	6,626	7,615
Females	22,737	0	62	212	431	1,446	3,801	6,179	10,604
Larynx									
Males	71,999	0	0	90	219	1,822	10,381	21,280	38,207
Females	17,266	0	0	36	135	841	3,336	4,868	8,051
Lung & Bronchus									
Males	183,215	45	110	405	1,195	5,366	24,255	54,762	97,077
Females	219,111	23	71	367	1,387	7,548	29,898	59,021	120,797
Melanoma of the Skin									
Males	471,220	56	594	5,039	17,738	46,603	97,543	130,762	172,885
Females	489,011	68	878	10,864	34,091	73,459	115,119	113,905	140,628

^a U.S. 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2011 U.S. population estimates based on the average of 2010 and 2011 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.23 - continued
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2011^a
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	14,871	0	0	22	112	591	1,838	4,277	8,030
Females	2,899,726	0	47	2,669	35,679	215,380	550,120	795,921	1,299,910
Cervix									
Females	249,632	0	57	2,142	15,934	41,969	59,700	57,440	72,389
Corpus & Uterus, NOS									
Females	610,804	12	12	581	5,712	25,316	89,564	165,807	323,801
Ovary ^d									
Females	188,867	45	1,063	3,475	6,807	18,458	41,719	49,281	68,021
Prostate									
Males	2,707,821	34	47	113	332	20,758	235,407	776,168	1,674,962
Urinary Bladder									
Males	425,722	57	88	567	2,476	10,749	43,060	102,624	266,102
Females	145,796	56	46	219	1,015	4,094	13,710	30,955	95,700
Kidney & Renal Pelvis									
Males	212,703	1,458	2,458	2,630	5,254	18,386	41,683	61,520	79,315
Females	145,900	1,374	2,649	2,922	4,870	12,287	25,458	35,857	60,483
Hodgkin Lymphoma									
Males	96,100	186	2,493	9,242	16,547	23,764	22,436	14,185	7,247
Females	89,693	67	2,083	9,407	16,399	23,306	19,845	11,219	7,368
Non-Hodgkin Lymphoma									
Males	278,836	842	3,598	7,821	13,627	29,722	54,887	71,106	97,233
Females	252,083	519	1,758	4,663	8,663	21,422	42,382	60,220	112,455
Myeloma									
Males	45,388	0	6	99	490	3,274	8,260	14,917	18,342
Females	37,979	0	0	28	312	2,325	6,791	11,124	17,399
Leukemia									
Males	170,969	6,686	13,047	12,524	11,179	14,206	23,300	35,001	55,027
Females	131,831	5,729	10,353	11,342	9,836	10,094	15,793	22,736	45,948
Acute Lymphocytic Leuk									
Males	39,514	5,599	11,159	9,649	6,330	3,974	1,505	836	463
Females	32,723	4,763	8,785	8,138	5,414	3,188	1,137	816	483

^a U.S. 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2011 U.S. population estimates based on the average of 2010 and 2011 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 2.21
All Cancer Sites (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	4,594,732	3,153,349	2,206,528	1,398,664	813,745	479,491	11,551,569	12,998,655	398,504	13,397,159
	Males	2,373,342	1,598,257	1,052,027	624,794	286,933	153,864	5,723,802	6,188,275	82,761	6,271,036
	Females	2,221,390	1,555,092	1,154,501	773,870	526,812	325,627	5,827,767	6,810,380	315,743	7,126,123
White ^b	Both Sexes	3,873,922	2,735,983	1,936,873	1,246,445	737,398	436,723	9,972,290	11,288,173	349,111	11,637,284
	Males	1,992,129	1,379,781	918,148	557,144	262,877	141,937	4,916,050	5,343,834	77,440	5,421,274
	Females	1,881,793	1,356,202	1,018,725	689,301	474,521	294,786	5,056,240	5,944,339	271,671	6,216,010
Black ^b	Both Sexes	470,162	283,128	181,576	101,652	48,754	27,655	1,049,074	1,132,495	23,724	1,156,219
	Males	254,286	156,022	95,636	47,986	15,328	7,370	558,243	580,643	2,081	582,724
	Females	215,876	127,106	85,940	53,666	33,426	20,285	490,831	551,852	21,643	573,495
Asian/ Pacific Islander ^c	Both Sexes	138,281	85,330	53,166	+	+	+	302,774	+	+	+
	Males	59,359	34,925	20,074	+	+	+	123,408	+	+	+
	Females	78,922	50,405	33,092	+	+	+	179,366	+	+	+
Hispanic ^d	Both Sexes	310,615	194,217	121,869	+	+	+	688,386	+	+	+
	Males	148,698	91,411	54,197	+	+	+	319,300	+	+	+
	Females	161,917	102,806	67,672	+	+	+	369,086	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	3.2217%	0.0862%	0.1837%	0.3198%	0.7689%	1.8721%	4.2666%	9.2546%	15.5822%	17.3733%	3.1412%
	Males	3.2166%	0.0913%	0.1931%	0.3099%	0.6094%	1.2979%	3.4940%	9.9466%	20.1479%	24.9665%	3.5286%
	Females	3.2266%	0.0809%	0.1738%	0.3301%	0.9289%	2.4413%	5.0049%	8.6287%	11.9106%	12.9282%	2.9139%
White ^c	Both Sexes	3.5585%	0.0939%	0.2039%	0.3630%	0.8634%	2.0102%	4.4752%	9.7013%	16.4512%	17.9921%	3.3133%
	Males	3.5192%	0.0986%	0.2161%	0.3513%	0.6930%	1.4141%	3.6333%	10.2198%	20.8721%	25.3324%	3.6571%
	Females	3.5976%	0.0889%	0.1911%	0.3755%	1.0425%	2.6247%	5.3066%	9.2172%	12.8078%	13.6557%	3.1218%
Black ^c	Both Sexes	2.3888%	0.0629%	0.1262%	0.2021%	0.5298%	1.5203%	4.0049%	9.5579%	15.4970%	15.4242%	2.9575%
	Males	2.6684%	0.0685%	0.1238%	0.2028%	0.3818%	1.0725%	3.9383%	12.6281%	23.6878%	26.9148%	3.9709%
	Females	2.1343%	0.0572%	0.1286%	0.2014%	0.6589%	1.9168%	4.0627%	7.1323%	9.8116%	9.9271%	2.3336%
Asian/ Pacific Islander ^c	Both Sexes	2.0299%	0.0675%	0.1285%	0.1901%	0.4691%	1.3207%	3.0094%	5.6545%	9.4723%	12.3994%	2.0502%
	Males	1.7744%	0.0727%	0.1351%	0.1768%	0.3183%	0.6954%	1.8658%	5.0668%	11.3639%	17.6094%	2.0567%
	Females	2.2638%	0.0621%	0.1215%	0.2031%	0.6043%	1.8758%	3.9907%	6.1417%	8.0194%	9.1981%	2.1021%
Hispanic ^d	Both Sexes	1.3751%	0.0808%	0.1742%	0.2457%	0.5507%	1.3078%	2.9931%	6.4163%	11.1656%	13.2202%	2.2708%
	Males	1.2494%	0.0868%	0.1837%	0.2461%	0.4265%	0.8128%	2.1598%	6.5842%	15.0580%	19.8157%	2.5617%
	Females	1.5037%	0.0745%	0.1641%	0.2452%	0.6830%	1.8232%	3.7866%	6.2732%	8.2937%	9.2981%	2.1372%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 3.21
Brain and Other Nervous System (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	44,217	26,484	19,591	16,190	11,780	6,972	109,276	131,408	13,055	144,463
	Males	24,099	13,786	9,994	8,930	6,186	3,704	58,368	69,809	5,939	75,748
	Females	20,118	12,698	9,597	7,260	5,594	3,268	50,908	61,599	7,116	68,715
White ^b	Both Sexes	38,106	23,266	17,243	14,134	10,379	6,194	95,125	114,930	4,635	119,565
	Males	21,097	12,280	8,856	7,751	5,338	3,327	51,289	61,510	2,622 ⁱ	64,132
	Females	17,009	10,986	8,387	6,383	5,041	2,867	43,836	53,420	2,013 ⁱ	55,433
Black ^b	Both Sexes	3,667	1,963	1,492	1,230	984	581	8,655	10,362	+	+
	Males	1,684	903	743	706	600	288	4,232	5,106	+	+
	Females	1,983	1,060	749	524	384	293	4,423	5,256	+	+
Asian/ Pacific Islander ^c	Both Sexes	1,362	787	552	+	+	+	3,047	+	+	+
	Males	733	370	252	+	+	+	1,541	+	+	+
	Females	629	417	300	+	+	+	1,506	+	+	+
Hispanic ^d	Both Sexes	4,602	2,969	2,209	+	+	+	10,885	+	+	+
	Males	2,550	1,592	1,091	+	+	+	5,746	+	+	+
	Females	2,052	1,377	1,118	+	+	+	5,139	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0306%	0.0147%	0.0320%	0.0311%	0.0306%	0.0363%	0.0376%	0.0373%	0.0289%	0.0181%	0.0302%
	Males	0.0333%	0.0153%	0.0335%	0.0330%	0.0339%	0.0399%	0.0418%	0.0417%	0.0307%	0.0196%	0.0328%
	Females	0.0280%	0.0142%	0.0305%	0.0292%	0.0272%	0.0327%	0.0335%	0.0333%	0.0275%	0.0172%	0.0278%
White ^c	Both Sexes	0.0348%	0.0163%	0.0361%	0.0354%	0.0350%	0.0416%	0.0434%	0.0422%	0.0318%	0.0186%	0.0343%
	Males	0.0379%	0.0166%	0.0384%	0.0374%	0.0379%	0.0462%	0.0479%	0.0471%	0.0348%	0.0207%	0.0372%
	Females	0.0317%	0.0160%	0.0335%	0.0331%	0.0320%	0.0369%	0.0389%	0.0376%	0.0293%	0.0173%	0.0314%
Black ^c	Both Sexes	0.0192%	0.0104%	0.0225%	0.0222%	0.0191%	0.0215%	0.0187%	0.0229%	0.0145%	0.0183%	0.0190%
	Males	0.0193%	0.0118%	0.0216%	0.0221%	0.0209%	0.0209%	0.0206%	0.0219%	0.0090%	-	0.0189%
	Females	0.0191%	0.0088%	0.0234%	0.0223%	0.0175%	0.0221%	0.0171%	0.0236%	0.0183%	0.0210%	0.0190%
Asian/ Pacific Islander ^c	Both Sexes	0.0176%	0.0102%	0.0185%	0.0177%	0.0182%	0.0196%	0.0201%	0.0180%	0.0213%	0.0118%	0.0175%
	Males	0.0185%	0.0109%	0.0175%	0.0193%	0.0227%	0.0190%	0.0218%	0.0192%	0.0183%	0.0140%	0.0183%
	Females	0.0167%	0.0096%	0.0196%	0.0162%	0.0142%	0.0202%	0.0186%	0.0171%	0.0235%	0.0104%	0.0167%
Hispanic ^d	Both Sexes	0.0214%	0.0105%	0.0242%	0.0182%	0.0193%	0.0281%	0.0317%	0.0313%	0.0240%	0.0247%	0.0228%
	Males	0.0223%	0.0100%	0.0258%	0.0180%	0.0196%	0.0316%	0.0332%	0.0378%	0.0226%	0.0216%	0.0239%
	Females	0.0205%	0.0112%	0.0226%	0.0185%	0.0191%	0.0245%	0.0302%	0.0256%	0.0250%	0.0266%	0.0216%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 4.25
Cancer of the Breast (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	861,388	681,999	539,624	352,387	226,061	119,480	2,489,129	2,861,883	52,714	2,914,597
	Males	5,977	3,716	2,395	1,156	775	443	13,456	14,737	134	14,871
	Females	855,411	678,283	537,229	351,231	225,286	119,037	2,475,673	2,847,146	52,580	2,899,726
White ^b	Both Sexes	726,636	595,024	477,668	314,850	205,071	108,022	2,162,558	2,501,069	46,544	2,547,613
	Males	4,987	3,184	2,123	1,006	666	399	11,475	12,626	4 ⁱ	12,630
	Females	721,649	591,840	475,545	313,844	204,405	107,623	2,151,083	2,488,443	46,540	2,534,983
Black ^b	Both Sexes	89,371	57,364	40,770	25,643	14,540	8,189	216,841	240,800	+	+
	Males	795	429	194	86	109	36	1,541	1,662	+	+
	Females	88,576	56,935	40,576	25,557	14,431	8,153	215,300	239,138	3,993	243,131
Asian/ Pacific Islander ^c	Both Sexes	32,843	23,025	15,781	+	+	+	79,088	+	+	+
	Males	151	76	38	+	+	+	301	+	+	+
	Females	32,692	22,949	15,743	+	+	+	78,787	+	+	+
Hispanic ^d	Both Sexes	58,348	40,516	27,038	+	+	+	139,463	+	+	+
	Males	268	142	78	+	+	+	510	+	+	+
	Females	58,080	40,374	26,960	+	+	+	138,953	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.6995%	-	0.0001%	0.0061%	0.0850%	0.4724%	1.1895%	2.1765%	2.9663%	3.4430%	0.6663%
	Males	0.0076%	-	-	-	0.0005%	0.0027%	0.0085%	0.0257%	0.0464%	0.0671%	0.0083%
	Females	1.3735%	-	0.0002%	0.0124%	0.1696%	0.9381%	2.3183%	4.1219%	5.3144%	5.4192%	1.2223%
White ^c	Both Sexes	0.7707%	-	-	0.0060%	0.0836%	0.4761%	1.2281%	2.3039%	3.1836%	3.6666%	0.7001%
	Males	0.0083%	-	-	-	0.0004%	0.0022%	0.0076%	0.0267%	0.0501%	0.0716%	0.0086%
	Females	1.5296%	-	-	0.0123%	0.1710%	0.9647%	2.4332%	4.4294%	5.7661%	5.7904%	1.3002%
Black ^c	Both Sexes	0.4978%	-	-	0.0067%	0.0992%	0.4767%	1.0759%	1.8870%	2.5237%	2.8899%	0.5929%
	Males	0.0072%	-	-	-	-	0.0051%	0.0155%	0.0339%	0.0414%	0.0680%	0.0101%
	Females	0.9444%	-	-	0.0128%	0.1849%	0.8942%	1.9965%	3.3509%	4.2467%	4.2399%	1.0261%
Asian/ Pacific Islander ^c	Both Sexes	0.5218%	-	-	0.0060%	0.0816%	0.4548%	1.0681%	1.6229%	2.0067%	2.1857%	0.5109%
	Males	0.0043%	-	-	-	-	0.0020%	0.0074%	0.0109%	0.0263%	0.0341%	0.0048%
	Females	0.9955%	-	-	0.0117%	0.1539%	0.8567%	1.9783%	2.9593%	3.5278%	3.5077%	0.9160%
Hispanic ^d	Both Sexes	0.2815%	-	-	0.0039%	0.0622%	0.3214%	0.8555%	1.5098%	1.9869%	2.2772%	0.4605%
	Males	0.0020%	-	-	-	-	0.0013%	0.0042%	0.0144%	0.0238%	0.0282%	0.0042%
	Females	0.5677%	-	-	0.0082%	0.1282%	0.6547%	1.6660%	2.7842%	3.4354%	3.6146%	0.8359%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 5.15
Cancer of the Cervix Uteri (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>										
All Races ^b	39,578	33,185	35,754	32,230	28,157	20,702	147,326	211,309	38,323	249,632
White ^b	31,065	26,389	28,581	26,721	23,662	17,154	118,141	171,365	32,320	203,685
Black ^b	5,426	4,525	4,703	3,484	2,922	2,720	18,942	26,687	+	+
Asian/Pacific Islander ^c	2,147	1,764	1,663	+	+	+	6,706	+	+	+
Hispanic ^d	7,909	7,382	6,634	+	+	+	26,543	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f All Ages
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races ^c	0.0928%	-	-	0.0089%	0.0762%	0.1797%	0.1983%	0.1762%	0.1544%	0.0989%	0.0879%
White ^c	0.0966%	-	-	0.0106%	0.0875%	0.1982%	0.2039%	0.1663%	0.1407%	0.0802%	0.0913%
Black ^c	0.0789%	-	-	0.0055%	0.0612%	0.1402%	0.1892%	0.1900%	0.1722%	0.1539%	0.0819%
Asian/Pacific Islander ^c	0.0820%	-	-	0.0036%	0.0325%	0.1095%	0.1693%	0.2115%	0.2172%	0.1893%	0.0755%
Hispanic ^d	0.1082%	-	-	0.0093%	0.0806%	0.2353%	0.3394%	0.3373%	0.3350%	0.2624%	0.1410%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 6.28
Cancer of the Colon and Rectum (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	403,434	281,507	193,503	116,069	76,677	43,080	1,013,148	1,140,625	21,801	1,162,426
	Males	206,233	141,727	97,318	56,567	37,430	18,506	511,310	567,872	7,585	575,457
	Females	197,201	139,780	96,185	59,502	39,247	24,574	501,838	572,753	14,216	586,969
White ^b	Both Sexes	330,229	238,982	166,908	100,366	66,971	38,458	852,636	965,306	19,172	984,478
	Males	170,643	122,414	84,789	49,390	33,264	16,513	435,487	486,080	6,763	492,843
	Females	159,586	116,568	82,119	50,976	33,707	21,945	417,149	479,226	12,409	491,635
Black ^b	Both Sexes	47,635	27,646	17,838	10,579	6,858	3,315	105,375	115,742	1,633	117,375
	Males	22,021	11,691	7,815	4,628	2,708	1,308	46,939	50,729	515	51,244
	Females	25,614	15,955	10,023	5,951	4,150	2,007	58,436	65,013	1,118	66,131
Asian/ Pacific Islander ^c	Both Sexes	17,551	10,605	6,452	+	+	+	37,770	+	+	+
	Males	9,130	5,358	3,296	+	+	+	19,376	+	+	+
	Females	8,421	5,247	3,156	+	+	+	18,394	+	+	+
Hispanic ^d	Both Sexes	30,018	16,387	9,618	+	+	+	60,324	+	+	+
	Males	16,210	8,541	5,180	+	+	+	32,041	+	+	+
	Females	13,808	7,846	4,438	+	+	+	28,283	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.2903%	-	0.0003%	0.0054%	0.0256%	0.1062%	0.3440%	0.7582%	1.5438%	2.3444%	0.2837%
	Males	0.2977%	-	0.0003%	0.0051%	0.0261%	0.1120%	0.3789%	0.8947%	1.8340%	2.6447%	0.3269%
	Females	0.2832%	-	0.0002%	0.0058%	0.0251%	0.1005%	0.3106%	0.6346%	1.3104%	2.1686%	0.2494%
White ^c	Both Sexes	0.3037%	-	0.0003%	0.0059%	0.0263%	0.1036%	0.3285%	0.7322%	1.5490%	2.3641%	0.2806%
	Males	0.3116%	-	0.0003%	0.0054%	0.0267%	0.1078%	0.3660%	0.8720%	1.8444%	2.6569%	0.3241%
	Females	0.2958%	-	0.0003%	0.0063%	0.0258%	0.0992%	0.2915%	0.6017%	1.3055%	2.1911%	0.2450%
Black ^c	Both Sexes	0.2427%	-	-	0.0043%	0.0231%	0.1179%	0.4092%	0.9427%	1.6686%	2.2417%	0.3108%
	Males	0.2295%	-	-	0.0050%	0.0211%	0.1174%	0.4159%	1.0542%	1.8362%	2.4503%	0.3373%
	Females	0.2548%	-	-	0.0037%	0.0248%	0.1184%	0.4033%	0.8547%	1.5523%	2.1419%	0.2929%
Asian/ Pacific Islander ^c	Both Sexes	0.2633%	-	-	0.0035%	0.0229%	0.1028%	0.3556%	0.7371%	1.3975%	2.2217%	0.2700%
	Males	0.2792%	-	-	0.0031%	0.0247%	0.1217%	0.3940%	0.8672%	1.7024%	2.5992%	0.3187%
	Females	0.2488%	-	-	0.0038%	0.0212%	0.0859%	0.3226%	0.6293%	1.1632%	1.9897%	0.2334%
Hispanic ^d	Both Sexes	0.1208%	-	-	0.0032%	0.0205%	0.0772%	0.2672%	0.6349%	1.1851%	1.7389%	0.2212%
	Males	0.1258%	-	-	0.0031%	0.0220%	0.0812%	0.2891%	0.7714%	1.5159%	2.1220%	0.2676%
	Females	0.1157%	-	-	0.0033%	0.0190%	0.0730%	0.2464%	0.5186%	0.9411%	1.5111%	0.1869%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 7.15
Cancer of the Corpus and Uterus, NOS (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>										
All Races ^b	177,109	124,264	99,217	69,643	47,530	33,070	481,292	584,180	26,624	610,804
White ^b	152,262	110,498	90,408	64,493	44,310	31,252	427,975	524,760	22,558	547,318
Black ^b	14,632	7,552	4,342	2,630	1,743	874	29,530	32,651	789	33,440
Asian/Pacific Islander ^c	6,474	4,219	2,718	+	+	+	14,851	+	+	+
Hispanic ^d	12,122	7,994	4,986	+	+	+	27,828	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f All Ages
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races ^c	0.2613%	-	-	0.0031%	0.0297%	0.1156%	0.3846%	0.8965%	1.1480%	1.0079%	0.2319%
White ^c	0.2988%	-	-	0.0027%	0.0289%	0.1169%	0.4058%	0.9799%	1.2908%	1.1246%	0.2525%
Black ^c	0.1295%	-	-	0.0047%	0.0248%	0.0766%	0.2046%	0.5624%	0.7916%	0.5267%	0.1450%
Asian/Pacific Islander ^c	0.1856%	-	-	0.0038%	0.0319%	0.1311%	0.4046%	0.6524%	0.5828%	0.4767%	0.1676%
Hispanic ^d	0.1136%	-	-	0.0042%	0.0365%	0.1187%	0.2997%	0.6408%	0.7037%	0.5882%	0.1659%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 8.21
Cancer of the Esophagus (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	21,165	7,280	3,839	1,333	498	214	33,798	34,428	123	34,551
	Males	16,662	5,575	2,970	1,030	367	169	26,376	26,846	63	26,909
	Females	4,503	1,705	869	303	131	45	7,422	7,582	60	7,642
White ^b	Both Sexes	18,841	6,591	3,355	1,176	416	187	30,106	30,662	96	30,758
	Males	15,108	5,155	2,692	901	310	142	23,975	24,379	56	24,435
	Females	3,733	1,436	663	275	106	45	6,131	6,283	40	6,323
Black ^b	Both Sexes	1,710	501	364	121	65	28	2,723	2,791	31	2,822
	Males	1,063	275	206	97	52	28	1,655	1,721	9	1,730
	Females	647	226	158	24	13	0	1,068	1,070	22	1,092
Asian/ Pacific Islander ^c	Both Sexes	413	109	87	+	+	+	639	+	+	+
	Males	319	80	50	+	+	+	474	+	+	+
	Females	94	29	37	+	+	+	165	+	+	+
Hispanic ^d	Both Sexes	903	252	112	+	+	+	1,344	+	+	+
	Males	759	211	96	+	+	+	1,125	+	+	+
	Females	144	41	16	+	+	+	219	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted ^f All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0089%	-	-	-	0.0004%	0.0027%	0.0121%	0.0327%	0.0516%	0.0395%	0.0086%
	Males	0.0140%	-	-	-	0.0006%	0.0044%	0.0194%	0.0571%	0.0859%	0.0744%	0.0147%
	Females	0.0040%	-	-	-	0.0002%	0.0010%	0.0051%	0.0106%	0.0241%	0.0190%	0.0036%
White ^c	Both Sexes	0.0103%	-	-	-	0.0005%	0.0030%	0.0132%	0.0371%	0.0565%	0.0412%	0.0094%
	Males	0.0164%	-	-	-	0.0008%	0.0050%	0.0211%	0.0652%	0.0951%	0.0780%	0.0163%
	Females	0.0043%	-	-	-	-	0.0010%	0.0053%	0.0109%	0.0246%	0.0194%	0.0037%
Black ^c	Both Sexes	0.0058%	-	-	-	-	0.0017%	0.0124%	0.0269%	0.0430%	0.0257%	0.0070%
	Males	0.0074%	-	-	-	-	0.0025%	0.0155%	0.0380%	0.0635%	0.0540%	0.0103%
	Females	0.0043%	-	-	-	-	-	0.0097%	0.0181%	0.0287%	0.0121%	0.0047%
Asian/ Pacific Islander ^c	Both Sexes	0.0044%	-	-	-	-	0.0021%	0.0055%	0.0110%	0.0297%	0.0342%	0.0047%
	Males	0.0068%	-	-	-	-	0.0032%	0.0109%	0.0196%	0.0450%	0.0593%	0.0078%
	Females	0.0023%	-	-	-	-	0.0011%	-	0.0038%	0.0180%	0.0187%	0.0023%
Hispanic ^d	Both Sexes	0.0027%	-	-	-	0.0006%	0.0020%	0.0056%	0.0163%	0.0278%	0.0258%	0.0048%
	Males	0.0044%	-	-	-	0.0010%	0.0032%	0.0097%	0.0297%	0.0545%	0.0562%	0.0091%
	Females	0.0009%	-	-	-	-	-	0.0016%	0.0049%	0.0081%	0.0078%	0.0014%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 9.21
Hodgkin LymphomaEstimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	38,529	32,259	28,383	24,620	20,988	16,105	128,515	172,689	13,104	185,793
	Males	20,912	16,780	14,406	12,472	10,357	8,594	67,067	89,550	6,550	96,100
	Females	17,617	15,479	13,977	12,148	10,631	7,511	61,448	83,139	6,554	89,693
White ^b	Both Sexes	32,135	27,499	24,853	22,086	19,267	15,073	110,813	152,110	14,550	166,660
	Males	17,509	14,470	12,551	11,186	9,575	7,839	58,021	78,738	7,517	86,255
	Females	14,626	13,029	12,302	10,900	9,692	7,234	52,792	73,372	7,033	80,405
Black ^b	Both Sexes	4,833	3,583	2,799	2,133	1,343	851	13,748	15,982	1,003	16,985
	Males	2,622	1,622	1,453	1,099	587	675	6,946	8,372	707	9,079
	Females	2,211	1,961	1,346	1,034	756	176	6,802	7,610	296	7,906
Asian/ Pacific Islander ^c	Both Sexes	932	734	423	+	+	+	2,315	+	+	+
	Males	454	390	230	+	+	+	1,161	+	+	+
	Females	478	344	193	+	+	+	1,154	+	+	+
Hispanic ^d	Both Sexes	4,173	3,005	2,322	+	+	+	10,828	+	+	+
	Males	2,290	1,665	1,285	+	+	+	5,982	+	+	+
	Females	1,883	1,340	1,037	+	+	+	4,846	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0351%	0.0007%	0.0106%	0.0395%	0.0624%	0.0593%	0.0412%	0.0327%	0.0313%	0.0238%	0.0350%
	Males	0.0373%	0.0009%	0.0117%	0.0388%	0.0611%	0.0627%	0.0482%	0.0399%	0.0360%	0.0278%	0.0372%
	Females	0.0329%	0.0005%	0.0093%	0.0402%	0.0636%	0.0559%	0.0346%	0.0261%	0.0275%	0.0214%	0.0331%
White ^c	Both Sexes	0.0399%	0.0007%	0.0116%	0.0448%	0.0724%	0.0675%	0.0466%	0.0367%	0.0357%	0.0270%	0.0400%
	Males	0.0422%	0.0009%	0.0130%	0.0430%	0.0706%	0.0708%	0.0534%	0.0447%	0.0408%	0.0314%	0.0420%
	Females	0.0376%	0.0005%	0.0101%	0.0468%	0.0743%	0.0642%	0.0398%	0.0292%	0.0315%	0.0244%	0.0381%
Black ^c	Both Sexes	0.0311%	0.0007%	0.0097%	0.0306%	0.0596%	0.0569%	0.0404%	0.0317%	0.0256%	0.0135%	0.0321%
	Males	0.0327%	-	0.0088%	0.0327%	0.0569%	0.0608%	0.0504%	0.0358%	0.0320%	0.0159%	0.0344%
	Females	0.0297%	-	0.0107%	0.0286%	0.0619%	0.0534%	0.0317%	0.0283%	0.0212%	0.0124%	0.0302%
Asian/ Pacific Islander ^c	Both Sexes	0.0132%	-	0.0053%	0.0213%	0.0215%	0.0191%	0.0117%	0.0097%	0.0095%	0.0078%	0.0126%
	Males	0.0138%	-	0.0067%	0.0240%	0.0201%	0.0187%	0.0125%	0.0122%	0.0102%	0.0092%	0.0134%
	Females	0.0126%	-	0.0038%	0.0186%	0.0227%	0.0194%	0.0110%	0.0076%	0.0090%	0.0069%	0.0119%
Hispanic ^d	Both Sexes	0.0214%	0.0011%	0.0100%	0.0263%	0.0336%	0.0329%	0.0283%	0.0329%	0.0373%	0.0204%	0.0236%
	Males	0.0233%	0.0012%	0.0127%	0.0271%	0.0331%	0.0357%	0.0355%	0.0369%	0.0389%	0.0423%	0.0263%
	Females	0.0195%	0.0009%	0.0072%	0.0254%	0.0341%	0.0299%	0.0214%	0.0295%	0.0361%	0.0074%	0.0212%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 10.15
Kaposi SarcomaEstimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	7,194	5,578	5,671	5,046	1,392	430	23,872	25,504	+	+
	Males	6,655	5,284	5,439	4,842	1,369	396	22,579	24,134	+	+
	Females	539	294	232	204	23	34	1,293	1,370	+	+
White ^b	Both Sexes	4,740	4,115	4,237	4,257	1,088	369	17,633	18,977	+	+
	Males	4,353	3,896	4,073	4,087	1,088	346	16,692	17,970	+	+
	Females	387	219	164	170	0	23	941	1,007	+	+
Black ^b	Both Sexes	1,980	1,234	1,078	520	122	26	4,840	4,973	+	+
	Males	1,874	1,177	1,056	486	122	26	4,620	4,753	+	+
	Females	106	57	22	34	0	0	220	220	+	+
Asian/ Pacific Islander ^c	Both Sexes	118	97	79	+	+	+	339	+	+	+
	Males	118	91	79	+	+	+	333	+	+	+
	Females	0	6	0	+	+	+	6	+	+	+
Hispanic ^d	Both Sexes	1,227	1,113	1,058	+	+	+	4,149	+	+	+
	Males	1,093	1,071	1,017	+	+	+	3,916	+	+	+
	Females	134	42	41	+	+	+	233	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0084%	-	-	0.0012%	0.0058%	0.0205%	0.0188%	0.0111%	0.0093%	0.0145%	0.0081%
	Males	0.0162%	-	-	0.0021%	0.0114%	0.0403%	0.0381%	0.0224%	0.0177%	0.0275%	0.0160%
	Females	0.0007%	-	-	-	0.0002%	0.0008%	0.0004%	0.0009%	0.0026%	0.0069%	0.0006%
White ^c	Both Sexes	0.0085%	-	-	0.0008%	0.0049%	0.0203%	0.0194%	0.0117%	0.0100%	0.0153%	0.0080%
	Males	0.0163%	-	-	0.0014%	0.0093%	0.0390%	0.0387%	0.0232%	0.0189%	0.0288%	0.0156%
	Females	0.0008%	-	-	-	-	0.0009%	0.0003%	0.0009%	0.0026%	0.0073%	0.0007%
Black ^c	Both Sexes	0.0118%	-	-	0.0038%	0.0156%	0.0342%	0.0236%	0.0125%	0.0064%	0.0089%	0.0124%
	Males	0.0239%	-	-	0.0072%	0.0329%	0.0719%	0.0497%	0.0263%	0.0114%	-	0.0257%
	Females	0.0008%	-	-	-	-	-	-	-	-	-	0.0009%
Asian/ Pacific Islander ^c	Both Sexes	0.0021%	-	-	-	0.0011%	0.0047%	0.0050%	0.0023%	0.0041%	0.0044%	0.0020%
	Males	0.0043%	-	-	-	0.0024%	0.0097%	0.0107%	0.0051%	0.0094%	0.0092%	0.0044%
	Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic ^d	Both Sexes	0.0083%	-	-	0.0011%	0.0065%	0.0252%	0.0228%	0.0153%	0.0179%	0.0385%	0.0111%
	Males	0.0155%	-	-	0.0020%	0.0127%	0.0481%	0.0462%	0.0309%	0.0318%	0.0659%	0.0212%
	Females	0.0010%	-	-	-	-	0.0015%	-	0.0020%	0.0076%	0.0222%	0.0017%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 11.21
Cancer of the Kidney And Renal Pelvis (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	149,969	87,553	48,081	29,936	17,014	9,310	319,656	347,916	10,687	358,603
	Males	91,313	52,407	27,681	17,630	9,319	5,384	191,436	206,735	5,968	212,703
	Females	58,656	35,146	20,400	12,306	7,695	3,926	128,220	141,181	4,719	145,900
White ^b	Both Sexes	124,730	75,017	41,056	26,270	15,289	8,156	270,771	295,946	12,063	308,009
	Males	76,289	44,953	24,231	15,726	8,541	4,825	163,388	177,362	5,351	182,713
	Females	48,441	30,064	16,825	10,544	6,748	3,331	107,383	118,584	6,712	125,296
Black ^b	Both Sexes	17,603	8,902	5,336	2,748	1,415	943	34,918	37,454	2,008	39,462
	Males	10,344	5,292	2,520	1,364	618	439	19,696	20,746	686	21,432
	Females	7,259	3,610	2,816	1,384	797	504	15,222	16,708	1,322	18,030
Asian/ Pacific Islander ^c	Both Sexes	3,691	1,937	928	+	+	+	6,954	+	+	+
	Males	2,265	1,178	550	+	+	+	4,236	+	+	+
	Females	1,426	759	378	+	+	+	2,718	+	+	+
Hispanic ^d	Both Sexes	13,867	7,357	3,621	+	+	+	26,557	+	+	+
	Males	7,930	4,232	1,969	+	+	+	15,065	+	+	+
	Females	5,937	3,125	1,652	+	+	+	11,492	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0891%	0.0069%	0.0108%	0.0050%	0.0148%	0.0542%	0.1360%	0.2805%	0.4081%	0.3692%	0.0859%
	Males	0.1090%	0.0068%	0.0094%	0.0045%	0.0162%	0.0670%	0.1761%	0.3798%	0.5546%	0.5191%	0.1132%
	Females	0.0696%	0.0070%	0.0122%	0.0055%	0.0135%	0.0415%	0.0977%	0.1907%	0.2903%	0.2815%	0.0633%
White ^c	Both Sexes	0.0983%	0.0073%	0.0117%	0.0053%	0.0154%	0.0582%	0.1432%	0.2955%	0.4344%	0.3852%	0.0908%
	Males	0.1199%	0.0071%	0.0111%	0.0046%	0.0161%	0.0721%	0.1831%	0.3946%	0.5887%	0.5452%	0.1189%
	Females	0.0768%	0.0076%	0.0123%	0.0062%	0.0146%	0.0439%	0.1038%	0.2030%	0.3072%	0.2907%	0.0669%
Black ^c	Both Sexes	0.0775%	0.0079%	0.0122%	0.0068%	0.0163%	0.0543%	0.1511%	0.3156%	0.4219%	0.3511%	0.0913%
	Males	0.0926%	0.0092%	0.0072%	0.0069%	0.0196%	0.0658%	0.2021%	0.4398%	0.5601%	0.4616%	0.1194%
	Females	0.0638%	0.0066%	0.0174%	0.0068%	0.0135%	0.0441%	0.1068%	0.2174%	0.3260%	0.2982%	0.0704%
Asian/ Pacific Islander ^c	Both Sexes	0.0465%	0.0038%	0.0035%	0.0018%	0.0072%	0.0282%	0.0707%	0.1483%	0.2185%	0.2493%	0.0467%
	Males	0.0589%	0.0025%	0.0028%	0.0024%	0.0085%	0.0326%	0.0973%	0.2195%	0.3053%	0.3473%	0.0640%
	Females	0.0351%	0.0052%	0.0044%	0.0012%	0.0061%	0.0243%	0.0479%	0.0893%	0.1519%	0.1890%	0.0333%
Hispanic ^d	Both Sexes	0.0530%	0.0060%	0.0086%	0.0032%	0.0145%	0.0510%	0.1291%	0.2729%	0.4138%	0.4443%	0.0867%
	Males	0.0590%	0.0053%	0.0068%	0.0024%	0.0137%	0.0594%	0.1519%	0.3605%	0.5785%	0.5581%	0.1103%
	Females	0.0468%	0.0066%	0.0105%	0.0042%	0.0153%	0.0423%	0.1074%	0.1982%	0.2923%	0.3766%	0.0685%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 12.21
Cancer of the Larynx (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	31,318	19,778	14,256	9,873	6,099	4,176	76,595	87,652	1,613	89,265
	Males	25,568	15,842	11,344	7,903	4,912	3,343	61,732	70,687	1,312	71,999
	Females	5,750	3,936	2,912	1,970	1,187	833	14,863	16,965	301	17,266
White ^b	Both Sexes	25,970	16,812	12,353	8,571	5,198	3,762	64,875	74,536	1,269	75,805
	Males	21,154	13,465	9,921	6,848	4,288	3,059	52,301	60,256	1,030	61,286
	Females	4,816	3,347	2,432	1,723	910	703	12,574	14,280	239	14,519
Black ^b	Both Sexes	4,352	2,446	1,529	1,092	729	346	9,586	10,700	110	10,810
	Males	3,512	1,947	1,088	901	531	233	7,581	8,406	92	8,498
	Females	840	499	441	191	198	113	2,005	2,294	18	2,312
Asian/ Pacific Islander ^c	Both Sexes	511	361	224	+	+	+	1,214	+	+	+
	Males	456	312	198	+	+	+	1,067	+	+	+
	Females	55	49	26	+	+	+	147	+	+	+
Hispanic ^d	Both Sexes	1,904	1,027	745	+	+	+	4,036	+	+	+
	Males	1,651	844	588	+	+	+	3,380	+	+	+
	Females	253	183	157	+	+	+	656	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted ^f All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0201%	-	-	0.0003%	0.0008%	0.0052%	0.0264%	0.0718%	0.1201%	0.0969%	0.0195%
	Males	0.0332%	-	-	0.0004%	0.0009%	0.0072%	0.0417%	0.1258%	0.2265%	0.2161%	0.0358%
	Females	0.0073%	-	-	0.0002%	0.0007%	0.0031%	0.0118%	0.0230%	0.0345%	0.0271%	0.0066%
White ^c	Both Sexes	0.0222%	-	-	0.0003%	0.0009%	0.0058%	0.0284%	0.0737%	0.1270%	0.0997%	0.0205%
	Males	0.0363%	-	-	0.0004%	0.0009%	0.0080%	0.0443%	0.1267%	0.2367%	0.2200%	0.0369%
	Females	0.0082%	-	-	-	0.0009%	0.0036%	0.0128%	0.0243%	0.0365%	0.0287%	0.0071%
Black ^c	Both Sexes	0.0213%	-	-	-	0.0010%	0.0052%	0.0365%	0.1131%	0.1553%	0.1106%	0.0263%
	Males	0.0353%	-	-	-	0.0019%	0.0068%	0.0578%	0.2074%	0.3109%	0.2720%	0.0511%
	Females	0.0086%	-	-	-	-	0.0038%	0.0180%	0.0385%	0.0474%	0.0334%	0.0093%
Asian/ Pacific Islander ^c	Both Sexes	0.0085%	-	-	-	-	0.0016%	0.0071%	0.0302%	0.0537%	0.0698%	0.0088%
	Males	0.0156%	-	-	-	-	0.0030%	0.0126%	0.0615%	0.1054%	0.1633%	0.0182%
	Females	0.0020%	-	-	-	-	-	0.0023%	0.0042%	0.0139%	0.0123%	0.0019%
Hispanic ^d	Both Sexes	0.0080%	-	-	0.0003%	0.0007%	0.0039%	0.0153%	0.0458%	0.0995%	0.0984%	0.0150%
	Males	0.0132%	-	-	-	0.0008%	0.0053%	0.0249%	0.0863%	0.2063%	0.2204%	0.0297%
	Females	0.0027%	-	-	-	-	0.0025%	0.0062%	0.0112%	0.0206%	0.0259%	0.0041%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 13.29
LeukemiaEstimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	115,823	74,234	43,360	28,669	16,988	10,694	265,824	297,129	5,671	302,800
	Males	66,545	42,601	24,570	16,036	9,539	5,550	151,719	168,443	2,526	170,969
	Females	49,278	31,633	18,790	12,633	7,449	5,144	114,105	128,686	3,145	131,831
White ^b	Both Sexes	100,301	65,805	38,527	25,841	15,211	9,730	233,818	262,198	4,749	266,947
	Males	57,669	37,933	21,855	14,604	8,651	5,127	133,872	149,222	1,783	151,005
	Females	42,632	27,872	16,672	11,237	6,560	4,603	99,946	112,976	2,966	115,942
Black ^b	Both Sexes	8,436	4,907	2,701	1,585	856	499	17,800	19,316	745	20,061
	Males	4,923	2,654	1,404	753	386	235	9,791	10,473	352	10,825
	Females	3,513	2,253	1,297	832	470	264	8,009	8,843	393	9,236
Asian/ Pacific Islander ^c	Both Sexes	2,958	1,820	1,176	+	+	+	6,662	+	+	+
	Males	1,702	986	676	+	+	+	3,720	+	+	+
	Females	1,256	834	500	+	+	+	2,942	+	+	+
Hispanic ^d	Both Sexes	11,187	6,994	4,455	+	+	+	25,370	+	+	+
	Males	6,457	3,950	2,431	+	+	+	14,442	+	+	+
	Females	4,730	3,044	2,024	+	+	+	10,928	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)								Age-Adjusted ^f		
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0760%	0.0320%	0.0597%	0.0368%	0.0265%	0.0408%	0.0773%	0.1615%	0.2685%	0.3153%	0.0753%
	Males	0.0883%	0.0342%	0.0654%	0.0401%	0.0298%	0.0493%	0.0948%	0.2097%	0.3581%	0.4093%	0.0927%
	Females	0.0641%	0.0298%	0.0538%	0.0334%	0.0232%	0.0325%	0.0606%	0.1178%	0.1964%	0.2602%	0.0607%
White ^c	Both Sexes	0.0869%	0.0357%	0.0686%	0.0417%	0.0293%	0.0427%	0.0843%	0.1794%	0.3005%	0.3476%	0.0835%
	Males	0.1004%	0.0377%	0.0747%	0.0453%	0.0328%	0.0508%	0.1036%	0.2314%	0.4000%	0.4484%	0.1023%
	Females	0.0734%	0.0336%	0.0622%	0.0377%	0.0256%	0.0343%	0.0653%	0.1309%	0.2185%	0.2882%	0.0673%
Black ^c	Both Sexes	0.0403%	0.0174%	0.0257%	0.0232%	0.0176%	0.0317%	0.0501%	0.1007%	0.1685%	0.1791%	0.0459%
	Males	0.0460%	0.0202%	0.0282%	0.0233%	0.0200%	0.0395%	0.0608%	0.1306%	0.2166%	0.2424%	0.0567%
	Females	0.0351%	0.0145%	0.0231%	0.0231%	0.0154%	0.0248%	0.0409%	0.0771%	0.1350%	0.1488%	0.0380%
Asian/ Pacific Islander ^c	Both Sexes	0.0402%	0.0271%	0.0482%	0.0244%	0.0201%	0.0335%	0.0411%	0.0640%	0.0984%	0.1124%	0.0412%
	Males	0.0470%	0.0295%	0.0548%	0.0272%	0.0220%	0.0430%	0.0464%	0.0763%	0.1302%	0.1686%	0.0498%
	Females	0.0339%	0.0247%	0.0412%	0.0216%	0.0185%	0.0250%	0.0364%	0.0538%	0.0740%	0.0779%	0.0342%
Hispanic ^d	Both Sexes	0.0497%	0.0359%	0.0680%	0.0396%	0.0269%	0.0362%	0.0528%	0.0978%	0.1455%	0.1624%	0.0571%
	Males	0.0557%	0.0398%	0.0743%	0.0447%	0.0322%	0.0434%	0.0592%	0.1194%	0.1739%	0.2068%	0.0667%
	Females	0.0436%	0.0319%	0.0614%	0.0339%	0.0212%	0.0288%	0.0467%	0.0793%	0.1245%	0.1360%	0.0486%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 13.30
Acute Lymphocytic Leukemia

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	16,260	13,209	11,408	9,489	7,668	6,025	51,988	68,728	3,509	72,237
	Males	9,051	7,551	6,191	5,281	4,261	3,012	28,925	37,729	1,785	39,514
	Females	7,209	5,658	5,217	4,208	3,407	3,013	23,063	30,999	1,724	32,723
White ^b	Both Sexes	13,645	11,332	9,992	8,237	6,726	5,465	44,665	59,741	+	+
	Males	7,482	6,533	5,388	4,677	3,754	2,752	24,848	32,841	+	+
	Females	6,163	4,799	4,604	3,560	2,972	2,713	19,817	26,900	+	+
Black ^b	Both Sexes	1,384	888	726	679	444	268	3,734	4,539	+	+
	Males	884	471	385	295	225	140	2,071	2,461	+	+
	Females	500	417	341	384	219	128	1,663	2,078	+	+
Asian/ Pacific Islander ^c	Both Sexes	768	661	464	+	+	+	2,239	+	+	+
	Males	432	370	295	+	+	+	1,266	+	+	+
	Females	336	291	169	+	+	+	973	+	+	+
Hispanic ^d	Both Sexes	5,096	3,281	2,589	+	+	+	12,682	+	+	+
	Males	3,025	1,865	1,393	+	+	+	7,311	+	+	+
	Females	2,071	1,416	1,196	+	+	+	5,371	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0169%	0.0269%	0.0512%	0.0249%	0.0073%	0.0047%	0.0039%	0.0042%	0.0031%	0.0019%	0.0174%
	Males	0.0193%	0.0288%	0.0562%	0.0281%	0.0090%	0.0052%	0.0042%	0.0048%	0.0032%	0.0028%	0.0193%
	Females	0.0146%	0.0248%	0.0459%	0.0214%	0.0056%	0.0041%	0.0037%	0.0037%	0.0030%	0.0014%	0.0154%
White ^c	Both Sexes	0.0191%	0.0303%	0.0597%	0.0292%	0.0088%	0.0051%	0.0042%	0.0045%	0.0030%	0.0018%	0.0200%
	Males	0.0216%	0.0323%	0.0649%	0.0330%	0.0108%	0.0057%	0.0044%	0.0052%	0.0034%	0.0023%	0.0220%
	Females	0.0167%	0.0283%	0.0541%	0.0252%	0.0068%	0.0045%	0.0041%	0.0040%	0.0026%	0.0015%	0.0179%
Black ^c	Both Sexes	0.0083%	0.0127%	0.0200%	0.0125%	0.0026%	0.0023%	0.0018%	0.0017%	-	-	0.0076%
	Males	0.0096%	0.0157%	0.0220%	0.0133%	0.0028%	0.0025%	-	-	-	-	0.0083%
	Females	0.0071%	0.0096%	0.0178%	0.0118%	0.0025%	0.0022%	0.0022%	-	-	-	0.0068%
Asian/ Pacific Islander ^c	Both Sexes	0.0122%	0.0227%	0.0388%	0.0146%	0.0043%	0.0041%	0.0036%	0.0033%	0.0041%	0.0029%	0.0132%
	Males	0.0145%	0.0239%	0.0448%	0.0169%	0.0054%	0.0044%	0.0053%	0.0037%	-	0.0077%	0.0149%
	Females	0.0101%	0.0215%	0.0326%	0.0123%	0.0034%	0.0038%	0.0020%	0.0030%	0.0059%	-	0.0114%
Hispanic ^d	Both Sexes	0.0245%	0.0303%	0.0605%	0.0278%	0.0095%	0.0066%	0.0058%	0.0077%	0.0047%	-	0.0208%
	Males	0.0278%	0.0337%	0.0660%	0.0326%	0.0115%	0.0085%	0.0057%	0.0066%	-	-	0.0231%
	Females	0.0211%	0.0267%	0.0547%	0.0226%	0.0073%	0.0046%	0.0060%	0.0086%	0.0045%	-	0.0182%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 14.21
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	33,682	7,065	2,521	1,025	580	430	44,415	45,468	474	45,942
	Males	24,703	4,859	1,647	591	231	212	31,870	32,312	77	32,389
	Females	8,979	2,206	874	434	349	218	12,545	13,156	397	13,553
White ^b	Both Sexes	24,394	5,153	1,799	771	440	348	32,185	33,019	333	33,352
	Males	18,015	3,649	1,149	444	157	151	23,289	23,620	210 ⁱ	23,830
	Females	6,379	1,504	650	327	283	197	8,896	9,399	123 ⁱ	9,522
Black ^b	Both Sexes	5,180	717	175	57	84	37	6,164	6,283	+	+
	Males	3,811	363	105	12	35	27	4,314	4,364	+	+
	Females	1,369	354	70	45	49	10	1,850	1,919	+	+
Asian/ Pacific Islander ^c	Both Sexes	3,492	1,064	448	+	+	+	5,105	+	+	+
	Males	2,458	720	314	+	+	+	3,566	+	+	+
	Females	1,034	344	134	+	+	+	1,539	+	+	+
Hispanic ^d	Both Sexes	4,938	935	317	+	+	+	6,375	+	+	+
	Males	3,569	649	188	+	+	+	4,530	+	+	+
	Females	1,369	286	129	+	+	+	1,845	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0163%	0.0028%	0.0024%	0.0010%	0.0017%	0.0063%	0.0340%	0.0583%	0.0633%	0.0413%	0.0152%
	Males	0.0235%	0.0034%	0.0024%	0.0013%	0.0021%	0.0089%	0.0552%	0.0924%	0.0887%	0.0634%	0.0227%
	Females	0.0093%	0.0022%	0.0024%	0.0007%	0.0013%	0.0037%	0.0138%	0.0274%	0.0428%	0.0284%	0.0085%
White ^c	Both Sexes	0.0134%	0.0031%	0.0025%	0.0010%	0.0013%	0.0049%	0.0296%	0.0471%	0.0417%	0.0269%	0.0119%
	Males	0.0195%	0.0037%	0.0026%	0.0012%	0.0014%	0.0066%	0.0476%	0.0749%	0.0582%	0.0422%	0.0177%
	Females	0.0075%	0.0026%	0.0024%	0.0007%	0.0012%	0.0031%	0.0119%	0.0211%	0.0281%	0.0178%	0.0066%
Black ^c	Both Sexes	0.0135%	-	0.0009%	0.0015%	0.0022%	0.0043%	0.0351%	0.0758%	0.0511%	0.0263%	0.0146%
	Males	0.0198%	-	0.0012%	0.0019%	0.0028%	0.0062%	0.0555%	0.1266%	0.0746%	0.0415%	0.0228%
	Females	0.0077%	-	-	-	0.0017%	0.0026%	0.0174%	0.0357%	0.0347%	0.0191%	0.0081%
Asian/ Pacific Islander ^c	Both Sexes	0.0340%	0.0026%	0.0033%	0.0007%	0.0033%	0.0156%	0.0557%	0.1115%	0.1920%	0.1436%	0.0343%
	Males	0.0492%	0.0039%	0.0025%	0.0012%	0.0052%	0.0243%	0.0978%	0.1769%	0.2762%	0.2101%	0.0525%
	Females	0.0202%	-	0.0040%	-	0.0016%	0.0079%	0.0195%	0.0573%	0.1273%	0.1027%	0.0197%
Hispanic ^d	Both Sexes	0.0127%	0.0035%	0.0024%	0.0008%	0.0009%	0.0069%	0.0438%	0.0786%	0.0746%	0.0636%	0.0193%
	Males	0.0178%	0.0044%	0.0027%	0.0009%	0.0007%	0.0103%	0.0727%	0.1258%	0.0974%	0.1028%	0.0293%
	Females	0.0075%	0.0027%	0.0022%	0.0008%	0.0011%	0.0034%	0.0162%	0.0383%	0.0577%	0.0403%	0.0109%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 15.27
Cancer of the Lung And Bronchus (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	231,126	77,607	41,694	22,634	12,124	6,214	376,088	395,186	7,140	402,326
	Males	107,276	34,010	17,823	10,198	5,209	2,765	170,515	179,129	4,086	183,215
	Females	123,850	43,597	23,871	12,436	6,915	3,449	205,573	216,057	3,054	219,111
White ^b	Both Sexes	196,046	67,584	36,611	19,757	10,797	5,441	322,693	339,549	2,910	342,459
	Males	90,417	29,237	15,323	8,775	4,505	2,411	144,794	152,256	1,359	153,615
	Females	105,629	38,347	21,288	10,982	6,292	3,030	177,899	187,293	1,551	188,844
Black ^b	Both Sexes	24,928	7,416	3,703	2,194	998	611	38,499	40,198	261	40,459
	Males	11,660	3,506	1,755	1,068	548	275	18,124	19,014	137	19,151
	Females	13,268	3,910	1,948	1,126	450	336	20,375	21,184	124	21,308
Asian/ Pacific Islander ^c	Both Sexes	7,365	2,011	1,020	+	+	+	10,778	+	+	+
	Males	3,791	964	548	+	+	+	5,503	+	+	+
	Females	3,574	1,047	472	+	+	+	5,275	+	+	+
Hispanic ^d	Both Sexes	8,687	2,559	1,275	+	+	+	13,118	+	+	+
	Males	4,093	1,156	606	+	+	+	6,151	+	+	+
	Females	4,594	1,403	669	+	+	+	6,967	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.1014%	0.0001%	0.0004%	0.0016%	0.0056%	0.0244%	0.1083%	0.3249%	0.6490%	0.6211%	0.1004%
	Males	0.0941%	-	0.0004%	0.0015%	0.0050%	0.0212%	0.1016%	0.3324%	0.6722%	0.7046%	0.1036%
	Females	0.1085%	-	0.0003%	0.0017%	0.0061%	0.0276%	0.1148%	0.3181%	0.6303%	0.5722%	0.0985%
White ^c	Both Sexes	0.1099%	0.0001%	0.0004%	0.0017%	0.0057%	0.0246%	0.1066%	0.3316%	0.6769%	0.6405%	0.1031%
	Males	0.0991%	-	0.0004%	0.0016%	0.0049%	0.0214%	0.0976%	0.3283%	0.6772%	0.7051%	0.1032%
	Females	0.1206%	-	0.0004%	0.0019%	0.0065%	0.0279%	0.1155%	0.3346%	0.6767%	0.6023%	0.1038%
Black ^c	Both Sexes	0.0871%	-	-	0.0016%	0.0062%	0.0314%	0.1539%	0.3995%	0.6713%	0.4987%	0.1092%
	Males	0.0863%	-	-	0.0019%	0.0076%	0.0262%	0.1533%	0.4525%	0.7743%	0.6605%	0.1236%
	Females	0.0878%	-	-	0.0013%	0.0050%	0.0360%	0.1544%	0.3575%	0.5998%	0.4213%	0.0999%
Asian/ Pacific Islander ^c	Both Sexes	0.0750%	-	-	0.0006%	0.0048%	0.0190%	0.0843%	0.2351%	0.4880%	0.5551%	0.0781%
	Males	0.0798%	-	-	-	0.0038%	0.0181%	0.0890%	0.2764%	0.5883%	0.7127%	0.0926%
	Females	0.0707%	-	-	-	0.0057%	0.0199%	0.0804%	0.2009%	0.4109%	0.4582%	0.0676%
Hispanic ^d	Both Sexes	0.0262%	-	-	0.0011%	0.0034%	0.0128%	0.0439%	0.1481%	0.3303%	0.3435%	0.0499%
	Males	0.0241%	-	-	0.0008%	0.0034%	0.0115%	0.0434%	0.1493%	0.3697%	0.3995%	0.0543%
	Females	0.0284%	-	-	0.0015%	0.0035%	0.0141%	0.0443%	0.1471%	0.3011%	0.3103%	0.0471%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 16.21
Melanoma of the Skin (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	282,028	208,237	155,229	107,755	79,776	53,373	771,284	924,397	35,834	960,231
	Males	151,313	105,989	77,824	52,347	35,218	22,168	395,832	459,686	11,534	471,220
	Females	130,715	102,248	77,405	55,408	44,558	31,205	375,452	464,711	24,300	489,011
White ^b	Both Sexes	270,630	201,635	149,586	103,407	76,509	51,407	742,777	889,700	33,779	923,479
	Males	145,767	102,885	74,988	50,377	34,052	21,320	382,176	443,630	10,979	454,609
	Females	124,863	98,750	74,598	53,030	42,457	30,087	360,601	446,070	22,800	468,870
Black ^b	Both Sexes	1,123	658	572	428	265	124	2,836	3,280	102	3,382
	Males	477	277	262	240	59	25	1,269	1,378	43	1,421
	Females	646	381	310	188	206	99	1,567	1,902	59	1,961
Asian/ Pacific Islander ^c	Both Sexes	686	539	383	+	+	+	1,760	+	+	+
	Males	327	228	159	+	+	+	785	+	+	+
	Females	359	311	224	+	+	+	975	+	+	+
Hispanic ^d	Both Sexes	4,793	3,797	2,672	+	+	+	12,743	+	+	+
	Males	2,004	1,289	965	+	+	+	4,783	+	+	+
	Females	2,789	2,508	1,707	+	+	+	7,960	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.1944%	0.0003%	0.0030%	0.0287%	0.0978%	0.2026%	0.3399%	0.5105%	0.6446%	0.7067%	0.1870%
	Males	0.2051%	0.0002%	0.0027%	0.0185%	0.0694%	0.1656%	0.3440%	0.6304%	0.9005%	1.0955%	0.2132%
	Females	0.1841%	0.0003%	0.0033%	0.0393%	0.1263%	0.2393%	0.3359%	0.4020%	0.4388%	0.4790%	0.1701%
White ^c	Both Sexes	0.2525%	0.0002%	0.0037%	0.0377%	0.1288%	0.2611%	0.4308%	0.6328%	0.7985%	0.8403%	0.2345%
	Males	0.2642%	-	0.0033%	0.0242%	0.0888%	0.2094%	0.4296%	0.7704%	1.1032%	1.2982%	0.2620%
	Females	0.2408%	0.0003%	0.0042%	0.0523%	0.1709%	0.3144%	0.4321%	0.5044%	0.5474%	0.5698%	0.2179%
Black ^c	Both Sexes	0.0061%	-	-	-	0.0042%	0.0067%	0.0087%	0.0198%	0.0310%	0.0431%	0.0075%
	Males	0.0059%	-	-	-	0.0030%	0.0043%	0.0093%	0.0224%	0.0414%	0.0730%	0.0089%
	Females	0.0063%	-	-	-	0.0052%	0.0087%	0.0081%	0.0177%	0.0238%	0.0288%	0.0069%
Asian/ Pacific Islander ^c	Both Sexes	0.0112%	-	0.0010%	0.0023%	0.0068%	0.0143%	0.0162%	0.0286%	0.0328%	0.0449%	0.0111%
	Males	0.0106%	-	-	0.0024%	0.0068%	0.0089%	0.0147%	0.0316%	0.0452%	0.0506%	0.0113%
	Females	0.0119%	-	-	0.0021%	0.0068%	0.0191%	0.0176%	0.0262%	0.0233%	0.0414%	0.0111%
Hispanic ^d	Both Sexes	0.0256%	-	0.0011%	0.0046%	0.0173%	0.0409%	0.0694%	0.1006%	0.1261%	0.1694%	0.0379%
	Males	0.0188%	-	0.0012%	0.0020%	0.0102%	0.0251%	0.0507%	0.0845%	0.1385%	0.2372%	0.0341%
	Females	0.0325%	-	0.0011%	0.0075%	0.0248%	0.0574%	0.0872%	0.1143%	0.1169%	0.1290%	0.0430%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 17.13
MesotheliomaEstimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	3,004	609	314	305	182	129	4,268	4,663	+	+
	Males	2,172	324	129	83	27	69	2,708	2,829	+	+
	Females	832	285	185	222	155	60	1,560	1,834	+	+
White ^b	Both Sexes	2,767	517	266	268	179	117	3,855	4,210	+	+
	Males	2,014	278	117	70	24	57	2,479	2,572	+	+
	Females	753	239	149	198	155	60	1,376	1,638	+	+
Black ^b	Both Sexes	167	61	36	24	0	12	289	325	+	+
	Males	108	38	12	12	0	12	170	195	+	+
	Females	59	23	24	12	0	0	119	130	+	+
Asian/ Pacific Islander ^c	Both Sexes	41	20	7	+	+	+	70	+	+	+
	Males	23	10	4	+	+	+	39	+	+	+
	Females	18	10	3	+	+	+	31	+	+	+
Hispanic ^d	Both Sexes	216	52	41	+	+	+	315	+	+	+
	Males	139	34	36	+	+	+	210	+	+	+
	Females	77	18	5	+	+	+	105	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted ^f All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0011%	-	-	-	0.0002%	0.0004%	0.0013%	0.0034%	0.0063%	0.0071%	0.0011%
	Males	0.0015%	-	-	-	-	0.0004%	0.0015%	0.0047%	0.0097%	0.0140%	0.0017%
	Females	0.0008%	-	-	-	0.0004%	0.0004%	0.0012%	0.0023%	0.0035%	0.0030%	0.0007%
White ^c	Both Sexes	0.0014%	-	-	-	0.0002%	0.0005%	0.0015%	0.0040%	0.0074%	0.0075%	0.0013%
	Males	0.0018%	-	-	-	-	0.0005%	0.0017%	0.0054%	0.0116%	0.0145%	0.0019%
	Females	0.0010%	-	-	-	0.0004%	0.0005%	0.0013%	0.0026%	0.0039%	0.0033%	0.0008%
Black ^c	Both Sexes	0.0005%	-	-	-	-	-	0.0012%	-	0.0035%	0.0051%	0.0007%
	Males	0.0006%	-	-	-	-	-	-	-	-	-	0.0010%
	Females	0.0005%	-	-	-	-	-	-	-	-	-	0.0005%
Asian/ Pacific Islander ^c	Both Sexes	0.0005%	-	-	-	-	-	-	0.0017%	0.0019%	0.0041%	0.0005%
	Males	0.0006%	-	-	-	-	-	-	0.0028%	-	0.0092%	0.0007%
	Females	0.0004%	-	-	-	-	-	-	-	-	-	0.0004%
Hispanic ^d	Both Sexes	0.0006%	-	-	-	-	-	0.0017%	0.0038%	0.0041%	0.0095%	0.0011%
	Males	0.0008%	-	-	-	-	-	0.0027%	0.0059%	0.0064%	0.0166%	0.0018%
	Females	0.0004%	-	-	-	-	-	-	0.0020%	-	-	0.0006%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 18.21
Myeloma (Invasive)Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h	
Race	Sex											
All Races ^b	Both Sexes	52,505	19,437	6,635	2,874	1,182	364	81,808	83,118	249	83,367	
	Males	28,573	10,613	3,696	1,542	657	181	44,601	45,299	89	45,388	
	Females	23,932	8,824	2,939	1,332	525	183	37,207	37,819	160	37,979	
White ^b	Both Sexes	39,107	15,225	5,082	2,239	858	261	61,917	62,852	0	62,852	
	Males	22,192	8,621	3,067	1,175	478	129	35,188	35,686	0 ⁱ	35,686	
	Females	16,915	6,604	2,015	1,064	380	132	26,729	27,166	0 ⁱ	27,166	
Black ^b	Both Sexes	10,804	3,489	1,263	509	249	93	16,141	16,442	+	+	
	Males	5,039	1,630	496	300	150	52	7,504	7,679	+	+	
	Females	5,765	1,859	767	209	99	41	8,637	8,763	+	+	
Asian/ Pacific Islander ^c	Both Sexes	1,330	383	189	+	+	+	1,967	+	+	+	
	Males	697	189	90	+	+	+	1,005	+	+	+	
	Females	633	194	99	+	+	+	962	+	+	+	
Hispanic ^d	Both Sexes	3,822	1,341	501	+	+	+	5,803	+	+	+	
	Males	2,086	731	300	+	+	+	3,194	+	+	+	
	Females	1,736	610	201	+	+	+	2,609	+	+	+	

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race	Sex											
All Races ^c	Both Sexes	0.0234%	-	-	0.0003%	0.0020%	0.0115%	0.0341%	0.0798%	0.1222%	0.1082%	0.0226%
	Males	0.0258%	-	-	0.0005%	0.0024%	0.0137%	0.0386%	0.0950%	0.1467%	0.1369%	0.0271%
	Females	0.0211%	-	-	-	0.0015%	0.0093%	0.0297%	0.0661%	0.1025%	0.0913%	0.0190%
White ^c	Both Sexes	0.0229%	-	-	0.0004%	0.0018%	0.0106%	0.0311%	0.0745%	0.1167%	0.0986%	0.0211%
	Males	0.0262%	-	-	0.0006%	0.0023%	0.0133%	0.0359%	0.0918%	0.1444%	0.1331%	0.0262%
	Females	0.0196%	-	-	-	0.0013%	0.0079%	0.0263%	0.0582%	0.0938%	0.0783%	0.0168%
Black ^c	Both Sexes	0.0371%	-	-	-	0.0035%	0.0242%	0.0683%	0.1688%	0.2258%	0.2297%	0.0455%
	Males	0.0359%	-	-	-	0.0044%	0.0237%	0.0735%	0.1866%	0.2477%	0.2359%	0.0489%
	Females	0.0381%	-	-	-	0.0028%	0.0247%	0.0638%	0.1547%	0.2107%	0.2268%	0.0429%
Asian/ Pacific Islander ^c	Both Sexes	0.0134%	-	-	-	0.0010%	0.0052%	0.0206%	0.0423%	0.0744%	0.0834%	0.0136%
	Males	0.0141%	-	-	-	0.0012%	0.0073%	0.0233%	0.0475%	0.0866%	0.0898%	0.0156%
	Females	0.0128%	-	-	-	-	0.0034%	0.0184%	0.0380%	0.0651%	0.0795%	0.0120%
Hispanic ^d	Both Sexes	0.0116%	-	-	0.0006%	0.0019%	0.0099%	0.0289%	0.0637%	0.1252%	0.0873%	0.0202%
	Males	0.0125%	-	-	0.0009%	0.0023%	0.0124%	0.0347%	0.0781%	0.1410%	0.1025%	0.0238%
	Females	0.0106%	-	-	-	0.0014%	0.0073%	0.0233%	0.0514%	0.1135%	0.0783%	0.0171%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 19.25
Non-Hodgkin LymphomaEstimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h	
Race	Sex											
All Races ^b	Both Sexes	203,992	138,548	82,649	46,499	26,613	14,965	478,424	522,113	8,806	530,919	
	Males	109,031	71,416	42,880	24,393	14,018	8,043	251,155	274,206	4,630	278,836	
	Females	94,961	67,132	39,769	22,106	12,595	6,922	227,269	247,907	4,176	252,083	
White ^b	Both Sexes	176,587	121,718	73,137	41,335	23,890	13,662	418,719	458,521	9,171	467,692	
	Males	94,865	62,935	38,025	21,688	12,655	7,313	220,566	241,606	5,180	246,786	
	Females	81,722	58,783	35,112	19,647	11,235	6,349	198,153	216,915	3,991	220,906	
Black ^b	Both Sexes	16,393	11,194	5,958	3,101	1,826	801	37,209	39,725	715	40,440	
	Males	8,305	5,551	3,045	1,601	926	456	18,751	20,065	299	20,364	
	Females	8,088	5,643	2,913	1,500	900	345	18,458	19,660	416	20,076	
Asian/ Pacific Islander ^c	Both Sexes	5,757	3,689	2,092	+	+	+	12,532	+	+	+	
	Males	3,044	1,904	1,117	+	+	+	6,533	+	+	+	
	Females	2,713	1,785	975	+	+	+	5,999	+	+	+	
Hispanic ^d	Both Sexes	15,678	8,999	5,398	+	+	+	32,329	+	+	+	
	Males	8,083	4,633	2,885	+	+	+	16,751	+	+	+	
	Females	7,595	4,366	2,513	+	+	+	15,578	+	+	+	

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)								Age-Adjusted ^f		
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races ^c	Both Sexes	0.1349%	0.0032%	0.0117%	0.0244%	0.0455%	0.0974%	0.1924%	0.3624%	0.5763%	0.6536%	0.1311%
	Males	0.1439%	0.0039%	0.0156%	0.0294%	0.0537%	0.1144%	0.2225%	0.4110%	0.6473%	0.7399%	0.1504%
	Females	0.1260%	0.0025%	0.0077%	0.0191%	0.0373%	0.0806%	0.1637%	0.3184%	0.5192%	0.6031%	0.1143%
White ^c	Both Sexes	0.1516%	0.0033%	0.0121%	0.0257%	0.0477%	0.1031%	0.2075%	0.3920%	0.6328%	0.6985%	0.1412%
	Males	0.1608%	0.0042%	0.0164%	0.0311%	0.0564%	0.1210%	0.2388%	0.4395%	0.7091%	0.7841%	0.1612%
	Females	0.1423%	0.0024%	0.0076%	0.0200%	0.0385%	0.0847%	0.1766%	0.3477%	0.5699%	0.6479%	0.1232%
Black ^c	Both Sexes	0.0835%	0.0024%	0.0121%	0.0239%	0.0472%	0.0935%	0.1607%	0.2708%	0.3385%	0.3185%	0.0953%
	Males	0.0896%	0.0024%	0.0154%	0.0302%	0.0554%	0.1105%	0.1894%	0.3229%	0.3191%	0.3524%	0.1073%
	Females	0.0779%	0.0025%	0.0086%	0.0180%	0.0401%	0.0785%	0.1358%	0.2297%	0.3520%	0.3023%	0.0852%
Asian/ Pacific Islander ^c	Both Sexes	0.0828%	0.0036%	0.0090%	0.0174%	0.0299%	0.0604%	0.1147%	0.2145%	0.3575%	0.4777%	0.0835%
	Males	0.0895%	0.0039%	0.0115%	0.0199%	0.0347%	0.0680%	0.1318%	0.2465%	0.4247%	0.5732%	0.0977%
	Females	0.0768%	0.0032%	0.0064%	0.0148%	0.0257%	0.0537%	0.1001%	0.1880%	0.3059%	0.4190%	0.0724%
Hispanic ^d	Both Sexes	0.0646%	0.0032%	0.0089%	0.0184%	0.0328%	0.0729%	0.1482%	0.2770%	0.4412%	0.5261%	0.1014%
	Males	0.0657%	0.0040%	0.0115%	0.0216%	0.0390%	0.0810%	0.1665%	0.2893%	0.4502%	0.5719%	0.1098%
	Females	0.0635%	0.0025%	0.0061%	0.0148%	0.0262%	0.0645%	0.1307%	0.2664%	0.4345%	0.4989%	0.0939%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 20.27
Cancer of the Oral Cavity and Pharynx (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	104,979	63,968	41,453	25,800	16,755	11,048	240,509	272,072	9,519	281,591
	Males	73,503	43,500	26,936	16,080	9,991	6,396	162,690	181,087	5,404	186,491
	Females	31,476	20,468	14,517	9,720	6,764	4,652	77,819	90,985	4,115	95,100
White ^b	Both Sexes	90,294	55,546	35,905	22,683	14,785	9,685	208,193	235,908	8,152	244,060
	Males	63,742	38,449	23,635	14,379	9,020	5,752	142,627	159,182	4,640	163,822
	Females	26,552	17,097	12,270	8,304	5,765	3,933	65,566	76,726	3,512	80,238
Black ^b	Both Sexes	7,910	4,687	2,906	1,706	1,069	791	17,519	19,657	442	20,099
	Males	5,302	2,827	1,705	852	483	319	10,795	11,685	174	11,859
	Females	2,608	1,860	1,201	854	586	472	6,724	7,972	268	8,240
Asian/ Pacific Islander ^c	Both Sexes	3,954	2,474	1,683	+	+	+	8,925	+	+	+
	Males	2,546	1,502	991	+	+	+	5,514	+	+	+
	Females	1,408	972	692	+	+	+	3,411	+	+	+
Hispanic ^d	Both Sexes	5,606	3,052	1,946	+	+	+	11,582	+	+	+
	Males	3,509	1,912	1,179	+	+	+	7,116	+	+	+
	Females	2,097	1,140	767	+	+	+	4,466	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)								Age-Adjusted ^f		
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0679%	0.0004%	0.0022%	0.0062%	0.0160%	0.0464%	0.1283%	0.2212%	0.2572%	0.2336%	0.0639%
	Males	0.0925%	0.0004%	0.0019%	0.0062%	0.0159%	0.0588%	0.1905%	0.3395%	0.3837%	0.3440%	0.0923%
	Females	0.0438%	0.0004%	0.0025%	0.0063%	0.0162%	0.0341%	0.0688%	0.1141%	0.1554%	0.1690%	0.0397%
White ^c	Both Sexes	0.0738%	0.0005%	0.0021%	0.0060%	0.0151%	0.0448%	0.1329%	0.2369%	0.2750%	0.2456%	0.0666%
	Males	0.1015%	0.0004%	0.0017%	0.0056%	0.0145%	0.0584%	0.1993%	0.3628%	0.4092%	0.3622%	0.0967%
	Females	0.0463%	0.0005%	0.0026%	0.0065%	0.0157%	0.0308%	0.0674%	0.1193%	0.1643%	0.1767%	0.0401%
Black ^c	Both Sexes	0.0409%	-	0.0024%	0.0066%	0.0131%	0.0353%	0.1011%	0.1604%	0.1793%	0.1137%	0.0461%
	Males	0.0538%	-	0.0025%	0.0091%	0.0125%	0.0378%	0.1467%	0.2528%	0.2642%	0.1721%	0.0652%
	Females	0.0292%	-	0.0023%	0.0042%	0.0137%	0.0330%	0.0614%	0.0873%	0.1204%	0.0857%	0.0315%
Asian/ Pacific Islander ^c	Both Sexes	0.0574%	-	0.0023%	0.0065%	0.0223%	0.0596%	0.1160%	0.1591%	0.1927%	0.1906%	0.0557%
	Males	0.0741%	-	0.0028%	0.0060%	0.0256%	0.0728%	0.1616%	0.2365%	0.2790%	0.2663%	0.0765%
	Females	0.0421%	-	0.0018%	0.0069%	0.0194%	0.0479%	0.0769%	0.0949%	0.1265%	0.1441%	0.0390%
Hispanic ^d	Both Sexes	0.0231%	0.0003%	0.0014%	0.0045%	0.0094%	0.0220%	0.0640%	0.1195%	0.1631%	0.1617%	0.0368%
	Males	0.0279%	-	0.0013%	0.0041%	0.0087%	0.0251%	0.0889%	0.1673%	0.2483%	0.2078%	0.0500%
	Females	0.0182%	-	0.0015%	0.0049%	0.0101%	0.0188%	0.0403%	0.0788%	0.1003%	0.1343%	0.0261%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 21.15
Cancer of the Ovary (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>										
All Races ^b	58,762	31,186	23,873	19,562	16,098	13,217	137,024	173,192	15,675	188,867
White ^b	50,895	27,153	20,890	17,129	14,675	11,819	119,489	151,815	13,345	165,160
Black ^b	4,161	2,084	1,566	1,401	914	883	9,341	11,883	1,518	13,401
Asian/Pacific Islander ^c	2,565	1,531	1,040	+	+	+	5,734	+	+	+
Hispanic ^d	5,381	2,750	1,715	+	+	+	10,980	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races ^c	0.0788%	0.0002%	0.0065%	0.0175%	0.0321%	0.0682%	0.1477%	0.2194%	0.2370%	0.1928%	0.0713%	
White ^c	0.0870%	0.0003%	0.0068%	0.0187%	0.0328%	0.0710%	0.1557%	0.2398%	0.2609%	0.2050%	0.0762%	
Black ^c	0.0417%	-	0.0044%	0.0118%	0.0261%	0.0458%	0.0864%	0.1155%	0.1403%	0.1392%	0.0442%	
Asian/Pacific Islander ^c	0.0692%	-	0.0079%	0.0168%	0.0338%	0.0705%	0.1529%	0.1779%	0.1634%	0.1366%	0.0632%	
Hispanic ^d	0.0447%	-	0.0089%	0.0202%	0.0318%	0.0546%	0.1123%	0.1660%	0.1883%	0.1847%	0.0591%	

- ^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census.
^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.
^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.
^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
 + Not available.

Table 22.21
Cancer of the Pancreas (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h	
Race	Sex											
All Races ^b	Both Sexes	32,756	5,171	2,500	1,355	787	360	41,977	43,238	300	43,538	
	Males	16,110	2,283	1,207	504	317	114	20,199	20,679	122	20,801	
	Females	16,646	2,888	1,293	851	470	246	21,778	22,559	178	22,737	
White ^b	Both Sexes	27,338	4,477	1,874	1,064	582	284	34,934	35,878	246	36,124	
	Males	13,448	2,074	998	418	266	77	17,033	17,408	108	17,516	
	Females	13,890	2,403	876	646	316	207	17,901	18,470	138	18,608	
Black ^b	Both Sexes	3,789	469	383	142	173	38	4,791	5,039	123	5,162	
	Males	1,965	133	129	0	52	26	2,227	2,316	54	2,370	
	Females	1,824	336	254	142	121	12	2,564	2,723	69	2,792	
Asian/ Pacific Islander ^c	Both Sexes	1,139	208	127	+	+	+	1,512	+	+	+	
	Males	511	75	49	+	+	+	651	+	+	+	
	Females	628	133	78	+	+	+	861	+	+	+	
Hispanic ^d	Both Sexes	2,013	443	193	+	+	+	2,777	+	+	+	
	Males	978	156	77	+	+	+	1,238	+	+	+	
	Females	1,035	287	116	+	+	+	1,539	+	+	+	

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races ^c	Both Sexes	0.0122%	-	0.0002%	0.0008%	0.0018%	0.0065%	0.0176%	0.0392%	0.0598%	0.0588%	0.0117%
	Males	0.0119%	-	-	0.0005%	0.0013%	0.0069%	0.0192%	0.0436%	0.0615%	0.0597%	0.0123%
	Females	0.0124%	-	0.0003%	0.0011%	0.0023%	0.0060%	0.0161%	0.0353%	0.0585%	0.0582%	0.0112%
White ^c	Both Sexes	0.0127%	-	0.0002%	0.0007%	0.0018%	0.0065%	0.0172%	0.0398%	0.0601%	0.0579%	0.0117%
	Males	0.0126%	-	-	0.0004%	0.0012%	0.0070%	0.0195%	0.0444%	0.0633%	0.0593%	0.0125%
	Females	0.0128%	-	-	0.0010%	0.0024%	0.0060%	0.0149%	0.0355%	0.0574%	0.0570%	0.0110%
Black ^c	Both Sexes	0.0109%	-	-	0.0009%	0.0021%	0.0081%	0.0229%	0.0442%	0.0610%	0.0568%	0.0130%
	Males	0.0104%	-	-	-	0.0016%	0.0096%	0.0250%	0.0461%	0.0615%	0.0540%	0.0135%
	Females	0.0114%	-	-	-	0.0026%	0.0068%	0.0212%	0.0426%	0.0606%	0.0581%	0.0126%
Asian/ Pacific Islander ^c	Both Sexes	0.0103%	-	-	0.0009%	0.0015%	0.0045%	0.0156%	0.0309%	0.0558%	0.0612%	0.0104%
	Males	0.0091%	-	-	-	0.0017%	0.0038%	0.0135%	0.0360%	0.0519%	0.0583%	0.0100%
	Females	0.0113%	-	-	0.0016%	0.0013%	0.0051%	0.0175%	0.0268%	0.0588%	0.0629%	0.0107%
Hispanic ^d	Both Sexes	0.0056%	-	-	0.0007%	0.0013%	0.0048%	0.0122%	0.0306%	0.0441%	0.0674%	0.0096%
	Males	0.0049%	-	-	-	-	0.0053%	0.0133%	0.0291%	0.0486%	0.0521%	0.0093%
	Females	0.0063%	-	-	0.0009%	0.0022%	0.0043%	0.0111%	0.0318%	0.0408%	0.0765%	0.0098%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.
^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.
^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
+ Not available.

Table 23.15
Cancer of the Prostate (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>										
All Races ^b	1,047,674	793,365	518,837	274,354	57,736	11,316	2,655,003	2,706,583	1,238	2,707,821
White ^b	842,834	661,840	439,478	239,979	52,556	10,463	2,203,073	2,250,165	1,432	2,251,597
Black ^b	147,861	103,144	63,749	28,209	4,112	753	344,435	348,058	120	348,178
Asian/Pacific Islander ^c	20,777	14,755	7,797	+	+	+	46,251	+	+	+
Hispanic ^d	58,787	42,509	23,815	+	+	+	134,478	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f All Ages
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races ^c	1.4779%	-	-	0.0003%	0.0012%	0.0848%	1.0293%	4.9999%	11.8211%	14.5189%	1.6885%
White ^c	1.5645%	-	-	0.0004%	0.0010%	0.0728%	0.9891%	4.9616%	11.9421%	14.3476%	1.6793%
Black ^c	1.6401%	-	-	-	0.0034%	0.2242%	1.9728%	8.4530%	17.5682%	20.0518%	2.6097%
Asian/Pacific Islander ^c	0.7034%	-	-	-	-	0.0224%	0.3432%	2.0078%	5.8919%	9.7833%	0.8715%
Hispanic ^d	0.5263%	-	-	-	0.0006%	0.0396%	0.6144%	3.3964%	9.4474%	12.7016%	1.3237%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 24.21
Cancer of the Stomach (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	36,380	15,961	8,793	5,341	3,445	1,880	67,193	72,734	1,301	74,035
	Males	21,307	9,112	4,940	2,857	1,695	928	38,531	41,394	733	42,127
	Females	15,073	6,849	3,853	2,484	1,750	952	28,662	31,340	568	31,908
White ^b	Both Sexes	27,127	11,759	6,308	4,008	2,583	1,398	49,735	53,965	1,047	55,012
	Males	16,656	6,894	3,597	2,015	1,269	737	29,399	31,649	570	32,219
	Females	10,471	4,865	2,711	1,993	1,314	661	20,336	22,316	477	22,793
Black ^b	Both Sexes	5,548	2,575	1,418	632	426	304	10,246	10,958	178	11,136
	Males	2,627	1,312	691	461	203	136	5,116	5,472	101	5,573
	Females	2,921	1,263	727	171	223	168	5,130	5,486	77	5,563
Asian/ Pacific Islander ^c	Both Sexes	3,276	1,630	990	+	+	+	6,415	+	+	+
	Males	1,819	884	575	+	+	+	3,578	+	+	+
	Females	1,457	746	415	+	+	+	2,837	+	+	+
Hispanic ^d	Both Sexes	5,404	2,059	1,152	+	+	+	9,196	+	+	+
	Males	3,052	1,141	571	+	+	+	4,997	+	+	+
	Females	2,352	918	581	+	+	+	4,199	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted ^f All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0240%	-	0.0001%	0.0005%	0.0033%	0.0118%	0.0297%	0.0641%	0.1253%	0.1711%	0.0235%
	Males	0.0277%	-	-	0.0005%	0.0033%	0.0125%	0.0365%	0.0833%	0.1684%	0.2281%	0.0303%
	Females	0.0204%	-	-	0.0006%	0.0034%	0.0111%	0.0232%	0.0467%	0.0906%	0.1376%	0.0182%
White ^c	Both Sexes	0.0204%	-	-	0.0006%	0.0030%	0.0104%	0.0247%	0.0526%	0.1012%	0.1261%	0.0189%
	Males	0.0241%	-	-	0.0005%	0.0031%	0.0113%	0.0307%	0.0710%	0.1386%	0.1673%	0.0247%
	Females	0.0167%	-	-	0.0007%	0.0029%	0.0095%	0.0188%	0.0355%	0.0703%	0.1018%	0.0142%
Black ^c	Both Sexes	0.0228%	-	-	-	0.0039%	0.0138%	0.0436%	0.0806%	0.1494%	0.1924%	0.0288%
	Males	0.0238%	-	-	-	0.0035%	0.0145%	0.0493%	0.0974%	0.1965%	0.2128%	0.0344%
	Females	0.0220%	-	-	-	0.0042%	0.0132%	0.0387%	0.0674%	0.1166%	0.1827%	0.0250%
Asian/ Pacific Islander ^c	Both Sexes	0.0455%	-	-	0.0006%	0.0045%	0.0174%	0.0463%	0.1172%	0.2500%	0.4618%	0.0475%
	Males	0.0528%	-	-	-	0.0039%	0.0173%	0.0609%	0.1474%	0.3308%	0.6405%	0.0622%
	Females	0.0388%	-	-	-	0.0050%	0.0175%	0.0338%	0.0922%	0.1879%	0.3519%	0.0369%
Hispanic ^d	Both Sexes	0.0184%	-	-	0.0008%	0.0050%	0.0152%	0.0392%	0.0871%	0.1849%	0.2411%	0.0332%
	Males	0.0196%	-	-	0.0010%	0.0054%	0.0147%	0.0432%	0.1116%	0.2482%	0.2914%	0.0411%
	Females	0.0172%	-	-	-	0.0045%	0.0158%	0.0355%	0.0663%	0.1382%	0.2112%	0.0273%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 25.15
Cancer of the Testis (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>										
All Races ^b	42,114	39,198	36,255	31,088	28,044	20,211	154,737	212,079	15,327	227,406
White ^b	38,604	37,033	34,094	29,566	27,142	19,521	145,211	200,534	14,651	215,185
Black ^b	1,309	889	944	594	320	325	3,794	4,697	166	4,863
Asian/Pacific Islander ^c	870	649	544	+	+	+	2,439	+	+	+
Hispanic ^d	6,416	4,519	3,407	+	+	+	16,352	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races ^c	0.0885%	0.0026%	0.0073%	0.0704%	0.1660%	0.2005%	0.1368%	0.0585%	0.0238%	0.0115%	0.0881%	
White ^c	0.1090%	0.0026%	0.0086%	0.0870%	0.2078%	0.2481%	0.1644%	0.0693%	0.0281%	0.0132%	0.1084%	
Black ^c	0.0187%	0.0016%	0.0025%	0.0138%	0.0315%	0.0416%	0.0378%	0.0167%	-	-	0.0193%	
Asian/Pacific Islander ^c	0.0293%	0.0042%	0.0042%	0.0269%	0.0497%	0.0579%	0.0455%	0.0192%	0.0077%	-	0.0282%	
Hispanic ^d	0.0638%	0.0029%	0.0102%	0.0750%	0.1429%	0.1291%	0.0731%	0.0280%	0.0110%	-	0.0642%	

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 26.21
Cancer of the Thyroid (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	174,786	117,871	76,909	54,925	41,306	31,258	433,781	527,919	38,789	566,708
	Males	38,665	25,267	16,501	12,249	8,454	6,424	94,464	114,267	8,821	123,088
	Females	136,121	92,604	60,408	42,676	32,852	24,834	339,317	413,652	29,968	443,620
White ^b	Both Sexes	150,376	103,251	67,741	48,798	37,221	28,251	378,604	463,565	35,135	498,700
	Males	34,346	22,675	14,957	11,190	7,774	5,810	84,812	102,959	7,429	110,388
	Females	116,030	80,576	52,784	37,608	29,447	22,441	293,792	360,606	27,706	388,312
Black ^b	Both Sexes	11,424	7,203	4,276	2,862	1,830	1,396	26,148	30,404	1,637	32,041
	Males	1,844	1,295	658	550	342	275	4,418	5,118	178	5,296
	Females	9,580	5,908	3,618	2,312	1,488	1,121	21,730	25,286	1,459	26,745
Asian/ Pacific Islander ^c	Both Sexes	8,426	5,356	3,410	+	+	+	19,246	+	+	+
	Males	1,551	917	610	+	+	+	3,417	+	+	+
	Females	6,875	4,439	2,800	+	+	+	15,829	+	+	+
Hispanic ^d	Both Sexes	17,678	11,227	6,911	+	+	+	39,901	+	+	+
	Males	2,946	1,951	1,112	+	+	+	6,642	+	+	+
	Females	14,732	9,276	5,799	+	+	+	33,259	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.1268%	0.0001%	0.0052%	0.0394%	0.1266%	0.2103%	0.2486%	0.2560%	0.2290%	0.1536%	0.1226%
	Males	0.0549%	-	0.0022%	0.0114%	0.0409%	0.0790%	0.1098%	0.1402%	0.1454%	0.1011%	0.0546%
	Females	0.1968%	-	0.0084%	0.0686%	0.2126%	0.3404%	0.3812%	0.3608%	0.2962%	0.1843%	0.1875%
White ^c	Both Sexes	0.1387%	0.0002%	0.0059%	0.0453%	0.1418%	0.2317%	0.2673%	0.2654%	0.2351%	0.1512%	0.1322%
	Males	0.0619%	-	0.0025%	0.0133%	0.0448%	0.0888%	0.1208%	0.1511%	0.1573%	0.1011%	0.0597%
	Females	0.2153%	-	0.0096%	0.0794%	0.2438%	0.3790%	0.4119%	0.3720%	0.2993%	0.1808%	0.2042%
Black ^c	Both Sexes	0.0604%	-	0.0016%	0.0123%	0.0544%	0.1052%	0.1287%	0.1725%	0.1483%	0.1152%	0.0658%
	Males	0.0213%	-	-	0.0031%	0.0152%	0.0350%	0.0447%	0.0723%	0.0760%	0.0860%	0.0258%
	Females	0.0961%	-	0.0023%	0.0210%	0.0885%	0.1674%	0.2017%	0.2518%	0.1985%	0.1292%	0.0991%
Asian/ Pacific Islander ^c	Both Sexes	0.1181%	-	0.0046%	0.0305%	0.1089%	0.1814%	0.2323%	0.2521%	0.2373%	0.1841%	0.1133%
	Males	0.0444%	-	0.0020%	0.0070%	0.0388%	0.0606%	0.0902%	0.1147%	0.1141%	0.1000%	0.0450%
	Females	0.1857%	-	0.0073%	0.0536%	0.1717%	0.2887%	0.3542%	0.3660%	0.3318%	0.2358%	0.1717%
Hispanic ^d	Both Sexes	0.0799%	-	0.0056%	0.0314%	0.0955%	0.1571%	0.2101%	0.2286%	0.2277%	0.1696%	0.1030%
	Males	0.0261%	-	0.0022%	0.0065%	0.0254%	0.0437%	0.0760%	0.1052%	0.1214%	0.0998%	0.0383%
	Females	0.1350%	-	0.0091%	0.0592%	0.1703%	0.2752%	0.3377%	0.3337%	0.3062%	0.2112%	0.1647%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 27.21
Cancer of the Urinary Bladder (Invasive and *In Situ*)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	200,849	136,192	91,719	59,753	36,263	22,082	497,175	560,445	11,073	571,518
	Males	152,753	101,789	67,760	44,341	26,750	15,913	372,925	418,562	7,160	425,722
	Females	48,096	34,403	23,959	15,412	9,513	6,169	124,250	141,883	3,913	145,796
White ^b	Both Sexes	183,533	126,659	86,005	56,476	34,544	21,106	460,827	521,241	8,660	529,901
	Males	140,477	95,090	63,718	41,827	25,429	15,192	347,031	390,556	5,497	396,053
	Females	43,056	31,569	22,287	14,649	9,115	5,914	113,796	130,685	3,163	133,848
Black ^b	Both Sexes	8,979	5,401	3,138	2,109	1,086	593	19,928	21,611	188	21,799
	Males	6,151	3,532	2,124	1,611	882	414	13,620	14,898	153	15,051
	Females	2,828	1,869	1,014	498	204	179	6,308	6,713	35	6,748
Asian/ Pacific Islander ^c	Both Sexes	3,647	2,174	1,216	+	+	+	7,652	+	+	+
	Males	2,807	1,660	870	+	+	+	5,813	+	+	+
	Females	840	514	346	+	+	+	1,839	+	+	+
Hispanic ^d	Both Sexes	7,543	4,539	2,837	+	+	+	16,403	+	+	+
	Males	5,620	3,416	2,101	+	+	+	12,204	+	+	+
	Females	1,923	1,123	736	+	+	+	4,199	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.1314%	0.0002%	0.0003%	0.0014%	0.0072%	0.0277%	0.1096%	0.3573%	0.7994%	1.1659%	0.1305%
	Males	0.2008%	0.0002%	0.0004%	0.0020%	0.0104%	0.0413%	0.1725%	0.5822%	1.3775%	2.2784%	0.2289%
	Females	0.0638%	0.0002%	0.0002%	0.0009%	0.0039%	0.0142%	0.0495%	0.1540%	0.3346%	0.5146%	0.0566%
White ^c	Both Sexes	0.1586%	0.0002%	0.0003%	0.0017%	0.0086%	0.0321%	0.1253%	0.4112%	0.9219%	1.2834%	0.1482%
	Males	0.2407%	-	0.0005%	0.0023%	0.0120%	0.0465%	0.1941%	0.6611%	1.5724%	2.5134%	0.2575%
	Females	0.0770%	0.0003%	-	0.0010%	0.0049%	0.0172%	0.0574%	0.1779%	0.3857%	0.5568%	0.0643%
Black ^c	Both Sexes	0.0471%	-	-	-	0.0038%	0.0157%	0.0625%	0.1803%	0.3628%	0.5344%	0.0629%
	Males	0.0683%	-	-	-	0.0070%	0.0260%	0.1014%	0.3022%	0.6038%	0.9550%	0.1071%
	Females	0.0278%	-	-	-	-	0.0065%	0.0288%	0.0840%	0.1955%	0.3332%	0.0335%
Asian/ Pacific Islander ^c	Both Sexes	0.0555%	-	-	-	0.0029%	0.0110%	0.0453%	0.1371%	0.3287%	0.6535%	0.0587%
	Males	0.0875%	-	-	-	0.0045%	0.0182%	0.0768%	0.2296%	0.5965%	1.2222%	0.1057%
	Females	0.0262%	-	-	-	0.0015%	0.0045%	0.0184%	0.0604%	0.1230%	0.3041%	0.0248%
Hispanic ^d	Both Sexes	0.0328%	-	-	0.0012%	0.0037%	0.0138%	0.0506%	0.1651%	0.3774%	0.6271%	0.0644%
	Males	0.0479%	-	-	0.0015%	0.0046%	0.0182%	0.0780%	0.2669%	0.6830%	1.2150%	0.1144%
	Females	0.0172%	-	-	0.0008%	0.0027%	0.0092%	0.0246%	0.0783%	0.1520%	0.2775%	0.0289%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 28.12
Childhood Cancer (Invasive, <20 Years old at diagnosis)

Estimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^f	Complete ^g
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	63,918	56,496	49,430	44,260	37,272	31,407	221,734	314,920	+	+
	Males	33,711	30,073	25,682	22,844	19,261	15,625	116,561	162,108	+	+
	Females	30,207	26,423	23,748	21,416	18,011	15,782	105,173	152,812	+	+
White ^b	Both Sexes	51,221	47,428	41,632	37,233	32,076	27,599	184,028	265,647	+	+
	Males	27,178	25,326	21,576	19,294	16,662	13,852	97,071	137,293	+	+
	Females	24,043	22,102	20,056	17,939	15,414	13,747	86,957	128,354	+	+
Black ^b	Both Sexes	7,930	5,674	4,812	4,506	3,300	2,583	23,560	31,387	+	+
	Males	4,094	2,923	2,498	2,250	1,573	1,197	12,070	15,552	+	+
	Females	3,836	2,751	2,314	2,256	1,727	1,386	11,490	15,835	+	+
Asian/ Pacific Islander ^c	Both Sexes	2,726	2,130	1,787	+	+	+	7,880	+	+	+
	Males	1,429	1,128	974	+	+	+	4,136	+	+	+
	Females	1,297	1,002	813	+	+	+	3,744	+	+	+
Hispanic ^d	Both Sexes	13,347	10,204	7,990	+	+	+	36,739	+	+	+
	Males	7,353	5,580	4,167	+	+	+	19,904	+	+	+
	Females	5,994	4,624	3,823	+	+	+	16,835	+	+	+

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census.

^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^{f g} (f) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <36 and >=36.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table 29.7
U.S. Childhood Cancer Survivors at January 1, 2011
Number of People Previously Diagnosed with Cancer as Children (Ages 0-19 Years) in the United States and Alive January 1, 2011
By International Classification of Childhood Cancer (ICCC)^a Selected Group and Subgroup, Age Group, and Sex

Site	Sex	Complete Prevalence Counts by Age at Prevalence							All Ages		
		0-19	20-29	30-39	40-49	50-59	60+	All Ages	Diagnosed between 1975-2010	Diagnosed before 1975	% ^b
									Count	Count	% ^b
All Sites	Both	114,469	96,086	63,670	56,072	39,055	19,149	388,501	314,744	73,757	19
	Male	61,300	50,413	31,574	25,285	16,475	9,976	195,023	162,249	32,774	17
	Female ^c	53,169	45,673	32,096	30,787	22,580	9,173	193,478	152,495	40,983	21
I Leukemia	Both	35,982	20,750	11,794	5,282	1,659	210	75,677	71,408	4,269	6
	Male	19,822	10,941	6,006	2,737	766	92	40,364	38,440	1,924	5
	Female	16,160	9,809	5,788	2,545	893	118	35,313	32,968	2,345	7
Acute lymphocytic leukemia ^d	Both	30,395	16,935	9,972	4,732	564	3	62,601	59,336	3,265	5
	Male	16,803	9,125	5,220	2,611	308	2	34,069	32,373	1,696	5
	Female	13,592	7,810	4,752	2,121	256	1	28,532	26,963	1,569	5
I(b) Acute myeloid leukemia	Both	3,992	2,528	1,185	690	245	4	8,644	8,145	499	6
	Male	2,075	1,183	498	321	137	3	4,217	3,952	265	6
	Female	1,917	1,345	687	369	108	1	4,427	4,193	234	5
II(a) Hodgkin lymphoma	Both	4,863	8,832	8,058	7,834	5,056	1,353	35,996	30,572	5,424	15
	Male	2,693	4,544	4,009	3,829	2,504	702	18,281	15,371	2,910	16
	Female	2,170	4,288	4,049	4,005	2,552	651	17,715	15,201	2,514	14
II(b,c,e) Non-Hodgkin lymphoma	Both	6,654	6,888	4,353	2,908	1,759	1,146	23,708	20,267	3,441	15
	Male	4,405	4,445	3,060	1,852	937	672	15,371	13,506	1,865	12
	Female	2,249	2,443	1,293	1,056	822	474	8,337	6,761	1,576	19
III CNS and misc intracranial & intraspinal neoplasms	Both	20,913	15,272	9,006	8,615	5,044	1,690	60,540	49,544	10,996	18
	Male	11,275	8,482	4,792	4,318	2,269	751	31,887	26,876	5,011	16
	Female	9,638	6,790	4,214	4,297	2,775	939	28,653	22,668	5,985	21
IV Neuroblastoma & other peripheral nervous cell tumor	Both	9,632	3,883	2,119	1,741	1,446	830	19,651	15,660	3,991	20
	Male	4,936	2,015	988	588	438	216	9,181	7,929	1,252	14
	Female	4,696	1,868	1,131	1,153	1,008	614	10,470	7,731	2,739	26
VI Renal tumors	Both	7,787	4,702	3,809	4,280	2,554	858	23,990	16,086	7,904	33
	Male	3,862	2,174	1,729	2,403	1,379	500	12,047	7,574	4,473	37
	Female	3,925	2,528	2,080	1,877	1,175	358	11,943	8,512	3,431	29
VIII Malignant bone tumors	Both	3,567	4,417	3,167	2,568	2,417	907	17,043	13,473	3,570	21
	Male	2,066	2,569	1,787	1,290	1,053	472	9,237	7,592	1,645	18
	Female	1,501	1,848	1,380	1,278	1,364	435	7,806	5,881	1,925	25
VIII(a) Osteosarcoma	Both	1,859	2,253	1,777	1,293	1,237	829	9,248	7,125	2,123	23
	Male	1,002	1,265	1,014	657	499	396	4,833	3,868	965	20
	Female	857	988	763	636	738	433	4,415	3,257	1,158	26
VIII(c) Ewing tumor & related sarcomas of bone	Both	1,347	1,467	972	728	358	148	5,020	4,420	600	12
	Male	819	872	487	336	197	43	2,754	2,506	248	9
	Female	528	595	485	392	161	105	2,266	1,914	352	16
IX Soft tissue & other extraosseous sarcomas	Both	6,799	7,126	4,166	4,913	4,267	4,670	31,941	21,477	10,464	33
	Male	3,724	3,812	2,131	2,424	1,995	2,608	16,694	11,350	5,344	32
	Female	3,075	3,314	2,035	2,489	2,272	2,062	15,247	10,127	5,120	34
X Germ cell & trophoblastic tumors & neoplasms of gonads	Both	5,233	7,869	5,634	6,245	6,641	6,817	38,439	24,486	13,953	36
	Male	2,749	5,143	3,461	3,322	3,301	3,136	21,112	14,507	6,605	31
	Female	2,484	2,726	2,173	2,923	3,340	3,681	17,327	9,979	7,348	42

^a International Classification of Childhood Cancer is based on ICD-O-3. Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of Childhood Cancer, Third Edition. Cancer. April 1, 2005; Vol 103, No. 7, pg 1457-1467.

^b Percent diagnosed before 1975 of all ages complete prevalence counts.

^c All sites females excludes breast cancer.

^d SEER site recode Acute Lymphocytic Leukemia was used instead of the ICCC classification lymphocytic leukemia (Ia) because of statistical modeling requirements.

For more information regarding the modeling of the prevalence estimates, see Mariotto et al. Long-Term Survivors of Childhood Cancers in the United States. Cancer Epidemiol Biomarkers Prev 2009, page 1034.