Table 5.9 Cancer of the Cervix Uteri (Invasive)

SEER^a Relative Survival (Percent) By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Tim	e																			
1-year	86.6	87.2	87.1	87.7	89.1	89.0	89.3	88.1	89.6	88.3	85.7	87.9	86.6	86.3	87.2	86.2	88.4	87.0	88.8	87.4
2-year	77.8	77.2	77.9	79.5	82.2	81.8	80.0	81.3	83.0	81.1	79.2	79.6	79.0	77.1	77.3	78.4	78.7	79.0	79.4	
3-year	73.2	72.7	73.2	75.1	78.8	77.9	75.6	76.7	78.4	76.4	75.5	74.9	75.7	73.1	72.3	75.5	74.6	75.1		
4-year	70.3	69.6	70.4	72.2	76.0	74.7	73.1	74.0	76.1	74.2	72.5	72.5	72.3	69.7	69.8	73.3	72.7			
5-year	68.3	67.6	68.6	70.3	74.9	73.5	71.6	72.5	74.4	72.1	70.5	69.8	70.1	67.3	68.0	71.5				
б-year	67.1	65.7	67.3	68.9	73.6	72.8	70.0	70.9	73.4	71.0	69.2	69.0	68.5	65.8	67.0					
7-year	66.0	64.1	66.1	68.0	72.4	71.6	69.5	69.7	72.9	70.2	67.8	68.1	67.0	64.9						
8-year	65.0	62.7	65.0	67.1	71.7	70.7	69.1	69.4	72.5	69.9	67.3	67.6	65.7							
9-year	64.3	62.1	64.3	66.2	70.7	69.2	67.5	68.8	72.1	68.8	66.8	67.2								
10-year	63.1	61.4	63.7	65.5	69.9	68.7	66.9	67.7	70.7	68.2	66.5									
11-year	62.4	60.7	63.0	64.9	69.6	68.1	65.9	67.1	69.1	67.7										
12-year	61.7	59.9	62.2	64.3	68.5	66.8	65.6	66.1	68.3											
13-year	61.1	59.4	61.6	63.8	67.0	66.2	65.3	64.7												
14-year	60.5	58.6	61.0	63.2	66.4	65.3	65.0													
15-year	59.8	57.8	60.3	62.6	66.1	64.5														
16-year	59.2	57.0	59.6	61.9	65.2															
17-year	58.6	56.1	58.8	61.4																
18-year	58.1	55.3	58.7																	
19-year	57.3	54.7	58.2																	
20-year	56.5	54.0	57.2																	

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

а