

Table 7.5
Cancer of the Corpus and Uterus, NOS (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Age

| | All Races, Females | | | White Females | | | Black Females | | |
|--------------------|--------------------|------|--------|---------------|------|--------|---------------|------|-------|
| | All | <50 | 50+ | All | <50 | 50+ | All | <50 | 50+ |
| Year of Diagnosis: | | | | | | | | | |
| 1975-2011 | 26.28 | 5.20 | 81.48 | 27.52 | 5.18 | 86.01 | 19.83 | 3.28 | 63.17 |
| 1975 | 35.47 | 6.48 | 111.37 | 37.32 | 6.81 | 117.22 | 19.91 | 3.64 | 62.51 |
| 1976 | 34.37 | 6.20 | 108.12 | 36.25 | 6.28 | 114.71 | 17.61 | 3.29 | 55.11 |
| 1977 | 31.68 | 5.75 | 99.58 | 33.17 | 5.91 | 104.56 | 20.06 | 3.52 | 63.37 |
| 1978 | 30.05 | 6.10 | 92.76 | 31.66 | 6.31 | 98.05 | 18.74 | 3.94 | 57.49 |
| 1979 | 27.90 | 5.31 | 87.06 | 29.25 | 5.32 | 91.90 | 16.96 | 3.94 | 51.06 |
| 1980 | 27.31 | 4.97 | 85.81 | 28.52 | 4.95 | 90.25 | 16.22 | - | 52.13 |
| 1981 | 27.28 | 5.62 | 83.98 | 28.34 | 5.45 | 88.26 | 17.42 | 2.86 | 55.52 |
| 1982 | 26.80 | 5.08 | 83.70 | 28.23 | 5.24 | 88.44 | 17.09 | 2.49 | 55.33 |
| 1983 | 26.51 | 4.86 | 83.21 | 27.78 | 4.91 | 87.66 | 18.81 | 3.37 | 59.25 |
| 1984 | 25.71 | 4.62 | 80.92 | 27.16 | 4.89 | 85.47 | 17.38 | - | 58.35 |
| 1985 | 25.24 | 4.63 | 79.20 | 26.56 | 4.83 | 83.46 | 17.41 | 2.23 | 57.17 |
| 1986 | 24.40 | 4.82 | 75.65 | 25.63 | 4.99 | 79.68 | 16.78 | 2.54 | 54.09 |
| 1987 | 24.71 | 5.00 | 76.33 | 26.12 | 5.28 | 80.68 | 15.77 | - | 51.79 |
| 1988 | 23.58 | 4.54 | 73.43 | 24.59 | 4.46 | 77.32 | 16.97 | 2.97 | 53.65 |
| 1989 | 24.43 | 4.49 | 76.66 | 25.50 | 4.78 | 79.77 | 19.36 | 1.78 | 65.40 |
| 1990 | 25.06 | 4.62 | 78.57 | 26.53 | 4.82 | 83.36 | 17.03 | 2.28 | 55.65 |
| 1991 | 24.64 | 4.37 | 77.70 | 25.99 | 4.48 | 82.34 | 17.48 | 3.12 | 55.09 |
| 1992 | 24.80 | 4.86 | 77.01 | 26.18 | 4.87 | 81.97 | 16.94 | 2.88 | 53.74 |
| 1993 | 24.29 | 5.19 | 74.31 | 25.68 | 5.14 | 79.45 | 16.97 | 3.48 | 52.27 |
| 1994 | 25.15 | 5.03 | 77.81 | 26.34 | 5.04 | 82.11 | 17.61 | 2.51 | 57.17 |
| 1995 | 25.33 | 4.71 | 79.33 | 26.50 | 4.63 | 83.78 | 18.82 | 2.75 | 60.90 |
| 1996 | 24.93 | 4.52 | 78.38 | 25.97 | 4.38 | 82.53 | 19.40 | 3.17 | 61.89 |
| 1997 | 25.89 | 5.05 | 80.46 | 27.35 | 5.04 | 85.78 | 16.47 | 2.46 | 53.15 |
| 1998 | 25.74 | 4.84 | 80.45 | 27.10 | 4.82 | 85.44 | 19.37 | 3.07 | 62.04 |
| 1999 | 25.33 | 5.19 | 78.08 | 26.79 | 5.06 | 83.66 | 17.70 | 2.97 | 56.27 |
| 2000 | 24.83 | 5.23 | 76.16 | 26.22 | 5.37 | 80.82 | 17.93 | 3.00 | 57.02 |
| 2001 | 25.29 | 5.12 | 78.10 | 26.73 | 4.98 | 83.69 | 19.34 | 4.07 | 59.34 |
| 2002 | 24.92 | 5.06 | 76.93 | 25.33 | 4.58 | 79.65 | 23.29 | 4.79 | 71.74 |
| 2003 | 24.36 | 5.31 | 74.25 | 25.54 | 5.30 | 78.57 | 20.05 | 3.68 | 62.90 |
| 2004 | 25.08 | 5.56 | 76.22 | 26.06 | 5.26 | 80.52 | 20.59 | 3.85 | 64.42 |
| 2005 | 24.97 | 5.55 | 75.83 | 25.84 | 5.40 | 79.38 | 23.12 | 4.12 | 72.87 |
| 2006 | 24.62 | 5.15 | 75.59 | 26.01 | 5.06 | 80.86 | 19.32 | 3.70 | 60.21 |
| 2007 | 25.26 | 5.84 | 76.09 | 26.08 | 5.89 | 78.96 | 22.42 | 3.51 | 71.94 |
| 2008 | 26.08 | 6.01 | 78.66 | 26.65 | 5.75 | 81.39 | 24.26 | 4.30 | 76.50 |
| 2009 | 27.35 | 5.99 | 83.30 | 27.97 | 5.92 | 85.72 | 25.16 | 3.72 | 81.32 |
| 2010 | 27.36 | 6.25 | 82.62 | 27.95 | 6.01 | 85.41 | 24.68 | 3.89 | 79.14 |
| 2011 | 27.05 | 5.54 | 83.37 | 27.22 | 5.30 | 84.61 | 28.27 | 4.00 | 91.81 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.