

Table 17.5
MesotheliomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time																				
1-year	34.8	37.7	35.7	35.9	35.8	41.9	35.3	37.4	31.2	40.5	40.5	31.7	46.6	39.6	41.2	44.2	42.1	44.1	44.8	33.7
2-year	15.9	18.7	16.5	15.7	13.8	25.1	16.9	21.2	17.9	14.6	24.0	15.6	22.3	16.6	20.5	28.1	18.2	19.2	25.3	
3-year	10.8	11.7	10.8	10.9	11.8	17.5	10.3	14.8	12.0	10.5	13.9	6.9	16.8	10.4	13.2	19.2	10.7	12.7		
4-year	9.2	8.4	8.0	8.2	8.9	11.4	7.8	12.6	9.7	7.1	11.4	4.7	13.5	8.5	10.2	14.4	6.2			
5-year	7.9	7.0	6.6	7.1	7.1	10.2	5.7	12.6	7.3	3.1	10.9	4.2	9.0	6.1	9.2	12.1				
6-year	6.7	5.7	5.2	6.0	6.1	10.0	4.6	11.4	6.9	1.3	9.9	3.7	8.0	6.1	9.2					
7-year	6.0	5.1	4.5	5.6	5.1	8.6	3.0	10.6	6.5	1.3	7.7	3.2	7.0	4.4						
8-year	5.6	4.8	4.0	5.0	5.1	6.5	3.0	8.5	6.5	0.7	7.7	3.2	5.7							
9-year	5.3	4.6	3.8	4.7	3.1	4.9	2.5	8.5	5.1	0.7	7.7	3.2								
10-year	5.2	4.3	3.4	4.3	3.1	4.4	2.5	7.7	5.1	0.7	6.7									
11-year	5.0	4.3	3.4	4.2	3.1	4.0	1.9	7.4	4.6	0.7										
12-year	4.9	4.1	3.4	3.8	2.5	3.5	1.9	5.6	4.6											
13-year	4.7	4.0	3.3	3.6	2.5	3.0	1.9	5.6												
14-year	4.7	4.0	3.3	3.6	2.5	3.0	1.9													
15-year	4.3	3.5	3.2	3.6	2.5	3.0														
16-year	4.1	3.5	3.2	3.6	2.5															
17-year	4.0	3.5	3.2	3.6																
18-year	4.0	3.5	3.2																	
19-year	4.0	3.5	3.2																	
20-year	4.0	3.5	3.1																	

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).