

Table 22.9
Cancer of the Pancreas (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Survival Time																					
1-year	14.2	15.9	17.0	18.5	17.9	19.0	19.9	19.5	22.3	21.8	23.8	24.9	22.8	23.1	26.7	25.8	28.0	29.7	29.6	30.3	31.2
2-year	5.9	6.2	6.8	8.6	8.2	9.3	9.9	8.7	10.5	10.9	11.4	11.9	11.3	11.8	12.9	13.1	13.8	15.0	15.6	16.6	
3-year	3.6	4.1	4.7	5.8	5.2	6.3	7.1	5.7	6.5	7.3	7.3	8.4	7.4	8.3	8.9	9.2	9.7	10.4	11.5		
4-year	3.0	3.3	3.8	4.8	3.9	4.8	5.6	4.5	5.6	6.1	6.1	6.8	5.6	6.4	7.1	7.9	8.2	8.4			
5-year	2.5	2.8	3.3	4.2	3.6	4.2	5.0	3.8	5.1	5.1	5.0	6.1	4.8	5.5	6.2	7.3	7.6				
6-year	2.3	2.5	2.9	4.0	3.2	3.8	4.5	3.7	4.6	4.6	4.7	5.7	4.4	4.9	5.7	6.7					
7-year	2.1	2.3	2.7	3.6	3.1	3.5	4.3	3.4	4.3	4.3	4.1	5.0	4.1	4.6	5.3						
8-year	1.9	2.1	2.5	3.4	2.7	3.4	4.1	3.1	3.9	3.8	3.9	4.7	3.9	4.3							
9-year	1.8	2.0	2.2	3.2	2.6	3.3	4.0	2.8	3.8	3.7	3.8	4.6	3.6								
10-year	1.8	1.9	2.2	3.0	2.6	3.2	3.9	2.7	3.6	3.6	3.4	4.4									
11-year	1.7	1.7	2.1	2.8	2.5	2.9	3.8	2.5	3.3	3.5	3.3										
12-year	1.7	1.7	2.1	2.7	2.3	2.8	3.6	2.4	3.1	3.1											
13-year	1.6	1.6	2.0	2.6	2.2	2.7	3.3	2.3	3.1												
14-year	1.6	1.6	1.9	2.5	1.8	2.7	3.1	1.9													
15-year	1.5	1.5	1.8	2.4	1.7	2.7	3.0														
16-year	1.5	1.4	1.7	2.2	1.7	2.2															
17-year	1.4	1.4	1.7	2.2	1.4																
18-year	1.3	1.3	1.7	2.2																	
19-year	1.3	1.3	1.6																		
20-year	1.2	1.3	1.6																		

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).