

Table 22.10
Cancer of the Pancreas (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Survival Time																					
1-year	14.1	15.0	16.7	17.8	17.0	19.9	21.1	19.3	23.0	21.9	23.3	24.8	24.3	25.1	25.7	27.2	28.3	31.1	29.2	29.8	32.4
2-year	5.8	5.8	6.2	8.5	7.2	10.8	9.9	9.2	9.8	11.3	12.1	11.1	11.3	12.5	11.8	13.7	13.7	15.9	16.3	16.2	
3-year	3.4	3.8	4.4	5.8	4.0	7.1	7.0	5.6	5.9	7.0	7.9	8.1	7.3	8.1	7.7	9.8	9.2	11.2	12.4		
4-year	3.0	3.0	3.3	4.7	3.4	5.8	5.6	4.6	5.5	5.9	6.3	6.8	5.3	5.7	5.5	8.1	7.9	9.5			
5-year	2.6	2.5	2.9	4.0	2.9	5.1	4.7	4.0	4.9	5.2	5.4	6.0	4.3	4.9	5.0	7.5	7.3				
6-year	2.4	2.2	2.4	3.8	2.6	4.7	3.9	3.8	4.0	4.9	5.0	5.2	3.9	4.2	4.6	6.9					
7-year	2.3	1.9	2.3	3.4	2.6	4.2	3.4	3.5	3.9	4.5	4.5	4.7	3.6	3.8	4.1						
8-year	2.0	1.7	2.1	3.1	2.4	4.2	3.4	3.3	3.8	3.9	4.1	4.3	3.5	3.5							
9-year	1.9	1.6	1.8	3.0	2.2	4.0	3.3	2.9	3.6	3.7	4.0	4.2	3.1								
10-year	1.8	1.5	1.7	2.7	2.1	3.8	3.2	2.9	3.6	3.5	3.8	4.1									
11-year	1.7	1.5	1.7	2.7	2.0	3.2	2.9	2.9	3.2	3.5	3.6										
12-year	1.6	1.5	1.7	2.6	1.8	3.0	2.8	2.6	3.1	2.9											
13-year	1.6	1.4	1.5	2.5	1.7	3.0	2.6	2.6	3.0												
14-year	1.6	1.4	1.5	2.4	1.2	2.9	2.4	2.4													
15-year	1.6	1.3	1.3	2.3	1.2	2.8	2.2														
16-year	1.6	1.3	1.3	2.1	1.2	2.0															
17-year	1.4	1.3	1.3	2.1	0.8																
18-year	1.3	1.3	1.2	2.1																	
19-year	1.2	1.3	1.2																		
20-year	1.1	1.2	1.2																		

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).