

Table 22.11
Cancer of the Pancreas (Invasive)
SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Survival Time																					
1-year	14.2	16.8	17.3	19.1	18.7	18.2	18.8	19.6	21.8	21.7	24.2	25.0	21.5	21.0	27.7	24.4	27.7	28.3	30.0	30.8	29.8
2-year	5.9	6.6	7.3	8.7	9.1	7.9	9.9	8.3	11.1	10.5	10.7	12.6	11.2	11.2	14.0	12.5	13.9	14.1	14.9	17.0	
3-year	3.8	4.4	5.1	5.8	6.2	5.5	7.2	5.8	7.1	7.6	6.7	8.7	7.5	8.5	10.1	8.6	10.2	9.5	10.6		
4-year	3.0	3.6	4.3	4.9	4.4	3.9	5.5	4.4	5.8	6.3	5.8	6.8	5.9	7.1	8.8	7.7	8.6	7.3			
5-year	2.4	3.2	3.7	4.4	4.2	3.4	5.2	3.7	5.2	5.0	4.6	6.3	5.3	6.1	7.5	7.0	7.9				
6-year	2.3	2.8	3.4	4.1	3.7	3.0	4.9	3.7	5.0	4.3	4.3	6.1	4.8	5.7	6.8	6.5					
7-year	2.0	2.7	3.1	3.8	3.4	2.8	4.9	3.4	4.6	3.9	3.7	5.2	4.5	5.4	6.6						
8-year	1.8	2.5	2.8	3.7	3.0	2.6	4.9	2.8	4.0	3.7	3.5	5.1	4.2	5.1							
9-year	1.8	2.3	2.7	3.4	3.0	2.5	4.6	2.5	3.9	3.6	3.4	4.9	4.1								
10-year	1.7	2.2	2.6	3.2	2.9	2.5	4.6	2.4	3.7	3.6	3.0	4.7									
11-year	1.7	2.0	2.5	2.9	2.9	2.5	4.6	2.1	3.4	3.6	2.9										
12-year	1.7	1.9	2.5	2.7	2.8	2.5	4.3	2.1	3.2	3.4											
13-year	1.6	1.8	2.4	2.7	2.6	2.3	3.9	1.9	3.1												
14-year	1.5	1.7	2.4	2.6	2.4	2.3	3.8	1.3													
15-year	1.3	1.6	2.3	2.3	2.1	2.3	3.7														
16-year	1.3	1.5	2.1	2.3	2.0	2.3															
17-year	1.3	1.5	2.1	2.3	1.9																
18-year	1.3	1.4	2.1	2.2																	
19-year	1.3	1.4	2.0																		
20-year	1.3	1.4	1.9																		

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).