

Table 28.2

Age-Adjusted SEER Cancer Incidence Rates^a, 1975-2012

By Primary Cancer Site

All Races, Males and Females

Year of Diagnosis

Age/Site	1975	1980	1985	1990	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
<u>Ages 0-14</u>																						
All Sites																						
All Races	11.6	12.9	14.6	14.3	14.1	14.8	14.1	15.3	14.5	15.4	15.8	15.6	13.2	15.3	16.7	14.5	14.8	15.9	16.1	16.6	15.9	15.7
Whites	12.1	13.2	15.6	14.4	14.9	15.0	14.7	16.1	15.6	16.6	17.0	17.0	14.0	16.1	17.7	15.2	15.6	16.2	16.8	17.3	17.0	16.0
Blacks	8.6	11.1	10.2	12.3	11.8	11.6	11.2	10.3	10.7	12.1	11.0	11.6	10.2	13.3	13.0	11.8	12.0	15.4	13.1	13.2	13.0	15.9
Bone & Joint	0.5	0.7	0.8	0.7	0.8	0.5	0.9	0.6	0.6	0.6	0.6	0.7	0.5	0.6	0.6	0.6	0.6	0.7	1.0	0.7	0.7	0.8
Brain & Other nervous	2.3	2.8	3.0	3.5	3.4	3.2	2.8	3.2	3.5	3.4	4.0	3.8	3.0	3.3	3.4	3.1	3.4	3.6	3.8	4.1	3.4	3.0
Hodgkin lymphoma	0.7	0.5	0.7	0.7	-	0.6	0.6	0.6	0.4	0.6	0.5	0.5	0.4	0.5	0.5	0.6	0.6	0.7	0.6	0.5	0.6	0.7
Kidney & Renal pelvis	0.7	0.8	0.9	0.8	0.6	1.0	1.0	1.0	0.7	1.0	0.9	0.9	0.7	0.7	0.9	0.6	0.7	1.0	0.8	0.9	0.9	0.6
Leukemia	3.3	4.0	4.6	4.5	4.5	4.7	4.3	4.8	4.7	4.6	4.4	4.8	4.1	5.0	5.4	4.6	4.2	4.8	4.8	4.9	5.1	4.7
Acute lymphocytic	2.2	3.1	3.5	3.5	3.7	3.6	3.5	3.9	3.6	3.5	3.5	4.0	3.2	3.8	4.1	3.5	3.6	3.6	3.8	3.9	4.3	3.7
Non-Hodgkin lymphoma	1.0	0.7	0.7	0.8	0.9	0.6	0.9	0.9	0.8	1.0	1.0	1.0	0.9	0.8	1.1	0.9	1.0	1.1	1.1	1.4	1.0	1.4
Soft tissue	0.9	0.9	0.9	0.7	1.0	1.1	1.0	1.1	0.9	1.0	1.2	1.0	0.8	1.0	1.5	0.9	1.1	0.9	1.0	1.0	1.1	1.3
<u>Ages 0-19</u>																						
All Sites																						
All Races	13.0	14.4	16.0	15.7	15.8	16.2	15.8	16.3	15.7	16.9	17.1	17.5	15.7	16.5	17.8	15.7	17.0	17.7	17.7	17.6	18.1	17.7
Whites	13.4	14.9	17.1	16.0	16.7	16.6	16.9	17.3	16.7	18.3	18.5	19.5	16.8	17.8	19.2	16.7	18.3	18.4	19.1	18.4	19.7	18.4
Blacks	10.8	10.9	10.6	13.3	12.4	12.6	11.1	11.5	11.3	12.9	11.7	11.4	11.6	13.2	13.2	12.4	13.0	15.2	12.9	14.2	13.2	16.1
Bone & Joint	0.7	0.9	1.0	0.8	0.9	0.7	1.2	0.8	0.7	0.8	0.9	1.0	0.8	0.7	0.9	0.9	0.8	1.0	1.1	0.7	0.9	1.0
Brain & Other nervous	2.1	2.5	2.7	3.2	3.1	3.0	2.6	2.8	3.0	3.1	3.5	3.4	2.9	3.0	3.0	2.8	3.1	3.2	3.3	3.6	3.0	2.9
Hodgkin lymphoma	1.5	1.2	1.4	1.3	0.9	1.4	1.3	1.4	1.1	1.2	1.1	1.2	1.0	1.1	0.9	1.1	1.4	1.4	1.3	1.3	1.3	1.5
Kidney & Renal pelvis	0.5	0.6	0.7	0.6	0.5	0.7	0.8	0.8	0.6	0.8	0.7	0.7	0.6	0.6	0.8	0.5	0.6	0.9	0.7	0.7	0.7	0.5
Leukemia	3.0	3.5	4.0	3.9	4.0	4.0	3.7	4.1	4.1	4.1	4.0	4.5	3.8	4.4	4.7	4.1	4.0	4.3	4.1	4.2	4.6	4.2
Acute lymphocytic	1.9	2.5	2.9	2.9	3.1	2.9	2.9	3.3	3.0	3.1	2.9	3.5	2.9	3.3	3.4	3.0	3.2	3.1	3.1	3.1	3.7	3.0
Non-Hodgkin lymphoma	1.0	0.8	1.0	1.0	1.1	1.0	1.1	1.0	0.9	1.3	1.1	1.2	1.3	1.1	1.3	1.2	1.3	1.3	1.4	1.5	1.2	1.4
Soft tissue	0.8	0.9	0.9	0.8	1.1	1.1	1.1	1.1	1.0	1.2	1.1	1.0	0.8	1.0	1.5	1.0	1.1	0.9	1.0	0.9	1.2	1.2

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Statistic could not be calculated. Rate based on less than 16 cases for the time interval.