

Table X-1

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1973-96

All Races, Males, All Ages

## Year of Diagnosis

SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
All Areas⊕	174	35	44	109	218	367	518	723	857	1017	1119	1107	1137	1082	978	952	817	532
All Areas Excluding SF-Oakland SMSA	130	24	21	35	51	104	153	196	234	352	420	426	414	494	478	434	414	293
San Francisco-Oakland SMSA	44	11	23	74	167	263	365	527	623	665	699	681	723	588	500	518	403	239
San Francisco City/County	17	6	14	61	140	226	313	442	515	547	536	529	570	432	366	380	301	165

AGE-ADJUSTED CANCER INCIDENCE RATES§

By SEER Location and Year of Diagnosis, 1973-96

All Races, Males, All Ages

## Year of Diagnosis

SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
All Areas⊕	0.3	0.4	0.4	0.9	1.7	2.8	3.8	5.1	6.0	6.9	7.5	7.2	7.4	6.9	6.1	5.9	5.1	3.2
All Areas Excluding SF-Oakland SMSA	0.3	0.3	0.3	0.4	0.5	1.0	1.4	1.7	2.0	2.9	3.4	3.4	3.2	3.8	3.6	3.2	3.1	2.1
San Francisco-Oakland SMSA	0.5	0.7	1.3	3.5	7.4	11.5	15.7	21.7	25.5	26.2	27.4	25.8	27.8	22.3	18.6	19.7	15.3	9.1
San Francisco City/County	0.6	1.5	3.3	12.4	26.8	43.1	59.4	79.9	93.1	97.9	97.8	95.0	104.4	79.5	66.4	71.2	56.5	32.0

§ SEER Program. Rates are per 100,000 males and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

⊕ All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

Table X-2

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1973-96

All Races, Males, Ages 20-54

## Year of Diagnosis

SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
All Areas⊕	18	5	20	75	179	328	476	683	802	946	1052	1046	1068	1002	922	898	747	487
All Areas Excluding SF-Oakland SMSA	14	<4	4	9	26	78	128	174	205	310	381	391	379	461	444	412	372	266
San Francisco-Oakland SMSA	4	4	16	66	153	250	348	509	597	636	671	655	689	541	478	486	375	221
San Francisco City/County	<4	4	13	59	134	219	300	429	497	528	519	513	548	401	353	358	283	153

AGE-ADJUSTED CANCER INCIDENCE RATES§

By SEER Location and Year of Diagnosis, 1973-96

All Races, Males, Ages 20-54

## Year of Diagnosis

SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
All Areas⊕	0.1	0.1	0.4	1.3	3.0	5.6	7.9	11.0	12.7	14.5	16.0	15.5	15.6	14.2	13.0	12.6	10.3	6.6
All Areas Excluding SF-Oakland SMSA	0.1	-	0.1	0.2	0.6	1.6	2.5	3.3	3.8	5.7	6.9	7.0	6.6	7.9	7.5	6.9	6.1	4.3
San Francisco-Oakland SMSA	0.1	0.5	1.9	6.9	15.0	24.8	34.1	47.6	55.5	56.5	59.5	56.2	59.6	44.9	40.1	41.1	31.5	18.6
San Francisco City/County	-	2.4	7.2	27.7	58.5	96.0	129.6	177.1	204.6	215.1	214.8	209.6	227.7	162.6	145.7	150.1	118.9	66.5

§ SEER Program. Rates are per 100,000 males aged 20-54 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

⊕ All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

Table X-3

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1973-96

All Races, Males, Ages 55+

## Year of Diagnosis

SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
All Areas⊕	156	30	24	34	39	38	42	38	55	69	66	61	69	79	56	54	70	45
All Areas Excluding SF-Oakland SMSA	116	23	17	26	25	26	25	21	29	42	39	35	35	32	34	22	42	27
San Francisco-Oakland SMSA	40	7	7	8	14	12	17	17	26	27	27	26	34	47	22	32	28	18
San Francisco City/County	16	<4	<4	<4	6	7	13	13	18	19	16	16	22	31	13	22	18	12

AGE-ADJUSTED CANCER INCIDENCE RATES§

By SEER Location and Year of Diagnosis, 1973-96

All Races, Males, Ages 55+


## Year of Diagnosis

SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
All Areas⊕	1.5	1.8	1.4	1.9	2.1	2.0	2.2	2.0	2.8	3.5	3.4	3.0	3.5	4.0	2.7	2.7	3.3	2.0
All Areas Excluding SF-Oakland SMSA	1.3	1.6	1.2	1.7	1.6	1.7	1.6	1.3	1.7	2.5	2.4	2.0	2.0	1.9	1.9	1.2	2.3	1.4
San Francisco-Oakland SMSA	2.3	2.7	2.4	2.7	4.8	4.0	5.4	5.5	8.3	8.6	8.9	8.4	11.0	15.6	6.9	10.4	8.9	5.5
San Francisco City/County	2.9	-	-	-	8.4	9.0	18.7	19.0	25.7	27.5	24.7	24.4	32.8	49.6	18.8	34.2	27.7	17.6

§ SEER Program. Rates are per 100,000 males aged 55+ and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.


⊕ All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

# All Cancer Sites vs. Kaposi's Sarcoma SEER Incidence Rates, 1973-1996 Males, Ages 20-54




Age-adjusted to 1970 Standard

### San Francisco/Oakland Ages 20-54, All Males, 1973-1996



### All SEER excluding San Francisco/Oakland Ages 20-54, All Males, 1973-1996



\*Kaposi's excluded from Oral, Colorec., Lung, Melanoma, Testis, and NHL.

Figure X-2