Table I-2 51-YEAR TRENDS IN U.S. CANCER DEATH RATES^a

All Races, Males and Females

All Primary Cancer Sites Combined

						Total
				Annual		Percent
				Percent Change		Change
Age Group	1950	1975	2000	1950-1975	1975-2000	1950-2000
0-4	11.1	5.2	2.7	-2.8	-2.9	-76.0
5-14	6.7	4.8	2.5	-1.0	-2.8	-62.9
15-24	8.6	6.6	4.4	-0.7	-1.7	-49.1
25-34	20.4	15.1	9.9	-1.2	-1.4	-51.7
35-44	63.6	53.3	36.7	-0.4	-1.4	-42.3
45-54	174.2	178.0	127.4	0.2	-1.4	-26.9
55-64	391.3	420.2	366.6	0.3	-0.5	-6.3
65-74	710.0	778.7	816.2	0.4	0.2	15.0
75-84	1,167.2	1,160.9	1,335.0	0.0	0.5	14.4
85+	1,450.7	1,437.9	1,819.4	-0.2	0.8	25.4
All Ages	195.4	199.1	199.6	0.1	0.0	2.2

Lung and Bronchus Cancerb

				Annual Percent Change		Total Percent Change
Age Group	1950	1975	2000	1950-1975	1975-2000	1950-2000
0-4	-	-	-	-	-	-
5-14	-	-	_	-	_	-
15-24	0.2	0.1	-	-2.5	-0.9	-63.9
25-34	0.8	0.9	0.5	0.4	-1.6	-38.4
35-44	4.6	10.6	6.2	3.7	-2.5	34.4
45-54	20.2	48.8	31.6	3.6	-2.0	56.2
55-64	48.9	122.6	122.6	3.6	0.0	150.9
65-74	59.4	196.2	284.6	4.5	1.5	378.8
75-84	55.4	192.3	371.6	5.3	2.5	571.0
85+	42.3	135.5	302.7	5.1	3.1	615.8
All Ages	14.9	42.8	56.2	4.2	1.0	276.0

Source: NCHS public use data file.

Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based

on less than 10 cases for at least one year within the time interval.

Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year

Due to coding changes throughout the years, Lung and Bronchus includes trachea and pleura.

Table I-3

SUMMARY OF CHANGES IN CANCER INCIDENCE AND MORTALITY, 1950-2000 AND

5-YEAR RELATIVE SURVIVAL RATES, 1950-1999

Males and Females, By Primary Cancer Site

Table is temporarily unavailable.