

Table VIII-1
ESOPHAGUS CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2004
WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		Joinpoint Segment 1		Joinpoint Segment 2		Joinpoint Segment 3		Joinpoint Segment 4	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-2004	0.6*						
All Races	Male	1975-2004	0.8*						
All Races	Female	1975-2004	-0.2						
White	Male and Female	1975-1981	-0.5	1981-2004	1.6*				
White	Male	1975-2004	1.7*						
White	Female	1975-2004	0.2						
Black	Male and Female	1975-1986	1.0	1986-2004	-4.4*				
Black	Male	1975-1986	1.2	1986-2004	-4.9*				
Black	Female	1975-1980	9.7	1980-2004	-2.5*				
<u>SEER Delay-Adjusted Incidence^a</u>									
All Races	Male and Female	1975-2004	0.6*						
All Races	Male	1975-2004	0.8*						
All Races	Female	1975-2004	-0.2						
White	Male and Female	1975-1981	-0.6	1981-2004	1.7*				
White	Male	1975-2004	1.7*						
White	Female	1975-2004	0.2						
Black	Male and Female	1975-1986	1.0	1986-2004	-4.4*				
Black	Male	1975-1986	1.2	1986-2004	-4.8*				
Black	Female	1975-1980	9.7	1980-2004	-2.5*				
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-2001	0.7*	2001-2004	-0.6				
All Races	Male	1975-1985	0.7*	1985-1994	1.2*	1994-2004	0.4*		
All Races	Female	1975-2000	0.1	2000-2004	-1.4				
White	Male and Female	1975-1983	0.4*	1983-2001	1.7*	2001-2004	0.1		
White	Male	1975-1982	0.4	1982-2000	1.9*	2000-2004	0.5		
White	Female	1975-2004	0.4*						
Black	Male and Female	1975-1982	1.1*	1982-1993	-1.6*	1993-2004	-4.4*		
Black	Male	1975-1984	0.8	1984-1993	-1.9*	1993-2004	-4.7*		
Black	Female	1975-1990	-0.1	1990-2004	-3.1*				

Joinpoint Regression Program Version 3.0, April 2005, National Cancer Institute. (<http://srab.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

^b Trends are from the NCHS public use data file for the total US.

* The APC is significantly different from zero (p<.05).

Table VIII-2
ESOPHAGUS CANCER (Invasive)

AGE ADJUSTED SEER INCIDENCE^a RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	4.1	6.8	2.1	3.5	5.6	2.0	10.9	19.9	3.7
1976	4.3	6.9	2.3	3.7	5.9	2.1	10.7	16.9	5.6
1977	4.1	6.6	2.2	3.3	5.3	1.9	12.0	20.9	5.4
1978	4.1	7.0	1.9	3.4	5.7	1.7	13.0	24.4	4.3
1979	4.4	7.4	2.2	3.7	6.2	2.0	12.9	23.0	5.2
1980	4.3	6.9	2.3	3.5	5.7	1.9	12.3	18.8	7.4
1981	4.1	6.5	2.4	3.3	5.2	1.9	13.5	21.3	7.5
1982	4.3	6.8	2.3	3.5	5.5	2.0	13.6	22.7	6.4
1983	4.3	7.4	2.0	3.6	6.1	1.8	12.2	22.2	4.3
1984	4.2	6.7	2.3	3.5	5.6	1.9	12.0	19.9	6.0
1985	4.5	7.4	2.2	3.7	6.3	1.9	12.7	22.1	5.8
1986	4.6	7.5	2.5	3.9	6.2	2.2	14.1	24.0	6.3
1987	4.5	7.5	2.3	3.9	6.4	2.0	11.4	20.1	4.8
1988	4.4	7.3	2.2	3.9	6.3	2.0	11.2	19.5	5.0
1989	4.4	7.3	2.2	3.8	6.2	2.0	10.5	17.3	5.4
1990	4.8	8.3	2.2	4.2	7.2	1.9	12.5	21.4	5.8
1991	4.6	7.4	2.4	4.2	6.7	2.2	10.2	17.8	4.9
1992	4.6	8.0	2.0	4.2	7.4	1.8	10.6	18.0	5.3
1993	4.6	7.7	2.2	4.2	6.9	2.1	10.0	17.5	4.6
1994	4.5	7.7	2.1	4.2	7.2	1.9	9.2	14.7	5.1
1995	4.4	7.1	2.2	4.1	6.6	2.1	8.4	13.7	4.4
1996	4.8	8.1	2.2	4.4	7.5	2.0	9.7	16.4	4.8
1997	4.7	7.8	2.2	4.6	7.7	2.0	7.1	10.8	4.5
1998	4.7	8.2	2.0	4.5	7.9	1.9	7.8	12.5	4.2
1999	4.9	8.4	2.1	4.9	8.4	2.0	7.2	11.5	4.1
2000	4.8	8.0	2.2	4.7	8.0	2.0	7.0	10.6	4.4
2001	4.8	8.4	1.9	4.8	8.4	1.9	6.3	11.3	2.6
2002	4.6	7.9	2.0	4.6	8.0	1.8	6.3	10.0	3.5
2003	4.6	8.0	1.9	4.7	8.2	1.9	6.3	10.0	3.7
2004	5.2	8.7	2.3	5.3	9.0	2.3	6.0	8.8	3.9
1975-2004	4.5	7.6	2.2	4.1	6.9	2.0	9.7	16.3	4.8

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VIII-3
ESOPHAGUS CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH^a RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DEATH:									
1975	3.7	6.4	1.7	3.1	5.4	1.5	9.7	17.0	4.1
1976	3.7	6.4	1.8	3.1	5.3	1.5	10.2	17.7	4.3
1977	3.7	6.3	1.8	3.1	5.2	1.5	10.1	17.6	4.4
1978	3.8	6.4	1.8	3.2	5.4	1.6	10.0	17.6	4.3
1979	3.8	6.4	1.8	3.1	5.3	1.5	10.3	18.6	4.2
1980	3.8	6.6	1.8	3.2	5.5	1.5	10.6	18.3	4.9
1981	3.8	6.6	1.8	3.2	5.4	1.5	10.4	18.7	4.3
1982	3.8	6.5	1.8	3.2	5.4	1.6	10.5	18.6	4.5
1983	3.9	6.6	1.8	3.2	5.6	1.5	10.2	17.9	4.5
1984	3.9	6.7	1.9	3.3	5.6	1.6	10.6	18.8	4.6
1985	3.9	6.7	1.7	3.3	5.7	1.5	9.9	17.7	4.3
1986	3.9	6.8	1.8	3.3	5.8	1.6	10.2	18.0	4.5
1987	4.0	6.9	1.8	3.4	6.0	1.5	9.7	17.1	4.4
1988	4.0	7.0	1.8	3.5	6.0	1.6	10.0	18.4	3.9
1989	4.1	7.1	1.8	3.6	6.2	1.6	9.7	17.3	4.3
1990	4.1	7.2	1.8	3.6	6.3	1.5	9.3	16.3	4.4
1991	4.1	7.2	1.8	3.6	6.4	1.6	9.2	16.5	4.1
1992	4.2	7.3	1.8	3.7	6.6	1.6	8.6	15.1	4.1
1993	4.2	7.3	1.8	3.7	6.6	1.6	9.0	16.0	4.1
1994	4.3	7.6	1.8	3.9	6.9	1.6	8.6	15.3	3.9
1995	4.3	7.6	1.8	3.9	7.0	1.6	8.0	14.4	3.5
1996	4.3	7.5	1.9	4.0	7.1	1.7	7.8	13.6	3.7
1997	4.3	7.5	1.8	4.0	7.1	1.6	7.3	12.8	3.4
1998	4.4	7.7	1.8	4.1	7.3	1.7	7.1	12.2	3.6
1999	4.3	7.7	1.8	4.2	7.4	1.6	6.5	11.6	3.1
2000	4.4	7.7	1.8	4.3	7.6	1.7	6.4	10.9	3.3
2001	4.4	7.8	1.8	4.3	7.7	1.7	6.4	11.3	3.0
2002	4.4	7.8	1.7	4.3	7.8	1.6	5.9	10.2	2.9
2003	4.4	7.7	1.7	4.3	7.7	1.7	5.7	9.7	2.9
2004	4.4	7.7	1.7	4.3	7.8	1.6	5.4	9.1	2.9
1975-2004	4.1	7.2	1.8	3.7	6.5	1.6	8.5	14.9	3.8

^a NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VIII-4
ESOPHAGUS CANCER (Invasive)

SEER INCIDENCE^a AND U.S. DEATH^b RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER INCIDENCE</u>									
<u>AGE AT DIAGNOSIS</u>									
AGE-ADJUSTED RATES, 2000-2004									
All ages	4.6	7.9	2.0	4.6	8.0	1.9	6.2	10.4	3.2
Under 65	1.9	3.2	0.7	1.9	3.2	0.6	3.1	5.0	1.5
65 and over	23.4	40.4	11.3	23.8	41.2	11.3	27.8	47.7	15.1
All ages(IARC world std) ^c	3.0	5.1	1.2	3.0	5.2	1.2	4.4	7.2	2.2
AGE-SPECIFIC RATES, 2000-2004									
<1	-	-	-	-	-	-	-	-	-
1-4	-	-	-	-	-	-	-	-	-
5-9	-	-	-	-	-	-	-	-	-
10-14	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-
20-24	-	-	-	-	-	-	-	-	-
25-29	0.1	-	-	-	-	-	-	-	-
30-34	0.2	0.3	-	0.2	0.3	-	-	-	-
35-39	0.4	0.6	0.2	0.4	0.7	0.2	-	-	-
40-44	1.0	1.7	0.4	1.0	1.6	0.3	1.8	2.7	0.9
45-49	2.7	4.3	1.1	2.7	4.5	0.9	4.2	5.7	2.9
50-54	5.2	9.1	1.5	5.2	9.2	1.2	8.7	14.5	3.7
55-59	9.3	15.6	3.3	9.0	15.3	2.9	16.2	26.1	7.9
60-64	13.8	23.3	5.1	13.6	23.1	4.8	22.9	38.4	10.4
65-69	19.8	33.5	8.1	19.7	33.1	8.0	28.5	48.7	13.0
70-74	23.7	39.9	10.9	24.2	40.8	10.7	30.6	50.4	16.8
75-79	25.9	45.6	12.1	26.6	47.1	12.1	31.7	54.8	17.4
80-84	25.9	45.8	13.6	26.4	47.2	13.6	24.3	40.1	15.7
85+	23.4	41.5	15.2	24.0	42.6	15.7	18.3	36.5	11.0
<u>U.S. MORTALITY</u>									
<u>AGE AT DEATH</u>									
AGE-ADJUSTED RATES, 2000-2004									
All ages	4.4	7.8	1.8	4.3	7.7	1.7	6.0	10.2	3.0
Under 65	1.7	3.0	0.5	1.6	2.9	0.4	2.8	4.8	1.2
65 and over	22.9	40.8	10.3	22.8	40.8	10.0	27.7	47.9	15.2
All ages(IARC world std) ^c	2.8	5.0	1.0	2.8	4.9	1.0	4.1	7.0	2.0
AGE-SPECIFIC RATES, 2000-2004									
<1	-	-	-	-	-	-	-	-	-
1-4	-	-	-	-	-	-	-	-	-
5-9	-	-	-	-	-	-	-	-	-
10-14	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-
20-24	0.0	-	-	0.0	-	-	-	-	-
25-29	0.0	0.1	-	0.0	0.1	-	-	-	-
30-34	0.1	0.2	0.0	0.1	0.2	0.0	-	-	-
35-39	0.3	0.6	0.1	0.4	0.6	0.1	0.2	0.3	-
40-44	1.0	1.7	0.3	1.0	1.7	0.3	1.4	2.2	0.7
45-49	2.4	4.1	0.7	2.2	4.0	0.5	3.9	6.3	1.8
50-54	4.8	8.3	1.3	4.5	8.0	1.1	8.0	13.6	3.2
55-59	8.4	14.7	2.5	8.0	14.0	2.2	14.7	25.4	5.9
60-64	13.0	22.7	4.2	12.4	21.9	3.6	21.3	36.3	9.6
65-69	17.9	31.1	6.5	17.4	30.4	6.0	26.6	45.6	12.5
70-74	22.2	38.7	9.0	22.0	38.5	8.5	29.3	49.5	15.7
75-79	25.5	45.8	11.1	25.6	46.1	10.9	30.4	54.1	16.2
80-84	27.4	50.2	13.5	27.6	51.1	13.2	28.0	47.2	18.1
85+	25.4	46.6	16.3	26.0	47.7	16.6	21.8	39.6	14.9

^a SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^c Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VIII-5
ESOPHAGUS CANCER (Invasive)

SURVIVAL RATES, BY RACE, SEX, DIAGNOSIS YEAR, STAGE AND AGE

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
5-YR RELATIVE SURVIVAL RATES									
YEAR OF DIAGNOSIS:									
1960-1963 ^a	-	-	-	4	4	6	1	0	2
1970-1973 ^a	-	-	-	4	4	4	4	4	3
1975-1977 ^b	5.1	4.0	7.5	5.6	4.9	7.1	3.1	1.6	7.1
1978-1980 ^b	5.2	4.9	5.7	5.6	5.6	5.3	4.3	3.3	7.0
1981-1983 ^b	6.8	6.2	8.4	7.6	6.8	9.3	4.3	3.6	6.5
1984-1986 ^b	9.7	8.5	12.5	10.6	9.3	13.6	8.2	7.6	9.9
1987-1989 ^b	9.8	9.6	10.1	11.0	11.4	9.8	6.4	5.0	10.3
1990-1992 ^b	12.7	12.3	13.8	13.6	12.9	15.4	9.4	9.7	8.4
1993-1995 ^b	12.7	12.8	12.4	14.3	14.5	13.7	7.5	7.6	7.2
1996-2003 ^b	16.4 ^e	16.0 ^e	17.8 ^e	17.5 ^e	17.1 ^e	19.0 ^e	10.9 ^e	9.4 ^e	13.8
5-YR PERIOD SURVIVAL RATES^{cd}									
YEAR OF DIAGNOSIS:									
1996-2003	16.5	15.5	19.8	17.2	16.2	20.5	11.3	10.0	14.1
STAGE DISTRIBUTION (%) 1996-2003^c									
All Stages									
Number of cases	15,153	11,460	3,693	12,317	9,431	2,886	1,937	1,342	595
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
Localized	24	23	25	25	24	26	20	20	21
Regional	30	31	27	30	31	27	29	29	29
Distant	30	32	25	30	31	24	32	33	29
Unstaged	16	14	22	16	14	23	19	17	21
5-YR RELATIVE SURVIVAL RATES, 1996-2003^c									
AGE AT DIAGNOSIS:									
<45	21.0	20.0	25.7	20.9	19.7	29.6 ^f	18.4	17.3 ^f	18.3 ^f
45-54	17.2	17.0	18.3	19.3	19.1	19.0	10.3	8.5	16.1
55-64	16.6	15.6	20.9	17.5	16.4	23.3	11.2	10.4	12.5
65-74	15.8	15.5	16.9	16.9	16.6	18.1	6.6	5.7	8.2
75+	11.4	10.8	12.4	12.1	11.4	13.1	9.1	9.9	6.1
Under 65	17.3	16.6	20.8	18.5	17.8	22.9	11.6	10.3	14.5
65 and over	14.1	13.8	14.7	14.9	14.7	15.6	7.4	7.1	7.8
STAGE:									
All Stages	15.6	15.3	16.8	16.6	16.3	17.8	10.1	9.3	11.7
Localized	33.7	34.1	32.1	35.9	36.7	33.0	19.6	16.6	24.5
Regional	16.9	16.2	19.5	18.1	17.6	20.2	11.1	9.3	14.2
Distant	2.9	2.8	3.4	3.1	2.8	4.3	1.5	1.7	-
Unstaged	10.8	10.5	11.2	9.7	8.8	11.1	13.0	14.8	8.9

^a Rates are based on End Results data from a series of hospital registries and one population-based registry.

^b Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

^c Rates are based on follow-up of patients into 2004.

^d Rates are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2003. The remaining 13 SEER Areas contribute cases for the entire period 1996-2003. Rates are based on follow-up of patients into 2004.

^e Period survival provides a more up-to-date estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three calendar year blocks (2001-2003: 0-1 year survival), (2000-2002: 1-2 year survival), (1999-2001: 2-3 year survival), (1998-2000: 3-4 year survival), (1997-1999: 4-5 years survival).

^f The difference in rates between 1975-1977 and 1996-2003 is statistically significant ($p < .05$).

^g The standard error of the survival rate is between 5 and 10 percentage points.

⁻ Statistic could not be calculated due to fewer than 16 cases during the time period.

Table VIII-6
ESOPHAGUS CANCER (Invasive)

SURVIVAL RATES^a

By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975-	1980-																					
	1979	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
Relative Survival Rates (SEER^b)																							
1-year	27.6	31.4	35.9	35.0	35.8	36.3	38.7	39.4	41.7	42.0	38.4	38.8	41.6	43.6	41.5	41.8	44.8	47.7	47.6	42.7	48.3		
2-year	11.7	14.6	20.7	19.1	18.2	20.4	20.2	20.6	21.4	24.8	22.6	23.3	24.0	24.5	24.0	24.5	27.0	29.1	29.7	25.9			
3-year	7.8	10.3	13.4	14.4	12.0	14.0	14.5	14.8	15.3	19.8	16.4	18.4	17.7	18.9	17.6	18.9	20.9	24.4	23.2				
4-year	6.0	8.5	10.1	11.8	9.5	12.1	11.7	11.9	13.1	17.1	14.1	15.3	13.8	16.2	14.4	14.7	18.5	21.3					
5-year	4.9	7.2	9.0	10.5	8.9	10.0	10.3	10.4	12.2	15.4	12.5	13.5	12.1	15.5	12.6	13.0	17.5						
6-year	4.4	6.4	7.5	9.4	7.9	9.2	9.1	10.0	10.7	14.4	11.3	12.3	11.4	14.4	11.6	12.2							
7-year	4.0	5.8	7.0	8.7	7.1	8.7	8.4	9.0	10.2	13.0	10.6	11.3	10.9	13.6	11.2								
8-year	3.5	5.2	6.2	7.9	6.2	8.2	8.3	8.6	9.0	12.4	10.2	10.7	10.7	10.8	13.0								
9-year	3.3	4.8	5.7	7.7	5.5	7.0	7.6	7.9	8.9	11.8	9.3	10.5	9.9										
10-year	3.0	4.6	5.7	6.8	5.4	6.5	6.9	7.5	8.6	10.7	8.9	9.2											
11-year	2.9	4.3	5.2	6.3	5.4	6.0	5.8	7.4	8.6	9.5	8.0												
12-year	2.7	4.2	4.5	6.0	5.4	5.6	4.7	5.6	8.2	8.6													
13-year	2.6	4.0	4.5	5.7	4.1	5.6	4.4	5.6	8.2														
14-year	2.6	3.9	4.5	5.7	3.9	5.1	3.9	5.6															
15-year	2.6	3.9	4.5	5.5	3.8	4.6	3.7																
16-year	2.6	3.8	4.5	4.3	3.6	4.3																	
17-year	2.6	3.5	3.7	3.7	3.6																		
18-year	2.5	3.2	3.7	3.7																			
19-year	2.5	3.0	3.4																				
20-year	2.4	2.9																					

^a Survival rates are relative rates expressed as percents.

^b Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table VIII-7
ESOPHAGUS CANCER (Invasive)

SURVIVAL RATES^a

By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975-		1980-																			
	1979	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	
Relative Survival Rates (SEER^b)																						
1-year	25.3	30.7	34.8	34.2	34.1	34.4	38.3	41.4	41.3	41.4	39.6	38.3	41.2	44.9	41.9	41.9	44.4	48.9	48.8	44.0	48.0	
2-year	10.3	14.1	19.5	19.7	17.2	19.2	19.6	21.5	21.0	24.7	22.3	22.0	24.3	25.1	24.7	24.8	26.6	29.4	30.1	26.4		
3-year	6.8	9.4	11.2	15.6	12.4	12.2	14.4	14.8	15.5	19.8	16.7	18.2	17.8	18.7	18.5	18.9	20.8	24.2	23.1			
4-year	5.2	7.7	7.3	12.5	9.8	10.8	11.6	11.8	12.9	16.3	14.3	14.9	13.6	15.5	15.3	14.4	18.2	20.2				
5-year	4.3	6.2	7.1	11.2	9.1	9.4	10.0	10.1	12.2	14.4	12.6	13.4	12.3	14.4	13.4	12.2	17.4					
6-year	3.7	5.4	6.2	10.3	8.4	8.8	8.6	9.7	10.5	13.4	11.5	12.0	11.6	13.6	12.1	11.6						
7-year	3.4	4.8	5.7	9.4	7.2	8.5	7.8	8.7	10.5	12.1	11.2	11.2	11.3	12.7	11.9							
8-year	3.1	4.3	5.0	8.2	6.4	8.1	7.6	8.0	8.9	11.5	10.8	10.3	11.0	12.2								
9-year	2.9	4.0	4.9	8.1	6.0	6.7	7.4	7.3	8.7	11.1	9.7	9.9	10.6									
10-year	2.8	3.8	4.8	7.7	6.0	6.2	7.1	6.9	8.3	9.7	8.9	8.9										
11-year	2.6	3.7	4.8	7.0	6.0	5.7	6.3	6.8	8.3	8.2	8.3											
12-year	2.5	3.6	3.6	6.4	6.0	5.0	5.3	5.4	7.7	7.5												
13-year	2.4	3.6	3.6	5.9	4.4	5.0	4.9	5.4	7.7													
14-year	2.4	3.5	3.6	5.9	3.9	4.3	4.0	5.4														
15-year	2.4	3.5	3.6	5.7	3.7	3.5	4.0															
16-year	2.4	3.5	3.6	4.6	3.4	3.0																
17-year	2.4	3.5	2.4	3.6	3.2																	
18-year	2.4	3.2	2.4	3.5																		
19-year	2.4	3.1	1.7																			
20-year	2.4	3.0																				

^a Survival rates are relative rates expressed as percents.

^b Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table VIII-8
ESOPHAGUS CANCER (Invasive)

SURVIVAL RATES^a

By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975-	1980-																					
	1979	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
Relative Survival Rates (SEER^b)																							
1-year	33.4	33.3	39.1	37.0	40.1	41.4	39.8	33.0	42.6	43.9	35.4	40.5	42.9	39.8	40.4	41.5	46.5	43.9	43.0	38.5	49.3		
2-year	15.2	16.0	23.9	17.8	20.6	23.7	21.7	17.9	22.1	25.2	23.3	27.6	22.9	22.6	22.0	23.4	28.5	28.1	28.2	24.4			
3-year	10.2	12.2	19.1	11.6	11.0	18.6	14.6	14.7	14.9	19.7	15.4	19.0	17.3	19.7	15.1	18.7	21.1	25.1	23.5				
4-year	8.0	10.6	17.6	10.1	8.8	15.4	11.9	12.0	13.7	19.3	13.5	16.0	14.2	18.1	11.7	15.7	19.3	24.5					
5-year	6.6	9.5	13.9	8.8	8.0	11.4	10.7	11.1	12.3	18.1	12.3	13.6	11.2	18.1	10.6	15.6	17.8						
6-year	6.1	8.7	10.8	7.1	6.6	10.2	10.5	10.7	11.1	17.9	10.7	13.0	10.7	16.7	9.8	14.1							
7-year	5.3	8.4	10.4	6.9	6.6	9.3	9.7	9.9	9.5	15.9	8.8	11.6	9.8	16.1	8.7								
8-year	4.5	7.5	8.8	6.9	5.3	8.5	9.7	9.9	9.2	15.8	8.5	11.2	9.8	15.1									
9-year	4.2	6.7	7.9	6.6	4.2	7.4	8.2	9.3	9.2	14.1	8.3	11.2	7.9										
10-year	3.6	6.6	7.5	4.6	3.6	7.2	6.2	8.9	9.2	13.8	8.3	9.7											
11-year	3.4	5.6	6.3	4.6	3.6	6.8	4.6	8.9	9.2	13.7	7.2												
12-year	3.2	5.3	6.3	4.6	3.6	6.8	3.1	6.4	9.1	12.6													
13-year	2.9	4.9	6.3	4.6	3.5	6.8	3.1	5.9	9.0														
14-year	2.9	4.7	6.3	4.6	3.5	6.8	3.1	5.9															
15-year	2.9	4.7	6.3	4.6	3.5	6.8	2.8																
16-year	2.8	4.0	6.2	3.6	3.5	6.8																	
17-year	2.5	3.2	6.2	3.6	3.5																		
18-year	2.3	3.1	6.2	3.6																			
19-year	2.3	2.8	6.2																				
20-year	2.3	2.7																					

^a Survival rates are relative rates expressed as percents.

^b Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table VIII-9
ESOPHAGUS CANCER (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.50 (0.49, 0.51)	0.48 (0.47, 0.48)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.51 (0.50, 0.52)	0.48 (0.48, 0.49)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.51 (0.50, 0.52)	0.48 (0.48, 0.49)	
	30	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.09 (0.09, 0.09)	0.52 (0.51, 0.53)	0.49 (0.48, 0.49)	
	40	0.02 (0.02, 0.02)	0.09 (0.08, 0.09)	0.23 (0.22, 0.23)	0.52 (0.51, 0.53)	0.49 (0.49, 0.50)	
	50	0.07 (0.07, 0.07)	0.22 (0.21, 0.22)	0.39 (0.38, 0.40)	0.52 (0.50, 0.53)	0.49 (0.49, 0.50)	
	60	0.15 (0.15, 0.16)	0.34 (0.33, 0.35)	0.45 (0.44, 0.46)	0.47 (0.46, 0.49)	0.45 (0.45, 0.46)	
	70	0.21 (0.20, 0.22)	0.34 (0.33, 0.35)	- (- , -)	0.37 (0.36, 0.38)	0.36 (0.36, 0.37)	
	80	0.18 (0.17, 0.19)	- (- , -)	- (- , -)	0.22 (0.21, 0.23)	0.23 (0.23, 0.24)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.51 (0.50, 0.53)	0.48 (0.47, 0.49)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.52 (0.51, 0.53)	0.49 (0.48, 0.49)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.52 (0.51, 0.53)	0.49 (0.48, 0.49)	
	30	0.00 (0.00, 0.00)	0.02 (0.02, 0.02)	0.09 (0.08, 0.09)	0.53 (0.51, 0.54)	0.49 (0.49, 0.50)	
	40	0.02 (0.02, 0.02)	0.09 (0.08, 0.09)	0.23 (0.22, 0.24)	0.53 (0.52, 0.54)	0.49 (0.49, 0.50)	
	50	0.07 (0.07, 0.07)	0.22 (0.21, 0.22)	0.39 (0.38, 0.40)	0.52 (0.51, 0.54)	0.49 (0.49, 0.50)	
	60	0.15 (0.15, 0.16)	0.34 (0.33, 0.35)	0.45 (0.44, 0.47)	0.48 (0.47, 0.49)	0.46 (0.45, 0.46)	
	70	0.22 (0.21, 0.23)	0.35 (0.33, 0.36)	- (- , -)	0.38 (0.37, 0.39)	0.37 (0.36, 0.37)	
	80	0.18 (0.17, 0.19)	- (- , -)	- (- , -)	0.23 (0.22, 0.24)	0.23 (0.23, 0.24)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.52 (0.49, 0.56)	0.51 (0.49, 0.52)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.54 (0.50, 0.57)	0.52 (0.50, 0.53)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.03 (0.03, 0.04)	0.54 (0.51, 0.57)	0.52 (0.51, 0.54)	
	30	0.00 (0.00, 0.00)	0.03 (0.03, 0.04)	0.13 (0.12, 0.15)	0.55 (0.51, 0.58)	0.53 (0.51, 0.55)	
	40	0.03 (0.03, 0.03)	0.14 (0.12, 0.15)	0.32 (0.30, 0.34)	0.56 (0.52, 0.60)	0.54 (0.52, 0.56)	
	50	0.11 (0.10, 0.12)	0.31 (0.28, 0.33)	0.48 (0.45, 0.51)	0.56 (0.52, 0.60)	0.54 (0.52, 0.56)	
	60	0.22 (0.20, 0.24)	0.41 (0.38, 0.44)	0.49 (0.45, 0.52)	0.50 (0.46, 0.54)	0.49 (0.47, 0.51)	
	70	0.24 (0.21, 0.27)	0.34 (0.30, 0.37)	- (- , -)	0.35 (0.32, 0.39)	0.36 (0.34, 0.38)	
	80	0.16 (0.13, 0.19)	- (- , -)	- (- , -)	0.18 (0.15, 0.22)	0.21 (0.20, 0.23)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

- Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VIII-9 - continued
ESOPHAGUS CANCER (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
 Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
 Both Sexes, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer						Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	Percent (95% C.I.)		
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.36 (0.32, 0.41)	0.36 (0.32, 0.41)	0.30 (0.26, 0.35)		
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.36 (0.32, 0.41)	0.36 (0.32, 0.41)	0.30 (0.27, 0.35)		
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.01)	0.36 (0.32, 0.42)	0.36 (0.32, 0.42)	0.30 (0.27, 0.35)		
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.04 (0.04, 0.05)	0.36 (0.32, 0.42)	0.36 (0.32, 0.42)	0.30 (0.27, 0.35)		
	40	0.01 (0.01, 0.01)	0.04 (0.03, 0.05)	0.11 (0.10, 0.13)	0.36 (0.32, 0.42)	0.36 (0.32, 0.42)	0.31 (0.27, 0.35)		
	50	0.03 (0.03, 0.04)	0.11 (0.09, 0.12)	0.20 (0.17, 0.22)	0.36 (0.32, 0.41)	0.31 (0.27, 0.35)			
	60	0.08 (0.06, 0.09)	0.17 (0.15, 0.19)	0.29 (0.26, 0.33)	0.34 (0.30, 0.39)	0.29 (0.25, 0.34)			
	70	0.10 (0.09, 0.12)	0.24 (0.20, 0.27)	- (-, -)	0.29 (0.24, 0.34)	0.26 (0.22, 0.31)			
	80	0.17 (0.14, 0.20)	- (-, -)	- (-, -)	0.23 (0.18, 0.30)	0.21 (0.16, 0.27)			
American	0	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.41 (0.28, 0.71)	0.40 (0.32, 0.52)			
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.42 (0.29, 0.72)	0.41 (0.32, 0.53)			
	20	0.00 (0.00, 0.01)	0.00 (0.00, 0.01)	0.01 (0.00, 0.03)	0.42 (0.29, 0.73)	0.41 (0.33, 0.54)			
	30	0.00 (0.00, 0.01)	0.01 (0.00, 0.03)	0.04 (0.02, 0.08)	0.43 (0.30, 0.74)	0.42 (0.33, 0.55)			
	40	0.01 (0.00, 0.03)	0.04 (0.02, 0.08)	0.19 (0.12, 0.28)	0.44 (0.30, 0.76)	0.43 (0.34, 0.56)			
	50	0.03 (0.01, 0.06)	0.18 (0.11, 0.28)	0.37 (0.25, 0.52)	0.44 (0.31, 0.78)	0.44 (0.34, 0.57)			
	60	0.16 (0.10, 0.26)	0.36 (0.24, 0.52)	0.44 (0.30, 0.65)	0.44 (0.30, 0.80)	0.43 (0.33, 0.58)			
	70	0.23 (0.14, 0.38)	0.33 (0.19, 0.54)	- (-, -)	0.33 (0.19, 0.73)	0.37 (0.26, 0.54)			
	80	0.13 (0.04, 0.34)	- (-, -)	- (-, -)	0.13 (0.04, 0.68)	0.28 (0.16, 0.52)			
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.41 (0.37, 0.47)	0.31 (0.29, 0.33)			
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.42 (0.38, 0.47)	0.31 (0.29, 0.34)			
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.42 (0.38, 0.47)	0.32 (0.29, 0.34)			
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.05 (0.04, 0.05)	0.42 (0.38, 0.48)	0.32 (0.30, 0.34)			
	40	0.01 (0.01, 0.01)	0.04 (0.04, 0.05)	0.13 (0.11, 0.14)	0.43 (0.38, 0.48)	0.32 (0.30, 0.35)			
	50	0.03 (0.03, 0.04)	0.12 (0.11, 0.14)	0.25 (0.22, 0.27)	0.43 (0.38, 0.48)	0.32 (0.30, 0.35)			
	60	0.09 (0.08, 0.10)	0.22 (0.20, 0.25)	0.34 (0.31, 0.37)	0.41 (0.36, 0.47)	0.31 (0.28, 0.33)			
	70	0.15 (0.13, 0.17)	0.28 (0.25, 0.31)	- (-, -)	0.36 (0.31, 0.42)	0.27 (0.24, 0.29)			
	80	0.17 (0.14, 0.21)	- (-, -)	- (-, -)	0.28 (0.22, 0.35)	0.20 (0.17, 0.24)			

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VIII-9 - continued
ESOPHAGUS CANCER (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.78 (0.76, 0.80)	0.75 (0.74, 0.76)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.79 (0.77, 0.81)	0.77 (0.76, 0.77)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.04 (0.03, 0.04)	0.79 (0.77, 0.81)	0.77 (0.76, 0.78)	
	30	0.00 (0.00, 0.01)	0.04 (0.03, 0.04)	0.15 (0.14, 0.15)	0.80 (0.78, 0.82)	0.78 (0.77, 0.79)	
	40	0.03 (0.03, 0.03)	0.14 (0.14, 0.15)	0.38 (0.36, 0.39)	0.81 (0.79, 0.83)	0.79 (0.78, 0.80)	
	50	0.12 (0.11, 0.12)	0.36 (0.35, 0.37)	0.63 (0.61, 0.64)	0.81 (0.79, 0.83)	0.79 (0.78, 0.80)	
	60	0.26 (0.25, 0.27)	0.55 (0.53, 0.57)	0.71 (0.69, 0.73)	0.75 (0.73, 0.77)	0.73 (0.72, 0.74)	
	70	0.35 (0.33, 0.36)	0.54 (0.52, 0.56)	- (-, -)	0.58 (0.56, 0.60)	0.59 (0.58, 0.60)	
	80	0.30 (0.28, 0.31)	- (-, -)	- (-, -)	0.36 (0.34, 0.38)	0.38 (0.37, 0.39)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.80 (0.78, 0.82)	0.77 (0.76, 0.78)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.81 (0.79, 0.83)	0.78 (0.77, 0.79)	
	20	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.04 (0.03, 0.04)	0.81 (0.79, 0.83)	0.78 (0.77, 0.79)	
	30	0.00 (0.00, 0.01)	0.04 (0.03, 0.04)	0.15 (0.14, 0.16)	0.82 (0.80, 0.84)	0.79 (0.78, 0.80)	
	40	0.03 (0.03, 0.03)	0.15 (0.14, 0.15)	0.38 (0.37, 0.39)	0.83 (0.81, 0.85)	0.80 (0.79, 0.81)	
	50	0.12 (0.11, 0.12)	0.36 (0.35, 0.37)	0.64 (0.62, 0.66)	0.83 (0.80, 0.85)	0.80 (0.79, 0.81)	
	60	0.26 (0.25, 0.27)	0.56 (0.54, 0.58)	0.73 (0.71, 0.75)	0.76 (0.74, 0.78)	0.74 (0.73, 0.75)	
	70	0.36 (0.34, 0.37)	0.56 (0.54, 0.58)	- (-, -)	0.60 (0.58, 0.62)	0.60 (0.59, 0.61)	
	80	0.30 (0.29, 0.32)	- (-, -)	- (-, -)	0.37 (0.35, 0.39)	0.39 (0.38, 0.40)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.76 (0.70, 0.82)	0.74 (0.72, 0.77)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.78 (0.72, 0.84)	0.76 (0.74, 0.79)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.04 (0.03, 0.05)	0.79 (0.73, 0.85)	0.77 (0.74, 0.80)	
	30	0.00 (0.00, 0.01)	0.04 (0.03, 0.05)	0.20 (0.18, 0.22)	0.81 (0.75, 0.87)	0.79 (0.76, 0.81)	
	40	0.04 (0.03, 0.05)	0.20 (0.18, 0.23)	0.50 (0.46, 0.54)	0.83 (0.77, 0.90)	0.81 (0.78, 0.84)	
	50	0.17 (0.15, 0.20)	0.49 (0.45, 0.53)	0.73 (0.68, 0.79)	0.84 (0.78, 0.91)	0.82 (0.79, 0.85)	
	60	0.36 (0.33, 0.41)	0.65 (0.59, 0.71)	0.75 (0.69, 0.83)	0.77 (0.70, 0.85)	0.75 (0.72, 0.79)	
	70	0.38 (0.33, 0.43)	0.52 (0.46, 0.59)	- (-, -)	0.54 (0.48, 0.62)	0.55 (0.52, 0.59)	
	80	0.26 (0.20, 0.33)	- (-, -)	- (-, -)	0.30 (0.24, 0.40)	0.33 (0.30, 0.37)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VIII-9 - continued
ESOPHAGUS CANCER (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Males, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.54 (0.47, 0.63)	0.47 (0.40, 0.55)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.54 (0.47, 0.63)	0.47 (0.41, 0.55)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.54 (0.47, 0.63)	0.47 (0.41, 0.56)	
	30	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.07 (0.05, 0.08)	0.55 (0.47, 0.64)	0.48 (0.41, 0.56)	
	40	0.01 (0.01, 0.02)	0.07 (0.05, 0.08)	0.19 (0.16, 0.22)	0.55 (0.48, 0.64)	0.48 (0.41, 0.56)	
	50	0.05 (0.04, 0.07)	0.18 (0.15, 0.21)	0.33 (0.29, 0.37)	0.54 (0.47, 0.64)	0.48 (0.41, 0.56)	
	60	0.13 (0.11, 0.16)	0.28 (0.24, 0.33)	0.47 (0.40, 0.53)	0.51 (0.44, 0.61)	0.46 (0.39, 0.54)	
	70	0.17 (0.14, 0.21)	0.37 (0.31, 0.44)	- (-, -)	0.42 (0.35, 0.52)	0.39 (0.32, 0.49)	
	80	0.26 (0.20, 0.34)	- (-, -)	- (-, -)	0.33 (0.25, 0.45)	0.31 (0.23, 0.42)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.61 (0.40, 1.29)	0.62 (0.48, 0.86)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.62 (0.41, 1.31)	0.63 (0.49, 0.88)	
	20	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.02 (0.00, 0.06)	0.63 (0.41, 1.33)	0.64 (0.50, 0.89)	
	30	0.00 (0.00, 0.03)	0.02 (0.00, 0.06)	0.07 (0.03, 0.14)	0.65 (0.42, 1.36)	0.66 (0.51, 0.91)	
	40	0.02 (0.00, 0.05)	0.07 (0.03, 0.14)	0.31 (0.19, 0.49)	0.67 (0.43, 1.41)	0.68 (0.53, 0.95)	
	50	0.06 (0.02, 0.13)	0.31 (0.18, 0.50)	0.57 (0.36, 0.85)	0.68 (0.44, 1.46)	0.70 (0.54, 0.98)	
	60	0.28 (0.15, 0.47)	0.56 (0.35, 0.86)	0.69 (0.43, 1.09)	0.69 (0.43, 1.54)	0.70 (0.53, 1.01)	
	70	0.33 (0.17, 0.61)	0.49 (0.25, 0.91)	- (-, -)	0.49 (0.25, 1.50)	0.61 (0.42, 0.98)	
	80	0.23 (0.06, 0.73)	- (-, -)	- (-, -)	0.23 (0.06, 1.76)	0.48 (0.26, 1.02)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.66 (0.58, 0.76)	0.48 (0.44, 0.52)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.66 (0.58, 0.77)	0.48 (0.45, 0.53)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.02 (0.01, 0.02)	0.67 (0.59, 0.77)	0.49 (0.45, 0.53)	
	30	0.00 (0.00, 0.00)	0.02 (0.01, 0.02)	0.08 (0.07, 0.09)	0.67 (0.59, 0.78)	0.49 (0.45, 0.54)	
	40	0.02 (0.01, 0.02)	0.08 (0.06, 0.09)	0.23 (0.20, 0.26)	0.68 (0.60, 0.79)	0.50 (0.46, 0.54)	
	50	0.06 (0.05, 0.07)	0.22 (0.19, 0.25)	0.42 (0.37, 0.46)	0.69 (0.60, 0.79)	0.50 (0.46, 0.55)	
	60	0.17 (0.14, 0.19)	0.38 (0.33, 0.43)	0.56 (0.50, 0.63)	0.67 (0.58, 0.78)	0.48 (0.44, 0.53)	
	70	0.24 (0.21, 0.29)	0.45 (0.39, 0.52)	- (-, -)	0.57 (0.48, 0.70)	0.41 (0.37, 0.47)	
	80	0.29 (0.23, 0.37)	- (-, -)	- (-, -)	0.47 (0.35, 0.63)	0.30 (0.25, 0.37)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VIII-9 - continued
ESOPHAGUS CANCER (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.25 (0.24, 0.26)	0.22 (0.22, 0.23)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.26 (0.25, 0.27)	0.22 (0.22, 0.23)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.26 (0.25, 0.27)	0.22 (0.22, 0.23)	
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.26 (0.25, 0.27)	0.22 (0.22, 0.23)	
	40	0.01 (0.01, 0.01)	0.03 (0.03, 0.03)	0.09 (0.08, 0.09)	0.26 (0.25, 0.27)	0.23 (0.22, 0.23)	
	50	0.02 (0.02, 0.03)	0.08 (0.08, 0.09)	0.17 (0.16, 0.18)	0.26 (0.25, 0.27)	0.23 (0.22, 0.23)	
	60	0.06 (0.06, 0.07)	0.15 (0.14, 0.16)	0.22 (0.21, 0.23)	0.24 (0.23, 0.26)	0.22 (0.21, 0.22)	
	70	0.10 (0.10, 0.11)	0.18 (0.17, 0.19)	- (-, -)	0.21 (0.20, 0.22)	0.19 (0.18, 0.19)	
	80	0.10 (0.10, 0.11)	- (-, -)	- (-, -)	0.14 (0.13, 0.15)	0.14 (0.13, 0.14)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.25 (0.24, 0.26)	0.21 (0.21, 0.22)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.25 (0.24, 0.27)	0.22 (0.21, 0.22)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.25 (0.24, 0.27)	0.22 (0.21, 0.22)	
	30	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.03 (0.02, 0.03)	0.26 (0.24, 0.27)	0.22 (0.21, 0.22)	
	40	0.01 (0.00, 0.01)	0.03 (0.02, 0.03)	0.08 (0.08, 0.09)	0.26 (0.24, 0.27)	0.22 (0.21, 0.22)	
	50	0.02 (0.02, 0.02)	0.08 (0.07, 0.08)	0.17 (0.16, 0.18)	0.26 (0.24, 0.27)	0.22 (0.21, 0.22)	
	60	0.06 (0.05, 0.06)	0.15 (0.14, 0.16)	0.22 (0.21, 0.23)	0.24 (0.23, 0.26)	0.21 (0.21, 0.22)	
	70	0.10 (0.10, 0.11)	0.18 (0.17, 0.19)	- (-, -)	0.21 (0.20, 0.22)	0.19 (0.18, 0.19)	
	80	0.10 (0.10, 0.11)	- (-, -)	- (-, -)	0.14 (0.13, 0.15)	0.14 (0.13, 0.14)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.31 (0.28, 0.35)	0.30 (0.28, 0.31)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.32 (0.28, 0.35)	0.30 (0.29, 0.32)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.02 (0.02, 0.03)	0.32 (0.28, 0.36)	0.30 (0.29, 0.32)	
	30	0.00 (0.00, 0.00)	0.02 (0.02, 0.03)	0.08 (0.06, 0.09)	0.32 (0.28, 0.36)	0.31 (0.29, 0.32)	
	40	0.02 (0.01, 0.03)	0.08 (0.06, 0.09)	0.16 (0.14, 0.19)	0.32 (0.29, 0.36)	0.31 (0.29, 0.33)	
	50	0.06 (0.05, 0.07)	0.15 (0.13, 0.17)	0.26 (0.23, 0.29)	0.32 (0.28, 0.36)	0.31 (0.29, 0.33)	
	60	0.10 (0.08, 0.12)	0.22 (0.19, 0.25)	0.27 (0.24, 0.31)	0.28 (0.24, 0.32)	0.29 (0.27, 0.31)	
	70	0.14 (0.11, 0.17)	0.20 (0.17, 0.24)	- (-, -)	0.22 (0.18, 0.26)	0.23 (0.21, 0.25)	
	80	0.10 (0.07, 0.13)	- (-, -)	- (-, -)	0.11 (0.08, 0.15)	0.15 (0.13, 0.17)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VIII-9 - continued
ESOPHAGUS CANCER (INVASIVE)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.20 (0.16, 0.28)	0.16 (0.12, 0.22)	
Islander	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.20 (0.16, 0.28)	0.16 (0.12, 0.22)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.00, 0.01)	0.21 (0.16, 0.28)	0.16 (0.12, 0.22)	
	30	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.02 (0.01, 0.03)	0.21 (0.16, 0.28)	0.16 (0.12, 0.22)	
	40	0.00 (0.00, 0.01)	0.02 (0.01, 0.03)	0.05 (0.03, 0.06)	0.20 (0.16, 0.28)	0.16 (0.12, 0.22)	
	50	0.01 (0.01, 0.02)	0.04 (0.03, 0.06)	0.09 (0.07, 0.11)	0.20 (0.15, 0.28)	0.16 (0.12, 0.22)	
	60	0.03 (0.02, 0.04)	0.08 (0.06, 0.10)	0.15 (0.12, 0.19)	0.19 (0.15, 0.27)	0.15 (0.11, 0.22)	
	70	0.05 (0.03, 0.07)	0.13 (0.10, 0.17)	- (-, -)	0.17 (0.12, 0.26)	0.15 (0.11, 0.22)	
	80	0.10 (0.07, 0.14)	- (-, -)	- (-, -)	0.15 (0.10, 0.25)	0.13 (0.08, 0.22)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.22 (0.10, 0.71)	0.20 (0.11, 0.37)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.22 (0.10, 0.72)	0.20 (0.11, 0.38)	
	20	0.00 (0.00, 0.03)	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.22 (0.10, 0.73)	0.20 (0.11, 0.38)	
	30	0.00 (0.00, 0.03)	0.01 (0.00, 0.04)	0.01 (0.00, 0.05)	0.22 (0.10, 0.73)	0.20 (0.11, 0.39)	
	40	0.01 (0.00, 0.04)	0.01 (0.00, 0.06)	0.07 (0.02, 0.17)	0.23 (0.10, 0.75)	0.21 (0.11, 0.39)	
	50	0.01 (0.00, 0.04)	0.06 (0.02, 0.16)	0.19 (0.08, 0.37)	0.23 (0.10, 0.76)	0.21 (0.11, 0.40)	
	60	0.06 (0.02, 0.16)	0.19 (0.08, 0.39)	0.23 (0.10, 0.49)	0.23 (0.10, 0.80)	0.20 (0.11, 0.41)	
	70	0.15 (0.05, 0.35)	0.20 (0.07, 0.47)	- (-, -)	0.20 (0.07, 0.85)	0.18 (0.08, 0.42)	
	80	0.06 (0.00, 0.34)	- (-, -)	- (-, -)	0.06 (0.00, 0.93)	0.16 (0.04, 0.49)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.21 (0.17, 0.27)	0.16 (0.14, 0.19)	
	10	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.21 (0.17, 0.27)	0.17 (0.14, 0.20)	
	20	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.00, 0.01)	0.21 (0.17, 0.28)	0.17 (0.14, 0.20)	
	30	0.00 (0.00, 0.00)	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.21 (0.17, 0.28)	0.17 (0.14, 0.20)	
	40	0.00 (0.00, 0.01)	0.01 (0.01, 0.02)	0.04 (0.03, 0.05)	0.21 (0.17, 0.28)	0.17 (0.14, 0.20)	
	50	0.01 (0.01, 0.01)	0.03 (0.02, 0.05)	0.10 (0.08, 0.12)	0.21 (0.17, 0.28)	0.17 (0.14, 0.20)	
	60	0.03 (0.02, 0.04)	0.09 (0.07, 0.11)	0.16 (0.13, 0.20)	0.21 (0.16, 0.28)	0.17 (0.14, 0.20)	
	70	0.07 (0.05, 0.09)	0.15 (0.12, 0.19)	- (-, -)	0.20 (0.15, 0.27)	0.16 (0.13, 0.19)	
	80	0.10 (0.07, 0.14)	- (-, -)	- (-, -)	0.16 (0.11, 0.25)	0.14 (0.11, 0.18)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VIII-10
ESOPHAGUS CANCER (Invasive)

SEER INCIDENCE AND U.S. MORTALITY
AGE-ADJUSTED RATES AND TRENDS^a
By Race/Ethnicity and Sex

SEER Incidence

RACE/ETHNICITY	Rate 2000-2004 ^b Rate per 100,000 persons			Trend 1995-2004 ^c APC (%)		
	Total	Males	Females	Total	Males	Females
	All Races	4.6	7.9	2.0	0.2	0.5
White	4.6	8.0	1.9	1.3*	1.7*	0.0
White Hispanic ^d	3.1	5.5	1.2	-1.2	-0.9	-1.7
White Non-Hispanic ^d	4.8	8.3	2.1	1.7*	2.0*	0.3
Black	6.2	10.4	3.2	-4.6*	-4.9*	-3.7*
Asian/Pacific Islander	2.5	4.2	1.2	-1.9	-2.4	1.5
Amer Ind/Alaska Nat ^e	4.1	7.0	-	-	-	-
Hispanic ^d	3.0	5.4	1.2	-1.1	-0.9	-1.3

U.S. Mortality^f

RACE/ETHNICITY	Rate 2000-2004 Rate per 100,000 persons			Trend 1995-2004 APC (%)		
	Total	Males	Females	Total	Males	Females
	All Races	4.4	7.8	1.8	0.3*	0.4*
White	4.3	7.7	1.7	1.2*	1.3*	0.2
White Hispanic ^d	2.5	4.4	1.0	-2.1*	-2.5*	-1.7
White Non-Hispanic ^d	4.5	7.9	1.7	1.5*	1.5*	0.4
Black	6.0	10.2	3.0	-4.2*	-4.7*	-2.9*
Asian/Pacific Islander	1.8	3.1	0.8	-2.6	-2.1	-3.8
Amer Ind/Alaska Nat						
Total U.S.	2.9	5.2	1.2	0.9	0.9	-
CHSDA Counties	3.8	6.7	1.5	4.5*	4.6*	-
Non-CHSDA Counties	1.9	3.1	-	-	-	-
Hispanic ^d	2.4	4.2	1.0	-2.2*	-2.5*	-1.8

- The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Incidence data used in calculating the rates are from the 17 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^c Incidence data used in calculating the trends are from the 13 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).

^d Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. The 2000-2004 Hispanic and Non-Hispanic death rates exclude deaths from Minnesota, New Hampshire and North Dakota. The 1995-2004 Hispanic and Non-Hispanic mortality trends exclude deaths from Maine, Minnesota, New Hampshire, North Dakota, and Oklahoma.

^e Incidence data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^f Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

* The APC is significantly different from zero ($p < .05$).

Table VIII-11
ESOPHAGUS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE RATES^a
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
SEER INCIDENCE RATES^a, 2000-2004									
REGISTRY									
Atlanta & Rural Georgia	4.8	7.9	2.4	4.1	7.0	1.8	6.4	10.7	3.4
Atlanta	4.7	7.7	2.3	4.1	7.0	1.8	6.1	10.2	3.3
Rural Georgia	7.4	11.7	-	4.3	-	-	13.0	22.0	-
California	4.1	6.9	1.9	4.3	7.3	1.9	5.0	8.1	2.8
Greater Bay Area	4.0	6.6	1.9	4.4	7.4	1.9	5.1	7.5	3.2
San Francisco-Oakland	4.0	6.7	1.9	4.4	7.4	2.0	5.0	7.5	3.1
San Jose-Monterey	3.9	6.5	1.8	4.2	7.3	1.8	-	-	-
Los Angeles	3.7	6.2	1.9	3.8	6.4	1.8	5.1	8.3	2.9
Greater California	4.3	7.4	1.9	4.5	7.6	1.9	4.8	8.5	2.1
Connecticut	5.6	9.6	2.6	5.6	9.5	2.5	7.4	12.0	4.1
Detroit	5.6	9.5	2.6	5.2	9.0	2.2	7.3	12.0	4.0
Hawaii	3.6	6.4	1.2	3.8	6.5	-	-	-	-
Iowa	5.1	9.2	1.8	5.2	9.3	1.8	-	-	-
Kentucky	5.0	9.0	1.8	4.9	8.7	1.7	7.9	14.6	3.3
Louisiana	5.0	8.8	2.0	4.6	8.2	1.7	6.0	10.7	2.7
New Jersey	5.1	8.6	2.3	4.9	8.4	2.2	7.2	12.7	3.5
New Mexico	4.1	7.2	1.5	4.3	7.4	1.6	-	-	-
Seattle-Puget Sound	5.2	8.7	2.3	5.3	9.0	2.3	4.6	-	-
Utah	3.2	5.8	1.0	3.3	5.9	1.0	-	-	-
9 SEER Areas ^b	4.8	8.2	2.1	4.8	8.3	2.0	6.4	10.1	3.6
11 SEER Areas ^b	4.5	7.7	2.0	4.6	7.9	2.0	6.0	9.6	3.5
13 SEER Areas ^b	4.5	7.7	2.0	4.6	7.9	2.0	6.1	9.7	3.5
17 SEER Areas ^b	4.6	7.9	2.0	4.6	8.0	1.9	6.2	10.4	3.2

^a Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130)
^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.
The SEER 11 areas comprise the SEER 9 areas plus San Jose-Monterey and Los Angeles.
The SEER 13 areas comprise the SEER 11 areas plus the Alaska Native Registry and Rural Georgia.
The SEER 17 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VIII-12
ESOPHAGUS CANCER (Invasive)

AGE-ADJUSTED SEER DEATH RATES^a
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER DEATH RATES^a, 2000-2004</u>									
REGISTRY									
Atlanta & Rural Georgia	4.1	6.9	2.0	3.4	5.7	1.6	6.0	10.9	2.8
Atlanta	4.0	6.8	1.9	3.4	5.7	1.6	5.7	10.7	2.5
Rural Georgia	6.8	10.3	-	4.1	-	-	12.0	17.8	-
California	3.8	6.5	1.7	4.0	6.9	1.7	4.5	7.1	2.6
Greater Bay Area	3.6	6.0	1.7	4.0	6.7	1.8	4.9	7.0	3.3
San Francisco-Oakland	3.7	6.2	1.8	4.0	6.8	1.8	5.0	7.1	3.4
San Jose-Monterey	3.4	5.6	1.7	3.8	6.3	1.8	-	-	-
Los Angeles	3.3	5.6	1.5	3.4	5.9	1.5	4.4	7.1	2.6
Greater California	4.1	7.0	1.7	4.3	7.4	1.8	4.1	7.3	1.9
Connecticut	4.5	8.0	1.9	4.4	7.8	1.7	6.2	10.2	3.3
Detroit	5.0	9.2	1.8	5.0	9.4	1.7	5.0	8.9	2.3
Hawaii	2.7	5.1	0.8	3.5	6.3	-	-	-	-
Iowa	4.6	8.2	1.7	4.6	8.2	1.7	-	-	-
Kentucky	4.4	8.3	1.4	4.3	8.0	1.4	6.9	13.5	-
Louisiana	4.5	8.2	1.7	4.2	7.7	1.5	5.5	10.0	2.3
New Jersey	4.3	7.6	1.9	4.3	7.5	1.8	5.6	10.2	2.4
New Mexico	3.9	7.1	1.3	4.0	7.3	1.3	-	-	-
Seattle-Puget Sound	4.8	8.4	1.9	5.0	8.8	1.9	4.2	-	-
Utah	2.9	5.5	0.8	3.0	5.6	0.8	-	-	-
9 SEER Areas ^b	4.2	7.5	1.7	4.3	7.7	1.6	5.2	8.8	2.7
11 SEER Areas ^b	4.0	7.0	1.6	4.1	7.3	1.6	5.0	8.3	2.7
13 SEER Areas ^b	4.0	7.0	1.6	4.1	7.3	1.6	5.1	8.4	2.7
17 SEER Areas ^b	4.1	7.2	1.7	4.2	7.4	1.6	5.2	9.0	2.5
Total U.S.	4.4	7.8	1.8	4.3	7.7	1.7	6.0	10.2	3.0

^a NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 11 areas comprise the SEER 9 areas plus San Jose-Monterey and Los Angeles.

The SEER 13 areas comprise the SEER 11 areas plus the Alaska Native Registry and Rural Georgia.

The SEER 17 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VIII-13
ESOPHAGUS CANCER (INVASIVE)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2000-2004

Males and Females

<u>State</u>	<u>Rate</u>	<u>SE</u>	<u>Rank</u>	<u>PD</u>	<u>State</u>	<u>Rate</u>	<u>SE</u>	<u>Rank</u>	<u>PD</u>
TOTAL U.S.	4.4	0.02							
High Five States									
District of Columbia	6.8 ^b	0.49	(01)	54.5 ^c					
New Hampshire	5.7 ^b	0.30	(02)	29.5 ^c					
Wyoming	5.7	0.48	(03)	29.5 ^c					
Maine	5.6 ^b	0.27	(04)	27.3 ^c					
Massachusetts	5.3 ^b	0.12	(05)	20.5 ^c					
Low Five States									
Kansas	3.8 ^b	0.17	(47)	-13.6					
California	3.8 ^b	0.05	(48)	-13.6					
Arkansas	3.6 ^b	0.16	(49)	-18.2 ^c					
Utah	2.9 ^b	0.19	(50)	-34.1 ^c					
Hawaii	2.7 ^b	0.20	(51)	-38.6 ^c					
State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.1	0.13	(40)	-6.8	Montana	4.9	0.31	(15)	11.4
Alaska	5.1	0.56	(07)	15.9 ^c	Nebraska	4.5	0.22	(23)	2.3
Arizona	4.1	0.12	(39)	-6.8	Nevada	4.6	0.22	(21)	4.5
Arkansas	3.6 ^b	0.16	(49)	-18.2 ^c	New Hampshire	5.7 ^b	0.30	(02)	29.5 ^c
California	3.8 ^b	0.05	(48)	-13.6	New Jersey	4.3	0.10	(33)	-2.3
Colorado	4.1	0.15	(41)	-6.8	New Mexico	3.9	0.21	(45)	-11.4
Connecticut	4.5	0.16	(24)	2.3	New York	4.4	0.07	(30)	0.0
Delaware	5.2	0.35	(06)	18.2 ^c	North Carolina	4.1	0.10	(38)	-6.8
District of Columbia	6.8 ^b	0.49	(01)	54.5 ^c	North Dakota	4.0	0.34	(43)	-9.1
Florida	4.2 ^b	0.06	(37)	-4.5	Ohio	5.1 ^b	0.09	(10)	15.9 ^c
Georgia	4.5	0.11	(26)	2.3	Oklahoma	3.9 ^b	0.15	(44)	-11.4
Hawaii	2.7 ^b	0.20	(51)	-38.6 ^c	Oregon	5.1 ^b	0.17	(08)	15.9 ^c
Idaho	4.3	0.26	(31)	-2.3	Pennsylvania	4.0 ^b	0.08	(14)	11.4
Illinois	4.7	0.09	(20)	6.8	Rhode Island	5.0	0.29	(12)	13.6
Indiana	4.8	0.12	(18)	9.1	South Carolina	5.1 ^b	0.16	(11)	15.9 ^c
Iowa	4.6	0.17	(22)	4.5	South Dakota	4.3	0.32	(35)	-2.3
Kansas	3.8 ^b	0.17	(47)	-13.6	Tennessee	4.0	0.12	(42)	-9.1
Kentucky	4.4	0.15	(29)	0.0	Texas	3.8 ^b	0.07	(46)	-13.6
Louisiana	4.5	0.14	(25)	2.3	Utah	2.9 ^b	0.19	(50)	-34.1 ^c
Maine	5.6 ^b	0.27	(04)	27.3 ^c	Vermont	4.8	0.38	(16)	9.1
Maryland	4.8	0.14	(17)	9.1	Virginia	4.3	0.11	(34)	-2.3
Massachusetts	5.3 ^b	0.12	(05)	20.5 ^c	Washington	4.7	0.13	(19)	6.8
Michigan	4.9 ^b	0.10	(13)	11.4	West Virginia	4.5	0.20	(28)	2.3
Minnesota	4.5	0.14	(27)	2.3	Wisconsin	5.1 ^b	0.13	(09)	15.9 ^c
Mississippi	4.2	0.17	(36)	-4.5	Wyoming	5.7	0.48	(03)	29.5 ^c
Missouri	4.3	0.12	(32)	-2.3					

^a NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

Table VIII-14
ESOPHAGUS CANCER (INVASIVE)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2000-2004

Males

<u>State</u>	<u>Rate</u>	<u>SE</u>	<u>Rank</u>	<u>PD</u>	<u>State</u>	<u>Rate</u>	<u>SE</u>	<u>Rank</u>	<u>PD</u>
TOTAL U.S.	7.8	0.04							
High Five States									
District of Columbia	12.1 ^b	1.02	(01)	55.1 ^c					
New Hampshire	10.1 ^b	0.61	(02)	29.5 ^c					
Maine	10.0 ^b	0.55	(03)	28.2 ^c					
Massachusetts	9.8 ^b	0.26	(04)	25.6 ^c					
Wyoming	9.4	0.95	(05)	20.5 ^c					
Low Five States									
Texas	6.7 ^b	0.13	(47)	-14.1					
California	6.5 ^b	0.10	(48)	-16.7 ^c					
Arkansas	6.1 ^b	0.31	(49)	-21.8 ^c					
Utah	5.5 ^b	0.38	(50)	-29.5 ^c					
Hawaii	5.1 ^b	0.41	(51)	-34.6 ^c					
State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.4	0.27	(38)	-5.1	Montana	8.2	0.61	(21)	5.1
Alaska	8.6	1.12	(14)	10.3	Nebraska	8.3	0.46	(18)	6.4
Arizona	7.2	0.24	(41)	-7.7	Nevada	7.8	0.42	(30)	0.0
Arkansas	6.1 ^b	0.31	(49)	-21.8 ^c	New Hampshire	10.1 ^b	0.61	(02)	29.5 ^c
California	6.5 ^b	0.10	(48)	-16.7 ^c	New Jersey	7.6	0.20	(34)	-2.6
Colorado	7.3	0.31	(40)	-6.4	New Mexico	7.1	0.43	(45)	-9.0
Connecticut	8.0	0.32	(25)	2.6	New York	7.6	0.14	(33)	-2.6
Delaware	9.4	0.73	(06)	20.5 ^c	North Carolina	7.2	0.21	(44)	-7.7
District of Columbia	12.1 ^b	1.02	(01)	55.1 ^c	North Dakota	7.4	0.68	(37)	-5.1
Florida	7.2 ^b	0.12	(42)	-7.7	Ohio	9.1 ^b	0.19	(08)	16.7 ^c
Georgia	7.9	0.23	(26)	1.3	Oklahoma	7.2	0.30	(43)	-7.7
Hawaii	5.1 ^b	0.41	(51)	-34.6 ^c	Oregon	9.0 ^b	0.34	(09)	15.4 ^c
Idaho	7.4	0.51	(36)	-5.1	Pennsylvania	8.8 ^b	0.17	(12)	12.8
Illinois	8.3	0.18	(17)	6.4	Rhode Island	8.4	0.58	(16)	7.7
Indiana	8.6	0.26	(13)	10.3	South Carolina	9.0 ^b	0.32	(10)	15.4 ^c
Iowa	8.2	0.34	(23)	5.1	South Dakota	7.7	0.64	(32)	-1.3
Kansas	6.8	0.34	(46)	-12.8	Tennessee	7.4	0.25	(39)	-5.1
Kentucky	8.3	0.31	(19)	6.4	Texas	6.7 ^b	0.13	(47)	-14.1
Louisiana	8.2	0.30	(22)	5.1	Utah	5.5 ^b	0.38	(50)	-29.5 ^c
Maine	10.0 ^b	0.55	(03)	28.2 ^c	Vermont	7.8	0.74	(29)	0.0
Maryland	8.4	0.28	(15)	7.7	Virginia	7.4	0.23	(35)	-5.1
Massachusetts	9.8 ^b	0.26	(04)	25.6 ^c	Washington	8.2	0.26	(20)	5.1
Michigan	9.0 ^b	0.20	(11)	15.4 ^c	West Virginia	7.9	0.42	(28)	1.3
Minnesota	7.9	0.27	(27)	1.3	Wisconsin	9.1 ^b	0.27	(07)	16.7 ^c
Mississippi	8.1	0.37	(24)	3.8	Wyoming	9.4	0.95	(05)	20.5 ^c
Missouri	7.7	0.24	(31)	-1.3					

^a NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

Table VIII-15
ESOPHAGUS CANCER (INVASIVE)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2000-2004

Females

<u>State</u>	<u>Rate</u>	<u>SE</u>	<u>Rank</u>	<u>PD</u>	<u>State</u>	<u>Rate</u>	<u>SE</u>	<u>Rank</u>	<u>PD</u>
TOTAL U.S.	1.8	0.01			Montana	2.0	0.27	(08)	11.1
High Five States					Nebraska	1.7	0.18	(35)	-5.6
District of Columbia	2.9	0.43	(01)	61.1 ^c	Nevada	1.7	0.18	(27)	-5.6
Wyoming	2.6	0.44	(02)	44.4 ^c	New Hampshire	2.1	0.24	(05)	16.7 ^c
Vermont	2.3	0.36	(03)	27.8 ^c	New Jersey	1.9	0.08	(22)	5.6
Rhode Island	2.3	0.26	(04)	27.8 ^c	New Mexico	1.3	0.16	(47)	-27.8 ^c
New Hampshire	2.1	0.24	(05)	16.7 ^c	New York	2.0 ^b	0.06	(12)	11.1
Low Five States					North Carolina	1.8	0.09	(23)	0.0
New Mexico	1.3	0.16	(47)	-27.8 ^c	North Dakota	1.3	0.25	(48)	-27.8 ^c
North Dakota	1.3	0.25	(48)	-27.8 ^c	Ohio	2.0 ^b	0.08	(07)	11.1
Oklahoma	1.2 ^b	0.11	(49)	-33.3 ^c	Oklahoma	1.2 ^b	0.11	(49)	-33.3 ^c
Utah	0.8 ^b	0.13	(50)	-55.6 ^c	Oregon	1.9	0.14	(17)	5.6
Hawaii	0.8 ^b	0.15	(51)	-55.6 ^c	Pennsylvania	2.0	0.07	(14)	11.1
Alabama	1.6	0.11	(39)	-11.1	Rhode Island	2.3	0.26	(04)	27.8 ^c
Alaska	2.1	0.50	(06)	16.7 ^c	South Carolina	2.0	0.13	(09)	11.1
Arizona	1.6	0.10	(37)	-11.1	South Dakota	1.5	0.25	(42)	-16.7 ^c
Arkansas	1.6	0.14	(40)	-11.1	Tennessee	1.5	0.09	(43)	-16.7 ^c
California	1.7	0.04	(34)	-5.6	Texas	1.6	0.06	(38)	-11.1
Colorado	1.5	0.12	(41)	-16.7 ^c	Utah	0.8 ^b	0.13	(50)	-55.6 ^c
Connecticut	1.9	0.13	(21)	5.6	Vermont	2.3	0.36	(03)	27.8 ^c
Delaware	2.0	0.29	(11)	11.1	Virginia	1.8	0.10	(26)	0.0
District of Columbia	2.9	0.43	(01)	61.1 ^c	Washington	1.9	0.11	(18)	5.6
Florida	1.7	0.05	(30)	-5.6	West Virginia	1.7	0.17	(28)	-5.6
Georgia	1.9	0.10	(20)	5.6	Wisconsin	1.9	0.11	(19)	5.6
Hawaii	0.8 ^b	0.15	(51)	-55.6 ^c	Wyoming	2.6	0.44	(02)	44.4 ^c
Idaho	1.6	0.22	(36)	-11.1					
Illinois	2.0	0.07	(15)	11.1					
Indiana	1.8	0.10	(24)	0.0					
Iowa	1.7	0.13	(32)	-5.6					
Kansas	1.4	0.13	(44)	-22.2 ^c					
Kentucky	1.4	0.11	(45)	-22.2 ^c					
Louisiana	1.7	0.12	(31)	-5.6					
Maine	2.0	0.21	(13)	11.1					
Maryland	2.0	0.12	(10)	11.1					
Massachusetts	1.9	0.10	(16)	5.6					
Michigan	1.8	0.08	(25)	0.0					
Minnesota	1.7	0.11	(29)	-5.6					
Mississippi	1.3 ^b	0.13	(46)	-27.8 ^c					
Missouri	1.7	0.10	(33)	-5.6					

^a NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

Table VIII-16
ESOPHAGUS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2004
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 ^e	0 to <29 ^e	>=29 ^g	Complete ^h
<u>Race/Sex</u>									
All Races ^b	17,196	4,268	1,939	764	288	23,242	24,565	220	24,785
Males	13,312	3,238	1,367	449	216	17,785	18,653	110	18,763
Females	3,884	1,030	572	315	72	5,457	5,912	110	6,022
White ^b	15,061	3,783	1,578	642	256	20,301	21,406	182	21,588
Males	11,838	2,941	1,095	385	198	15,765	16,529	103	16,632
Females	3,223	842	483	257	58	4,536	4,877	79	4,956
Black ^b	1,725	409	308	112	29	2,407	2,609	56	2,665
Males	1,151	236	231	64	15	1,596	1,697	18	1,715
Females	574	173	77	48	14	811	912	38	950
Asian/Pacific Islander ^c	266	76	+	+	+	386	+	+	+
Males	201	57	+	+	+	292	+	+	+
Females	65	19	+	+	+	94	+	+	+
Hispanic ^d	551	154	+	+	+	804	+	+	+
Males	462	110	+	+	+	656	+	+	+
Females	89	44	+	+	+	148	+	+	+

Estimated prevalence percent^a on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)								Age-Adjusted ^f All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	
<u>Race/Sex</u>										
All Races ^c	0.0061%	-	-	0.0001%	0.0005%	0.0022%	0.0104%	0.0254%	0.0376%	0.0263%
Males	0.0094%	-	-	-	0.0008%	0.0035%	0.0172%	0.0436%	0.0642%	0.0465%
Females	0.0029%	-	-	-	-	0.0010%	0.0040%	0.0091%	0.0175%	0.0153%
White ^c	0.0068%	-	-	0.0001%	0.0006%	0.0023%	0.0108%	0.0269%	0.0407%	0.0270%
Males	0.0105%	-	-	-	0.0010%	0.0037%	0.0185%	0.0464%	0.0699%	0.0473%
Females	0.0031%	-	-	-	-	0.0009%	0.0033%	0.0090%	0.0182%	0.0161%
Black ^c	0.0053%	-	-	-	-	0.0029%	0.0149%	0.0317%	0.0322%	0.0242%
Males	0.0069%	-	-	-	-	0.0037%	0.0197%	0.0467%	0.0526%	0.0390%
Females	0.0038%	-	-	-	-	0.0022%	0.0109%	0.0199%	0.0189%	0.0171%
Asian/Pacific Islander ^c	0.0031%	-	-	-	-	0.0009%	0.0043%	0.0105%	0.0204%	0.0229%
Males	0.0048%	-	-	-	-	0.0017%	0.0058%	0.0209%	0.0328%	0.0470%
Females	0.0015%	-	-	-	-	-	0.0030%	-	0.0115%	0.0069%
Hispanic ^d	0.0017%	-	-	-	0.0004%	0.0010%	0.0048%	0.0124%	0.0220%	0.0142%
Males	0.0027%	-	-	-	0.0006%	0.0015%	0.0079%	0.0229%	0.0446%	0.0246%
Females	0.0007%	-	-	-	-	-	0.0018%	0.0037%	0.0059%	0.0083%

^a US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^{f g h} Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^(g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). ^(h)Complete prevalence is obtained by summing 0 to <29 and >=29.

⁻ Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

⁺ Not available.

Table VIII-17

ESOPHAGUS CANCER (Invasive)

Percent Distribution and Counts by Histology among Histologically Confirmed Cases, 2001-2004
Both Sexes by Race

Histology	All Races		White		Black	
	Count	Percent	Count	Percent	Count	Percent
Adenocarcinoma	6,774	54.0%	6,439	61.6%	166	11.6%
Adenocarcinoma, NOS (8140)	5,950	47.5%	5,662	54.1%	144	10.1%
Mucinous and mucin-producing adenocarcinoma (8480-8481)	224	1.8%	209	2.0%	-	-
Other adenocarcinomas ^a	600	4.8%	568	5.4%	-	-
Squamous cell carcinoma	4,860	38.8%	3,240	31.0%	1,180	82.7%
Squamous cell carcinoma (8070-8077)	4,798	38.3%	3,192	30.5%	1,171	82.1%
Other specified squamous carcinomas (8050-8069, 8078-8089, 8094)	62	0.5%	48	0.5%	-	-
Other malignant histologies	901	7.2%	781	7.5%	80	5.6%
Total	12,535	100.0%	10,460	100.0%	1,426	100.0%

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Percents may not sum to 100 due to rounding.

^a Statistic not shown due to fewer than 16 cases during the time period.

^a Includes histologies 8141-8389, 8401, 8408, 8410, 8411, 8413, 8441, 8450, 8460, 8470-8482, 8490, 8500, 8503, 8504, 8510, 8520, 8525, 8530, 8571-8574, 8576, 9070, 9110.

SEER Incidence, Delay Adjusted Incidence and US Death Rates^a Esophagus Cancer, by Race and Sex




Figure VIII-1

^a Source: SEER 9 areas and NCHS public use data file for the total US. Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines and APCs are calculated using the Joinpoint Regression Program Version 3.0, April 2005, National Cancer Institute.


The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

SEER Incidence and US Death Rates^a

Esophagus Cancer, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2004
and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2004



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from NCHS public use data file for the total US.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 3.0, April 2005, National Cancer Institute.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from Connecticut, Maine, Maryland, Minnesota, New Hampshire, New York, North Dakota, Oklahoma, and Vermont.

Figure VIII-2