

Table XXXI-1
BENIGN BRAIN TUMORS

A Comparison of Benign/Borderline and Malignant Brain Tumors
Counts, Percents and Age-Adjusted Incidence Rates^a by WHO Histology Grouping, 2004

| WHO Histology Grouping | Benign/Borderline | | | Malignant | | |
|---|-------------------|---------|------|-----------|---------|------|
| | Count | Percent | Rate | Count | Percent | Rate |
| BRAIN | | | | | | |
| Diffuse astrocytoma (protoplasma, fibrillary) | - | - | - | 64 | 1.2% | 0.1 |
| Anaplastic astrocytoma | - | - | - | 301 | 5.8% | 0.4 |
| Glioblastoma | - | - | - | 2,324 | 44.4% | 3.1 |
| Pilocytic astrocytoma | - | - | - | 201 | 3.8% | 0.3 |
| Unique astrocytoma variants | 34 | 0.4% | 0.0 | 21 | 0.4% | 0.0 |
| Oligodendroglioma | - | - | - | 232 | 4.4% | 0.3 |
| Anaplastic oligodendroglioma | - | - | - | 104 | 2.0% | 0.1 |
| Ependymoma/anaplastic ependymoma | - | - | - | 94 | 1.8% | 0.1 |
| Ependymoma variants | - | - | - | - | - | - |
| Mixed glioma | - | - | - | 128 | 2.4% | 0.2 |
| Astrocytoma, NOS | - | - | - | 325 | 6.2% | 0.4 |
| Glioma, NOS | - | - | - | 241 | 4.6% | 0.3 |
| Choroid plexus | 36 | 0.5% | 0.0 | - | - | - |
| Neuroepithelial | - | - | - | - | - | - |
| Neuronal/glial, neuronal and mixed | 103 | 1.3% | 0.1 | - | - | - |
| Embryonal/primitive/medulloblastoma | - | - | - | 147 | 2.8% | 0.2 |
| Nerve sheath | 112 | 1.5% | 0.1 | - | - | - |
| Meningioma | 534 | 6.9% | 0.7 | 28 | 0.5% | 0.0 |
| Hemangioma and hemangioblastoma | 130 | 1.7% | 0.2 | - | - | - |
| Germ cell tumors, cysts, and heterotopias | - | - | - | 23 | 0.4% | 0.0 |
| Chordoma/chondrosarcoma | - | - | - | - | - | - |
| Craniopharyngioma | - | - | - | - | - | - |
| Neoplasm, unspecified | 131 | 1.7% | 0.2 | 208 | 4.0% | 0.3 |
| Other histologies ^b | - | - | - | - | - | - |
| MENINGES | | | | | | |
| Neuroepithelial | - | - | - | - | - | - |
| Nerve sheath | - | - | - | - | - | - |
| Meningioma | 3,282 | 42.7% | 4.4 | 62 | 1.2% | 0.1 |
| Hemangioma and hemangioblastoma | - | - | - | - | - | - |
| Chordoma/chondrosarcoma | - | - | - | - | - | - |
| Neoplasm, unspecified | - | - | - | - | - | - |
| Other histologies ^b | - | - | - | - | - | - |
| CRANIAL NERVES AND OTHER CNS | | | | | | |
| Glioblastoma | - | - | - | - | - | - |
| Pilocytic astrocytoma | - | - | - | - | - | - |
| Astrocytoma, NOS | - | - | - | - | - | - |
| Glioma, NOS | - | - | - | 21 | 0.4% | 0.0 |
| Neuronal/glial, neuronal and mixed | - | - | - | - | - | - |
| Nerve sheath | 970 | 12.6% | 1.3 | - | - | - |
| Meningioma | - | - | - | - | - | - |
| Hemangioma and hemangioblastoma | - | - | - | - | - | - |
| Germ cell tumors, cysts, and heterotopias | - | - | - | - | - | - |
| Chordoma/chondrosarcoma | - | - | - | - | - | - |
| Neoplasm, unspecified | - | - | - | - | - | - |
| Other histologies ^b | - | - | - | - | - | - |

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Count and percent may not sum to total due to suppression and rounding.

- Statistic not shown due to fewer than 16 cases during the time period.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

The histologic groupings exclude lymphomas, mesotheliomas, and Kaposi Sarcoma.

^b Includes histologies not included in current groupings.

Table XXXI-1 - continued
BENIGN BRAIN TUMORS

A Comparison of Benign/Borderline and Malignant Brain Tumors
 Counts, Percents and Age-Adjusted Incidence Rates^a by WHO Histology Grouping, 2004

| WHO Histology Grouping | Benign/Borderline | | | Malignant | | |
|---|-------------------|---------------|-------------|--------------|---------------|------------|
| | Count | Percent | Rate | Count | Percent | Rate |
| TUMORS OF THE SELLAR REGION | | | | | | |
| Glioblastoma | - | - | - | - | - | - |
| Pilocytic astrocytoma | - | - | - | - | - | - |
| Ependymoma/anaplastic ependymoma | - | - | - | - | - | - |
| Glioma, NOS | - | - | - | - | - | - |
| Pineal parenchymal | - | - | - | 19 | 0.4% | 0.0 |
| Embryonal/primitive/medulloblastoma | - | - | - | - | - | - |
| Nerve sheath | - | - | - | - | - | - |
| Meningioma | - | - | - | - | - | - |
| Germ cell tumors, cysts, and heterotopias | - | - | - | 32 | 0.6% | 0.0 |
| Chordoma/chondrosarcoma | - | - | - | - | - | - |
| Pituitary | 1,641 | 21.3% | 2.2 | - | - | - |
| Craniopharyngioma | 87 | 1.1% | 0.1 | - | - | - |
| Neoplasm, unspecified | 29 | 0.4% | 0.0 | - | - | - |
| Other histologies ^b | - | - | - | - | - | - |
| SPINAL CORD | | | | | | |
| Diffuse astrocytoma (protoplasma, fibrillary) | - | - | - | - | - | - |
| Anaplastic astrocytoma | - | - | - | - | - | - |
| Glioblastoma | - | - | - | - | - | - |
| Pilocytic astrocytoma | - | - | - | 16 | 0.3% | 0.0 |
| Unique astrocytoma variants | - | - | - | - | - | - |
| Ependymoma/anaplastic ependymoma | - | - | - | 96 | 1.8% | 0.1 |
| Ependymoma variants | 61 | 0.8% | 0.1 | - | - | - |
| Mixed glioma | - | - | - | - | - | - |
| Astrocytoma, NOS | - | - | - | 16 | 0.3% | 0.0 |
| Glioma, NOS | - | - | - | - | - | - |
| Neuronal/glial, neuronal and mixed | - | - | - | - | - | - |
| Embryonal/primitive/medulloblastoma | - | - | - | - | - | - |
| Nerve sheath | 143 | 1.9% | 0.2 | - | - | - |
| Meningioma | 29 | 0.4% | 0.0 | - | - | - |
| Hemangioma and hemangioblastoma | 24 | 0.3% | 0.0 | - | - | - |
| Chordoma/chondrosarcoma | - | - | - | - | - | - |
| Neoplasm, unspecified | - | - | - | - | - | - |
| Other histologies ^b | - | - | - | - | - | - |
| SPINAL MENINGES | | | | | | |
| Ependymoma/anaplastic ependymoma | - | - | - | - | - | - |
| Neuronal/glial, neuronal and mixed | - | - | - | - | - | - |
| Nerve sheath | 20 | 0.3% | 0.0 | - | - | - |
| Meningioma | 207 | 2.7% | 0.3 | - | - | - |
| Hemangioma and hemangioblastoma | - | - | - | - | - | - |
| Neoplasm, unspecified | - | - | - | - | - | - |
| Other histologies ^b | - | - | - | - | - | - |
| TOTAL | 7,695 | 100.0% | 10.2 | 5,233 | 100.0% | 7.0 |

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

- Count and percent may not sum to total due to suppression and rounding.
 - Statistic not shown due to fewer than 16 cases during the time period.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
 The histologic groupings exclude lymphomas, mesotheliomas, and Kaposi Sarcoma.

^b Includes histologies not included in current groupings.

Table XXXI-2
BENIGN BRAIN TUMORS

Age-Adjusted SEER Incidence Rates^a for Benign Brain Tumors, 2004
By Race, Sex and Age

Brain and Other Nervous System(Benign/Borderline)
Excluding Endocrine Glands and Related Structures

| Age | All Races | | | Whites | | | Blacks | | |
|----------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| All Ages | 7.9 | 5.8 | 9.7 | 7.9 | 5.8 | 9.6 | 7.9 | 5.2 | 9.9 |
| <20 | 0.8 | 0.9 | 0.8 | 0.8 | 0.9 | 0.8 | 0.7 | - | - |
| 20-44 | 4.0 | 3.1 | 4.8 | 3.9 | 3.2 | 4.7 | 3.9 | 2.4 | 5.2 |
| 45-64 | 12.5 | 8.7 | 16.1 | 12.6 | 8.7 | 16.3 | 11.0 | 6.6 | 14.7 |
| 65-74 | 22.9 | 16.8 | 28.0 | 22.9 | 16.9 | 27.9 | 24.3 | 16.5 | 30.0 |
| 75-84 | 30.7 | 21.8 | 36.8 | 30.2 | 21.3 | 36.3 | 36.5 | 30.5 | 40.2 |
| 85+ | 34.3 | 25.1 | 38.6 | 34.6 | 26.3 | 38.5 | 37.1 | - | 48.8 |

Brain and Other Nervous System(Benign/Borderline)
Including Endocrine Glands and Related Structures

| Age | All Races | | | Whites | | | Blacks | | |
|----------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| All Ages | 10.2 | 8.1 | 12.1 | 10.1 | 8.0 | 11.9 | 11.7 | 8.9 | 13.8 |
| <20 | 1.3 | 1.2 | 1.4 | 1.3 | 1.2 | 1.4 | 1.2 | - | 1.4 |
| 20-44 | 6.2 | 4.7 | 7.7 | 6.1 | 4.9 | 7.5 | 6.7 | 4.5 | 8.7 |
| 45-64 | 15.9 | 12.3 | 19.3 | 15.7 | 12.1 | 19.2 | 16.7 | 13.1 | 19.7 |
| 65-74 | 28.3 | 23.3 | 32.4 | 27.6 | 22.9 | 31.6 | 35.7 | 28.8 | 40.6 |
| 75-84 | 36.1 | 29.5 | 40.6 | 35.1 | 28.6 | 39.6 | 48.3 | 44.6 | 50.7 |
| 85+ | 36.5 | 28.2 | 40.4 | 36.6 | 29.7 | 39.8 | 43.8 | - | 56.4 |

Source: SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

- Rate not shown due to fewer than 16 cases during the time period.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Table XXXI-3

Brain Tumor Histology Groupings^a

| Site Group | ICD-0-3 Morphology |
|--|---|
| TUMORS OF NEUROEPITHELIAL TISSUE | |
| Diffuse astrocytoma (protoplasma, fibrillary) | 9410, 9420 |
| Anaplastic astrocytoma | 9401, 9411 |
| Glioblastoma | 9440, 9441, 9442/3 |
| Pilocytic astrocytoma | 9421 |
| Unique astrocytoma variants | 9383, 9384, 9424 |
| Oligodendroglioma | 9450 |
| Anaplastic oligodendroglioma | 9451, 9460 |
| Ependymoma/anaplastic ependymoma | 9391, 9392, 9393 |
| Ependymoma variants | 9394 |
| Mixed glioma | 9382 |
| Astrocytoma, NOS | 9400 |
| Glioma, NOS | 9380 |
| Choroid plexus | 9390 |
| Neuroepithelial | 9381, 9423, 9430, 9444 |
| Benign and Malignant neuronal/glial, neuronal and mixed | 8680/1, 8680/3, 8681, 8682, 8690, 8693, 9412, 9413, 9442/1, 9490-9493, 9500, 9505, 9506, 9522, 9523 |
| Pineal parenchymal | 9360, 9361, 9362 |
| Embryonal/primitive/medulloblastoma | 8963, 9363, 9364, 9470-9474, 9501-9503, 9508 |
| TUMORS OF CRANIAL AND SPINAL NERVES | |
| Nerve sheath, benign and malignant | 9540, 9541, 9550, 9560, 9561, 9570, 9571 |
| TUMORS OF THE MENINGES | |
| Meningioma, benign and malignant | 9530-9534, 9537-9539 |
| Other mesenchymal, benign and malignant | 8324, 8728, 8800-8806, 8810, 8815, 8824, 8830, 8831, 8850, 8851, 8857, 8861, 8890, 8897, 8900, 8910, 8920, 8990, 9040, 9150, 9180, 9210, 9241, 9260, 9480 |
| Hemangioblastoma and hemangioblastoma | 9120-9123, 9125, 9130, 9131, 9133, 9140, 9161, 9535 |
| GERM CELL TUMORS AND CYSTS | |
| Germ cell tumors, cysts, and heterotopias | 8020, 9060, 9061, 9064, 9065, 9070-9072, 9080-9085, 9100 |
| LOCAL EXTENSIONS FROM REGIONAL TUMORS | |
| Chordoma/chondrosarcoma | 9220, 9231, 9240, 9370-9373 |
| TUMORS OF THE SELLAR REGION | |
| Pituitary | 8022, 8040, 8140, 8146, 8246, 8260, 8270, 8271, 8272, 8280, 8281, 8290, 8300, 8310, 8323, 8333, 8334, 9582 |
| Craniopharyngioma | 9350, 9351, 9352 |
| UNCLASSIFIED TUMORS | |
| Neoplasm, unspecified | 8000-8005, 8010, 8013, 8021 |
| All other | any remaining histologies not included in above categories |

Source: Modified from CBTRUS (2005). Statistical Report: Primary Brain Tumors in the United States, 1998-2002. Published by the Central Brain Tumor Registry of the United States.

^a The histologic groupings exclude lymphomas, mesotheliomas, and Kaposi Sarcoma.