

Table VII-4
CORPUS AND UTERUS, NOS CANCER (Invasive)

SEER INCIDENCE^a AND U.S. DEATH^b RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	Incidence ^a			US Death ^b		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<u>AGE-ADJUSTED RATES, 2000-2004</u>						
AGE AT DIAGNOSIS/DEATH:						
All ages	23.2	24.3	19.6	4.1	3.9	7.1
Under 65	14.5	15.1	10.4	1.4	1.3	2.2
65 and over	83.5	87.4	82.9	22.8	21.4	40.8
All ages(IARC world std) ^c	17.0	17.7	14.0	2.6	2.4	4.5
<u>AGE-SPECIFIC RATES, 2000-2004</u>						
AGE AT DIAGNOSIS/DEATH:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	0.3	0.3	-	-	-	-
25-29	1.3	1.3	1.4	0.1	0.0	-
30-34	3.4	3.3	3.0	0.2	0.2	0.3
35-39	6.8	6.6	5.6	0.4	0.4	0.5
40-44	12.8	13.1	8.3	0.9	0.9	1.2
45-49	23.1	23.5	13.3	1.8	1.7	2.1
50-54	42.5	45.1	25.2	3.7	3.6	5.1
55-59	66.9	70.7	44.7	6.7	6.4	9.9
60-64	81.8	85.6	76.1	11.2	10.2	21.5
65-69	89.1	93.9	89.1	16.0	14.0	34.3
70-74	86.5	91.0	88.8	20.5	18.8	41.6
75-79	86.6	90.6	88.9	23.9	22.6	42.7
80-84	81.8	85.3	72.0	28.7	27.7	45.9
85+	61.4	62.9	59.2	34.0	33.6	44.2

^a SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^c Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.