
# Trends in SEER Incidence & US Death Rates by Primary Cancer Site 1997-2006

#### Trends in SEER Incidence Rates

#### Trends in US Cancer Death Rates

Figure 1.4



Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia) and US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. For sex-specific cancer sites, the population was limited to the population of the appropriate sex.

Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

<sup>\*</sup> The Annual Percent Change is significantly different from zero (p<.05).

<sup>&</sup>lt;sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

#### Trends in SEER Incidence Rates by Primary Cancer Site 1997-2006





Figure 1.5

Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). For sex-specific cancer sites, the population was limited to the population of the appropriate sex. Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

<sup>\*</sup> The Annual Percent Change is significantly different from zero (p<.05).

<sup>&</sup>lt;sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

### Trends in US Death Rates by Primary Cancer Site 1997-2006



Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. For sex-specific cancer sites, the population was limited to the population of the appropriate sex.

<sup>\*</sup> Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). The Annual Percent Change is significantly different from zero (p<.05).

### Trends in SEER Incidence Rates by Primary Cancer Site 1997-2006




Figure 1.7


Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).

Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

The Annual Percent Change is significantly different from zero (p<.05).


Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

### Trends in US Death Rates by Primary Cancer Site 1997-2006



Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). \* The Annual Percent Change is significantly different from zero (p<.05).

## Trends in US Death Rates, 1997-2006 by Primary Cancer Site All Ages, by Race



Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).