

Table 1.22
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2006^a
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	5,168,889	17,479	39,370	75,164	144,948	331,732	720,142	1,194,219	2,645,836
Females	6,216,003	14,832	33,518	81,003	213,538	597,268	1,130,077	1,333,585	2,812,181
Oral Cavity & Pharynx									
Males	157,250	23	492	1,318	3,276	14,407	37,003	40,095	60,638
Females	87,223	101	501	1,668	3,489	8,222	15,686	18,110	39,446
Esophagus									
Males	20,742	0	0	17	130	1,166	4,085	6,436	8,907
Females	6,861	0	0	12	50	303	909	1,550	4,037
Stomach									
Males	36,725	0	5	86	649	2,426	5,764	8,519	19,274
Females	27,397	11	23	91	491	1,896	3,966	4,776	16,143
Colon & Rectum									
Males	536,944	11	46	961	5,116	22,614	69,355	116,858	321,983
Females	567,158	0	115	1,190	4,881	20,121	58,565	96,908	385,378
Liver & Intrahep									
Males	16,484	415	403	334	382	1,391	5,714	3,821	4,025
Females	8,445	424	329	318	328	843	1,678	1,630	2,896
Pancreas									
Males	14,979	0	35	69	255	1,190	3,251	4,246	5,933
Females	16,201	0	44	114	434	1,327	2,738	4,093	7,452
Larynx									
Males	73,244	0	0	11	232	2,513	10,523	19,979	39,986
Females	18,326	0	0	46	132	1,174	3,010	4,737	9,227
Lung & Bronchus									
Males	171,522	45	46	345	1,118	6,174	24,269	49,093	90,433
Females	193,474	0	32	372	1,441	8,500	25,665	51,689	105,775
Melanoma of the Skin									
Males	367,925	72	692	5,217	16,598	46,626	83,425	86,050	129,247
Females	390,763	84	980	10,591	32,264	68,250	93,268	73,818	111,509

^a U.S. 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2006 U.S. population estimates based on the average of 2005 and 2006 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.22 - continued
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2006^a
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	13,132	0	0	23	75	542	2,078	3,322	7,093
Females	2,533,193	0	104	2,335	35,060	206,965	500,267	614,770	1,173,691
Cervix									
Females	248,166	11	92	2,502	18,006	43,602	57,740	48,934	77,278
Corpus & Uterus, NOS									
Females	572,603	0	55	543	4,924	23,372	76,468	124,512	342,730
Ovary ^d									
Females	176,007	45	891	3,473	7,019	20,547	37,874	39,554	66,605
Prostate									
Males	2,177,975	45	92	69	349	16,618	187,672	552,329	1,420,801
Urinary Bladder									
Males	388,965	23	143	717	2,390	12,796	41,822	85,952	245,121
Females	138,531	33	68	414	1,120	4,555	13,137	26,664	92,539
Kidney & Renal Pelvis									
Males	156,166	1,489	2,424	2,363	4,140	13,996	31,742	40,436	59,577
Females	109,902	1,565	2,655	2,595	3,885	10,133	17,631	24,085	47,353
Hodgkin Lymphoma									
Males	82,450	129	2,058	8,923	16,247	22,575	18,055	9,050	5,412
Females	77,557	101	1,477	10,135	16,381	21,291	15,813	6,811	5,549
Non-Hodgkin Lymphoma									
Males	217,143	663	3,555	6,616	12,138	26,624	42,988	48,865	75,694
Females	202,390	360	1,529	3,938	7,562	18,662	35,213	43,070	92,057
Myeloma									
Males	32,483	0	0	51	549	2,188	6,759	9,771	13,165
Females	26,720	0	0	23	222	1,661	4,882	7,103	12,830
Leukemia									
Males	130,746	6,418	11,449	10,565	9,207	10,991	17,345	23,135	41,637
Females	101,110	5,023	9,672	9,353	7,857	7,821	11,507	14,611	35,265
Acute Lymphocytic Leuk									
Males	32,255	5,310	9,965	8,164	5,276	1,976	781	520	263
Females	26,601	4,202	8,131	6,920	4,273	1,699	654	442	281

^a U.S. 2006 cancer prevalence counts are based on 2006 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2006 U.S. population estimates based on the average of 2005 and 2006 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.