

Table 5.6
Cancer of the Cervix Uteri (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	Incidence ^a			US Death ^b		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<u>Age-Adjusted Rates, 2002-2006</u>						
Age at Diagnosis/Death:						
All ages	8.2	8.1	10.4	2.5	2.2	4.6
Under 65	7.6	7.6	8.7	1.9	1.7	3.0
65 and over	12.6	11.3	22.4	6.4	5.5	15.0
All ages (IARC world std) ^c	6.5	6.4	8.0	1.8	1.6	3.2
<u>Age-Specific Rates, 2002-2006</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	0.1	-	-	-	-	-
20-24	1.6	1.5	2.1	0.1	0.1	-
25-29	5.5	5.8	5.8	0.5	0.5	1.0
30-34	11.0	12.0	9.7	1.4	1.4	1.9
35-39	13.4	14.4	12.0	2.3	2.3	3.3
40-44	15.3	15.2	16.9	3.4	3.1	5.4
45-49	14.3	13.7	16.9	4.1	3.8	6.7
50-54	13.5	13.0	16.8	4.5	4.2	7.3
55-59	13.5	12.7	18.0	5.2	4.8	8.2
60-64	13.7	12.3	22.4	5.3	4.8	10.1
65-69	14.7	13.3	22.8	5.9	5.1	12.4
70-74	13.2	12.2	19.9	5.9	5.1	13.6
75-79	12.0	10.2	23.6	6.2	5.3	14.8
80-84	10.9	9.7	23.2	7.2	6.3	16.9
85+	10.0	8.5	23.9	8.0	6.7	22.1

^a SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.