

Table 1.22
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2007^a
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	5,353,054	17,384	39,980	78,051	145,588	336,123	749,038	1,258,920	2,727,970
Females	6,360,682	15,217	33,503	84,509	213,851	604,412	1,155,959	1,395,096	2,858,136
Oral Cavity & Pharynx									
Males	161,112	34	462	1,460	3,306	14,446	38,574	42,305	60,526
Females	88,254	112	478	1,499	3,730	8,328	16,027	18,737	39,343
Esophagus									
Males	21,782	0	0	16	110	1,155	4,158	7,094	9,250
Females	6,947	0	0	12	71	215	906	1,589	4,154
Stomach									
Males	37,297	0	5	29	621	2,421	5,852	9,147	19,223
Females	28,342	5	11	106	575	2,139	3,970	4,882	16,654
Colon & Rectum									
Males	540,636	11	46	1,079	5,377	23,458	70,527	118,887	321,251
Females	571,857	0	109	1,421	4,924	21,129	59,895	98,155	386,224
Liver & Intrahep									
Males	18,644	408	525	363	435	1,451	6,731	4,464	4,268
Females	9,109	411	346	356	427	842	1,852	1,835	3,041
Pancreas									
Males	16,057	0	23	93	227	1,283	3,502	4,577	6,353
Females	16,936	0	53	137	418	1,208	3,029	4,061	8,031
Larynx									
Males	72,246	0	0	24	215	2,470	10,579	19,435	39,524
Females	18,192	0	0	57	77	1,144	3,131	4,429	9,355
Lung & Bronchus									
Males	172,739	34	75	375	1,109	6,021	23,958	50,028	91,141
Females	197,878	11	55	332	1,269	8,266	26,154	53,184	108,608
Melanoma of the Skin									
Males	385,054	94	750	5,271	16,607	46,961	86,015	93,408	135,947
Females	408,229	72	991	11,270	32,943	70,027	96,409	80,585	115,933

^a U.S. 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2007 U.S. population estimates based on the average of 2006 and 2007 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.22 - continued
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2007^a
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	13,326	0	0	29	93	497	2,014	3,353	7,340
Females	2,591,855	0	81	2,530	35,087	208,053	505,550	645,055	1,195,498
Cervix									
Females	247,180	11	81	2,348	17,537	43,343	58,279	48,897	76,684
Corpus & Uterus, NOS									
Females	575,108	0	67	452	5,116	23,123	78,436	129,723	338,192
Ovary ^d									
Females	177,162	84	949	3,571	6,786	20,117	37,716	40,600	67,339
Prostate									
Males	2,276,112	50	69	80	387	17,453	197,739	585,651	1,474,683
Urinary Bladder									
Males	395,480	11	131	702	2,434	12,104	42,058	87,203	250,837
Females	139,756	33	46	342	1,153	4,415	13,116	27,208	93,443
Kidney & Renal Pelvis									
Males	164,839	1,413	2,428	2,472	4,088	14,777	33,714	43,718	62,228
Females	116,651	1,441	2,633	2,809	3,988	10,792	19,272	25,537	50,178
Hodgkin Lymphoma									
Males	84,583	162	2,193	9,019	16,237	22,676	18,661	9,862	5,771
Females	79,690	107	1,397	10,426	16,314	21,812	16,409	7,293	5,934
Non-Hodgkin Lymphoma									
Males	226,855	689	3,434	7,073	12,028	27,188	44,816	51,557	80,070
Females	211,470	393	1,596	4,130	7,622	18,650	37,239	45,364	96,476
Myeloma									
Males	33,522	0	5	40	488	2,380	6,786	9,878	13,944
Females	28,120	0	0	23	215	1,671	5,234	7,536	13,441
Leukemia									
Males	137,398	6,452	11,601	11,047	9,522	11,671	18,275	24,790	44,041
Females	106,874	5,226	9,722	9,893	8,060	8,514	12,311	15,613	37,535
Acute Lymphocytic Leuk									
Males	33,394	5,340	10,132	8,470	5,396	2,320	905	538	291
Females	27,389	4,366	8,155	7,282	4,309	1,895	706	455	220

^a U.S. 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2007 U.S. population estimates based on the average of 2006 and 2007 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.