

Table 1.9  
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex  
Whites, 2001-2010

Site	Incidence <sup>a</sup>			US Mortality <sup>b</sup>		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
All Sites	-0.6*	-1.1*	-0.3	-1.4*	-1.7*	-1.3*
Oral Cavity & Pharynx:						
Lip	0.5*	0.6*	0.1	-0.8*	-0.8*	-1.3*
Tongue	-4.1*	-4.7*	-2.5*	-3.8*	-5.1*	-1.3
Salivary gland	2.3*	2.6*	1.3*	-0.2	-0.1	-0.6
Floor of mouth	1.1	0.6	1.4	-0.3	-0.2	-0.9
Gum & other oral cavity	-3.2*	-4.1*	-1.4	-6.5*	-6.9*	-5.6*
Nasopharynx	-1.0*	-1.2*	-0.7	-1.7*	-2.1*	-1.4
Tonsil	-0.3	-0.7	0.0	-0.6	-0.5	-1.3
Oropharynx	3.8*	4.2*	1.6*	1.3*	1.9*	-1.3
Hypopharynx	2.0	2.4	0.2	1.8*	1.8*	1.2
Other oral cavity & pharynx	-2.5*	-2.4*	-3.6*	-3.5*	-3.2*	-5.1*
Digestive System:	-3.3*	-2.9	-5.3*	-2.2*	-2.2*	-2.7*
Esophagus	-1.1*	-1.2*	-1.2*	-1.1*	-1.1*	-1.3*
Stomach	-0.3	-0.2	-1.7*	0.0	0.1	-0.9*
Small intestine	-0.9*	-1.3*	-0.7	-3.0*	-3.4*	-2.7*
Colon & Rectum:	2.3*	2.3*	2.3*	-0.3	-0.2	-0.6
Colon	-2.8*	-3.1*	-2.5*	-3.0*	-3.1*	-3.0*
Rectum	-3.0*	-3.3*	-2.8*	-	-	-
Anus, anal canal & anorectum	-2.3*	-2.7*	-1.9*	-	-	-
Liver & intrahepatic	2.6*	2.0*	3.1*	3.3*	3.5*	3.3*
bile duct	4.2*	4.5*	3.0*	2.5*	2.6*	1.9*
Gallbladder	-0.7	-1.2	-0.4	-1.5*	-1.2*	-1.6*
Other biliary	0.9*	1.0	0.5	-3.0*	-3.3*	-2.8*
Pancreas	1.0*	0.8*	1.1*	0.6*	0.5*	0.6*
Retroperitoneum	-1.5	-1.3	-2.1	-2.5*	-1.9	-3.3*
Peritoneum, omentum &	-0.2	0.8	-0.2	0.9	0.9	1.0
mesentery	1.7*	2.2*	1.3	-1.0	0.0	-2.0
Respiratory System:	-1.4*	-2.1*	-0.6*	-1.6*	-2.4*	-0.9*
Nose, nasal cavity &	0.3	0.7	-0.5	-0.7	-0.6	-0.9
middle ear	-1.8*	-2.0*	-1.8*	-2.0*	-2.4*	-1.8*
Larynx	-1.4*	-2.2*	-0.6*	-1.6*	-2.4*	-0.8*
Lung & bronchus	-1.5	-0.9	-	-7.1*	-7.1*	-7.9*
Pleura	-0.4	-1.1	0.7	-4.2*	-4.6*	-4.0*
Trachea & other						
respiratory organs						
Bones & joints	-0.1	0.2	-0.7	-0.2	-0.2	-0.1
Soft tissue (including heart)	1.2*	1.1*	1.3*	0.6*	1.2*	-0.1
Skin (excl. basal & squamous):	1.5*	1.8*	1.2*	0.7*	1.0*	-0.1
Melanoma of the skin	1.6*	1.8*	1.3*	0.7*	1.0*	0.0
Other non-epithelial skin	1.4*	2.0*	0.4	0.8	1.0*	-0.4
Breast	-1.0*	1.2	-0.8*	-2.2*	-1.2*	-2.0*
Breast ( <i>in situ</i> )	0.7	3.0	0.9	-	-	-

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population  
(19 age groups - Census P25-1130).

<sup>a</sup> SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

<sup>b</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

\* The APC is significantly different from zero ( $p<.05$ ).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.9 - continued  
 SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex  
 Whites, 2001-2010

Site	Incidence <sup>a</sup>			US Mortality <sup>b</sup>		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
Female Genital System:	-0.4	-	-0.2	-1.3*	-	-1.0*
Cervix uteri	-1.5*	-	-1.4*	-1.5*	-	-1.3*
Corpus uteri	0.4	-	0.7*	-0.9*	-	-0.5
Uterus, NOS	0.1	-	0.4	0.8*	-	1.3*
Ovary <sup>c</sup>	-1.9*	-	-1.7*	-2.0*	-	-1.8*
Vagina	0.3	-	0.6	-0.7	-	-0.4
Vulva	1.1*	-	1.3*	0.8*	-	1.3*
Other female genital system	5.1*	-	5.1*	0.4	-	0.7
Male Genital System:	-1.9*	-2.4*	-	-2.5*	-3.2*	-
Prostate	-2.0*	-2.5*	-	-2.5*	-3.3*	-
Testis	0.7*	0.7*	-	0.3	0.2	-
Penis	0.3	-0.2	-	-0.3	-0.5	-
Other male genital system	0.5	0.3	-	3.6	3.1	-
Urinary System:	0.6*	0.3	0.7	-0.3*	-0.2*	-0.8*
Urinary bladder	-0.6*	-0.7*	-0.9*	0.3*	0.2	-0.4
Kidney & renal pelvis	2.3*	2.2*	2.3*	-1.0*	-1.0*	-1.2*
Ureter	-0.2	-0.2	-0.5	0.4	0.1	0.2
Other urinary system	2.2*	1.5	2.8	2.2	3.2	0.2
Eye & Orbit	-0.4	-1.1	0.3	1.0	0.1	1.8
Brain & Nervous System: <sup>d</sup>	-0.2	-0.1	-0.4	-0.4	-0.4	-0.4
Brain	-0.1	-0.1	-0.2	-	-	-
Cranial nerves & other nervous system	-1.8*	-1.1	-2.4	-	-	-
Endocrine System:	6.0*	5.0*	6.4*	0.1	0.2	0.2
Thyroid	6.4*	5.8*	6.6*	1.0*	1.4*	0.8
Other endocrine & thymus	0.6	-0.1	1.2	-1.2*	-1.7*	-1.0
Lymphoma:	0.1	0.2	-0.1	-2.7*	-2.6*	-3.0*
Hodgkin lymphoma	-0.1	0.0	-0.2	-2.4*	-2.1*	-2.7*
Non-Hodgkin lymphoma	0.1	0.2	0.0	-2.7*	-2.6*	-3.1*
Myeloma	0.4	0.7*	-0.1	-1.8*	-1.5*	-2.3*
Leukemia:	-0.5*	-0.8*	-0.4	-0.9*	-0.9*	-1.2*
Lymphocytic:	-0.6	-0.9*	-0.3	-1.3*	-1.4*	-1.4*
Acute lymphocytic	1.1*	0.9	1.5*	-0.9*	-1.1*	-0.6
Chronic lymphocytic	-1.2*	-1.5*	-1.1	-1.5*	-1.5*	-1.8*
Other lymphocytic	-1.1	-0.9	-2.0	-0.8	-1.2	-1.0
Myeloid & Monocytic:	0.0	-0.2	0.0	-0.4*	-0.3	-0.9*
Acute myeloid	0.1	-0.3	0.3	0.4*	0.5*	-0.1
Chronic myeloid	0.2	0.3	-0.1	-6.7*	-6.0*	-7.5*
Acute monocytic	-1.2	-0.2	-2.3	-3.3*	-4.0*	-2.9
Other myeloid & monocytic	-1.4	-0.7	-2.5	0.3	-0.2	0.5
Other leukemia:	-4.0*	-4.9*	-3.1*	-1.4*	-1.5*	-1.4*
Other acute leukemia	-5.7*	-6.1*	-5.5*	-3.7*	-4.2*	-3.3*
Aleukemic, subleukemic & NOS	-2.7*	-4.0*	-1.4	0.3	0.4	0.0
Kaposi Sarcoma <sup>e</sup>	-3.5*	-3.4*	-4.5*	-	-	-
Mesothelioma <sup>e</sup>	-0.8	-1.0	-0.5	-	-	-
Ill-defined & unspecified	-2.7*	-2.5*	-3.0*	-1.9*	-1.9*	-2.2*

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>a</sup> SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

<sup>b</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

<sup>c</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

<sup>d</sup> Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

<sup>e</sup> Trend not shown for mortality. Category did not exist in mortality coding until 1999.

\* The APC is significantly different from zero ( $p<.05$ ).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.