

Table 1.21
U.S. and SEER Death Rates by Primary Cancer Site and Race/Ethnicity, 2006-2010

Site		Total United States ^a							SEER 18 Areas ^{ab}						
		Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e	Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e
All Sites	Both Sexes	176.4	175.8	210.3	160.4	108.8	121.8	179.8	170.9	172.7	210.5	137.6	115.2	123.2	178.4
	Male	215.3	213.1	276.6	191.0	132.4	152.1	217.3	206.1	206.7	272.7	159.9	140.7	150.7	212.9
	Female	149.7	149.8	171.2	139.0	92.1	101.2	153.6	146.9	149.2	173.6	121.8	97.4	104.5	154.6
Oral Cavity & Pharynx	Both Sexes	2.5	2.4	3.0	2.3	2.0	1.5	2.5	2.5	2.4	3.0	2.2	2.1	1.5	2.6
	Male	3.8	3.6	5.2	3.4	3.0	2.5	3.7	3.8	3.7	5.2	3.3	3.1	2.4	3.8
	Female	1.4	1.4	1.4	1.5	1.2	0.8	1.4	1.4	1.4	1.4	-	1.3	0.8	1.5
Esophagus	Both Sexes	4.3	4.4	4.4	3.6	1.8	2.3	4.6	3.9	4.1	4.3	2.8	1.9	2.2	4.3
	Male	7.6	7.8	7.7	6.1	3.1	4.3	8.1	6.8	7.2	7.3	4.4	3.3	4.1	7.6
	Female	1.6	1.6	2.1	1.6	0.8	0.8	1.6	1.6	1.5	2.2	1.6	0.9	0.7	1.6
Stomach	Both Sexes	3.5	3.1	6.7	5.6	6.6	5.7	2.8	4.1	3.5	7.0	6.5	6.8	6.3	3.0
	Male	4.9	4.2	9.8	8.1	8.7	7.6	3.9	5.5	4.8	9.9	8.9	8.9	8.4	4.2
	Female	2.5	2.2	4.7	3.8	5.1	4.4	1.9	3.0	2.5	5.2	4.5	5.4	4.8	2.1
Colon & Rectum	Both Sexes	16.4	15.9	22.8	16.9	11.2	12.7	16.1	15.9	15.6	22.9	15.9	11.7	12.3	15.9
	Male	19.6	19.1	28.7	18.7	13.1	16.1	19.2	19.0	18.6	28.1	18.8	13.9	15.7	18.9
	Female	13.9	13.4	19.0	15.4	9.7	10.2	13.6	13.6	13.2	19.5	13.8	10.0	9.7	13.6
Liver & Intrahepatic Bile Duct	Both Sexes	5.6	5.2	7.4	9.3	9.7	8.5	4.8	6.1	5.4	7.6	9.3	9.8	8.6	4.9
	Male	8.3	7.6	11.8	13.2	14.4	12.3	7.1	8.9	7.9	11.9	12.8	14.6	12.2	7.2
	Female	3.4	3.2	4.1	6.1	6.0	5.4	2.9	3.7	3.4	4.3	6.3	6.0	5.6	3.0
Pancreas	Both Sexes	10.9	10.8	13.7	9.2	7.6	8.6	10.9	10.9	10.9	13.9	9.9	8.3	9.2	11.0
	Male	12.5	12.5	15.3	10.1	8.3	9.6	12.7	12.4	12.5	15.4	10.6	9.1	10.1	12.7
	Female	9.6	9.4	12.5	8.6	7.1	7.7	9.5	9.6	9.5	12.6	9.2	7.6	8.5	9.7
Larynx	Both Sexes	1.1	1.1	2.0	1.0	0.4	0.8	1.1	1.0	1.0	1.7	-	0.5	0.7	1.0
	Male	2.0	1.9	3.9	2.1	0.8	1.7	1.9	1.8	1.8	3.3	-	0.9	1.5	1.8
	Female	0.4	0.4	0.7	-	0.1	0.2	0.4	0.4	0.4	0.7	-	0.1	0.2	0.4
Lung & Bronchus	Both Sexes	49.5	50.2	53.5	40.2	25.5	21.3	52.6	45.4	46.6	53.4	29.6	27.0	20.2	50.1
	Male	63.5	63.2	78.5	49.6	35.5	31.3	65.7	57.3	57.4	76.5	36.4	38.0	28.6	61.2
	Female	39.2	40.4	37.2	33.1	18.4	14.1	42.7	36.6	38.5	38.1	24.6	19.0	14.2	41.9
Melanoma of the Skin	Both Sexes	2.7	3.1	0.4	1.1	0.4	0.8	3.4	2.6	3.1	0.4	-	0.4	0.8	3.5
	Male	4.1	4.6	0.5	1.7	0.4	1.1	5.0	3.9	4.6	0.6	-	0.5	1.0	5.2
	Female	1.7	2.0	0.4	0.8	0.3	0.6	2.1	1.6	2.0	0.3	-	0.3	0.7	2.2
Breast	Female	22.6	22.1	30.8	15.5	11.5	14.8	22.7	22.7	22.7	31.3	14.0	12.8	14.9	23.7

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 18 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

^c Rates for American Indian/Alaska Native (AI/AN) are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Asian/Pacific Islander.

^e Hispanic (Hisp) and White Non-Hispanic (W-NHisp) are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Mortality data for Hispanics and White Non-Hispanics do not include cases from the District of Columbia, North Dakota, and South Carolina.

- Statistic could not be calculated due to less than 16 cases in the time interval.

Table 1.21 - continued
U.S. and SEER Death Rates by Primary Cancer Site and Race/Ethnicity, 2006-2010

Site		Total United States ^a							SEER 18 Areas ^{ab}						
		Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e	Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e
Cervix	Female	2.4	2.2	4.2	3.5	1.9	2.9	2.1	2.4	2.2	4.0	3.1	1.9	3.1	2.0
Corpus & Uterus, NOS	Female	4.3	4.0	7.4	3.2	2.6	3.3	4.0	4.3	4.1	7.1	2.5	2.9	3.4	4.1
Ovary	Female	8.1	8.4	6.7	7.1	4.8	5.8	8.6	8.2	8.7	6.9	6.9	5.0	6.3	9.0
Prostate	Male	23.0	21.2	50.9	20.7	10.1	19.2	21.3	23.1	22.0	49.6	17.5	10.9	19.7	22.1
Testis	Male	0.2	0.3	0.1	-	0.1	0.3	0.3	0.2	0.3	0.1	-	0.1	0.3	0.3
Urinary Bladder	Both Sexes	4.4	4.6	3.7	2.4	1.7	2.4	4.8	4.3	4.6	3.8	2.0	1.8	2.3	4.8
	Male	7.7	8.1	5.5	4.1	2.8	4.0	8.4	7.4	8.0	5.8	3.1	3.0	3.9	8.5
	Female	2.2	2.2	2.6	1.2	0.9	1.3	2.3	2.2	2.2	2.7	-	1.0	1.3	2.4
Kidney & Renal	Both Sexes	4.0	4.1	3.9	6.7	2.0	3.5	4.1	3.8	3.9	3.9	6.2	2.1	3.6	3.9
	Male	5.8	5.9	5.7	9.5	3.0	5.1	5.9	5.5	5.7	5.8	8.5	3.2	5.2	5.7
Pelvis	Female	2.6	2.6	2.6	4.4	1.2	2.3	2.6	2.4	2.5	2.5	4.4	1.4	2.4	2.5
Brain & Nervous System	Both Sexes	4.3	4.6	2.5	2.6	1.9	2.8	4.8	4.2	4.7	2.6	1.8	2.0	3.0	5.0
	Male	5.2	5.6	3.0	2.8	2.3	3.3	5.8	5.2	5.7	3.1	1.9	2.5	3.4	6.1
	Female	3.5	3.8	2.1	2.3	1.6	2.4	3.9	3.4	3.8	2.2	1.6	1.7	2.6	4.0
Thyroid	Both Sexes	0.5	0.5	0.5	0.5	0.7	0.6	0.5	0.5	0.5	0.4	-	0.8	0.6	0.5
	Male	0.5	0.5	0.4	-	0.5	0.5	0.5	0.5	0.5	0.3	-	0.6	0.5	0.5
	Female	0.5	0.5	0.6	-	0.9	0.7	0.5	0.6	0.5	0.5	-	1.0	0.7	0.5
Hodgkin Lymphoma	Both Sexes	0.4	0.4	0.4	-	0.2	0.4	0.4	0.4	0.4	0.4	-	0.2	0.4	0.4
	Male	0.5	0.5	0.4	-	0.2	0.5	0.5	0.5	0.5	0.5	-	0.2	0.5	0.5
	Female	0.3	0.3	0.3	-	0.1	0.3	0.3	0.3	0.3	0.3	-	0.1	0.3	0.3
Non-Hodgkin Lymphoma	Both Sexes	6.4	6.7	4.6	4.8	4.1	5.3	6.8	6.3	6.7	4.6	3.6	4.5	5.6	6.7
	Male	8.2	8.5	5.9	5.4	5.2	6.5	8.6	8.0	8.4	6.1	3.7	5.7	6.8	8.5
	Female	5.1	5.3	3.6	4.3	3.4	4.4	5.4	5.0	5.3	3.5	3.4	3.7	4.6	5.4
Myeloma	Both Sexes	3.4	3.1	6.4	2.9	1.7	2.8	3.1	3.3	3.1	6.4	1.8	1.9	2.9	3.1
	Male	4.3	4.0	7.9	3.6	2.3	3.5	4.1	4.2	4.0	7.9	2.3	2.5	3.6	4.0
	Female	2.7	2.5	5.4	2.4	1.3	2.3	2.5	2.6	2.4	5.5	1.5	1.5	2.5	2.4
Leukemia	Both Sexes	7.1	7.3	6.1	4.8	3.9	4.9	7.4	6.9	7.2	6.2	3.8	4.1	4.9	7.4
	Male	9.5	9.8	8.2	7.0	5.0	6.1	10.0	9.2	9.7	8.2	5.1	5.2	6.2	9.9
	Female	5.3	5.5	4.8	3.3	3.1	4.0	5.5	5.2	5.4	4.9	2.9	3.3	4.0	5.5

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 18 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

^c Rates for American Indian/Alaska Native (AI/AN) are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Asian/Pacific Islander.

^e Hispanic (Hisp) and White Non-Hispanic (W-NHisp) are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Mortality data for Hispanics and White Non-Hispanics do not include cases from the District of Columbia, North Dakota, and South Carolina.

- Statistic could not be calculated due to less than 16 cases in the time interval.