

Table 1.23  
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2011<sup>a</sup>  
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	6,271,036	19,001	41,846	85,407	155,919	356,211	883,826	1,658,545	3,070,283
Females	7,126,123	16,447	36,376	88,223	226,162	638,879	1,316,044	1,758,236	3,045,756
Oral Cavity & Pharynx									
Males	186,491	62	423	1,548	3,544	14,256	45,288	58,097	63,273
Females	95,100	90	522	1,483	3,689	8,554	18,160	23,692	38,911
Esophagus									
Males	26,909	0	0	22	144	1,109	4,697	9,747	11,190
Females	7,642	0	0	11	11	240	1,378	1,832	4,170
Stomach									
Males	42,127	0	29	62	603	2,502	7,119	11,218	20,593
Females	31,908	4	23	154	618	2,252	4,650	6,664	17,543
Colon & Rectum									
Males	575,457	11	69	1,276	5,758	25,233	82,732	139,236	321,141
Females	586,969	0	67	1,472	5,551	23,983	70,097	112,784	373,015
Liver & Intrahep									
Males	32,389	556	508	616	547	1,590	10,492	11,364	6,716
Females	13,553	396	521	359	403	889	3,016	3,463	4,506
Pancreas									
Males	20,801	23	12	130	276	1,653	4,466	6,626	7,615
Females	22,737	0	62	212	431	1,446	3,801	6,179	10,604
Larynx									
Males	71,999	0	0	90	219	1,822	10,381	21,280	38,207
Females	17,266	0	0	36	135	841	3,336	4,868	8,051
Lung & Bronchus									
Males	183,215	45	110	405	1,195	5,366	24,255	54,762	97,077
Females	219,111	23	71	367	1,387	7,548	29,898	59,021	120,797
Melanoma of the Skin									
Males	471,220	56	594	5,039	17,738	46,603	97,543	130,762	172,885
Females	489,011	68	878	10,864	34,091	73,459	115,119	113,905	140,628

<sup>a</sup> U.S. 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2011 U.S. population estimates based on the average of 2010 and 2011 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.23 - continued  
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2011<sup>a</sup>  
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages <sup>c</sup>	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	14,871	0	0	22	112	591	1,838	4,277	8,030
Females	2,899,726	0	47	2,669	35,679	215,380	550,120	795,921	1,299,910
Cervix									
Females	249,632	0	57	2,142	15,934	41,969	59,700	57,440	72,389
Corpus & Uterus, NOS									
Females	610,804	12	12	581	5,712	25,316	89,564	165,807	323,801
Ovary <sup>d</sup>									
Females	188,867	45	1,063	3,475	6,807	18,458	41,719	49,281	68,021
Prostate									
Males	2,707,821	34	47	113	332	20,758	235,407	776,168	1,674,962
Urinary Bladder									
Males	425,722	57	88	567	2,476	10,749	43,060	102,624	266,102
Females	145,796	56	46	219	1,015	4,094	13,710	30,955	95,700
Kidney & Renal Pelvis									
Males	212,703	1,458	2,458	2,630	5,254	18,386	41,683	61,520	79,315
Females	145,900	1,374	2,649	2,922	4,870	12,287	25,458	35,857	60,483
Hodgkin Lymphoma									
Males	96,100	186	2,493	9,242	16,547	23,764	22,436	14,185	7,247
Females	89,693	67	2,083	9,407	16,399	23,306	19,845	11,219	7,368
Non-Hodgkin Lymphoma									
Males	278,836	842	3,598	7,821	13,627	29,722	54,887	71,106	97,233
Females	252,083	519	1,758	4,663	8,663	21,422	42,382	60,220	112,455
Myeloma									
Males	45,388	0	6	99	490	3,274	8,260	14,917	18,342
Females	37,979	0	0	28	312	2,325	6,791	11,124	17,399
Leukemia									
Males	170,969	6,686	13,047	12,524	11,179	14,206	23,300	35,001	55,027
Females	131,831	5,729	10,353	11,342	9,836	10,094	15,793	22,736	45,948
Acute Lymphocytic Leuk									
Males	39,514	5,599	11,159	9,649	6,330	3,974	1,505	836	463
Females	32,723	4,763	8,785	8,138	5,414	3,188	1,137	816	483

<sup>a</sup> U.S. 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2011 U.S. population estimates based on the average of 2010 and 2011 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b</sup> Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

<sup>c</sup> Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.