Table 1.24 Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites by Race/Ethnicity Both Sexes

All Races			White			Black		
	Rateb	APC <sup>c</sup>		Rateb	APC <sup>c</sup>		Rateb	APC <sup>c</sup>
	2010-2014	2005-2014		2010-2014	2005-2014		2010-2014	2005-2014
All Sites	442.7	-1.1*	All Sites	451.8	-1.1*	All Sites	459.3	-1.5*
Breast	67.0	0.0	Breast	67.8	-0.1	Prostate <sup>f</sup>	80.7	-4.5*
Lung and Bronchus	55.8	-2.3*	Lung and Bronchus	57.3	-2.2*	Breast	71.1	0.5
Prostate <sup>f</sup>	54.7	-4.8*	Prostate <sup>f</sup>	52.1	-5.2*	Lung and Bronchus	63.0	-2.5*
Colon and Rectum	40.1	-2.8*	Colon and Rectum	39.3	-2.8*	Colon and Rectum	48.7	-3.3*
Melanoma of the Skin	22.3	1.1*	Melanoma of the Skin	26.6	1.1*	Kidney and Renal Pelvis	18.0	1.0
Urinary Bladder	19.8	-1.2*	Urinary Bladder	21.7	-1.2*	Pancreas	15.5	-0.5
Non-Hodgkin Lymphoma	19.5	-0.5*	Non-Hodgkin Lymphoma	20.4	-0.6*	Non-Hodgkin Lymphoma	14.6	-0.3
Kidney and Renal Pelvis	15.6	0.7	Kidney and Renal Pelvis		0.7	Corpus and Uterus, NOS <sup>f</sup>	14.0	2.3*
Thyroid	14.2	4.0*	Thyroid	14.9	3.9*	Myeloma	13.2	0.8
Leukemia	13.7	0.5*	Leukemia	14.5	0.4	Urinary Bladder	12.5	-0.9*
Corpus and Uterus, NOS <sup>f</sup>	13.6	1.2*	Corpus and Uterus, NOS <sup>f</sup>	13.8	0.9*	Leukemia	10.9	1.2*
Pancreas	12.5	0.5*	Pancreas	12.5	0.6*	Stomach	10.2	-3.0*
Oral Cavity and Pharynx	11.2	0.7*	Oral Cavity and Pharynx		1.0*	Liver & IBD <sup>g</sup>	10.1	2.5*
Liver & IBD <sup>g</sup>	8.6	2.6*	Liver & IBD <sup>g</sup>	7.8	3.4*	Oral Cavity and Pharynx	9.1	-1.5*
Stomach	7.3	-0.9*	Brain and ONS <sup>g</sup>	7.0	-0.5*	Thyroid	8.7	4.4*
Asian/Pacific	Islander		American Indian/Alaska Natived			Hispanic <sup>e</sup>		
	Rateb	APCc		Rate <sup>b</sup>	APC <sup>c</sup>		Rate <sup>b</sup>	APC <sup>c</sup>
		2005-2014			2005-2014			2005-2014
All Sites	298.8	-1.2*	All Sites	316.0	-1.4*	All Sites	340.6	-1.3*
Breast	54.1	0.9*	Breast	44.1	0.0	Breast	49.8	0.1
Lung and Bronchus	35.7	-1.8*	Colon and Rectum	41.1	-0.6	Prostate <sup>f</sup>	43.7	-5.6*
Colon and Rectum	33.8	-2.8*	Lung and Bronchus	37.4	-3.8*	Colon and Rectum	33.9	-2.5*
Prostate <sup>f</sup>	27.5	-6.4*	Prostate <sup>f</sup>	26.5	-6.3*	Lung and Bronchus	28.8	-2.5*
Thyroid	13.7	3.3*	Kidney and Renal Pelvis	17.2	-3.2*	Non-Hodgkin Lymphoma	17.8	-0.2
Non-Hodgkin Lymphoma	13.5	0.2	Liver & IBDg	13.3	2.2	Kidney and Renal Pelvis	15.4	0.6
Liver & IBD <sup>g</sup>	13.2	-2.2*	Non-Hodgkin Lymphoma	12.1	-2.0	Liver & IBDg	13.2	1.9*
Corpus and Uterus, NOSf	11.2	2.1*	Stomach	10.6	0.5	Thyroid	12.3	4.2*
Stomach	10.6	-3.2*	Corpus and Uterus, NOSf	10.5	0.8	Corpus and Uterus, NOSf	11.6	2.5*
Pancreas	10.0	0.5	Pancreas	9.6	-2.9	Pancreas	11.1	-0.2
Urinary Bladder	8.7	-1.9*	Thyroid	9.3	5.0*	Urinary Bladder	11.0	-1.7*
Kidney and Renal Pelvis	8.2	0.5	Oral Cavity and Pharynx	8.9	1.4	Leukemia	10.6	0.2
Leukemia	7.8	0.5	Urinary Bladder	8.5	-0.4	Stomach	10.5	-1.7*
Oral Cavity and Pharynx	7.7	-0.5	Leukemia	8.1	0.7	Oral Cavity and Pharynx	6.6	0.1
Ovary <sup>fh</sup>	5.2	-0.7	Ovary <sup>fh</sup>	5.1	-3.6	Myeloma	6.1	0.9

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

- Top 15 cancer sites selected based on 2010-2014 age-adjusted rates for the race/ethnic group.
- Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- The APC is the Annual Percent Change over the time interval.
- Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- d Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.
- Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. f
- The rates for sex-specific cancer sites are calculated using the population for both sexes combined.
- IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
- Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
- The APC is significantly different from zero (p<.05).
- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.25

Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites by Race/Ethnicity

Males

All Races			White			Black		
	Rate <sup>b</sup> 2010-2014	APC° 2005-2014		Rate <sup>b</sup> 2010-2014	APC <sup>c</sup> 2005-2014		Rate <sup>b</sup> 2010-2014	APC <sup>c</sup> 2005-2014
All Sites	492.4	-2.0*	All Sites	497.1	-2.0*	All Sites	549.7	-2.8*
Prostate	119.8	-5.2*	Prostate	112.8	-5.6*	Prostate	188.7	-4.8*
Lung and Bronchus	65.7	-2.9*	Lung and Bronchus	65.9	-2.9*	Lung and Bronchus	83.7	-3.1*
Colon and Rectum	46.0	-3.0*	Colon and Rectum	45.0	-3.1*	Colon and Rectum	56.4	-3.4*
Urinary Bladder	34.9	-1.4*	Urinary Bladder	38.1	-1.4*	Kidney and Renal Pelvis	25.0	0.7
Melanoma of the Skin	29.2	1.3*	Melanoma of the Skin	34.4	1.3*	Urinary Bladder	21.0	-0.6
Non-Hodgkin Lymphoma	23.7	-0.4*	Non-Hodgkin Lymphoma	24.8	-0.4*	Non-Hodgkin Lymphoma	17.6	-0.4
Kidney and Renal Pelvis	21.4	0.7	Kidney and Renal Pelvis	21.9	0.7	Pancreas	17.0	-0.4
Leukemia	17.6	0.4	Leukemia	18.5	0.2	Liver & IBD <sup>f</sup>	16.7	2.5*
Oral Cavity and Pharynx	16.9	0.9*	Oral Cavity and Pharynx	17.6	1.1*	Myeloma	15.9	0.8
Pancreas	14.2	0.6*	Pancreas	14.2	0.7*	Oral Cavity and Pharynx	14.2	-1.3*
Liver & IBD <sup>f</sup>	13.3	2.5*	Liver & IBD <sup>f</sup>	11.9	3.1*	Leukemia	14.0	1.5*
Stomach	9.9	-1.3*	Stomach	9.0	-0.9*	Stomach	13.6	-3.5*
Myeloma	8.3	1.2*	Brain and ONS <sup>f</sup>	8.3	-0.5*	Larynx	8.5	-2.6*
Brain and ONS <sup>f</sup>	7.5	-0.5*	Myeloma	7.8	1.3*	Esophagus	6.9	-4.9*
Esophagus	7.3	-1.3*	Esophagus	7.7	-0.7	Brain and ONS <sup>f</sup>	4.8	0.0
Asian/Pacific	: Islander		American Indian/Alaska Native <sup>d</sup>			Hispanic <sup>e</sup>		
	Rateb	APCc		Rateb	APC°		Rateb	APCc
		2005-2014			2005-2014		2010-2014	
All Sites	308.6	-2.5*	All Sites	331.4	-2.0*	All Sites	372.2	-2.6*
Prostate	62.9	-6.6*	Prostate	59.9	-6.5*	Prostate	98.3	-5.9*
Lung and Bronchus	46.4	-2.5*	Lung and Bronchus	46.9	-2.9*	Colon and Rectum	40.0	-3.1*
Colon and Rectum	40.2	-2.2*	Colon and Rectum	45.5	-0.5	Lung and Bronchus	35.3	-3.6*
Liver & IBD <sup>f</sup>	20.2	-2.0*	Kidney and Renal Pelvis		-1.8	Non-Hodgkin Lymphoma	20.7	-0.2
Non-Hodgkin Lymphoma	16.7	0.1	Liver & IBD <sup>f</sup>	18.7	2.3	Kidney and Renal Pelvis	20.2	0.1
Urinary Bladder	15.2	-2.1*	Urinary Bladder	14.9	-0.7	Liver & IBD <sup>f</sup>	19.7	1.5*
Stomach	14.0	-3.3*	Stomach	13.7	-0.7	Urinary Bladder	19.3	-1.7*
Kidney and Renal Pelvis		0.9	Non-Hodgkin Lymphoma	13.2	-3.8	Stomach	13.0	-2.9*
Oral Cavity and Pharynx		0.0	Oral Cavity and Pharynx		2.0	Leukemia	12.9	0.0
Pancreas	11.0	0.3	Pancreas	11.1	-	Pancreas	11.8	-0.4
Leukemia	9.7	0.7	Leukemia	9.8	1.4	Oral Cavity and Pharynx		0.3
Thyroid	6.6	5.1*	Testis	5.4	4.9	Myeloma	7.7	1.5
Myeloma	4.7	0.1	Esophagus	5.1	_	Brain and ONS <sup>f</sup>	5.7	-1.4*
Brain and ONS <sup>f</sup> Esophagus	4.4	0.4	Myeloma Melanoma of the Skin	5.0 4.3	-	Thyroid Testis	5.2 5.1	3.6* 2.6*

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

- a Top 15 cancer sites selected based on 2010-2014 age-adjusted rates for the race/ethnic group.
- Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- The APC is the Annual Percent Change over the time interval.
- Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- d Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.
- IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
- \* The APC is significantly different from zero (p<.05).
- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

- - - -

Table 1.26

Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites by Race/Ethnicity

Females

All Races			White			Black		
	Rate <sup>b</sup> 2010-2014	APC <sup>c</sup> 2005-2014		Rate <sup>b</sup> 2010-2014	APC <sup>c</sup> 2005-2014		Rate <sup>b</sup> 2010-2014	APC <sup>c</sup> 2005-2014
All Sites	408.7	-0.2*	All Sites	421.5	-0.2*	All Sites	397.4	-0.4*
Breast	124.9	0.2	Breast	127.7	0.0	Breast	125.1	0.6*
Lung and Bronchus	48.4	-1.7*	Lung and Bronchus	50.8	-1.6*	Lung and Bronchus	49.0	-1.7*
Colon and Rectum	35.1	-2.7*	Colon and Rectum	34.4	-2.6*	Colon and Rectum	43.2	-3.3*
Corpus and Uterus, NOS	25.7	1.4*	Corpus and Uterus, NOS	26.3	1.1*	Corpus and Uterus, NOS	24.8	2.5*
Thyroid	21.0	4.0*	Thyroid	22.4	4.0*	Pancreas	14.3	-0.7
Melanoma of the Skin	17.3	0.6	Melanoma of the Skin	20.9	0.7	Thyroid	13.0	4.8*
Non-Hodgkin Lymphoma	16.0	-0.6*	Non-Hodgkin Lymphoma	16.9	-0.7*	Kidney and Renal Pelvis	12.6	1.1
Ovary <sup>g</sup>	11.7	-1.8*	Ovary <sup>g</sup>	12.2	-2.0*	Non-Hodgkin Lymphoma	12.2	0.0
Pancreas	11.1	0.4*	Leukemia	11.3	0.4	Myeloma	11.4	0.7
Kidney and Renal Pelvis	10.7	0.5	Kidney and Renal Pelvis	11.0	0.5	Ovary <sup>g</sup>	9.4	-1.3*
Leukemia	10.7	0.4	Pancreas	11.0	0.5*	Leukemia	8.8	0.8
Urinary Bladder	8.4	-1.4*	Urinary Bladder	9.1	-1.3*	Cervix Uteri	8.7	-2.3*
Cervix Uteri	7.4	-1.6*	Cervix Uteri	7.4	-1.5*	Stomach	7.8	-2.7*
Oral Cavity and Pharynx	6.2	0.0	Oral Cavity and Pharynx	6.4	0.4	Urinary Bladder	6.8	-1.6*
Brain and ONS <sup>f</sup>	5.3	-0.6	Brain and ONS <sup>f</sup>	5.9	-0.4	Oral Cavity and Pharynx	5.1	-2.1*
Asian/Pacific	c Islander		American Indian/Alaska Natived			Hispanic <sup>e</sup>		
	Rateb	APC <sup>c</sup>		Rateb	APC <sup>c</sup>		Rateb	APC <sup>c</sup>
	2010-2014	2005-2014		2010-2014			2010-2014	2005-2014
All Sites	296.1	-0.2	All Sites	308.9	-0.8	All Sites	323.3	-0.1
Breast	98.5	1.0*	Breast	82.2	0.2	Breast	93.1	0.2
Colon and Rectum	28.8	-3.4*	Colon and Rectum	37.5	-0.4	Colon and Rectum	29.1	-2.0*
Lung and Bronchus	27.9	-0.8	Lung and Bronchus	30.4	-5.0*	Lung and Bronchus	24.2	-1.4*
Corpus and Uterus, NOS	20.5	2.2*	Corpus and Uterus, NOS	19.9	0.9	Corpus and Uterus, NOS	21.9	2.7*
Thyroid	19.9	2.8*	Thyroid	14.1	4.9*	Thyroid	19.3	4.4*
Non-Hodgkin Lymphoma	11.1	0.2	Kidney and Renal Pelvis	12.7	-4.5*	Non-Hodgkin Lymphoma	15.4	-0.1
Ovary <sup>g</sup>	9.5	-0.7	Non-Hodgkin Lymphoma	11.1	-0.5	Kidney and Renal Pelvis	11.5	1.0
Pancreas	9.2	0.7	Ovary <sup>g</sup>	9.5	-3.3	Ovary <sup>g</sup>	10.6	-1.6*
Stomach	8.0	-3.3*	Liver & IBD <sup>f</sup>	8.6	1.4	Pancreas	10.5	-0.1
Liver & IBD <sup>f</sup>	7.6	-2.5*	Pancreas	8.3	-4.4*	Cervix Uteri	9.1	-4.4*
Leukemia	6.3	0.2	Stomach	8.1	2.0	Leukemia	8.8	0.3
Cervix Uteri	6.1	-2.9*	Cervix Uteri	7.7	-0.5	Stomach	8.5	-0.3
Kidney and Renal Pelvis	5.5	0.0	Leukemia	6.8	0.2	Liver & IBD <sup>f</sup>	7.6	2.8*
Oral Cavity and Pharynx	5.0	-1.3	Oral Cavity and Pharynx	5.5	_	Myeloma	4.9	0.2
Urinary Bladder	3.9	-1.5	Myeloma	5.2	-	Urinary Bladder	4.9	-1.9*

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

- a Top 15 cancer sites selected based on 2010-2014 age-adjusted rates for the race/ethnic group.
- Parallel Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- The APC is the Annual Percent Change over the time interval.
- Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- d Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.
- Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.
- IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
- Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
- \* The APC is significantly different from zero (p<.05).
- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.27 Age-Adjusted U.S. Death Rates and Trends for the Top 15 Cancer Sites by Race/Ethnicity Both Sexes

All Races			White			Black		
	Rate <sup>b</sup> 2010-2014	APC <sup>c</sup> 2005-2014		Rate <sup>b</sup> 2010-2014	APC <sup>c</sup> 2005-2014		Rate <sup>b</sup> 2010-2014	APC <sup>c</sup> 2005-2014
All Sites	166.1	-1.5*	All Sites	166.2	-1.4*	All Sites	194.2	-2.0*
Lung and Bronchus	44.7	-2.5*	Lung and Bronchus	45.5	-2.4*	Lung and Bronchus	48.0	-2.8*
Colon and Rectum	14.8	-2.5*	Colon and Rectum	14.4	-2.4*	Colon and Rectum	20.0	-3.3*
Breast	11.8	-1.9*	Breast	11.4	-1.9*	Breast	17.1	-1.7*
Pancreas	10.9	0.1	Pancreas	10.8	0.2	Prostate <sup>f</sup>	15.2	-3.8*
Prostate <sup>f</sup>	8.1	-2.6*	Prostate <sup>f</sup>	7.5	-2.4*	Pancreas	13.4	-0.5*
Leukemia	6.8	-0.9*	Leukemia	7.1	-0.9*	Liver & IBD <sup>g</sup>	8.2	2.5*
Liver & IBD <sup>g</sup>	6.3	2.7*	Non-Hodgkin Lymphoma	6.1	-2.2*	Myeloma	6.3	-0.3
Non-Hodgkin Lymphoma	5.9	-2.3*	Liver & IBD <sup>g</sup>	5.8	2.9*	Stomach	5.9	-3.1*
Urinary Bladder	4.4	-0.1	Brain and ONS <sup>g</sup>	4.7	0.4*	Leukemia	5.7	-1.3*
Brain and ONS <sup>g</sup>	4.3	0.4	Urinary Bladder	4.6	0.1	Corpus and Uterus, NOSf	4.8	1.8*
Esophagus	4.1	-1.1*	Esophagus	4.3	-0.6*	Non-Hodgkin Lymphoma	4.3	-1.6*
Ovary <sup>f</sup>	4.1	-2.5*	Ovary <sup>f</sup>	4.2	-2.6*	Ovary <sup>f</sup>	3.8	-1.5*
Kidney and Renal Pelvis	3.9	-0.8*	Kidney and Renal Pelvis	4.0	-0.7*	Esophagus	3.7	-4.4*
Myeloma	3.3	-0.6*	Myeloma	3.1	-0.7*	Kidney and Renal Pelvis	3.7	-1.3*
Stomach	3.2	-2.3*	Melanoma of the Skin	3.1	-0.3	Urinary Bladder	3.6	-0.8
Asian/Pacifi	c Islander		American Indian/Alaska Native <sup>d</sup>			<u> Hispanic</u> e		
	Rateb	APC <sup>c</sup>		Rate <sup>b</sup>	APC <sup>c</sup>		Rateb	APC°
	2010-2014	2005-2014		2010-2014	2005-2014		2010-2014	2005-2014
All Sites	102.8	-1.4*	All Sites	152.4	-1.2*	All Sites	116.2	-1.3*
Lung and Bronchus	23.7	-1.7*	Lung and Bronchus	37.5	-1.5*	Lung and Bronchus	19.3	-2.5*
Colon and Rectum	10.3	-2.0*	Colon and Rectum	16.4	-0.3	Colon and Rectum	11.7	-2.0*
Liver & IBD <sup>g</sup>	9.7	-0.5	Liver & IBD <sup>g</sup>	10.5	3.0*	Liver & IBD <sup>g</sup>	9.1	1.2*
Pancreas	7.7	0.0	Pancreas	9.0	-1.2	Pancreas	8.5	-0.4
Breast	6.4	-0.7	Breast	7.9	-3.9*	Breast	8.0	-0.7*
Stomach	5.5	-4.2*	Prostate <sup>f</sup>	7.9	-2.0	Prostate <sup>f</sup>	6.6	-3.0*
Non-Hodgkin Lymphoma	4.0	-1.1*	Kidney and Renal Pelvis	6.3	-0.3	Stomach	5.3	-2.0*
Leukemia	3.9	-0.4	Stomach	5.4	-1.2	Non-Hodgkin Lymphoma	5.0	-1.5*
Prostate <sup>f</sup>	3.5	-3.3*	Leukemia	4.4	-1.9	Leukemia	4.9	-0.2
Ovary <sup>f</sup>	2.5	-2.2*	Non-Hodgkin Lymphoma	4.2	-2.7*	Kidney and Renal Pelvis	3.5	-0.9
Brain and ONS <sup>g</sup>	2.1	1.5	Ovary <sup>f</sup>	3.5	-2.5*	Ovary <sup>f</sup>	3.0	-2.0*
Oral Cavity and Pharynx	1.9	-1.2*	Esophagus	3.4	-1.7	Brain and ONS <sup>g</sup>	2.9	0.4
Kidney and Renal Pelvis	1.8	-1.1	Myeloma	3.0	0.3	Myeloma	2.8	-0.8
Urinary Bladder	1.7	0.1	Brain and ONS <sup>g</sup>	2.5	0.9	Urinary Bladder	2.3	-0.9
Esophagus	1.7	-1.7*	Urinary Bladder	2.3	0.4	Esophagus	2.2	-1.5

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

- a Top 15 cancer sites selected based on 2010-2014 age-adjusted rates for the race/ethnic group.
- Bates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- The APC is the Annual Percent Change over the time interval.
- Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.
- The rates for sex-specific cancer sites are calculated using the population for both sexes combined.
- IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
- \* The APC is significantly different from zero (p<.05).
- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.28

Age-Adjusted U.S. Death Rates and Trends for the Top 15 Cancer Sites by Race/Ethnicity

Males

White

Black

	Rate <sup>b</sup>	APC <sup>c</sup>		Rate <sup>b</sup>	APC <sup>c</sup>		Rate <sup>b</sup>	APC <sup>c</sup>
	2010-2014	2005-2014		2010-2014	2005-2014		2010-2014	2005-2014
All Sites	200.5	-1.8*	All Sites	199.8	-1.6*	All Sites	247.3	-2.7*
Lung and Bronchus	55.9	-3.1*	Lung and Bronchus	55.9	-3.0*	Lung and Bronchus	68.0	-3.5*
Prostate	20.1	-3.3*	Prostate	18.7	-3.1*	Prostate	42.0	-4.5*
Colon and Rectum	17.7	-2.6*	Colon and Rectum	17.2	-2.5*	Colon and Rectum	25.3	-3.1*
Pancreas	12.6	0.1	Pancreas	12.6	0.2*	Pancreas	15.0	-0.4
Liver & IBD <sup>f</sup>	9.2	2.7*	Leukemia	9.5	-0.9*	Liver & IBD <sup>f</sup>	13.0	2.3*
Leukemia	9.1	-1.0*	Liver & IBD <sup>f</sup>	8.5	2.8*	Stomach	8.6	-3.3*
Urinary Bladder	7.6	-0.2*	Urinary Bladder	8.1	-0.1	Myeloma	7.5	-1.0*
Non-Hodgkin Lymphoma	7.6	-2.0*	Non-Hodgkin Lymphoma	7.9	-2.0*	Leukemia	7.5	-1.7*
Esophagus	7.3	-1.2*	Esophagus	7.6	-0.7*	Esophagus	6.3	-4.9*
Kidney and Renal Pelvis	5.6	-0.6*	Kidney and Renal Pelvis	5.8	-0.4*	Non-Hodgkin Lymphoma	5.5	-1.7*
Brain and ONS <sup>f</sup>	5.3	0.4	Brain and ONS <sup>f</sup>	5.7	0.4*	Kidney and Renal Pelvis	5.5	-1.0
Stomach	4.4	-2.5*	Melanoma of the Skin	4.6	-0.2	Urinary Bladder	5.4	-0.7
Myeloma	4.2	-0.7*	Myeloma	4.0	-0.6*	Oral Cavity and Pharynx	5.0	-2.4*
Melanoma of the Skin	4.0	-0.4	Stomach	3.8	-2.5*	Larynx	3.4	-3.7*
Oral Cavity and Pharynx	3.8	0.2	Oral Cavity and Pharynx	3.8	0.7*	Brain and ONS <sup>f</sup>	3.2	1.0
Asian/Pacifi	c Islander		American Indian/Alaska Native <sup>d</sup>			Hispanic <sup>e</sup>		
	Rateb	APCc		Rateb	APC°		Rateb	APC <sup>c</sup>
	2010-2014	2005-2014		2010-2014	2005-2014		2010-2014	2005-2014
All Sites	122.7	-1.8*	All Sites	184.0	-1.0	All Sites	142.6	-1.7*
Lung and Bronchus	31.7	-2.6*	Lung and Bronchus	46.3	-1.4	Lung and Bronchus	27.3	-3.4*
Liver & IBD <sup>f</sup>	14.3	-0.4	Colon and Rectum	19.5	-0.3	Prostate	16.5	-3.5*
Colon and Rectum	12.4	-1.5*	Prostate	19.4	-2.7*	Colon and Rectum	15.0	-1.5*
Prostate	8.8	-3.3*	Liver & IBD <sup>f</sup>	14.9	3.6*	Liver & IBD <sup>f</sup>	13.1	1.2*
Pancreas	8.2	-0.6	Pancreas	9.8	-1.7	Pancreas	9.5	-0.5
Stomach	7.1	-4.6*	Kidney and Renal Pelvis	8.9	-1.1	Stomach	6.9	-2.4*
Leukemia	5.0	-0.6	Stomach	7.5	-2.4	Non-Hodgkin Lymphoma	6.2	-1.0*
Non-Hodgkin Lymphoma	4.9	-1.0	Leukemia	5.8	-2.7	Leukemia	6.1	-0.4
Oral Cavity and Pharynx	2.9	-0.2	Non-Hodgkin Lymphoma	5.6	0.1	Kidney and Renal Pelvis	4.9	-1.3
Urinary Bladder	2.9	0.6	Esophagus	5.6	-1.4	Esophagus	4.0	-1.7
Esophagus	2.9	-1.0	Oral Cavity and Pharynx	3.7	0.0	Urinary Bladder	3.9	-1.4
Kidney and Renal Pelvis	2.7	-0.5	Urinary Bladder	3.6	-2.1	Myeloma	3.4	-0.6
Brain and ONSf	2.4	0.5	Myeloma	3.3	-2.4	Brain and ONS <sup>f</sup>	3.4	0.3
Myeloma	2.1	-0.9	Brain and ONSf	3.0	0.1	Oral Cavity and Pharynx		-0.9*
Soft Tissue including H	leart 1.0	-1.5	Larynx	1.7	_	Larynx	1.6	-2.5*

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

- Top 15 cancer sites selected based on 2010-2014 age-adjusted rates for the race/ethnic group.
- Pates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- The APC is the Annual Percent Change over the time interval.
- Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- d Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.
- IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
- \* The APC is significantly different from zero (p<.05).

All Races

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.29 Age-Adjusted U.S. Death Rates and Trends for the Top 15 Cancer Sites by Race/Ethnicity Females

All Races			White			Black		
	Rateb	APC <sup>c</sup>		Rate <sup>b</sup>	APC <sup>c</sup>		Rateb	APC <sup>c</sup>
	2010-2014	2005-2014		2010-2014	2005-2014		2010-2014	2005-2014
All Sites	141.5	-1.4*	All Sites	141.9	-1.3*	All Sites	161.8	-1.5*
Lung and Bronchus	36.3	-1.8*	Lung and Bronchus	37.5	-1.7*	Lung and Bronchus	34.6	-2.0*
Breast	21.2	-1.7*	Breast	20.6	-1.7*	Breast	29.2	-1.6*
Colon and Rectum	12.4	-2.6*	Colon and Rectum	12.1	-2.5*	Colon and Rectum	16.5	-3.6*
Pancreas	9.5	0.0	Pancreas	9.4	0.1	Pancreas	12.1	-0.6*
Ovary	7.4	-2.3*	Ovary	7.7	-2.3*	Corpus and Uterus, NOS	8.1	2.1*
Leukemia	5.1	-1.1*	Leukemia	5.3	-1.0*	Ovary	6.4	-1.3*
Non-Hodgkin Lymphoma	4.6	-2.6*	Non-Hodgkin Lymphoma	4.8	-2.7*	Myeloma	5.5	0.2
Corpus and Uterus, NOS	4.6	1.5*	Corpus and Uterus, NOS	4.2	1.4*	Leukemia	4.6	-1.1*
Liver & IBD <sup>f</sup>	3.7	2.5*	Brain and ONS <sup>f</sup>	3.8	0.4	Liver & IBD <sup>f</sup>	4.5	2.7*
Brain and ONS <sup>f</sup>	3.5	0.3	Liver & IBD <sup>f</sup>	3.5	2.5*	Stomach	4.1	-3.3*
Myeloma	2.7	-0.7	Kidney and Renal Pelvis	2.5	-1.6*	Cervix Uteri	3.8	-2.1*
Kidney and Renal Pelvis	2.4	-1.6*	Myeloma	2.4	-0.9*	Non-Hodgkin Lymphoma	3.5	-1.4*
Stomach	2.3	-2.2*	Urinary Bladder	2.2	-0.5*	Urinary Bladder	2.5	-1.4*
Cervix Uteri	2.3	-0.8*	Cervix Uteri	2.1	-0.4	Kidney and Renal Pelvis		-1.9*
Urinary Bladder	2.2	-0.7*	Stomach	2.0	-2.0*	Brain and ONS <sup>f</sup>	2.1	1.2
Asian/Pacific	: Islander		American Indian/Alaska Native <sup>d</sup>			<u> Hispanic</u> e		
	Rateb	APC°		Rateb	APC°		Rate <sup>b</sup>	APC
		2005-2014			2005-2014			2005-2014
All Sites	88.8	-1.1*	All Sites	129.3	-1.5*	All Sites	97.7	-1.0*
Lung and Bronchus	18.0	-0.6*	Lung and Bronchus	30.8	-1.7*	Breast	14.4	-0.6*
Breast	11.3	-0.7	Breast	14.1	-3.6*	Lung and Bronchus	13.4	-1.3*
Colon and Rectum	8.8	-2.6*	Colon and Rectum	14.0	-0.7	Colon and Rectum	9.2	-2.7*
Pancreas	7.3	0.4	Pancreas	8.1	-1.4	Pancreas	7.7	-0.4
Liver & IBD <sup>f</sup>	6.1	-0.6	Liver & IBD <sup>f</sup>	6.9	1.7	Liver & IBD <sup>f</sup>	5.8	1.1*
Ovary	4.4	-2.0*	Ovary	6.2	-2.3*	Ovary	5.4	-1.8*
Stomach	4.3	-3.9*	Kidney and Renal Pelvis	4.2	0.7	Non-Hodgkin Lymphoma	4.1	-2.1*
Non-Hodgkin Lymphoma	3.2	-1.3*	Stomach	3.8	0.0	Stomach	4.1	-1.7*
Leukemia	3.0	-0.3	Corpus and Uterus, NOS	3.8	1.4	Leukemia	4.0	0.1
Corpus and Uterus, NOS	2.9	2.2*	Leukemia	3.4	-1.3	Corpus and Uterus, NOS	3.6	2.0*
Brain and ONSf	1.8	2.4	Non-Hodgkin Lymphoma	3.2	-5.7*	Cervix Uteri	2.6	-2.5*
Cervix Uteri	1.7	-1.9	Cervix Uteri	2.8	-1.9	Brain and ONS <sup>f</sup>	2.4	0.5
Myeloma	1.3	-2.5	Myeloma	2.7	2.5	Myeloma	2.3	-1.1
Kidney and Renal Pelvis	1.1	-2.1	Brain and ONS <sup>f</sup>	2.0	_	Kidney and Renal Pelvis	2.3	-0.7
Oral Cavity and Pharynx		-3.0*	Gallbladder	1.7	-5.1	Urinary Bladder	1.2	-0.5

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

- <sup>a</sup> Top 15 cancer sites selected based on 2010-2014 age-adjusted rates for the race/ethnic group.
- Pates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- The APC is the Annual Percent Change over the time interval.
- Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.
- IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
- \* The APC is significantly different from zero (p<.05).
- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.