Supplemental Materials #1: Rates and Trends for the Top 15 Cancer Sites By Sex and Race/Ethnicity for 1992-2000

SEER Cancer Incidence, 1992-2000 Top 15 Male Cancer Incidence Sites

Age-Adjusted Rates and Annual Percentage Change (APC)

All Races			White	s		Blacks		
	Rate	<u>APC</u>		Rate	APC		Rate	APC
All Sites	574.2	-1.7 *	All Sites	574.3	-1.8 *	All Sites	724.4	-1.9 *
Prostate	180.6	-3.1 *	Prostate	175.4	-3.4 *	Prostate	286.5	-2.3 *
Lung and Bronchus	84.5	-2.3 *	Lung and Bronchus	83.1	-2.3 *	Lung and Bronchus	126.0	-2.5 *
Colon and Rectum	65.0	-0.9 *	Colon and Rectum	64.9	-1.0 *	Colon and Rectum	73.2	-0.4
Urinary Bladder	36.1	-0.1	Urinary Bladder	39.6	0.0	Oral Cavity and Pharynx	21.8	-3.2 *
Non-Hodgkin Lymphoma	23.7	-0.5	Non-Hodgkin Lymphoma	24.8	-0.5	Urinary Bladder	20.4	0.2
Melanoma of the Skin	19.8	2.8 *	Melanoma of the Skin	23.1	3.1 *	Stomach	20.3	-2.0
Oral Cavity and Pharynx	16.9	-2.3 *	Leukemia	17.2	-1.6 *	Non-Hodgkin Lymphoma	19.0	-1.4
Leukemia	16.2	-1.5 *	Oral Cavity and Pharynx	16.7	-2.2 *	Pancreas	18.4	-1.0 *
Kidney and Renal Pelvis	15.1	1.1 *	Kidney and Renal Pelvis	15.5	1.3 *	Kidney and Renal Pelvis	17.7	1.0
Stomach	13.5	-1.9 *	Pancreas	12.4	-0.2	Myeloma	13.3	-2.1
Pancreas	12.7	-0.4	Stomach	11.7	-2.1 *	Leukemia	13.0	-0.9
Liver and Intrahepatic Bile Duct	8.3	3.7 *	Brain and Other Nervous System	8.5	-0.4	Larynx	13.0	-3.4 *
Brain and Other Nervous System	7.7	-0.6	Esophagus	7.3	1.8 *	Esophagus	12.9	-6.4 *
Esophagus	7.5	0.4	Liver and Intrahepatic Bile Duct	6.7	4.1 *	Liver and Intrahepatic Bile Duct	10.2	4.3 *
Testis	5.0	1.3 *	Testis	5.9	1.3 *	Other Non-Epithelial Skin	7.2	-10.9 *

Hispanics#			American Indian or Alaskan Natives			Asian or Pacific Islanders		
	Rate	<u>APC</u>		<u>Rate</u>	<u>APC</u>		Rate	<u>APC</u>
All Sites	429.7	-1.2 *	All Sites	285.1	-4.8 *	All Sites	401.3	-1.1 *
Prostate	141.8	-1.4 *	Prostate	62.9	-9.8 *	Prostate	106.1	-2.3
Colon and Rectum	48.4	0.9	Lung and Bronchus	51.7	-5.4	Lung and Bronchus	62.7	-0.7
Lung and Bronchus	48.3	-2.4 *	Colon and Rectum	40.5	-3.4	Colon and Rectum	57.6	-0.3
Non-Hodgkin Lymphoma	19.3	-1.2	Kidney and Renal Pelvis	15.4	-5.3	Stomach	24.2	-2.9 *
Stomach	19.0	-2.1	Stomach	14.4	~	Liver and Intrahepatic Bile Duct	20.6	0.7
Urinary Bladder	18.7	-0.6	Oral Cavity and Pharynx	13.0	~	Non-Hodgkin Lymphoma	16.7	0.3
Kidney and Renal Pelvis	14.0	0.3	Non-Hodgkin Lymphoma	8.7	~	Urinary Bladder	16.5	0.3
Liver and Intrahepatic Bile Duct	12.9	2.5	Urinary Bladder	8.4	~	Oral Cavity and Pharynx	13.2	-0.9
Leukemia	11.8	-0.1	Liver and Intrahepatic Bile Duct	8.0	~	Pancreas	11.1	-2.5
Pancreas	10.8	-1.1	Pancreas	7.8	~	Leukemia	10.0	0.4
Oral Cavity and Pharynx	10.6	-3.4 *	Esophagus	5.5	~	Kidney and Renal Pelvis	8.9	0.9
Myeloma	6.4	-1.4	Leukemia	5.0	~	Esophagus	5.4	0.0
Esophagus	6.1	0.0	Myeloma	4.3	~	Myeloma	4.3	-1.6
Brain and Other Nervous System	5.6	0.2	Testis	2.5	~	Brain and Other Nervous System	4.2	-1.6
Testis	3.4	1.6	Soft Tissue including Heart	1.5	~	Thyroid	3.8	-0.3

Rates are per 100,000 and age-adjusted to the 2000 U.S. standard by 5-year age groups.

Data Source - SEER 11 (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey and Los Angeles).

American Indians/Alaskan Natives also include cases diagnosed in Alaska.

Incidence data for Hispanics does not include cases from Detroit or Hawaii.

[~] Statistic could not be calculated. APC based on less than 10 cases for at least one year in within the time interval.

[^] Rate not displayed due to less than 25 cases.

^{*} The APC is significantly different from zero (p<0.05).

[#] Hispanics are not mutually exclusive from Whites, Blacks, Asian or Pacific Islanders, and American Indians or Alaskan Natives.