

Table XXVII-1

AGE-ADJUSTED CANCER SEER INCIDENCE RATES AND 22-YEAR TRENDS, 1973-94

By Primary Cancer Site

All Races, Males and Females

Site	Ages 0-14				Ages 0-19			
	Average 1973-74	Rate <sup>⊕</sup> 93-94	% Change 73-94	EAPC 73-94	Average 1973-74	Rate <sup>⊕</sup> 93-94	% Change 73-94	EAPC 73-94
All Sites								
All Races	12.8	14.2	10.5	0.8★	13.8	15.6	13.4	0.9★
Whites	13.3	14.3	7.6	0.7★	14.2	15.8	11.0	0.8★
Blacks	10.3	13.0	25.5	1.4★	11.1	13.6	22.5	1.1★
Bone & Joint	0.7	0.7	-0.7	0.6	0.8	1.0	16.6	0.9★
Brain & Other nervous	2.4	3.3	35.1	1.8★	2.2	2.9	28.8	1.7★
Hodgkin's disease	0.7	0.6	-20.6	-1.1★	1.3	1.3	-0.5	-0.2
Kidney & Renal pelvis	0.7	1.0	45.6	1.0	0.6	0.8	50.0	1.0★
Leukemias	4.3	3.8	-11.8	0.4	3.8	3.5	-9.6	0.4
Acute lymphocytic	2.7	2.9	4.9	1.0★	2.3	2.4	4.7	1.0★
Non-Hodgkin's lymphomas	0.8	0.9	16.6	0.4	0.8	1.1	35.9	1.2★
Soft tissue	0.7	0.7	5.3	0.4	0.8	0.7	-4.9	0.3

The EAPC is the Estimated Annual Percent Change over the time interval.

§ SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

⊕ The Average Rate is the Average Annual Rate over the specified two-year period.

\* The EAPC is significantly different from zero ( $p < .05$ ).

Table XXVII-2

AGE-ADJUSTED CANCER U.S. MORTALITY RATES\* AND 22-YEAR TRENDS, 1973-94  
 By Race and Primary Cancer Site  
 Males and Females

Race/Site	Ages 0-14				Ages 0-19			
	Average 1973-74	Rate <sup>⊕</sup> 93-94	% Change 73-94	EAPC 73-94	Average 1973-74	Rate <sup>⊕</sup> 93-94	% Change 73-94	EAPC 73-94
<u>All Races</u>								
All Sites	5.4	0.9	-46.0	-3.0*	5.6	0.9	-43.0	-2.7*
Bone & Joint	0.0	0.0	-	-	0.0	0.0	-	-
Brain & Other nervous	0.0	0.0	-	-	0.0	0.0	-	-
Hodgkin's disease	0.0	0.0	-	-	0.0	0.0	-	-
Kidney & Renal pelvis	0.0	0.0	-	-	0.0	0.0	-	-
Leukemias	0.0	0.0	-	-	0.0	0.0	-	-
Acute lymphocytic	0.0	0.0	-	-	0.0	0.0	-	-
Non-Hodgkin's lymphomas	0.6	0.1	-73.2	-6.8*	0.5	0.2	-60.5	-4.8*
Soft tissue	0.0	0.0	-	-	0.0	0.0	-	-
<u>Whites</u>								
All Sites	5.0	0.8	-49.0	-3.0*	5.0	0.8	-43.0	-2.8*
Bone & Joint	0.0	0.0	-	-	0.0	0.0	-	-
Brain & Other nervous	0.0	0.0	-	-	0.0	0.0	-	-
Hodgkin's disease	0.0	0.0	-	-	0.0	0.0	-	-
Kidney & Renal pelvis	0.0	0.0	-	-	0.0	0.0	-	-
Leukemias	0.0	0.0	-	-	0.0	0.0	-	-
Acute lymphocytic	0.0	0.0	-	-	0.0	0.0	-	-
Non-Hodgkin's lymphomas	0.0	0.0	-	-	0.0	0.0	-	-
Soft tissue	0.0	0.0	-	-	0.0	0.0	-	-
<u>Blacks</u>								
All Sites	4.0	0.8	-41.5	-2.6*	3.8	0.8	-38.0	-2.4*
Bone & Joint	0.0	0.0	-	-	0.0	0.0	-	-
Brain & Other nervous	0.0	0.0	-	-	0.0	0.0	-	-
Hodgkin's disease	0.0	0.0	-	-	0.0	0.0	-	-
Kidney & Renal pelvis	0.0	0.0	-	-	0.0	0.0	-	-
Leukemias	0.0	0.0	-	-	0.0	0.0	-	-
Acute lymphocytic	0.0	0.0	-	-	0.0	0.0	-	-
Non-Hodgkin's lymphomas	0.0	0.0	-	-	0.0	0.0	-	-
Soft tissue	0.0	0.0	-	-	0.0	0.0	-	-

The EAPC is the Estimated Annual Percent Change over the time interval.

\* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

⊕ The Average Rate is the Average Annual Rate over the specified two-year period.

\* The EAPC is significantly different from zero ( $p < .05$ ).

Table XXVII-3

AGE-ADJUSTED CANCER SEER INCIDENCE RATES\$, 1973-94

By Primary Cancer Site

All Races, Males and Females

## Year of Diagnosis

<u>Age/Site</u>	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
<u>Ages 0-14</u>																						
All Sites																						
All Races	12.7	13.0	11.3	12.9	12.7	13.0	12.9	12.7	12.2	12.8	12.8	13.7	14.3	14.0	14.0	13.2	14.8	13.9	14.9	13.4	14.5	13.8
Whites	13.0	13.5	11.8	13.2	13.4	13.5	13.5	12.8	12.2	12.9	13.4	14.1	15.3	14.6	14.4	13.7	15.2	13.9	15.7	13.6	14.5	14.0
Blacks	10.5	10.2	8.5	11.1	8.5	9.8	10.4	11.0	13.0	11.8	9.9	10.3	9.9	10.5	11.4	11.8	13.6	12.3	11.7	12.4	12.8	13.2
Bone & Joint	0.8	0.6	0.5	0.5	0.6	0.7	0.8	0.7	0.6	0.6	0.7	0.7	0.8	0.7	0.7	0.6	0.6	0.7	0.9	0.6	0.6	0.8
Brain & Other nervous	2.3	2.6	2.3	2.4	3.0	2.7	2.8	2.8	2.5	2.4	2.2	2.8	2.9	3.4	3.1	3.3	3.2	3.4	3.5	3.2	3.4	3.2
Hodgkin's disease	0.9	0.6	0.7	0.8	0.7	0.7	0.9	0.5	0.6	0.5	0.7	0.7	0.7	0.5	0.7	0.6	0.8	0.7	0.6	0.5	0.6	0.6
Kidney & Renal pelvis	0.7	0.7	0.6	0.9	0.7	0.9	0.9	0.8	0.8	0.9	1.1	0.7	0.8	0.9	0.8	0.7	1.1	0.7	0.9	0.7	0.9	1.1
Leukemias	3.8	4.8	3.2	4.0	3.8	3.8	3.6	3.9	3.9	3.7	3.6	4.4	4.5	4.0	4.4	4.0	4.6	4.3	4.4	3.9	4.1	3.5
Acute lymphocytic	2.4	3.1	2.1	2.9	3.0	3.0	2.7	3.0	3.1	2.9	2.9	3.4	3.4	3.2	3.3	3.2	3.8	3.3	3.4	3.0	2.9	2.8
Non-Hodgkin's lymph	0.6	0.9	1.0	0.8	0.8	1.0	0.8	0.7	0.7	0.9	0.9	0.9	0.7	1.0	1.1	0.7	0.8	0.8	0.9	0.8	0.8	1.0
Soft tissue	0.8	0.7	0.8	0.7	0.7	0.8	0.6	0.9	0.7	0.9	0.9	0.7	0.8	0.9	0.8	0.6	0.9	0.8	1.0	0.7	0.6	0.9
<u>Ages 0-19</u>																						
All Sites																						
All Races	13.6	13.9	12.8	14.1	14.1	14.4	14.5	14.1	13.7	14.4	14.4	15.3	15.7	15.6	15.2	15.0	16.4	15.5	16.1	15.9	15.9	15.4
Whites	14.1	14.3	13.2	14.7	14.7	15.0	15.1	14.6	14.0	14.9	15.2	15.9	16.7	16.4	16.0	15.5	17.0	15.7	17.1	16.4	15.9	15.7
Blacks	10.3	11.8	10.5	10.7	10.4	10.8	12.6	11.0	12.9	11.8	11.1	12.1	10.4	12.0	11.2	12.9	14.0	13.0	12.4	12.7	13.3	13.8
Bone & Joint	0.9	0.8	0.7	0.6	0.7	0.9	0.9	0.9	1.0	0.8	0.8	0.9	1.0	0.9	1.0	0.8	0.9	0.8	1.0	0.9	0.9	1.0
Brain & Other nervous	2.1	2.3	2.1	2.2	2.9	2.5	2.5	2.4	2.3	2.2	2.2	2.5	2.6	3.2	2.9	3.0	2.9	3.1	3.1	3.2	2.9	2.8
Hodgkin's disease	1.4	1.1	1.5	1.4	1.4	1.5	1.6	1.2	1.2	1.5	1.4	1.5	1.4	1.3	1.4	1.3	1.7	1.3	1.4	1.4	1.3	1.3
Kidney & Renal pelvis	0.6	0.5	0.5	0.7	0.6	0.7	0.7	0.6	0.6	0.7	0.8	0.6	0.7	0.7	0.7	0.6	0.8	0.6	0.7	0.5	0.8	0.9
Leukemias	3.5	4.2	3.0	3.6	3.4	3.4	3.2	3.4	3.6	3.4	3.4	3.9	4.0	3.6	3.7	3.5	4.1	3.7	4.0	3.6	3.6	3.3
Acute lymphocytic	2.0	2.7	1.9	2.4	2.5	2.5	2.3	2.4	2.7	2.5	2.5	2.9	2.8	2.7	2.7	2.7	3.3	2.8	2.9	2.5	2.5	2.4
Non-Hodgkin's lymph	0.7	0.9	1.0	0.9	0.8	1.0	0.9	1.0	1.0	1.1	1.0	1.0	1.1	1.0	1.0	1.0	1.0	1.1	1.0	0.9	0.9	1.3
Soft tissue	0.7	0.8	0.8	0.7	0.7	0.9	0.7	0.9	0.7	1.0	1.0	0.7	0.9	0.9	0.7	0.9	0.8	1.0	0.7	0.7	0.7	0.7

§ SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-4

AGE-ADJUSTED CANCER U.S. MORTALITY RATES\*, 1973-94

By Primary Cancer Site

All Races, Males and Females

## Year of Diagnosis

<u>Age/Site</u>	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
<u>Ages 0-14</u>																						
All Sites	5.5	5.3	4.9	4.9	4.7	4.3	4.4	4.3	4.2	4.2	4.1	3.6	3.6	3.5	3.4	3.3	3.3	3.1	3.1	3.0	3.0	2.8
Bone & Joint	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1
Brain & Other nervous	1.0	1.1	1.0	1.0	0.8	0.9	0.9	0.9	1.0	0.9	0.8	0.8	0.9	0.7	0.8	0.9	0.9	0.9	0.8	0.8	0.8	
Hodgkin's disease	0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Kidney & Renal pelvis	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	
Leukemias	2.4	2.3	2.1	2.1	2.0	1.9	1.8	1.7	1.6	1.6	1.5	1.3	1.4	1.3	1.3	1.2	1.2	1.1	1.1	1.1	1.1	
Acute lymphocytic	1.3	1.3	1.2	1.2	1.1	1.0	1.0	0.9	1.0	0.9	0.8	0.8	0.8	0.7	0.7	0.6	0.6	0.5	0.5	0.5	0.4	
Non-Hodgkin's lymph	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.1	
Soft tissue	0.6	0.5	0.5	0.6	0.6	0.5	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	
<u>Ages 0-19</u>																						
All Sites	5.8	5.5	5.1	5.1	5.0	4.5	4.6	4.6	4.4	4.5	4.3	3.9	3.8	3.8	3.6	3.5	3.5	3.4	3.4	3.3	3.3	3.2
Bone & Joint	0.4	0.4	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Brain & Other nervous	0.9	1.0	0.9	0.9	0.9	0.8	0.9	0.8	0.8	0.9	0.8	0.8	0.8	0.8	0.7	0.8	0.8	0.8	0.7	0.8	0.7	
Hodgkin's disease	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	
Kidney & Renal pelvis	0.2	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	
Leukemias	2.3	2.2	2.0	2.1	2.0	1.8	1.8	1.8	1.6	1.6	1.5	1.4	1.4	1.3	1.4	1.2	1.2	1.2	1.2	1.1	1.1	
Acute lymphocytic	1.2	1.1	1.0	1.1	1.0	0.9	1.0	0.9	0.9	0.9	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.5	0.6	0.5	0.5	
Non-Hodgkin's lymph	0.5	0.5	0.4	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	
Soft tissue	0.5	0.5	0.5	0.5	0.5	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	

\* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-5

AGE-ADJUSTED CANCER U.S. MORTALITY RATES\*, 1973-94

By Primary Cancer Site

Whites, Males and Females

## Year of Diagnosis

<u>Age/Site</u>	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
<u>Ages 0-14</u>																						
All Sites	5.6	5.4	5.0	5.0	4.8	4.3	4.5	4.4	4.2	4.3	4.1	3.7	3.6	3.6	3.5	3.3	3.4	3.2	3.2	3.1	3.1	2.8
Bone & Joint	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1
Brain & Other nervous	1.0	1.1	1.0	1.0	0.8	0.9	0.9	0.9	1.0	0.9	0.9	0.9	0.9	0.9	0.8	0.8	0.9	0.9	0.9	0.8	0.8	0.8
Hodgkin's disease	0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Kidney & Renal pelvis	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Leukemias	2.5	2.4	2.3	2.3	2.1	1.9	1.9	1.8	1.7	1.6	1.6	1.4	1.4	1.4	1.4	1.2	1.2	1.1	1.1	1.2	1.1	1.0
Acute lymphocytic	1.4	1.4	1.2	1.3	1.2	1.1	1.1	1.0	1.0	0.9	0.9	0.8	0.8	0.8	0.7	0.7	0.6	0.6	0.6	0.5	0.6	0.5
Non-Hodgkin's lymph	0.5	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2
Soft tissue	0.6	0.5	0.5	0.6	0.6	0.5	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1
<u>Ages 0-19</u>																						
All Sites	5.9	5.6	5.2	5.2	5.1	4.6	4.7	4.7	4.4	4.5	4.4	4.0	3.9	3.9	3.7	3.6	3.6	3.4	3.5	3.4	3.3	3.2
Bone & Joint	0.4	0.4	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2
Brain & Other nervous	0.9	1.0	0.9	0.9	0.9	0.8	0.9	0.8	0.8	0.9	0.9	0.8	0.8	0.8	0.7	0.8	0.8	0.8	0.8	0.8	0.8	0.7
Hodgkin's disease	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0
Kidney & Renal pelvis	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Leukemias	2.4	2.3	2.1	2.2	2.1	1.9	1.9	1.8	1.7	1.7	1.6	1.4	1.4	1.4	1.4	1.3	1.2	1.2	1.2	1.2	1.2	1.1
Acute lymphocytic	1.3	1.2	1.1	1.1	1.1	1.0	1.0	1.0	0.9	0.9	0.8	0.8	0.8	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.5	0.5
Non-Hodgkin's lymph	0.5	0.5	0.5	0.4	0.4	0.3	0.4	0.3	0.4	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2
Soft tissue	0.5	0.5	0.5	0.5	0.5	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2

\* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-6

AGE-ADJUSTED CANCER U.S. MORTALITY RATES\*, 1973-94

By Primary Cancer Site

Blacks, Males and Females

## Year of Diagnosis

<u>Age/Site</u>	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
<u>Ages 0-14</u>																						
All Sites	5.2	4.4	4.5	4.2	4.2	4.1	4.0	3.8	4.2	4.0	3.9	3.5	3.3	3.4	3.0	3.2	3.2	3.0	3.1	2.7	2.7	2.9
Bone & Joint	0.2	0.2	0.3	0.3	0.2	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Brain & Other nervous	1.2	1.1	0.9	1.0	1.0	0.8	0.9	0.9	1.0	0.8	0.8	0.7	0.6	0.9	0.7	1.0	0.8	0.7	0.8	0.7	0.6	0.8
Hodgkin's disease	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.1
Kidney & Renal pelvis	0.4	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.1
Leukemias	1.8	1.6	1.6	1.6	1.4	1.5	1.3	1.2	1.6	1.5	1.2	1.3	1.2	1.1	1.1	1.0	0.9	1.0	0.9	1.0	0.8	0.9
Acute lymphocytic	0.7	0.7	0.8	0.8	0.7	0.6	0.6	0.6	0.8	0.7	0.7	0.7	0.7	0.7	0.6	0.5	0.4	0.5	0.5	0.4	0.4	0.3
Non-Hodgkin's lymph	0.3	0.3	0.4	0.2	0.3	0.4	0.3	0.2	0.2	0.2	0.2	0.1	0.2	0.3	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1
Soft tissue	0.6	0.4	0.4	0.5	0.5	0.5	0.3	0.1	0.3	0.3	0.3	0.2	0.2	0.1	0.3	0.2	0.2	0.3	0.1	0.2	0.2	0.3
<u>Ages 0-19</u>																						
All Sites	5.6	4.8	4.7	4.5	4.7	4.5	4.3	4.2	4.5	4.3	3.8	3.7	3.5	3.7	3.6	3.5	3.5	3.4	3.4	3.0	3.1	3.3
Bone & Joint	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2
Brain & Other nervous	1.1	1.0	0.9	0.9	0.9	0.7	0.9	0.8	0.9	0.8	0.7	0.7	0.6	0.8	0.7	0.9	0.7	0.7	0.7	0.7	0.7	0.8
Hodgkin's disease	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1
Kidney & Renal pelvis	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.1
Leukemias	1.8	1.5	1.6	1.5	1.4	1.5	1.4	1.4	1.5	1.5	1.1	1.3	1.1	1.1	1.2	1.0	1.1	1.0	1.1	0.9	1.0	1.0
Acute lymphocytic	0.7	0.6	0.7	0.7	0.7	0.6	0.6	0.6	0.7	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.3	0.3	0.3
Non-Hodgkin's lymph	0.5	0.3	0.4	0.3	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.3	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2
Soft tissue	0.5	0.4	0.5	0.5	0.5	0.5	0.3	0.2	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.3

\* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-7

AGE-ADJUSTED CANCER SEER INCIDENCES AND U.S. MORTALITY\* RATES, 1990-94

By Primary Cancer Site

All Races, Males and Females

Site	Ages 0-14						Ages 0-19					
	Incidence			Mortality			Incidence			Mortality		
	Total	Males	Females									
<b>All Sites</b>												
All Races	14.1	15.0	13.2	3.0	3.3	2.7	15.8	16.5	15.0	3.3	3.7	2.9
Whites	14.4	15.3	13.4	3.1	3.4	2.7	16.2	17.1	15.2	3.4	3.8	2.9
Blacks	12.5	12.7	12.3	2.9	3.1	2.6	13.0	12.8	13.3	3.2	3.5	3.0
Bone & Joint	0.7	0.7	0.8	0.1	0.1	0.1	0.9	1.0	0.8	0.2	0.2	0.2
Brain & Other nervous	3.3	3.6	3.1	0.8	0.9	0.8	3.0	3.3	2.7	0.8	0.8	0.7
Hodgkin's disease	0.6	0.6	0.5	0.0	0.0	0.0	1.3	1.3	1.4	0.1	0.1	0.0
Kidney & Renal pelvis	0.9	0.9	0.8	0.1	0.1	0.1	0.7	0.7	0.7	0.1	0.1	0.1
Leukemias	4.0	4.2	3.8	1.1	1.2	0.9	3.6	3.9	3.4	1.1	1.3	0.9
Acute lymphocytic	3.1	3.3	2.9	0.5	0.6	0.4	2.6	2.9	2.3	0.5	0.6	0.4
Non-Hodgkin's lymphomas	0.9	1.2	0.5	0.2	0.2	0.1	1.1	1.4	0.7	0.2	0.3	0.1
Soft tissue	0.8	0.8	0.8	0.2	0.2	0.1	0.8	0.8	0.8	0.2	0.2	0.2

<sup>§</sup> SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

\* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

Table XXVII-8

5-YEAR RELATIVE SURVIVAL RATES (Percent)

By Selected Primary Cancer Site and Year Of Diagnosis

All Races, Males and Females

Site	Ages 0-14					Ages 0-19				
	1974-76	1977-79	1980-82	1983-85	1986-93	1974-76	1977-79	1980-82	1983-85	1986-93
<b>All Sites</b>										
All Races	55.6	61.4	64.6	67.8	71.8S	58.4	63.9	66.4	69.7	73.4S
Whites	55.6	61.9	65.1	69.3	72.6S	58.7	64.4	67.3	71.3	74.7S
Blacks	56.8	57.4	61.2	57.2	66.3S	56.6	59.9	61.2	58.7	64.4S
Bone & Joint	53.7†	51.8†	54.1†	58.5†	63.5S	50.3	50.9	49.8	53.7	63.1S
Brain & Other nervous	54.4	56.5	55.3	62.0	61.0S	55.2	56.7	56.8	63.4	63.6S
Hodgkin's disease	79.0	83.4	90.8	89.6	93.5S	85.4	87.9	87.1	88.1	91.0S
Leukemias	44.9	58.8	61.9	61.3	72.2S	40.6	53.2	57.4	58.1	67.8S
Acute lymphocytic	52.8	66.8	70.3	69.4	79.8S	48.8	62.7	66.5	66.8	76.4S
Acute myeloid	14.1	26.2†	21.1†	32.0†	37.2S	16.2	25.4	21.7	33.3†	38.0S
Neuroblastoma	52.2	53.2	52.8	54.0	64.5S	51.7	53.2	53.0	53.3	63.0S
Non-Hodgkin's lymphomas	44.6	50.7	61.8	70.4	73.0S	44.0	49.7	59.4	66.4	72.3S
Soft tissue	60.3	68.6	65.0	76.4	72.6S	61.8	64.6	64.5	74.0	67.9
Wilms' tumor	74.3	77.3	85.9	86.4	91.7S	73.6	76.8	85.3	86.7	91.4S

Rates are from the SEER Program. They are based on data from population-based registries in Connecticut, New Mexico, Utah, Iowa, Hawaii, Atlanta, Detroit, Seattle-Puget Sound and San Francisco-Oakland. Rates are based on follow-up of patients through 1994.

S The difference in rates between 1974-76 and 1986-93 is statistically significant ( $p<.05$ ).

† The standard error of the survival rate is between 5 and 10 percentage points.

† The standard error of the survival rate is greater than 10 percentage points.

Table XXVII-9

**CHILDHOOD CANCER**  
**SEER Incidence, U.S. Mortality and 5-Year Relative Survival Rates**  
**For Top Cancer Sites<sup>⊕</sup> By 5-Year Age Groups**  
**All Races, Males and Females**

Age/Site	SEER Incidences				US Mortality*				Survivals	
	1990-94 Num Cases	1990-94 Rate	1973-94 PC	1973-94 EAPC	1990-94 Num Cases	1990-94 Rate	1973-94 PC	1973-94 EAPC	1974-76	1986-93
<b>Ages 0-4</b>										
All Sites	1,871	20.2	17.3	0.9★	2,967	3.1	-46.9	-3.0★	56.1↑	72.0SS
Soft tissue	119	1.3	10.1	0.0000	140	0.1	-86.7	-10.5★	57.7↑	76.0SS
Kidney and Renal pelvis	186	2.0	10.1	0.0000	92	0.1	-67.3	-6.0★	74.2	89.1SS
Eye and Orbit	136	1.5	10.0	0.4	59	0.1	-64.7	-3.3★	89.2	92.1SS
Brain and Other nervous	358	3.9	46.9	2.5★	731	0.8	-20.4	-1.3★	51.2	53.0SS
Other endocrine	123	1.3	-	-	552	0.6	-	-	32.8↑	57.1SS
Leukemias	622	6.7	-10.5	0.2	974	1.0	-54.2	-3.7★	50.6	75.2SS
Acute lymphocytic	493	5.3	-10.6	0.8★	327	0.3	-68.7	-5.6★	57.9	82.8
<b>Ages 5-9</b>										
All Sites	943	10.8	9.0	0.6★	2,831	3.1	-49.0	-3.3★	55.5	71.6S
Brain and Other nervous	292	3.4	29.3	1.4★	899	1.0	-23.3	-0.9★	54.4	63.4
Leukemias	308	3.5	-13.3	0.6	1,008	1.1	-65.4	-5.3★	46.7	74.2SS
Acute lymphocytic	240	2.8	-8.6	1.2	537	0.6	-71.5	-6.3★	53.4	80.5S
<b>Ages 10-14</b>										
All Sites	1,043	12.2	3.2	0.7★	2,616	2.9	-41.5	-2.5★	54.8↑	71.6SS
Bone and Joint	115	1.3	-10.7	0.4	252	0.3	-46.9	-3.1★	52.8↑	63.3SS
Brain and Other nervous	248	2.9	30.2	1.6★	647	0.7	-23.2	-1.3★	57.8	70.2SS
Hodgkin's disease	109	1.3	-19.9	-0.84	38	0.0	-66.9	-5.1★	75.9	95.0SS
Leukemias	197	2.0	-15.9	-0.7	983	1.1	-48.7	-3.0★	29.0	57.0SS
Acute lymphocytic	134	1.6	-15.9	1.2	482	0.5	-50.2	-3.4★	37.2↑	65.9
<b>Ages 15-19</b>										
All Sites	1,669	20.8	20.1	1.1★	3,682	4.2	-35.1	-2.1★	63.8	76.8SS
Bone and Joint	128	1.6	45.5	1.0000	438	0.5	-38.9	-2.6★	46.6↑	62.0
Melanomas of skin	126	1.4	66.0	0.6000	44	0.1	-48.1	-3.9★	76.7↑	94.0
Ovary(females)	196	2.4	66.0	0.6000	40	0.1	-70.3	-6.4★	69.0↑	90.0
Testis(males)	130	3.1	55.5	1.0000	49	0.0	-78.0	-8.5★	55.4↑	93.0
Brain and Other nervous	167	2.1	-7.2	-0.9	547	0.6	-14.5	-0.9★	58.4↑	75.0
Thyroid	116	1.4	-7.2	-0.9	-	-	-	-	100.0	97.0
Hodgkin's disease	285	3.6	14.4	1.0000	139	0.2	-70.9	-4.7★	88.8↑	89.0
Non-Hodgkin's lymphomas	130	1.6	88.0	0.2000	324	0.4	-45.4	-2.5★	42.6↑	70.8SS
Leukemias	197	1.6	88.0	0.2000	1,216	1.0	-32.0	-1.9★	21.5↑	41.0SS
Acute lymphocytic	194	1.2	88.0	0.2000	518	0.6	-31.1	-1.6★	21.2↑	45.7

The PC is the Percent Change over the time interval.

The EAPC is the Estimated Annual Percent Change over the time interval.

Incidence and mortality rates are per 100,000. Survival rates are expressed as percents.

SEER Program.

NCHS public use tape.

The EAPC is significantly different from zero ( $p < .05$ ).

The difference in rates between 1974-76 and 1986-93 is statistically significant ( $p < .05$ ).

The standard error of the survival rate is between 5 and 10 percentage points.

The standard error of the survival rate is greater than 10 percentage points.

Sites were chosen based on more than 100 SEER incident cases between 1990-94.

Statistic could not be calculated.

Note:

\*\*

★

S

↑

↑↑

⊕

-

Table XXVII-10  
ALL SITES (Invasive)

INCIDENCE, MORTALITY, AND SURVIVAL RATES

By Year of Diagnosis/Death

All Races, Males and Females

Ages 0-14

Year of Diagnosis/Death

	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
<b>Incidence</b>																						
SEER	12.7	13.0	11.3	12.9	12.7	13.0	12.9	12.7	12.2	12.8	12.8	13.7	14.3	14.0	14.0	13.2	14.8	13.9	14.9	13.4	14.5	13.8
<b>Mortality</b>																						
U.S.	5.5	5.3	4.9	4.9	4.7	4.3	4.4	4.3	4.2	4.2	4.1	3.6	3.6	3.5	3.4	3.3	3.3	3.1	3.1	3.0	3.0	2.8
SEER	5.8	5.9	5.0	4.7	4.9	4.3	4.6	4.2	4.1	4.3	4.5	3.6	4.0	4.1	3.8	3.6	2.8	3.2	3.4	2.9	3.1	2.6
<b>Relative Survival Rates (SEER)</b>																						
1-year	76.3	75.7	79.8	80.5	82.6	81.9	81.7	84.0	83.9	85.9	84.9	86.1	87.1	85.6	85.6	88.5	88.8	89.3	90.9	91.1	88.3	
2-year	63.4	63.1	66.6	70.1	71.4	73.7	72.7	74.4	74.4	76.2	76.6	76.2	78.0	76.2	76.4	80.1	80.0	81.5	82.0	82.0	84.1	
3-year	57.8	57.7	61.0	63.7	65.4	69.3	66.5	68.2	69.9	71.2	72.2	71.2	73.8	72.3	71.6	76.4	75.8	77.2	78.2			
4-year	53.8	55.4	58.5	60.0	62.3	65.6	62.7	64.4	65.5	69.7	69.5	67.1	70.5	70.7	69.7	73.2	73.1	74.7				
5-year	50.8	53.3	55.4	57.8	59.9	63.4	61.0	62.1	64.1	67.7	67.3	66.3	69.6	68.3	69.1	72.1	71.5					
6-year	49.7	51.3	53.1	56.4	58.6	61.6	59.3	60.5	63.0	66.1	66.3	65.5	68.2	67.3	67.7	71.6						
7-year	48.2	50.3	51.3	55.1	57.6	61.0	58.1	59.7	61.6	65.5	66.0	64.4	67.0	67.0	66.1							
8-year	47.3	49.9	50.2	54.7	57.1	60.3	56.9	58.8	60.8	65.0	65.3	63.4	65.7	66.6								
9-year	46.9	49.0	49.9	54.4	56.8	59.7	56.8	57.9	60.1	64.7	65.2	62.8	65.0									
10-year	46.1	48.9	49.3	54.1	56.6	59.3	56.4	57.4	59.7	64.0	64.7	62.2										
11-year	45.6	48.6	49.3	54.1	56.1	58.8	55.7	57.3	58.9	63.6	64.3											
12-year	45.0	48.4	48.8	53.5	56.0	58.5	55.4	56.9	58.1	63.1												
13-year	44.4	48.3	48.8	53.2	55.9	58.5	55.1	56.8	58.1													
14-year	44.1	48.2	48.8	53.0	55.2	57.7	54.7	56.0														
15-year	43.5	48.0	48.8	52.9	55.0	57.4	53.9															
16-year	43.3	47.5	48.8	52.2	54.3	56.8																
17-year	43.2	47.4	48.4	51.7	54.1																	
18-year	42.8	46.8	48.0	51.3																		
19-year	42.8	46.8	48.0																			
20-year	42.8	46.7																				
21-year	42.8																					

Note: Incidence and mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Survival rates are relative rates expressed as percents.

Table XXVII-11  
ALL SITES (Invasive)

INCIDENCE, MORTALITY, AND SURVIVAL RATES

By Year of Diagnosis/Death

All Races, Males and Females

Ages 0-19

Year of Diagnosis/Death

	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
<b>Incidence</b>																						
SEER	13.6	13.9	12.8	14.1	14.1	14.4	14.5	14.1	13.7	14.4	14.4	15.3	15.7	15.6	15.2	15.0	16.4	15.5	16.1	15.9	15.9	15.4
<b>Mortality</b>																						
U.S.	5.8	5.5	5.1	5.1	5.0	4.5	4.6	4.6	4.4	4.5	4.3	3.9	3.8	3.8	3.6	3.5	3.5	3.4	3.4	3.3	3.3	3.2
SEER	5.8	5.9	5.4	5.2	4.9	4.4	5.0	4.4	4.4	4.7	4.8	3.8	4.2	4.3	3.9	3.8	3.1	3.6	3.6	3.4	3.5	3.1
<b>Relative Survival Rates (SEER)</b>																						
1-year	77.1	77.7	81.4	82.4	84.0	83.2	84.0	86.2	84.9	85.9	86.9	87.4	87.7	88.1	87.9	90.0	89.7	90.3	91.5	90.4	89.5	
2-year	65.6	64.3	69.2	73.0	74.0	73.9	74.3	76.6	74.8	77.7	79.3	77.8	78.5	79.0	79.5	82.6	81.3	82.7	83.4	82.7	83.2	
3-year	60.5	58.8	64.1	67.5	68.6	69.5	69.2	70.3	70.1	72.5	74.9	73.2	75.0	75.5	74.8	78.4	77.5	78.2	79.4			
4-year	56.9	56.6	61.3	63.9	65.3	66.5	65.8	67.9	66.3	70.5	72.0	69.1	72.3	73.6	72.3	75.6	74.1	75.7				
5-year	54.2	54.3	58.5	62.2	63.2	64.5	64.0	65.9	64.9	68.6	69.6	68.1	71.4	71.6	71.7	74.6	72.7					
6-year	52.7	52.5	56.1	60.5	61.9	63.0	62.0	64.5	63.5	67.3	68.3	66.6	71.4	71.6	71.7	74.6	72.7					
7-year	51.2	51.5	54.8	59.3	61.2	62.4	60.9	63.9	62.6	66.6	67.6	65.7	68.9	70.3	69.2							
8-year	50.6	50.7	54.0	58.7	60.6	61.8	59.4	63.3	61.9	66.0	66.5	64.5	67.9	69.9								
9-year	50.4	49.8	53.4	58.3	59.9	60.9	59.4	62.7	61.4	65.1	66.0	63.8	67.2									
10-year	49.8	49.7	52.8	57.8	59.3	60.8	59.2	62.2	61.0	64.4	65.6	63.2										
11-year	49.3	49.4	52.8	57.5	58.9	60.3	58.6	61.9	60.2	64.0	65.4											
12-year	48.8	49.2	52.3	57.0	58.9	60.0	58.0	61.8	59.4	63.7												
13-year	48.4	48.6	52.3	56.8	58.8	59.8	57.7	61.5	59.4													
14-year	47.6	48.5	52.2	56.6	58.2	59.3	57.5	60.6														
15-year	47.0	48.2	52.1	56.6	57.7	59.1	56.9															
16-year	46.8	47.9	52.1	56.0	57.1	58.5																
17-year	46.2	47.3	51.8	55.6	57.1																	
18-year	45.7	46.8	51.3	55.4																		
19-year	45.6	46.7	51.3																			
20-year	45.3	46.6																				
21-year	45.3																					

Note: Incidence and mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Survival rates are relative rates expressed as percents.