

Table V-1
CERVIX UTERI CANCER (Invasive)

TRENDS IN SEER INCIDENCE AND U.S. MORTALITY USING THE JOINPOINT REGRESSION PROGRAM, 1973-1997
WITH UP TO THREE JOINPOINTS BY RACE AND AGE

	All Races		Whites		Blacks	
	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence</u>						
Age - All Ages	1973-81	-4.8★	1973-82	-5.0★	1973-97	-4.0★
	1981-97	-1.1★	1982-90	0.4		
			1990-97	-2.7★		
Age - Under 50	1973-81	-4.3★	1973-81	-4.7★	1973-97	-3.8★
	1981-97	-0.4	1981-89	1.4		
			1989-97	-1.8★		
Age - 50 and Over	1973-82	-4.9★	1973-85	-4.9★	1973-97	-4.1★
	1982-97	-1.7★	1985-91	0.9		
			1991-97	-4.5★		
<u>U.S. Cancer Mortality</u>						
Age - All Ages	1973-82	-4.7★	1973-83	-4.5★	1973-79	-5.8★
	1982-97	-1.5★	1983-97	-1.2★	1979-97	-2.8★
Age - Under 50	1973-79	-5.8★	1973-79	-5.4★	1973-81	-6.0★
	1979-97	-0.8★	1979-97	-0.4★	1981-97	-2.1★
Age - 50 and Over	1973-83	-4.7★	1973-84	-4.8★	1973-79	-5.2★
	1983-97	-1.9★	1984-97	-1.6★	1979-97	-2.9★

The APC is the Annual Percent Change based on rates age-adjusted to the 1970 U.S. standard population using the Joinpoint Regression Program.

* The APC is significantly different from zero ($p < .05$).

Table V-2
CERVIX UTERI CANCER (Invasive)

SEER INCIDENCES AND U.S. MORTALITY* RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	Incidence§			US Mortality*		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<u>AGE-SPECIFIC RATES, 1993-97</u>						
AGE AT DIAGNOSIS/DEATH:						
0-4	0.0	0.0	0.0	0.0	0.0	0.0
5-9	0.0	0.0	0.0	0.0	0.0	0.0
10-14	0.0	0.0	0.0	0.0	0.0	0.0
15-19	0.3	0.3	0.4	0.0	0.0	0.0
20-24	2.5	2.5	3.4	0.2	0.2	0.4
25-29	7.6	8.2	5.2	0.9	0.8	1.3
30-34	11.7	12.2	11.1	1.9	1.9	2.9
35-39	14.2	13.8	14.5	3.1	2.8	5.2
40-44	14.4	12.9	20.0	4.2	3.7	7.6
45-49	16.7	15.1	21.8	5.3	4.7	10.0
50-54	15.8	13.8	22.9	6.2	5.4	11.8
55-59	15.6	12.7	28.3	6.5	5.6	12.9
60-64	16.2	14.1	24.3	7.3	6.2	15.1
65-69	15.8	13.4	27.8	7.7	6.7	17.1
70-74	14.4	13.0	26.8	8.2	7.1	20.0
75-79	15.0	12.5	34.0	8.7	7.5	22.1
80-84	14.5	11.1	48.5	10.3	8.8	27.7
85+	13.3	10.2	47.9	12.0	10.3	32.8
<u>AGE-ADJUSTED RATES, 1993-97</u>						
AGE AT DIAGNOSIS/DEATH:						
All ages	7.8	7.1	11.4	2.7	2.4	5.6
Under 65	7.0	6.5	9.1	2.1	1.8	3.9
65 and over	14.9	12.6	32.6	8.7	7.5	21.2
All ages (world std.)⊕	7.4	6.8	10.2	2.4	2.1	4.8

§ or * SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population, except where noted.
NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population, except where noted.

⊕ Rates are per 100,000 and are age-adjusted to the world standard population.

Table V-3
CERVIX UTERI CANCER (Invasive)

SEER INCIDENCES AND U.S. MORTALITY* RATES, AGE-ADJUSTED RATES, BY RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1973	14.2	8.7	31.2	12.8	7.7	28.8	30.0	18.1	66.8
1974	12.7	7.6	28.7	11.6	7.0	26.0	25.2	12.0	65.7
1975	12.4	7.3	28.2	11.1	6.7	24.6	27.9	13.2	73.2
1976	11.9	7.2	26.6	10.8	6.7	23.6	24.6	12.0	63.6
1977	10.9	6.4	24.7	9.8	6.7	22.4	22.9	12.4	55.4
1978	10.5	6.3	23.5	9.2	5.8	20.8	19.9	9.4	52.3
1979	10.6	6.6	23.1	9.1	5.8	19.7	23.5	12.0	59.3
1980	10.2	6.1	22.8	9.0	5.5	20.4	19.0	10.0	46.8
1981	9.0	5.5	19.6	8.0	5.0	17.3	19.0	10.1	46.5
1982	8.9	5.4	19.7	7.9	5.1	16.5	17.8	7.6	49.0
1983	8.8	5.6	18.7	8.1	5.2	16.9	15.3	8.9	35.0
1984	8.9	6.1	18.7	8.3	5.7	16.4	17.2	9.5	41.0
1985	8.5	5.3	18.4	7.6	5.1	15.3	15.9	6.8	43.7
1986	8.0	5.9	18.6	8.0	5.5	16.0	15.4	8.1	38.1
1987	8.3	5.4	17.4	7.5	5.2	14.5	15.4	6.6	42.6
1988	8.9	6.0	17.7	8.0	5.6	15.1	15.5	8.0	38.6
1989	8.9	5.9	18.0	8.3	5.9	15.6	13.2	5.9	35.7
1990	8.9	6.1	17.7	8.4	5.9	16.0	13.9	7.2	34.5
1991	8.4	5.3	18.0	7.7	5.1	15.7	13.5	6.9	34.0
1992	8.3	5.6	16.5	7.9	5.5	15.2	11.2	7.3	23.1
1993	8.2	5.5	16.4	7.7	5.2	14.7	11.2	6.6	25.6
1994	8.0	5.3	16.4	7.4	5.2	14.1	11.8	5.3	31.7
1995	7.5	5.1	14.8	6.6	4.8	12.3	11.4	6.1	27.9
1996	7.9	5.6	15.1	7.1	5.3	12.7	11.3	5.9	27.7
1997	7.5	5.2	14.8	6.7	4.9	12.2	11.5	6.2	27.6
1993-97	7.8	5.3	15.5	7.1	5.1	13.2	11.4	6.0	28.1
<u>U.S. MORTALITY RATES</u>									
<u>YEAR OF DEATH:</u>									
1973	5.2	2.0	15.0	4.4	1.6	12.9	13.1	5.0	37.9
1974	5.0	1.9	14.6	4.3	1.6	12.9	11.7	4.6	33.6
1975	4.6	1.8	13.4	3.9	1.4	11.6	11.1	4.2	32.1
1976	4.5	1.6	13.4	3.9	1.4	11.5	10.7	3.6	32.4
1977	4.1	1.5	12.3	3.5	1.2	10.6	10.1	3.7	30.0
1978	4.0	1.6	11.6	3.4	1.3	10.0	9.6	3.5	28.3
1979	3.8	1.4	11.2	3.3	1.2	9.8	8.8	3.3	25.8
1980	3.7	1.5	10.6	3.1	1.2	9.1	8.9	3.4	25.8
1981	3.6	1.4	10.5	3.1	1.2	9.0	8.2	2.7	24.9
1982	3.4	1.3	9.8	2.9	1.1	8.3	8.1	2.7	24.6
1983	3.3	1.3	9.6	2.8	1.1	8.1	8.1	2.9	24.3
1984	3.3	1.3	9.2	2.8	1.1	7.8	7.6	2.9	22.1
1985	3.1	1.3	8.9	2.7	1.1	7.6	7.3	2.5	22.3
1986	3.2	1.3	8.9	2.7	1.1	7.5	7.7	3.0	22.2
1987	3.0	1.3	8.3	2.6	1.1	7.1	6.8	2.7	19.6
1988	3.0	1.3	8.3	2.6	1.1	7.0	6.7	2.2	20.6
1989	3.0	1.3	8.2	2.5	1.1	6.8	7.1	2.4	21.5
1990	3.0	1.3	8.3	2.6	1.2	7.1	6.4	2.5	18.5
1991	2.9	1.3	8.0	2.5	1.1	6.7	6.4	2.3	18.8
1992	2.9	1.3	7.9	2.4	1.1	6.6	6.7	2.5	19.6
1993	2.8	1.2	7.7	2.4	1.1	6.6	6.3	2.3	18.7
1994	2.8	1.2	7.7	2.5	1.1	6.8	5.6	2.1	16.2
1995	2.7	1.2	7.2	2.3	1.1	6.2	5.7	2.2	16.5
1996	2.7	1.2	7.4	2.4	1.1	6.4	5.2	1.9	15.4
1997	2.6	1.2	7.1	2.3	1.1	6.2	5.4	2.1	15.3
1993-97	2.7	1.2	7.4	2.4	1.1	6.4	5.6	2.1	16.4

SEER Program. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.
 NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Table V-4
CERVIX UTERI CANCER (Invasive)

SURVIVAL RATES, BY RACE, DIAGNOSIS YEAR, STAGE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
5-YR RELATIVE SURVIVAL RATES									
YEAR OF DIAGNOSIS:									
1960-63 ^a	-	-	-	58	-	-	47	-	-
1970-73 ^a	-	-	-	64	-	-	61	-	-
1974-76 ^b	69.1	81.4	58.0	69.8	82.3	58.6	63.5	75.2	53.4
1977-79 ^b	68.4	78.1	59.7	69.6	79.8	60.5	62.2	71.5	52.9
1980-82 ^b	67.4	79.1	56.1	68.0	80.5	55.7	61.2	72.0	51.1
1983-85 ^b	68.6	78.2	57.6	70.3	80.4	58.5	60.2	69.7	50.3
1986-88 ^b	68.8	78.6	56.7	71.7	81.4	58.5	55.3	63.9	47.4
1989-96 ^b	69.9	78.3S	58.4	71.6S	80.9	58.1	58.6	63.4S	52.7
1989-96 ^b									
ALL STAGES	69.9	78.3	58.4	71.6	80.9	58.1	58.6	63.4	52.7
LOCALIZED	91.5	93.1	88.1	92.4	94.5	87.5	86.1	86.7	84.3
REGIONAL	48.6	51.4	46.4	49.6	53.9	46.1	37.2	34.5	39.8
DISTANT	12.6	20.0	8.8	14.5	23.2	9.7	7.4	6.5 [†]	5.6
UNSTAGED	60.3	75.1	46.3	59.5	79.2	42.2	60.9 [†]	63.1 [†]	58.2 [†]
STAGE DISTRIBUTION (%) 1989-96^b									
ALL STAGES									
Number of cases	8,593	4,834	3,759	6,502	3,766	2,736	1,239	649	590
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
LOCALIZED	53	65	38	55	67	39	45	53	35
REGIONAL	32	24	42	31	23	42	36	31	41
DISTANT	7	5	11	7	5	11	9	6	13
UNSTAGED	7	6	9	7	5	9	10	10	11
5-YR RELATIVE SURVIVAL RATES, 1989-96^b									
AGE AT DIAGNOSIS:									
<45	80.6	-	-	83.4	-	-	65.6	-	-
45-54	68.1	-	-	69.3	-	-	53.2	-	-
55-64	63.4	-	-	62.5	-	-	58.4	-	-
65-74	54.6	-	-	54.3	-	-	53.5 [†]	-	-
75+	44.2	-	-	44.0	-	-	40.8 [†]	-	-
Under 65	74.4	-	-	76.5	-	-	61.2	-	-
65 and over	50.7	-	-	50.3	-	-	48.3	-	-

^a Rates are based on End Results data from a series of hospital registries and one population-based registry.
^b Rates are from the SEER Program. They are based on data from population-based registries in Connecticut, New Mexico, Utah, Iowa, Hawaii, Atlanta, Detroit, Seattle-Puget Sound and San Francisco-Oakland. Rates are based on follow-up of patients through 1997.

The difference in rates between 1974-76 and 1989-96 is statistically significant ($p<.05$).

The standard error of the survival rate is between 5 and 10 percentage points.

The standard error of the survival rate is greater than 10 percentage points.

Statistic could not be calculated.

S
↑
-

Table V-5
CERVIX UTERI CANCER (Invasive)

INCIDENCE, MORTALITY, AND SURVIVAL RATES

By Year of Diagnosis/Death

All Races, Females

Year of Diagnosis/Death

	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
Incidence																						
SEER	11.9	10.9	10.5	10.6	10.2	9.0	8.9	8.8	9.2	8.5	9.0	8.3	8.9	8.9	8.9	8.4	8.3	8.2	8.0	7.5	7.9	7.5
Mortality																						
U.S.	4.5	4.1	4.0	3.8	3.7	3.6	3.4	3.3	3.3	3.1	3.2	3.0	3.0	3.0	3.0	2.9	2.9	2.8	2.8	2.7	2.7	2.6
SEER	4.1	3.5	3.6	3.2	2.9	3.1	2.9	2.8	2.7	2.7	2.6	2.4	2.5	2.7	2.3	2.4	2.3	2.2	2.5	2.0	2.1	1.8
Relative Survival Rates (SEER)																						
1-year	88.3	87.4	86.3	87.2	86.8	86.3	87.3	89.2	88.5	86.9	86.0	87.5	87.4	89.2	88.2	88.9	87.3	89.6	87.7	89.2	87.9	
2-year	78.8	78.5	77.4	77.9	77.1	77.2	75.6	78.6	78.9	75.8	76.8	77.8	79.0	80.8	79.5	79.9	78.0	79.8	79.9	80.5		
3-year	74.3	73.7	73.3	72.0	73.2	73.1	70.0	74.2	74.1	71.0	71.9	72.4	75.5	75.8	75.5	74.5	73.8	75.2	75.8			
4-year	71.9	72.0	69.9	69.0	70.2	69.9	67.7	71.0	71.1	68.5	69.0	69.3	73.7	73.0	73.2	70.9	70.0	72.1				
5-year	70.0	69.9	68.4	67.1	68.5	67.5	66.2	69.8	69.3	66.6	66.4	67.2	72.7	71.0	72.1	68.7	67.3					
6-year	69.2	69.2	67.1	65.7	67.4	66.3	64.0	67.6	67.1	65.7	65.8	66.5	70.9	69.5	70.8	66.9						
7-year	68.5	68.2	65.6	65.0	65.1	63.8	63.4	66.4	65.7	64.3	64.5	65.6	70.1	68.4	69.8							
8-year	68.0	68.0	64.6	64.2	64.2	62.4	62.9	64.7	64.4	63.2	63.6	65.3	69.2	67.2								
9-year	67.0	67.0	63.9	64.0	63.2	62.3	62.7	64.7	63.2	62.3	63.6	65.0	68.5									
10-year	66.3	66.2	63.7	62.4	61.9	62.2	62.6	64.4	63.1	61.6	63.1	64.8										
11-year	65.7	65.5	63.3	62.4	61.5	62.2	62.4	63.6	62.3	61.1	62.6											
12-year	64.6	64.9	63.1	62.3	60.7	61.9	61.7	62.5	61.7	60.0												
13-year	64.2	64.9	62.5	62.3	60.7	61.6	60.4	62.2	61.0													
14-year	63.4	64.6	61.4	62.3	60.1	59.8	59.8	61.7														
15-year	63.1	64.4	60.7	61.9	60.1	58.4	58.6															
16-year	63.1	64.1	60.4	61.8	58.9	58.1																
17-year	62.9	63.7	59.6	61.6	58.0																	
18-year	62.9	62.5	59.1	61.3																		
19-year	62.2	62.0	58.1																			
20-year	61.6	61.2																				
21-year	60.1																					

Note: Incidence and mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.

Survival rates are relative rates expressed as percents.

Table V-6
CERVIX UTERI CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed With Cancer and
Lifetime Risk (Percent) of Dying From Cancer

9 SEER Areas, 1995-97

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.05	0.78
10	0.00	0.05	0.18	0.78
20	0.05	0.17	0.33	0.78
30	0.13	0.28	0.43	0.74
40	0.16	0.30	0.44	0.62
50	0.15	0.29	0.39	0.47
60	0.15	0.26	0.32	0.34

Lifetime Risk of Being Diagnosed = 0.78%
Lifetime Risk of Dying = 0.23%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.05	0.69
10	0.00	0.05	0.18	0.69
20	0.05	0.18	0.31	0.69
30	0.13	0.26	0.38	0.65
40	0.14	0.26	0.37	0.52
50	0.12	0.24	0.33	0.39
60	0.12	0.22	0.27	0.28

Lifetime Risk of Being Diagnosed = 0.69%
Lifetime Risk of Dying = 0.21%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.04	1.15
10	0.00	0.04	0.17	1.17
20	0.04	0.17	0.37	1.17
30	0.12	0.33	0.55	1.14
40	0.21	0.43	0.64	1.04
50	0.24	0.46	0.64	0.87
60	0.24	0.45	0.62	0.70

Lifetime Risk of Being Diagnosed = 1.15%
Lifetime Risk of Dying = 0.42%

Table V-7
CERVIX UTERI CANCER (Invasive)
AGE-ADJUSTED SEER INCIDENCE AND U.S. MORTALITY RATES
By Race/Ethnicity and Sex

Incidenceⁱ

RACE/ETHNICITY	Rate 1990-1997	Trend 1990-1997
	Rate per 100,000 persons	EAPC (%)
	Females	Females
All Races	8.9	-2.3*
White	8.4	-2.8*
White Hispanic	16.2	-4.2*
White Non-Hispanic	7.1	-2.8*
Black	11.7	-3.0*
Asian/Pacific Islander	10.2	0.3
American Indian	6.4	-
Hispanic [⊗]	15.3	-4.4*

Mortality[♦]

RACE/ETHNICITY	Rate 1990-1997	Trend 1990-1997
	Rate per 100,000 persons	EAPC (%)
	Females	Females
All Races	2.8	-1.8*
White	2.4	-1.4*
White Hispanic	3.7	-0.3
White Non-Hispanic	2.3	-1.4*
Black	5.9	-3.4*
Asian/Pacific Islander	2.7	-1.5*
American Indian	3.6	-8.3*
Hispanic [⊗]	3.4	-0.6

- The EAPC is the Estimated Annual Percent Change over the time interval.
- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
i Incidence data are from the 11 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, and Los Angeles). Unless specified, other tables use data from 9 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

⊗ Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians. For incidence, all 11 SEER areas are included. For mortality, information is included for all states except Connecticut, Oklahoma, Louisiana, and New Hampshire.

♦ Mortality data are analyzed from a public-use file provided by the National Center for Health Statistics (NCHS).

* The EAPC is significantly different from zero ($p < .05$).

Table V-8
CERVIX UTERI CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE RATES\$ AND SEER MORTALITY RATES*
By Registry, Race and Age

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES\$, 1993-97</u>									
REGISTRY									
11 registries	8.6	5.7	17.6	8.0	5.7	15.2	11.2	6.0	27.5
9 registries	7.8	5.3	15.5	7.1	5.1	13.2	11.4	6.0	28.1
San Francisco-Oakland	8.0	5.2	16.5	7.2	5.2	13.3	9.0	5.1	21.2
Connecticut	7.2	5.0	14.2	6.7	4.8	12.6	12.6	6.4	31.9
Metropolitan Detroit	8.5	5.8	16.9	7.2	5.2	13.4	12.5	7.0	29.6
Hawaii	8.1	5.3	17.0	6.6	4.9	11.9	-	-	-
Iowa	7.3	5.6	12.6	7.1	5.5	12.2	-	-	-
New Mexico	8.5	5.7	17.4	8.4	5.7	16.9	-	-	-
Seattle-Puget Sound	7.1	5.1	13.3	6.8	5.1	12.1	6.7	4.7	12.8
Utah	6.7	4.6	13.4	6.7	4.7	12.9	-	-	-
Metropolitan Atlanta	9.2	5.7	20.2	7.6	5.3	14.7	12.7	5.6	34.8
San Jose-Monterey	8.6	5.2	18.9	7.8	5.3	15.6	9.6	4.3	25.7
Los Angeles	11.2	7.0	24.0	11.2	7.6	22.1	10.9	5.9	26.2
<u>SEER MORTALITY RATES*, 1993-97</u>									
REGISTRY									
11 registries	2.3	1.0	6.4	2.1	1.0	5.7	4.3	1.7	12.2
9 registries	2.1	0.9	5.7	1.9	0.8	5.2	4.0	1.6	11.4
San Francisco-Oakland	2.0	0.9	5.4	1.8	0.9	4.6	3.4	1.4	9.7
Connecticut	1.8	0.9	4.7	1.7	0.8	4.3	4.1	1.8	11.3
Metropolitan Detroit	2.4	1.1	6.3	2.0	0.9	5.5	3.9	2.0	9.9
Hawaii	1.6	0.8	4.1	1.6	0.5	4.9	-	-	-
Iowa	2.1	0.9	5.7	2.1	0.9	5.6	-	-	-
New Mexico	2.5	1.1	6.7	2.4	1.0	6.6	-	-	-
Seattle-Puget Sound	2.0	0.8	5.7	1.9	0.8	5.3	3.0	1.6	7.4
Utah	2.1	0.9	5.7	2.1	0.9	5.7	-	-	-
Metropolitan Atlanta	2.5	0.8	7.6	1.6	0.6	4.8	5.4	1.4	17.8
San Jose-Monterey	2.2	0.8	6.8	1.9	0.8	5.5	4.3	0.4	16.5
Los Angeles	3.1	1.3	8.5	2.9	1.4	7.7	4.8	1.9	13.7

Note: San Jose-Monterey and Los Angeles are new SEER areas as of 1992 and their data are only used in tables where indicated.
 SEER Program. Incidence rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.
 NCHS public use tape. Mortality rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.
 Statistic could not be calculated or populations are inadequate.

Table V-9
CERVIX UTERI CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER MORTALITY* RATES BY STATE, 1993-97

All Races, Females

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
TOTAL U.S.	2.7	0.02			Montana	2.1*	0.27	(42)	-22.2
High Five States					Nebraska	2.1*	0.21	(40)	-22.2
District of Columbia	4.3*	0.49	(01)	59.3	Nevada	2.9	0.26	(22)	7.4
Delaware	4.0*	0.43	(02)	48.1	New Hampshire	2.5	0.28	(32)	-7.4
Kentucky	3.6*	0.17	(03)	33.3	New Jersey	2.9	0.11	(20)	7.4
West Virginia	3.6*	0.24	(04)	33.3	New Mexico	2.5	0.23	(33)	-7.4
Louisiana	3.6*	0.17	(05)	33.3	New York	2.9	0.07	(17)	7.4
Low Five States					North Carolina	2.9	0.11	(19)	7.4
South Dakota	1.7*	0.29	(47)	-37.0	North Dakota	2.6	0.39	(29)	-3.7
Minnesota	1.7*	0.11	(48)	-37.0	Ohio	2.7	0.09	(26)	0.0
Hawaii	1.6*	0.22	(49)	-40.7	Oklahoma	3.0	0.17	(15)	11.1
Alaska	1.5*	0.36	(50)	-44.4	Oregon	2.3	0.15	(37)	-14.8
Wyoming	1.5*	0.32	(51)	-44.4	Pennsylvania	2.7	0.08	(28)	0.0
State	Rate	SE	Rank	PD	Rhode Island	2.7	0.31	(25)	0.0
Alabama	3.1	0.15	(13)	14.8	South Carolina	3.4*	0.17	(09)	25.9
Alaska	1.5*	0.36	(50)	-44.4	South Dakota	1.7*	0.29	(47)	-37.0
Arizona	2.3	0.14	(35)	-14.8	Tennessee	3.5*	0.15	(07)	29.6
Arkansas	3.2*	0.20	(10)	18.5	Texas	3.1	0.08	(12)	14.8
California	2.6	0.06	(30)	-3.7	Utah	2.1*	0.22	(39)	-22.2
Colorado	2.1*	0.14	(43)	-22.2	Vermont	3.4*	0.44	(08)	25.9
Connecticut	1.8*	0.13	(46)	-33.3	Virginia	2.7	0.12	(27)	0.0
Delaware	4.0*	0.43	(02)	48.1	Washington	2.0*	0.11	(45)	-25.9
District of Columbia	4.3*	0.49	(01)	59.3	West Virginia	3.6*	0.24	(04)	33.3
Florida	2.9	0.08	(16)	7.4	Wisconsin	2.3	0.13	(36)	-14.8
Georgia	3.0	0.12	(14)	11.1	Wyoming	1.5*	0.32	(51)	-44.4
Hawaii	1.6*	0.22	(49)	-40.7					
Idaho	2.1*	0.26	(38)	-22.2					
Illinois	2.9	0.09	(21)	7.4					
Indiana	2.9	0.13	(18)	7.4					
Iowa	2.1*	0.16	(41)	-22.2					
Kansas	2.5	0.19	(31)	-7.4					
Kentucky	3.6*	0.17	(03)	33.3					
Louisiana	3.6*	0.17	(05)	33.3					
Maine	3.2*	0.29	(11)	18.5					
Maryland	2.8	0.14	(24)	3.7					
Massachusetts	2.0*	0.11	(44)	-25.9					
Michigan	2.4	0.09	(34)	-11.1					
Minnesota	1.7*	0.11	(48)	-37.0					
Mississippi	3.6*	0.21	(06)	33.3					
Missouri	2.9	0.13	(23)	7.4					

* NCHS public use tape. Rates are per 100,000 and are age-adjusted to the 1970 U.S. standard population.
SE Standard error of the rate.


PD Percent difference between state rate and total U.S. rate.

* Absolute difference between state rate and total U.S. rate is 15% or more.

⊕ Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

Figure V-1

SEER Incidence and U.S. Mortality Rates⁺ Cervix Uteri By Race




⁺ Rates are age-adjusted to the 1970 U.S. standard million population. Regression lines are calculated using the Joinpoint Regression Program.


Figure V-2

SEER Incidence and U.S. Mortality Rates⁺ Cervix Uteri, By Age and Race


White - Under 50 Years of Age




Black - Under 50 Years of Age



White - Ages 50 and Over



Black - Ages 50 and Over



• Incidence × Mortality

⁺ Rates are age-adjusted to the 1970 U.S. standard million population. Regression lines are calculated using the Joinpoint Regression Program.