

Table X-1

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1973-99

All Races, Males, All Ages

Year of Diagnosis

SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
All Areas [⊕]	174	35	44	109	219	369	518	722	857	1019	1125	1111	1137	1083	981	966	836	565	263	192	184
All Areas but SF-Oakland	130	23	21	35	52	106	153	196	233	353	421	429	415	494	481	444	428	307	149	121	114
San Francisco-Oakland SMSA	44	12	23	74	167	263	365	526	624	666	704	682	722	589	500	522	408	258	114	71	70
San Francisco City/County	17	7	14	61	140	226	313	441	515	548	537	530	569	432	366	383	307	178	74	43	39

AGE-ADJUSTED CANCER INCIDENCE RATES[§]

By SEER Location and Year of Diagnosis, 1973-99

All Races, Males, All Ages

Year of Diagnosis

SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
All Areas [⊕]	0.4	0.6	0.5	1.2	2.2	3.6	4.9	6.6	7.6	8.9	9.6	9.3	9.4	8.8	7.8	7.6	6.5	4.4	2.1	1.5	1.5
All Areas but SF-Oakland	0.4	0.4	0.3	0.6	0.7	1.3	1.8	2.2	2.5	3.8	4.3	4.4	4.1	4.8	4.5	4.1	4.0	2.9	1.4	1.2	1.1
San Francisco-Oakland SMSA	0.7	1.1	1.5	4.5	9.5	14.7	19.6	27.6	32.6	33.4	35.1	33.2	35.4	28.2	23.5	24.9	19.4	12.3	5.4	3.4	3.3
San Francisco City/County	0.9	2.1	4.1	15.8	34.1	55.0	75.0	102.6	120.0	125.5	125.3	122.5	132.5	99.8	83.6	89.4	71.7	41.9	17.1	9.8	9.0

[§] SEER 9 areas. Rates are per 100,000 males and are age-adjusted to the 2000 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

[⊕] All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

Table X-2

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1973-99

All Races, Males, Ages 20-54

Year of Diagnosis

SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
All Areas [⊕]	18	6	20	75	179	329	477	682	803	949	1058	1050	1068	1004	924	911	765	517	229	158	145
All Areas but SF-Oakland	14	<4	4	9	26	79	128	174	205	312	382	394	380	462	446	421	385	277	127	98	90
San Francisco-Oakland SMSA	4	5	16	66	153	250	349	508	598	637	676	656	688	542	478	490	380	240	102	60	55
San Francisco City/County	<4	5	13	59	134	219	300	428	497	529	520	514	547	401	353	361	288	166	70	39	33

AGE-ADJUSTED CANCER INCIDENCE RATES§

By SEER Location and Year of Diagnosis, 1973-99

All Races, Males, Ages 20-54

Year of Diagnosis

SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
All Areas [⊕]	0.1	0.1	0.4	1.5	3.3	6.1	8.7	12.2	14.1	16.2	17.7	17.3	17.3	15.8	14.3	14.1	11.6	7.8	3.5	2.4	2.2
All Areas but SF-Oakland	0.1	-	0.1	0.2	0.6	1.7	2.7	3.6	4.1	6.3	7.5	7.6	7.1	8.7	8.2	7.7	6.9	4.9	2.3	1.8	1.6
San Francisco-Oakland SMSA	0.1	0.6	2.0	7.6	16.4	27.2	37.3	52.9	62.0	63.3	66.6	63.1	66.0	50.4	44.3	45.7	35.3	22.3	9.3	5.4	4.9
San Francisco City/County	-	2.9	7.6	30.6	64.4	106.2	143.0	197.9	230.3	241.0	240.2	236.2	251.8	181.4	160.1	166.3	132.5	77.2	32.1	17.5	14.7

§ SEER 9 areas. Rates are per 100,000 males aged 20-54 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

⊕ All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

Table X-3

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1973-99

All Races, Males, Ages 55+

Year of Diagnosis

SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
All Areas [⊕]	156	29	24	34	39	39	41	39	54	68	66	61	69	78	57	55	71	48	34	34	39
All Areas but SF-Oakland	116	22	17	26	25	27	25	22	28	41	39	35	35	31	35	23	43	30	22	23	24
San Francisco-Oakland SMSA	40	7	7	8	14	12	16	17	26	27	27	26	34	47	22	32	28	18	12	11	15
San Francisco City/County	16	<4	<4	<4	6	7	13	13	18	19	16	16	22	31	13	22	19	12	4	4	6

AGE-ADJUSTED CANCER INCIDENCE RATES§

By SEER Location and Year of Diagnosis, 1973-99

All Races, Males, Ages 55+

Year of Diagnosis


SEER Location	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
All Areas [⊕]	1.8	2.3	1.6	2.3	2.5	2.5	2.4	2.3	2.8	3.7	3.4	3.3	3.8	3.9	2.8	2.6	3.4	2.4	1.7	1.6	1.7
All Areas but SF-Oakland	1.6	2.0	1.4	2.2	1.9	2.0	2.0	1.7	1.9	2.9	2.5	2.5	2.4	1.9	2.1	1.3	2.5	1.9	1.3	1.3	1.3
San Francisco-Oakland SMSA	2.9	3.8	2.5	3.1	5.9	4.9	4.6	5.1	7.7	7.8	8.4	7.8	11.3	14.3	6.5	9.5	8.3	5.2	3.7	3.3	4.0
San Francisco City/County	4.0	-	-	-	8.8	9.3	16.4	17.6	23.4	24.1	22.9	21.0	31.4	43.2	16.8	30.0	26.0	15.8	5.1	5.0	7.5

§ SEER 9 areas. Rates are per 100,000 males aged 55+ and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

⊕ All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

Figure X-1

All Cancer Sites vs. Kaposi's Sarcoma SEER Incidence Rates⁺, 1973-1999 Males, Ages 20-54



Source: SEER 9 areas. Rates are age-adjusted to the 2000 U.S. standard million population by 5-year age groups.
Regression lines are calculated using the Joinpoint Regression Program.






Figure X-2

Source: SEER 9 areas. Rates are age-adjusted to the 2000 U.S. standard million population by 5-year age groups.

* Kaposi's sarcoma excluded from Oral, Colorectal, Lung, Melanoma, Testis, and NHL[^].

[^] Non-Hodgkin Lymphoma.